

Wheatgrass Extract Ameliorates Hypoxia-induced Mucin Gene Expression in A549 cells

Ju hwan Sim, Moon-Hee Choi¹, Hyun-Jae Shin¹, Ji-Eun Lee,

Department of Otorhinolaryngology-Head and Neck Surgery, Chosun University College of Medicine, ¹Department of Biochemical and Polymer Engineering, Chosun University, Gwangju, Korea

Submitted: 30-12-2015

Revised: 14-02-2016

Published: 06-01-2017

ABSTRACT


Background: Wheatgrass is known to have antioxidant, antiaging, and anti-inflammatory effect. However, its protective effect against hypoxia is not yet evaluated. **Objective:** In this study, we evaluated the protective and anti-inflammatory effect of wheatgrass against the hypoxia in airway epithelial cells. **Materials and Methods:** A549 human lung adenocarcinoma cells were incubated in a hypoxic condition (CO₂ 5%/O₂ 1%) for 24 hr in the presence of different concentration of wheatgrass 50, 75, 100, and 150 µg/mL, and the magnitude of each immunologic response produced by the A549 cells was compared. The mRNA expression level of mucin gene (MUC), 5A, 5B, 8, GM-CSF, TNF-α, and VEGF were evaluated by using real-time polymerase chain reaction. The MUC proteins level before and after knocking out the hypoxia-inducible factor (hif)-1α via short interfering (si) RNA transfection were assessed by immunoblot analysis. Accordingly, the involved cell signaling pathway was evaluated by immunoblot analysis. **Results:** The inflammatory cytokines (GM-CSF, TNF-α) and the expressions of MUC 5A, 5B, and 8 were augmented by hypoxia. The augmented MUC expression was decreased by the wheatgrass extract administration. Hif-1α gene expression after hypoxia exposure was decreased by wheatgrass. Knockdown of hif-1α by siRNA reduced the mucin gene expression and which was more enhanced by wheatgrass extract. **Conclusion:** These results suggest that wheatgrass may be useful in the treatment of sinonasal disease by inhibiting mucus hypersecretion in airway epithelium.

Key words: Airway epithelium, hypoxia, mucin, wheatgrass, VEGF

SUMMARY

- Wheatgrass extract decreases the hypoxia-induced MUC 5A, 5B and 8 expression.
- Hif-1α gene expression after hypoxia exposure was decreased by wheatgrass.

- Wheatgrass inhibits p44/42 phosphorylation in hypoxia-exposed airway epithelial cells.



Abbreviations used:

A549: human lung adenocarcinoma cells, GM-CSF: granulocyte-macrophage colony stimulating factor, HIF: hypoxia inducible factor, IL: interleukin, MUC: mucin, MTT: 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide, TNF: tumor necrosis factors, VEGF: vascular endothelial growth factor, si RNA: short interfering RNA

Correspondence:

Dr. Ji-Eun Lee,
Department of Otorhinolaryngology- Head &
Neck Surgery
College of Medicine,
Chosun University, 365 Plimundaero, Dong-gu,
Gwangju, South Korea.
E-mail: jeviolin@hanmail.net, jeviolins@gmail.com
DOI: 10.4103/0973-1296.197660

Access this article online

Website: www.phcog.com

Quick Response Code:



INTRODUCTION

The airway epithelium is lined by mucus, which is a mixture of water, glycoprotein, proteins, and lipids. Mucin is one of the components of mucus and has been implicated in several airway diseases. In the sinus, mucin gene (MUC) 5AC and MUC 5B are known to be predominant and their expression is controlled by various mediators such as interleukin (IL)-1β, IL-6, IL-13, IL-17 or tumor necrosis factor-α.^[1-3] Recently, one study demonstrated anoxia upregulates MUC5AC by the hypoxia-inducible factor-1α (hif-1α) signaling pathway in human nasal epithelium.^[4]

The roles of hypoxia in sinus, in terms of cause of inflammation, are supposed to be exist generally as at least two different ways. One way is to induce an epithelial–mesenchymal transition, which is basic feature of nasal polyps,^[5] and the other is to overexpress the mucin gene. Thus, hif-1α can be a promising target for the prevention and treatment sinus inflammatory disease.

Wheatgrass refers to the young grass of the common monocot wheat plant "*Triticum aestivum*". Its consumption in the Western world began in

the 1930s. Today, wheatgrass is quickly becoming one of the most widely used supplemental health foods and is available in many health food stores as fresh produce, tablets, frozen juice, and powder. Wheatgrass contains vitamins, minerals, enzymes, amino acids, polysaccharides, and large amounts (70%) of chlorophyll. Several papers have indicated that wheatgrass has antitumor activities,^[6] antioxidant properties,^[7] and a therapeutic effect on distal ulcerative colitis.^[8] In addition, wheatgrass may help prevent some disorders, including diabetes and heart disease.^[9] Considering that hypoxia has been getting an attention as one of the

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: Sim Jh, Choi MH, Shin HJ, Lee JE. Wheatgrass extract ameliorates hypoxia-induced mucin gene expression in A549 cells. *Phcog Mag* 2017;13:7-12.

pathogenic mechanism of sinusitis, mucus hypersecretion induced by hypoxia could be a new therapeutic target of sinonasal disease. In this study, we have aimed to evaluate the protective and anti-inflammatory effect of wheatgrass against the hypoxia in airway epithelial cells. We found that wheatgrass extract inhibits hif-1 α expression and downregulates hif-1 α induced mucin overexpression in airway epithelial cells. This suggests that wheatgrass may have potential therapeutic relevance.

MATERIALS AND METHODS

SP600125 (JNK inhibitor), PD980599 (ERK inhibitor), and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) were obtained from Sigma–Aldrich (Munich, Germany). The other chemicals used were of the purest grade available from Sigma (St Louis, MO).

Preparation of wheat sprout sample

The wheatgrass sprout was harvested from a commercial planting located Gwang-ju, Korea in 2015.

Extracts preparation

After grinding dried wheat sprouts in a mixer, they were extracted by 80% ethanol for three days in shaking incubators. Extracting conditions were at 25 °C, 200 rpm. They were concentrated by the rotary evaporator. We first extracted wheat sprout with 80% ethanol, and the ethanol extracts were fractionated with ethyl acetate or water. Ethyl acetate layer was further separated with n-butanol and water. Then, the butanol fraction was separated with hexane and water.

Cell cultures

A549 human lung adenocarcinoma cells were maintained in RPMI 1640 (Gibco BRL, Gnd Island, NY) media supplemented with 1% penicillin/streptomycin (gibco/Invitrogen, Carlsbad, CA) and 10% fetal calf serum (Capricorn) (w/v). The cells were grown to 60% confluence in 100 mm culture plates and kept at 37°C in a carbon-dioxide-enriched (95% air, 5% CO₂) humidified atmosphere.

Cells were pre-treated with different concentration of wheatgrass extract (50–150 μ g/mL) for 30 min and then incubated in modular incubator chamber (hypoxic, 5% CO₂/1% O₂) (Billups-Rotheberg, Del Mar, CA) or in normoxic condition (5% CO₂/20% O₂) for additional 24 hr.

Cell cytotoxicity assays

Cell proliferation was measured by a colorimetric assay using MTT. In brief, a549 cells cultured overnight on 96-well plates were administered media containing different concentrations of wheatgrass (50–150 μ g/mL) and incubated for 24 hr under hypoxic condition. 10 μ L of MTT stock solution (5 mg/mL) were added to each well, followed by incubation for an additional 4 hr. Blue formazans were eluted from cells by the addition of 100 μ L of DMSO with gentle shaking for 10 min at room temperature. Absorbances were measured at 570 nm using an enzyme-linked immunosorbent assay reader (Spectra MAX; Molecular Devices, Sunnyvale, CA).

Real-time polymerase chain reaction (qRT-PCR)

Total RNA was extracted from the cells using TRIzol reagent (Invitrogen) according to manufacturer's instructions, and reverse transcribed into cDNA with the Quantitect Reverse Transcription kit (Qiagen, Venlo, NLD). qRT-PCR analyses were performed using a 7500 FAST qRT-PCR System (Applied Biosystems, Foster City, CA). Each reaction mixture contained 10 μ L of SYBR® Green PCR Master Mix (Applied Biosystems), 4 pmol of forward and reverse primers each, and 1 μ L of cDNA in a final volume of 20 μ L. Reaction mixtures were incubated at 95 °C for 5 min to

Table 1: Primers for real-time polymerase chain reaction

Primers	
MUC 5AC	F: 5 ϵ -GGGACTTCTCCTACCAAT-3 ϵ R: 5 ϵ -TATATGGTGGATCCTGCA GGGTAG-3 ϵ
MUC 5B	F: 5 ϵ -CAC ATC CAC CCT TCC AAC-3 ϵ R: 5 ϵ -GGC TCA TTG TCG TCT CTG-3 ϵ
MUC 8	F: 5 ϵ -GAC AGG GTT TCT CCT CAT TG-3 ϵ R: 5 ϵ -CGT TTA TTC CAG CAC TGT TC-3 ϵ
VEGF	F: 5 ϵ -CTACCTCCACCATGCCAAGT -3 ϵ R: 5 ϵ -GCAGTAGCTGCGCTGATAGA -3 ϵ
TNF- α	F: 5'-AGACGCCACATCCCCTGACAA-3' R: 5'-AGACGGCGATGCGGCTGATG-3'
GAPDH	F: 5 ϵ -ATCATCCCTGCCTCTACTGG-3 ϵ R: 5 ϵ -GTCAGGTCCACCACTGACAC-3 ϵ

activate FastStartTaq DNA Polymerase, followed by amplification for 40 cycles. Data were analyzed using Sequence Detection Software version 1.9.1 (Applied Biosystems). Target mRNA expression was normalized to glyceraldehyde-3-phosphate dehydrogenase (GAPDH) expression, and calculated using the comparative Ct method. Primers constructed were shown in Table 1.

si RNA transfection

Cells were transfected with short interfering (si) RNA corresponding to human hif-1 α or control siRNA using Lipofectamine RNAiMAX (Invitrogen) according to the manufacturer's protocol. The si-RNA was #1068432V for hif-1 α (Bioneer, Daejeon, Korea). After incubation for 4 h, media were changed with complete medium containing 10% serum and antibiotics.

Nuclear protein extract

The nuclear protein was extracted by using the NE-PER® Nuclear and Cytoplasmic Extraction Reagents (Pierce, Rockford, IL) according to the manufacturer's instruction.

Immunoblot analysis

After incubation in hypoxic condition, media was removed and cells were washed with phosphate-buffered saline (10 mM, pH 7.4). Then, cells were lysed with lysis buffer (50 mM Tris pH 7.7, 150 mM NaCl, 1% NP-40, 5 mM EGTA, 50 mM glycerophosphate, 20 mM NaF, 1 mM Na₃VO₄, 2 mM phenylmethyl sulfonyl fluoride, 10 mg/mL leupeptin, and 10 mg/mL aprotinin) and incubated for 20 min at 4°C. After sonication briefly, the cells were centrifuged at 13,000 g for 10 min at 4°C. The supernatant that contained the total cell lysate was collected. Protein concentration of the lysates was measured by Bio-Rad Protein Assay (Bio-Rad, Hercules, CA). Equal amounts of protein were mixed with sample buffer, boiled for 5 min, and separated by electrophoresis on 10%-12% Tris-HCl gels. The protein content of the gels was transferred to a PVDF membrane (Amersham, Buckinghamshire, UK), and the membranes were blocked with TBS-T (20 mM Tris, 500 nM NaCl, with 0.1% Tween-20) containing 5% (w/v) skim milk for 1 hr at room temperature. The membrane was incubated overnight at 4°C with a specific primary antibody of MUC 8 (Sigma-Aldrich, St. Louis, MO), Hif-1 α (Novus, Littleton, Co) p38, phospho-p38, p42/44, phospho-p42/44 (Cell Signaling Technology, Danvers, MA), and GAPDH (Santa Cruz Biotechnology, Dallas, TX) followed by peroxidase-conjugated antimouse IgG or antirabbit IgG (Jackson Immuno Research, West Grove, PA). The membranes were developed using the enhanced chemiluminescent analysis system (SuperSignal®

West Pico Chemiluminescent Substrate, Pierce, Waltham, MA) and the signal was captured on an image reader (LAS4000; Fuji Photo Film, Tokyo, Japan). Results were obtained from three independent experiments.

Statistical analysis

All data are expressed as mean standard deviation (SD). Statistical analysis was performed using student's *t*-test of variance. $p < 0.05$ for the null hypothesis was accepted as indicating a statistically significant difference. Statistical analyses were performed using SPSS for Windows (Ver. 12.0, SPSS Inc., Chicago, IL).

RESULTS

Non-cytotoxic effects of wheatgrass extract on airway epithelial cells

To examine the effect of wheatgrass, we used MTT assays to measure the viabilities of cells that had been exposed to different concentration of wheatgrass for 24 hr under hypoxic condition. Wheatgrass per se did not show toxicity of cells at various concentration of 50–150 $\mu\text{g/mL}$ [Figure 1].

Wheatgrass extract decreases mucin gene expression induced by hypoxia exposure

We investigate the effects of wheatgrass extract on MUC gene production in hypoxia-exposed airway epithelial cells. The mRNA expression

of MUC 5AC and MUC8 was increased when cells were exposed to hypoxia for 24 hr [Figure 2]. To elucidate the effect of wheatgrass, we treated wheatgrass extract at various concentration of 50, 75, 100,




Figure 1: MTT assay to determine cytotoxic effects of hypoxia and wheatgrass. WG, wheatgrass. Wheatgrass per se did not have cytotoxic effect.





Figure 2: Hypoxia induces the expression of MUC mRNA in A549 cells. WG, wheatgrass; VEGF, vascular endothelial growth factor hypoxia-induced MUC mRNA or VEGF, TNF- α was decreased by wheatgrass administration in a dose-dependent pattern.



and 150 µg/mL to cells 30 min before hypoxia exposure. There after, cells were incubated in a hypoxic condition for 24 hr. MUC 5AC and MUC 8 mRNA expression were increased after hypoxia exposure, and decreased after wheatgrass administration. It was confirmed at protein level for MUC 8 gene. Hypoxia-induced MUC production was inhibited by wheatgrass in a dose-dependent manner which was confirmed by western blotting Figure 3a. Control GAPDH was constitutively expressed and was not affected by wheatgrass treatment.


Inhibition of hif-1α activation by wheatgrass extract in hypoxia-exposed airway epithelial cells

Hypoxia is known to induce MUC5AC production via hif-1α signaling pathway.^[4] The expression of hif-1α was increased when cells were exposed to hypoxia for 24 hr. The increased hif-1α after hypoxia exposure was decreased dose-dependently by the wheatgrass administration [Figure 3b]. These results showed that wheatgrass extract decreased hif-1α gene expression, which was known to be involved in MUC gene expression.

Knockdown of hif-1α reduced the production of mucin gene expression

Next, to further evaluate the effect of wheatgrass extract on hif-1α expression, we transiently knock downed hif-1α gene by using si RNA-hif-1α. After si RNA-hif-1α transfection, cells were incubated in a hypoxic (5% CO₂/1% O₂) or normoxic condition for 24 hr. The transfection efficiency was confirmed at the protein level. As shown in [Figure 4], signal intensity of MUC 8 protein in cells treated with wheatgrass extract (150 µg/mL) for 30 min and then exposed to hypoxia

for 24 hr was similar with that in cells transiently hif-1α gene silenced and then exposed to hypoxia for 24 hr. Moreover, its signal was as low as the normoxic control when cells were transiently transfected and then treated with wheatgrass extract (150 µg/mL). These results suggest that hypoxia-induced increase in mucin gene expression in airway epithelial cells is dependent of hif-1α transcription factor and wheatgrass acts in a way decreasing hif-1α transcriptional factor, thus reduces the mucin gene expression.






Figure 5: JNK/ERK MAPK signal pathway is involved in the regulation of MUC 8 expression by the wheatgrass treatment. WG, wheatgrass. (a) The expression of MUC 8 was decreased by the treatment of SP600125 or PD98059. (b) The signal intensity of phosphorylation of p44/42 was decreased by the wheatgrass administration.

Inhibition of p44/42 (ERK1/2) phosphorylation by wheatgrass in hypoxia-exposed airway epithelial cells

To evaluate the downstream signal pathway for hif-1 α transcriptional factor, we used the SB230580 (a p38 inhibitor), SP600125 (a JNK inhibitor), and PD98059. PD98059 is a specific inhibitor of mitogen-activated protein kinase/extracellular signal-regulated kinase 1 (MEK-1), which is responsible for ERK1/2 (p44/42) activation. Cells treated with wheatgrass extract demonstrated a similar signal density of MUC 8 protein with cells treated the SP600125 or PD98059, whereas SB203580 did not show a definite inhibitory effect [Figure 5a]. These observations indicate a possible involvement of JNK or ERK in the wheatgrass-induced anti-hypoxic effect. Since p44/42 kinase is important for hif-1 α downstream pathway in airway epithelial cells and a possible target of wheatgrass extract, we further determined the role p44/42 in hypoxia-induced MUC gene production. So, cells were treated with wheatgrass extract and exposed hypoxia for indicated time [Figure 5b]. A kinetic study showed that hypoxia-induced phosphorylation of p44/42 peaked at 5 min, was maintained until 3 hr. When cells were treated with wheatgrass extract and exposed hypoxia for the same indicated time, the signal intensity of phosphorylation of p44/42 was decreased as shown in Figure 5b. These results suggest that the p44/42 kinase pathway plays an important role in the regulation of MUC gene production of wheatgrass extract in relation with hypoxia in airway epithelial cells.

DISCUSSION

Since hypoxia has been demonstrated to be involved in chronic rhinosinusitis and nasal polyp,^[5] targeting hif-1 α represents an important therapeutic strategy for sinonasal disease. We sought to identify a natural compound capable of inhibiting hypoxia-mediated inflammatory responses. Wheatgrass is known to have antioxidant, antiaging, and anti-inflammatory effect.^[7,8] However, its immune-modulatory effect is not yet evaluated. In this study, we evaluated the protective effect of wheatgrass against hypoxia in airway epithelial cells. We demonstrated that wheatgrass extract inhibits the p44/42 (ERK1/2) pathway in hypoxia-exposed airway epithelial cells and that decreases MUC gene expression in response to hypoxia.

Hif-1 α is important for maintaining oxygen homeostasis by transcriptional activation of erythropoietin, vascular endothelial growth factor, and transferrin.^[4] Several lines of evidence demonstrated that hif-1 α plays crucial roles not only in cancers but also in benign lesions.^[10,11] Hif-1 α is induced in normal organ under systemic hypoxia.^[12,13] Under hypoxia conditions, stable hif-1 α dimerizes with hif-1 β and binds to the hypoxia-response element for transcriptional activation.^[4] The expression of hif-1 α in nasal polyp was reported previously and it was also noticed that hif-1 α is expressed in nasal epithelium.^[14,15] Moreover, ERK activity is required for hif-1 α transcriptional factor activation. ERK has been reported to be involved in hif-1 α -mediated transcription,^[16]

indicating that ERK could regulate mucin gene expression by increasing the transactivation capacity of hif-1 α .

Wheatgrass extract contains chlorophyll, phenolic compounds, and flavonols. Since it was known to have anti-oxidative effect, it has been mainly studied in cancer-related topics.^[17,18] In this study, we found that wheatgrass extract acts similarly with PD98059 which selectively inhibits the ERK 1/2 pathway, and that decreases MUC gene expression in response to hypoxia.

In summary, these experiments demonstrate that wheatgrass inhibits hypoxia-induced MUC gene expression in airway epithelial cells. Based on our findings, the most likely mechanism that can account for this biological effect involves inhibition of the ERK 1/2 kinase pathway and hif-1 α activity. Further research such as *in vitro* study using primary human nasal epithelial cells or other airway epithelial cell lines, or *in vivo* study using animal model would be necessary for clinical implication of wheatgrass in sinonasal disease.

CONCLUSION

Considering the importance of hif-1 α in sinonasal disease, wheatgrass may be useful in the treatment of sinonasal disease by inhibiting mucus hypersecretion in airway epithelium.

Acknowledgement

This work was supported by a grant from the Clinical Medicine Research Institute of the Chosun University Hospital (2014).

Financial support and sponsorship

Nil

Conflicts of interest

There are no conflict of interest.

REFERENCES

1. Koo JS, Kim YD, Jetten AM, Belloni P, Nettekheim P. Overexpression of mucin genes induced by interleukin-1 beta, tumor necrosis factor-alpha, lipopolysaccharide, and neutrophil elastase is inhibited by a retinoic acid receptor alpha antagonist. *Exp Lung Res* 2002;28:315-32.
2. Chen Y, Thai P, Zhao YH, Ho YS, DeSouza MM, Wu R. Stimulation of airway mucin gene expression by interleukin (IL)-17 through IL-6 paracrine/autocrine loop. *J Biol Chem* 2003;278:17036-43.
3. Evans CM, Williams OW, Tuvim MJ, Nigam R, Mixides GP, Blackburn MR. *et al.* Mucin is produced by clara cells in the proximal airways of antigen-challenged mice. *Am J Respir Cell Mol Biol* 2004;31:382-94.
4. Kim YJ, Cho HJ, Shin WC, Song HA, Yoon JH, Kim CH. Hypoxia-mediated mechanism of MUC5AC production in human nasal epithelia and its implication in rhinosinusitis. *PLoS One* 2014;9:e98136.
5. Shin HW, Cho K, Kim DW, Han DH, Khalmuratova R, Kim SW. *et al.* Hypoxia-inducible factor 1 mediates nasal polyposis by inducing epithelial-to-mesenchymal transition. *Am J Respir Crit Care Med* 2012;185:944-54.
6. Alitheen NB, Oon CL, Keong YS, Chuan TK, Li HK, Yong HW. Cytotoxic effects of commercial wheatgrass and fiber towards human acute promyelocytic leukemia cells (HL60). *Pak J Pharm Sci* 2011;24:243-50.
7. Das A, Raychaudhuri U, Chakraborty R. Effect of freeze drying and oven drying on antioxidant properties of fresh wheatgrass. *Int J Food Sci Nutr* 2012;63:718-21.
8. Ben-Arye E, Goldin E, Wengrower D, Stamper A, Kohn R, Berry E. Wheat grass juice in the treatment of active distal ulcerative colitis: A randomized double-blind placebo-controlled trial. *Scand J Gastroenterol* 2002;37:444-9.
9. Shermer M. Wheatgrass juice and folk medicine. *Sci Am* 2008;299:42.
10. Rankin EB, Rha J, Unger TL, Wu CH, Shutt HP, Johnson RS. *et al.* Hypoxia-inducible factor-2 regulates vascular tumorigenesis in mice. *Oncogene* 2008;27:5354-8.
11. Adunka O, Gstoettner W, Knecht R, Kierner AC. Expression of hypoxia inducible factor 1 alpha and Von Hippel Lindau protein in human middle ear cholesteatoma. *Laryngoscope* 2003;113:1210-5.
12. Stroka DM, Burkhardt T, Desbaillets I, Wenger RH, Neil DA, Bauer C. *et al.* HIF-1 is expressed in normoxic tissue and displays an organ-specific regulation under systemic hypoxia. *FASEB J* 2001;15:2445-53.
13. Wiesener MS, Jurgensen JS, Rosenberger C, Scholze CK, Horstrup JH, Warnecke C. *et al.* Widespread hypoxia-inducible expression of hif-2alpha in distinct cell populations of different organs. *FASEB J* 2003;17:271-3.
14. Chien CY, Tai CF, Ho KY, Kuo WR, Chai CY, Hsu YC. *et al.* Expression of hypoxia-inducible factor 1alpha in the nasal polyps by real-time RT-PCR and immunohistochemistry. *Otolaryngol Head Neck Surg* 2008;139:206-10.
15. Hsu YC, Kuo WR, Chen YY, Tai CF, Tsai CJ, Wang LF. Increased expression of hypoxia-inducible factor 1alpha in the nasal polyps. *Am J Otolaryngol* 2007;28:379-83.
16. Movafagh S, Crook S, Vo K. Regulation of hypoxia-inducible factor-1a by reactive oxygen species: new developments in an old debate. *J Cell Biochem* 2015;116:696-703.
17. Shakya G, Balasubramanian S, Rajagopalan R. Methanol extract of wheatgrass induces G1 cell cycle arrest in a p53-dependent manner and down regulates the expression of cyclin D1 in human laryngeal cancer cells-an *in vitro* and *in silico* approach. *Pharmacogn Mag* 2015;11:S139-47.
18. Chiu LC, Kong CK, Ooi VE. The chlorophyllin induced cell cycle arrest and apoptosis in human breast cancer MCF 7 cells is associated with ERK deactivation and Cyclin D1 depletion. *Int J Mol Med* 2005;16:735-40.