



Published in final edited form as:

Am J Prev Med. 2017 January ; 52(1 Suppl 1): S63–S76. doi:10.1016/j.amepre.2016.09.026.

Systemic Inflammation in Midlife: Race, Socioeconomic Status, and Perceived Discrimination

Irena Stepanikova, PhD^{1,2}, Lori Brand Bateman, PhD³, and Gabriela R. Oates, PhD³

¹Department of Sociology, University of Alabama at Birmingham, Birmingham, Alabama

²Research Centre for Toxic Compounds in the Environment, Masaryk University, Brno, Czech Republic

³School of Medicine, Division of Preventive Medicine, University of Alabama at Birmingham, Birmingham, Alabama

Abstract

Introduction—This study investigates social determinants of systemic inflammation, focusing on race, SES, and perceived discrimination.

Methods—Data on 884 white and 170 black participants were obtained from the Survey of Midlife in the U.S., a cross-sectional observational study combining survey measures, anthropometry, and biomarker assay. Data, collected in 2004–2009, were analyzed in 2016. Main outcome measures were fasting blood concentrations of C-reactive protein, interleukin 6, fibrinogen, and E-selectin. For each biomarker, series of multivariate linear regression models were estimated for the pooled sample and separately for blacks and whites. Full models included social determinants; psychological, lifestyle, and health factors; and demographic covariates.

Results—Bivariate analyses indicated higher concentrations of all inflammation markers among blacks compared with whites ($p < 0.001$). In fully adjusted models using the pooled sample, racial differences persisted for interleukin 6 ($p < 0.001$) and fibrinogen ($p < 0.01$). For E-selectin and C-reactive protein, racial differences were explained after adjusting for covariates. Education was linked to lower fibrinogen concentration ($p < 0.05$) in the fully adjusted model and C-reactive protein concentration ($p < 0.01$) after adjusting for demographic factors and income. Lifetime perceived discrimination was related to higher concentrations of fibrinogen ($p < 0.05$) in the fully adjusted model, and higher concentrations of E-selectin and interleukin 6 ($p < 0.05$) after adjusting for socioeconomic status (SES) and demographic factors.

Conclusions—This study clarifies the contributions of race, SES, and perceived discrimination to inflammation. It suggests that inflammation-reducing interventions should focus on blacks and individuals facing socioeconomic disadvantages, especially low education.

Address correspondence to: Irena Stepanikova, PhD, Department of Sociology, University of Alabama at Birmingham, 1401 University Boulevard, Birmingham AL 35294-1152. irena@uab.edu.

SUPPLEMENTAL MATERIAL

Supplemental materials associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.amepre.2016.09.026>.

This article is part of a supplement issue titled Social Determinants of Health: An Approach to Health Disparities Research.

INTRODUCTION

Systemic inflammation has received attention as a preventable factor in chronic conditions such as hypertension,¹ cardiovascular disease,^{2,3} insulin resistance,⁴ Type 2 diabetes,^{3,5-7} and cancer.⁸⁻¹⁰ Smoking,¹¹ alcohol consumption,¹² sedentary lifestyle,^{13,14} and obesity,^{4,15,16} are established factors in inflammation. Recent research, however, indicates that social determinants are as important—if not more important—as health behaviors for shaping health.^{17,18} In fact, social determinants affect both the biological processes and health lifestyles of individuals.

Key social determinants of health include SES and race/ethnicity.¹⁹⁻²¹ Higher inflammation levels among racial/ethnic minorities, especially blacks,²²⁻²⁴ and individuals with lower SES^{25,26} have been reported, but several investigations do not corroborate these findings.^{1,27} Others argue that the role of SES in inflammation varies with SES measures²⁸ and racial/ethnic background.²⁹

Perceived discrimination (PD) has been suggested as another social factor with relevance to inflammation. PD has been associated with inflammation among young adults,³⁰ midlife adults,³¹ low-income black youths,³² and older blacks,³³ although the Dallas Heart Study showed no relationship between PD and inflammation among blacks, Hispanics, and whites.³⁴ In other studies, the link between PD and inflammation was limited to specific subpopulations, including women anticipating a racial threat³⁵ and non-obese women.³⁶

One important limitation of most prior research on inflammation is the lack of a theoretically grounded framework. To address this limitation, this study proposes a conceptual model of social determinants of inflammation (Figure 1) informed by fundamental cause theory (FCT), a sociological perspective. FCT postulates social determinants as key causes shaping health outcomes through multiple pathways that can evolve dynamically across life stages and historic periods in response to societal and technologic changes.³⁷⁻³⁹

Using the FCT, the proposed model specifies that SES, race, and PD act as key social determinants of inflammation. SES is a multidimensional construct consisting of income, education, and occupational prestige⁴⁰; this study focuses more specifically on income and education as dimensions strongly associated with health outcomes in the U.S.^{41,42} In the conceptual model, minority race contributes to lower SES²⁰ (Arrow 1) and to higher PD, which is also linked to lower SES⁴³⁻⁴⁷ (Arrows 2 and 3). Proximal factors are those through which social determinants influence inflammation. Stress, an established mechanism leading to poor health,⁴⁸⁻⁵⁰ is linked to all three social determinants specified in this model, with minorities, persons of lower SES, and those exposed to discrimination experiencing higher stress (Arrows 4-6). Stress harms health through overactivation of the biological stress response system, which may directly lead to increased inflammation (Arrow 7); furthermore, it contributes to unhealthy lifestyle and poorer psychological health (Arrows 8-10), which are also implicated in inflammation (Arrows 11-13).^{11-14,51-54}

Figure 1 illustrates the complexity of relations among social determinants and inflammation. In addition to effects through proximal factors, the model allows for direct effects of

unspecified mechanisms (Arrows 14–16). Reciprocal relationships reflect yet-unknown causal direction. Because of the model's complexity, comprehensive evaluation is outside the scope of this study. Instead, the study focuses on evaluating select components using three hypotheses:

1. Inflammation levels are higher among blacks compared with whites.
2. Inflammation levels decrease with higher SES.
3. Inflammation levels are higher among persons reporting PD.

This study sequentially evaluates the contribution of each group of proximal factors as suggested by the conceptual model, using a hierarchy-of-effects approach.⁵⁵ In addition to testing for the pooled sample, inflammation is modeled separately for blacks and whites because factors contributing to inflammation may vary by race/ethnicity.^{29,30,51–54}

METHODS

Data Sample

Data were obtained from the Survey of Midlife in the U.S., an ongoing national survey using a random-digit-dial sample representative of non-institutionalized English-speaking residents of 48 contiguous U.S. states who are aged 35 years.⁵⁶ The present study was limited to 1,054 participants in the biomarker sub-study that collected biological specimens (whites, $n=884$; blacks, $n=170$). Sub-study participants are similar to the national sample on age, sex, race, marital status, income, and health characteristics (subjective health, chronic conditions, activities of daily living, exercise, alcohol consumption, health insurance, physician visits) but are more educated and less likely to smoke.⁵⁷ Data collection took place between 2004 and 2009. Biological specimens and anthropometry were collected by trained staff during an overnight clinic stay. Demographic, social, and psychological indicators were measured using mail surveys and telephone interviews.

Measures

Biomarkers of systemic inflammation included C-reactive protein (CRP), which is produced by hepatocytes in response to infection or injury⁵⁸; interleukin 6 (IL-6), a proinflammatory cytokine; fibrinogen, a blood clotting factor involved in the coagulation response to vascular injury⁵⁹; and soluble E-selectin, an endothelial adhesion molecule expressed as a result of endothelial damage.⁶⁰ Fibrinogen concentrations (mg/dL) and CRP concentrations (ug/mL) were measured in citrated plasma using immunoturbidometric assay. Soluble E-selectin concentrations (ng/mL) and IL-6 concentrations (pg/mL) were measured in serum using enzyme-linked immunosorbent assay. Standardized procedures were used for fasting blood samples collection and processing.

Race was self-reported and categorized as black and white. Dimensions of SES were years of education and annual household income from all sources, measured in U.S. dollars and log-transformed. Age in years and gender (woman=1, man=0) were also included.

The Daily Discrimination and Lifetime Discrimination scales⁶¹ were used to measure PD. Consistent with the argument that PD is harmful to health regardless of the reason (race/

ethnicity, gender, or others),⁴⁷ these scales measure PD experiences of any type. The Daily Discrimination scale asks respondents how often they experience each of nine types of discrimination (e.g., being treated with less courtesy, less respect, or receiving poorer service at restaurants because of race/ethnicity, gender, age, religion, physical appearance, sexual orientation, or other characteristics [*never*=1, *rarely*=2, *sometimes*=3, *often*=4]). The Daily Discrimination scale totals the responses; higher values indicate higher levels of perceived discrimination. The Lifetime Discrimination scale measures experiences of major discriminatory events in life domains including employment, education, health care, and housing. Examples include not being hired for a job or being prevented from renting or buying a home. Respondents are asked how many times in their lifetime they have experienced each event. Lifetime discrimination is calculated as a total of items for which respondents indicate experiencing the event at least once.

Measures of generalized anxiety and depressed affect were based on Wang et al.⁶² The generalized anxiety scale consists of ten items, for example: *How often over the past 12 months were you restless because of your worry?*, the scale totals items for which *most days* was chosen. For depressed affect, respondents were asked: *During the past 12 months, was there ever a time when you felt sad, blue, or depressed for 2 weeks or more in a row?* (*yes/no*), and an additional seven items, for example, *During two weeks in past 12 months, when you felt sad, blue, or depressed, did you feel more tired out or low on energy than is usual?* (*yes/no*). The depressed affect scale totals *yes* answers on these seven items. This study also used three scales representing Negative Emotionality in Multidimensional Personality Questionnaire⁶³: stress reactivity (three items; e.g., *minor setbacks sometimes irritate me too much*), aggression (four items; e.g., *when I get angry I am often ready to hit someone*), and alienation (three items; e.g., *I would be more successful if people did not make things difficult for me*). Items use a 1–4 response scale (false to true). A sum of responses is calculated for each scale.

Participants' BMI, a measure of weight status (calculated as kg/m²), was based on anthropometric data. Two dichotomous indicators for weekly strenuous physical activity and weekly moderate physical activity were included (Appendix 1, available online), as well as two indicators of smoking capturing whether respondents ever smoked cigarettes regularly and whether they currently smoked cigarettes regularly (both *yes/no*). Because of potential effects on inflammation, current preventive use of aspirin was included. Finally, models controlled for chronic conditions during the past 12 months that had prevalence $\geq 5\%$ in the sample and showed relationships with inflammation at $p < 0.10$ in preliminary analyses. These conditions included high blood pressure/hypertension (henceforth hypertension), diabetes/high blood sugar (henceforth diabetes), joint/bone diseases, persistent skin trouble, teeth trouble, and sleep problems.

Statistical Analysis

First, descriptive statistics for the pooled sample and by race were obtained. *T*-tests were used to compare blacks with whites on continuous variables, and chi-square tests were performed for categorical variables. Next, series of multivariate linear regression models of each inflammation marker were estimated; robust estimators accounted for deviations from

normality. Because CRP and IL-6 had skewed distributions, they were log-transformed for modeling purposes. Model 1 included race and demographic covariates (gender, age). Model 2 added income and education. Model 3 further added PD measures. Model 4 added psychological factors. Model 5 added lifestyle (smoking indicators, physical activity indicators, and BMI). Finally, Model 6 added health characteristics, including preventive use of aspirin and chronic conditions. After estimating multivariate models for the pooled sample, Models 2–6 were estimated separately by race.

RESULTS

As shown in Table 1, blacks had higher concentrations of all four biomarkers of inflammation compared with whites ($p < 0.001$). They had lower SES as indicated by fewer years of education and lower income, and scored higher on both measures of PD, as well as generalized anxiety, alienation, and BMI (p -values < 0.001). They were also more likely to smoke regularly ($p < 0.01$) and less likely to engage in weekly physical activity (vigorous, $p < 0.05$; moderate, $p < 0.001$). They had higher rates of diabetes, teeth problems (p -values < 0.001), joint/bone disease, and sleep problems (p -values < 0.01), but lower rates of hypertension ($p < 0.001$) and preventive aspirin use ($p < 0.05$). Demographically, blacks were younger ($p < 0.001$) and more commonly women ($p < 0.01$).

Table 2 summarizes results of multivariate models of inflammation markers for the pooled sample. Because of space limitations, only Models 2, 3, and 6 are displayed; Models 1, 4, and 5 appear in Appendix 2 (available online). In Model 3, which included race, SES, PD, and demographic covariates, blacks had higher concentrations of fibrinogen ($p > 0.001$), IL-6 ($p > 0.001$), CRP ($p < 0.01$), and E-selectin ($p < 0.05$), lending support to Hypothesis 1. In fully adjusted models (Model 6), higher levels of IL-6 ($p < 0.001$) and fibrinogen ($p < 0.01$) among blacks persisted, though the coefficients underwent attenuation. For E-selectin, the black–white difference was explained after including PD, SES, psychological characteristics, and lifestyle factors in Model 5 (Table 1); for CRP, the difference was explained when health factors were controlled in Model 6. Hypothesis 2, which argues that inflammation decreases with SES, was supported for education but not for income in the pooled sample. Individuals with higher education had lower concentrations of fibrinogen ($p < 0.05$ in Model 6) and CRP ($p < 0.01$ in Model 3); a trend toward lower E-selectin was evident as well ($p < 0.10$ in Model 3). Lifetime perceived discrimination was related to higher concentrations of fibrinogen ($p < 0.05$ in Model 6), E-selectin ($p < 0.05$ in Model 3), and IL-6 ($p < 0.05$ in Model 3). These results support Hypothesis 3, which predicts increased inflammation with PD.

Tables 3 and 4 shows multivariate model estimates separately for whites and blacks. Results for whites were similar to the pooled sample and generally supportive of Hypothesis 2; one exception concerns income, which showed a positive relation to E-selectin ($p < 0.05$ in Model 6). Models for blacks supported the hypothesized inverse relationships between education and CRP ($p < 0.05$, Model 3) and between income on E-selectin ($p < 0.01$, Model 6). Hypothesis 3, however, was not supported for blacks, except for a marginally significant relationship between fibrinogen and lifetime perceived discrimination in Model 3 ($p < 0.10$).

BMI showed a positive relationship with three inflammation markers in the pooled sample (Table 2, Model 6), including fibrinogen ($p<0.05$), CRP ($p<0.001$), and IL-6 ($p<0.001$). Among whites, CRP concentrations decreased with moderate physical activity, and IL-6 concentrations decreased with vigorous physical activity (p -values <0.05 , Model 6, Table 3).

Among psychological factors, general anxiety was associated with higher CRP ($p<0.05$, Model 6, Table 4) and lower E-selectin ($p<0.01$) in blacks; there were also trends toward increased E-selectin with higher depressed affect and lower alienation in this racial group (p -values <0.10). Among whites, stress reactivity was linked to lower IL-6 ($p<0.05$, Model 6, Table 3), whereas alienation was linked to higher E-selectin ($p<0.05$, Model 6, Table 3). E-selectin concentrations further increased among blacks and whites who reported having diabetes ($p<0.05$, Model 6, Tables 3 and 4); among whites, diabetes was further linked to higher CRP ($p<0.01$, Model 6, Table 3). In whites, hypertension was inversely associated with IL-6 ($p<0.01$, Model 6, Table 3) and marginally with CRP ($p<0.10$); an inverse relationship between hypertension and CRP was also evident among blacks ($p<0.01$, Model 6, Table 4).

In models for the pooled sample (Table 2), older individuals had higher fibrinogen ($p<0.001$) and IL-6 ($p<0.001$), but lower E-selectin ($p<0.01$). Older whites had higher CRP ($p<0.05$, Model 3, Table 3), but the opposite was true for blacks ($p<0.01$ in Model 6, Table 4). Older blacks also had lower E-selectin ($p<0.05$, Model 3, Table 4). White women had higher fibrinogen and CRP ($p<0.001$, Model 6, Table 3), but lower E-selectin (p -values <0.01) compared with white men. Black women had higher fibrinogen and CRP (p -values <0.01 , Model 6, Table 4), as well as higher IL-6 concentrations ($p<0.05$, Model 3, Table 4) compared with black men.

DISCUSSION

Among the examined social determinants (race, SES, and PD), race was most consistently linked to inflammation, with blacks showing higher levels of all examined biomarkers. For fibrinogen and IL-6, racial differences tended to persist in fully adjusted models, whereas for E-selectin and CRP, racial differences were explained after including covariates. Consistent with research highlighting the importance of education for health,⁴¹ inverse associations between inflammation markers and educational attainment were observed, especially among whites, and an inverse association between income and E-selectin was found for blacks. For whites, there was an increase—not a decrease—of E-selectin with income, suggesting that higher income may have protective effects for blacks but not for whites. PD, the third social determinant considered in this study, was related to increased concentrations of most biomarkers of inflammation, but only for lifetime discrimination, not daily discrimination. In supplementary analyses (data not shown), bivariate associations between daily discrimination and inflammation markers were statistically significant but dissipated after controlling for lifetime discrimination. The reasons for this are unclear. Daily discrimination represents relatively minor events, such as being given poor service in a restaurant. Nevertheless, chronic exposure to such experiences may influence responses to major lifetime discriminatory events and other race-related stressors once they occur, increasing

physiologic and perceived stress and consequently harming health. More research is needed to disentangle these processes.

Importantly, the hypotheses proposed in this study received weaker support for blacks than for whites. This could be because of fewer blacks in the sample and lower statistical power, but the possibility that different mechanisms contribute to inflammation among different racial populations cannot be ruled out, especially in light of prior studies, in which education, weight status, and depressive symptoms showed weaker associations with inflammation among blacks compared with whites.^{29,51–53,64} By contrast, the results of this study were less consistent with prior research, suggesting that the effects of PD on health are mediated by psychological factors. Few statistically significant relationships between psychological factors and inflammation emerged. The absence of positive associations between stress reactivity and inflammation biomarkers was especially surprising, given the known role of stress in physical and mental health. One explanation is that the measure of stress reactivity does not capture the level of stress exposure; instead, it measures a relatively stable personality characteristic representing how a person responds emotionally once stress has occurred.⁶³ Future research should assess stress exposure more directly to clarify its influence on inflammation.

Limitations

This study has a cross-sectional, observational design, which prevents assessing changes over time and causally interpreting results, though specifically for race and education, it seems unlikely that they might change in response to changes in inflammation among midlife individuals. Racial identification tends to be stable over the life course, and education is typically completed during young adulthood. Nevertheless, inflammation may reduce SES by contributing to poorer health, which may limit earnings, productivity, and career advancement. This study addressed two dimensions of PD but it did not capture institutional, implicit, and covert discrimination. Finally, this investigation focused on blacks and whites; in future studies, it will be important to assess inflammation among Latinos and Native Americans, who have high prevalence of cardiovascular disease⁶⁵ and diabetes⁶⁶ and may be at risk of perceived discrimination.

CONCLUSIONS

As systemic inflammation is implicated in many chronic diseases, evidence of the role of social determinants in inflammation highlights the social origins of chronic disease during midlife and informs scholarship seeking to pinpoint the processes leading to health disparities. Better understanding is the first step toward preventive interventions to reduce the health risks among vulnerable populations, including racial/ethnic minorities, individuals at risk for discrimination, and those facing socioeconomic disadvantages. Notably, this work suggests the importance of systemic interventions that address large-scale social determinants of health, including system-level factors that underlie discrimination and lead to disparities in income and education.

Acknowledgments

Publication of this article was supported by the National Institutes of Health. The findings and conclusions in this article are those of the authors and do not necessarily represent the official position of the National Institutes of Health.

This work was supported by a grant from the National Institute on Minority Health and Health Disparities (U54MD008176).

No financial disclosures were reported by the authors of this paper.

References

1. Savoia C, Schiffrin EL. Inflammation in hypertension. *Curr Opin Nephrol Hypertens*. 2006; 15(2): 152–158. <http://dx.doi.org/10.1097/01.mnh.0000203189.57513.76>. [PubMed: 16481882]
2. Everett BM, Pradhan AD, Solomon DH, et al. Rationale and design of the Cardiovascular Inflammation Reduction Trial: a test of the inflammatory hypothesis of atherothrombosis. *Am Heart J*. 2013; 166(2):199–207. <http://dx.doi.org/10.1016/j.ahj.2013.03.018>. [PubMed: 23895801]
3. Haffner SM. The metabolic syndrome: inflammation, diabetes mellitus, and cardiovascular disease. *Am J Cardiol*. 2006; 97(2):3–11. <http://dx.doi.org/10.1016/j.amjcard.2005.11.010>. [PubMed: 16675316]
4. Olefsky JM, Glass CK. Macrophages, inflammation, and insulin resistance. *Annu Rev Physiol*. 2010; 72:219–246. <http://dx.doi.org/10.1146/annurev-physiol-021909-135846>. [PubMed: 20148674]
5. Kolb H, Mandrup-Poulsen T. The global diabetes epidemic as a consequence of lifestyle-induced low-grade inflammation. *Diabetologia*. 2010; 53(1):10–20. <http://dx.doi.org/10.1007/s00125-009-1573-7>. [PubMed: 19890624]
6. Shiraiishi, FG., Stringuetta Belik, F., e Silva, O., et al. Inflammation, diabetes, and chronic kidney disease: role of aerobic capacity. *Exp Diabetes Res*. 2012. <http://dx.doi.org/10.1155/2012/750286>
7. Hu FB. Globalization of diabetes: the role of diet, lifestyle, and genes. *Diabetes Care*. 2011; 34(6): 1249–1257. <http://dx.doi.org/10.2337/dc11-0442>. [PubMed: 21617109]
8. Elinav E, Nowarski R, Thaiss CA, et al. Inflammation-induced cancer: crosstalk between tumours, immune cells and microorganisms. *Nat Rev Cancer*. 2013; 13(11):759–771. <http://dx.doi.org/10.1038/nrc3611>. [PubMed: 24154716]
9. Trinchieri G. Cancer and inflammation: an old intuition with rapidly evolving new concepts. *Annu Rev Immunol*. 2012; 30:677–706. <http://dx.doi.org/10.1146/annurev-immunol-020711-075008>. [PubMed: 22224761]
10. Kamp DW, Shacter E, Weitzman SA. Chronic inflammation and cancer: the role of the mitochondria. *Oncology*. 2011; 25(5):400–410 413. [PubMed: 21710835]
11. Arnsen Y, Shoenfeld Y, Amital H. Effects of tobacco smoke on immunity, inflammation and autoimmunity. *J Autoimmun*. 2010; 34(3):J258–J265. <http://dx.doi.org/10.1016/j.jaut.2009.12.003>. [PubMed: 20042314]
12. Wang HJ, Zakhari S, Jung MK. Alcohol, inflammation, and gut-liver-brain interactions in tissue damage and disease development. *World J Gastroenterol*. 2010; 16(11):1304–1313. <http://dx.doi.org/10.3748/wjg.v16.i11.1304>. [PubMed: 20238396]
13. Woods JA, Vieira VJ, Keylock KT. Exercise, inflammation, and innate immunity. *Immunol Allergy Clin North Am*. 2009; 29(2):381–393. <http://dx.doi.org/10.1016/j.iac.2009.02.011>. [PubMed: 19389588]
14. Beavers KM, Brinkley TE, Nicklas BJ. Effect of exercise training on chronic inflammation. *Clin Chim Acta*. 2010; 411(11–12):785–793. <http://dx.doi.org/10.1016/j.cca.2010.02.069>. [PubMed: 20188719]
15. Hajer GR, van Haften TW, Visseren FL. Adipose tissue dysfunction in obesity, diabetes, and vascular diseases. *Eur Heart J*. 2008; 29(24):2959–2971. <http://dx.doi.org/10.1093/eurheartj/ehn387>. [PubMed: 18775919]

16. Pischon N, Heng N, Bernimoulin J-P, et al. Obesity, inflammation, and periodontal disease. *J Dent Res.* 2007; 86(5):400–409. <http://dx.doi.org/10.1177/154405910708600503>. [PubMed: 17452558]
17. Marmot M, Allen JJ. Social determinants of health equity. *Am J Public Health.* 2014; 104(S4):S517–S519. <http://dx.doi.org/10.2105/AJPH.2014.302200>. [PubMed: 25100411]
18. Braveman PA, Egerter SA, Mockenhaupt RE. Broadening the focus: the need to address the social determinants of health. *Am J Prev Med.* 2011; 40(1):S4–S18. <http://dx.doi.org/10.1016/j.amepre.2010.10.002>. [PubMed: 21146778]
19. Braveman PA, Cubbin C, Egerter S, Williams DR, Pamuk E. Socioeconomic disparities in health in the United States: what the patterns tell us. *Am J Public Health.* 2010; 100(suppl 1):S186–S196. <http://dx.doi.org/10.2105/AJPH.2009.166082>. [PubMed: 20147693]
20. Williams DR, Mohammed SA, Leavell J, Collins C. Race, socioeconomic status, and health: complexities, ongoing challenges, and research opportunities. *Ann N Y Acad Sci.* 2010; 1186:69–101. <http://dx.doi.org/10.1111/j.1749-6632.2009.05339.x>. [PubMed: 20201869]
21. Frieden TR. CDC Health Disparities and Inequalities Report–United States, 2013. Foreword. *MMWR Surveill Summ.* 2013; 62:1–2.
22. Kelley-Hedgpeath A, Lloyd-Jones DM, Colvin A, et al. Ethnic differences in C-reactive protein concentrations. *Clin Chem.* 2008; 54(6):1027–1037. <http://dx.doi.org/10.1373/clinchem.2007.098996>. [PubMed: 18403563]
23. Khera A, McGuire DK, Murphy SA, et al. Race and gender differences in C-reactive protein levels. *J Am Coll Cardiol.* 2005; 46(3):464–469. <http://dx.doi.org/10.1016/j.jacc.2005.04.051>. [PubMed: 16053959]
24. Paalani M, Lee JW, Haddad E, Tonstad S. Determinants of inflammatory markers in a bi-ethnic population. *Ethn Dis.* 2011; 21(2):142. [PubMed: 21749016]
25. Pollitt RA, Kaufman JS, Rose KM, et al. Early-life and adult socioeconomic status and inflammatory risk markers in adulthood. *Eur J Epidemiol.* 2007; 22(1):55–66. <http://dx.doi.org/10.1007/s10654-006-9082-1>. [PubMed: 17225957]
26. Jousilahti P, Salomaa V, Rasi V, Vahtera E, Palosuo T. Association of markers of systemic inflammation, C reactive protein, serum amyloid A, and fibrinogen, with socioeconomic status. *J Epidemiol Community Health.* 2003; 57(9):730–733. <http://dx.doi.org/10.1136/jech.57.9.730>. [PubMed: 12933781]
27. Ford ES, Giles WH, Myers GL, et al. C-reactive protein concentration distribution among U.S. children and young adults: findings from the National Health and Nutrition Examination Survey, 1999–2000. *Clin Chem.* 2003; 49(8):1353–1357. <http://dx.doi.org/10.1373/49.8.1353>. [PubMed: 12881452]
28. Friedman EM, Herd P. Income, education, and inflammation: differential associations in a national probability sample (the MIDUS study). *Psychosom Med.* 2010; 72(3):290. <http://dx.doi.org/10.1097/PSY.0b013e3181cfe4c2>. [PubMed: 20100883]
29. Fuller-Rowell TE, Curtis DS, Doan SN, Coe CL. Racial disparities in the health benefits of educational attainment: a study of inflammatory trajectories among African American and white adults. *Psychosom Med.* 2015; 77(1):33–40. <http://dx.doi.org/10.1097/PSY.0000000000000128>. [PubMed: 25490696]
30. Cunningham TJ, Seeman TE, Kawachi I, et al. Racial/ethnic and gender differences in the association between self-reported experiences of racial/ethnic discrimination and inflammation in the CARDIA cohort of 4 U.S. communities. *Soc Sci Med.* 2012; 75(5):922–931. <http://dx.doi.org/10.1016/j.socscimed.2012.04.027>. [PubMed: 22682683]
31. Friedman EM, Williams DR, Singer BH, Ryff CD. Chronic discrimination predicts higher circulating levels of E-selectin in a national sample: the MIDUS study. *Brain Behav Immun.* 2009; 23(5):684–692. <http://dx.doi.org/10.1016/j.bbi.2009.01.002>. [PubMed: 19171188]
32. Goosby BJ, Malone S, Richardson EA, Cheadle JE, Williams DT. Perceived discrimination and markers of cardiovascular risk among low-income African American youth. *Am J Hum Biol.* 2015; 27(4):546–552. <http://dx.doi.org/10.1002/ajhb.22683>. [PubMed: 25753652]
33. Lewis TT, Aiello AE, Leurgans S, Kelly J, Barnes LL. Self-reported experiences of everyday discrimination are associated with elevated C-reactive protein levels in older African-American

- adults. *Brain Behav Immun*. 2010; 24(3):438–443. <http://dx.doi.org/10.1016/j.bbi.2009.11.011>. [PubMed: 19944144]
34. Albert MA, Ravenell J, Glynn RJ, et al. Cardiovascular risk indicators and perceived race/ethnic discrimination in the Dallas Heart Study. *Am Heart J*. 2008; 156(6):1103–1109. <http://dx.doi.org/10.1016/j.ahj.2008.07.027>. [PubMed: 19033005]
 35. Nuru-Jeter A, Chae DH, Price M, et al. Anticipatory racism threat and superwoman schema: elucidating the relationship between racial discrimination and chronic inflammation. *Circulation*. 2013; 128(22 suppl):A9550.
 36. Beatty Moody DL, Brown C, Matthews KA, Bromberger JT. Everyday discrimination prospectively predicts inflammation across 7–years in racially diverse midlife women: Study of Women’s Health across the Nation. *J Soc Issues*. 2014; 70(2):298–314. <http://dx.doi.org/10.1111/josi.12061>. [PubMed: 25342861]
 37. Link BG, Phelan J. Social conditions as fundamental causes of disease. *J Health Soc Behav*. 1995; 36:80–94. <http://dx.doi.org/10.2307/2626958>.
 38. Phelan JC, Link BG. Is racism a fundamental cause of inequalities in health? *Annu Rev Sociol*. 2015; 41:311–330. <http://dx.doi.org/10.1146/annurev-soc-073014-112305>.
 39. Phelan JC, Link BG, Tehranifar P. Social conditions as fundamental causes of health inequalities theory, evidence, and policy implications. *J Health Soc Behav*. 2010; 51(1 suppl):S28–S40. <http://dx.doi.org/10.1177/0022146510383498>. [PubMed: 20943581]
 40. Galobardes, B., Lynch, J., Smith, GD. Measuring socioeconomic position in health research; *Br Med Bull*. 2007. p. 81-82. <http://dx.doi.org/10.1093/bmb/ldm001>
 41. Mirowsky, J., Ross, CE. *Education, Social Status, and Health*. Livingston, NJ: Transaction Publishers; 2003.
 42. Lynch, J., Kaplan, GA. Socioeconomic position. In: Berkman, LF., Kawachi, I., editors. *Social Epidemiology*. New York: Oxford University Press; 2000. p. 13-35.
 43. Fuller-Rowell TE, Evans GW, Ong AD. Poverty and health: the mediating role of perceived discrimination. *Psychol Sci*. 2012; 23(7):734–739. <http://dx.doi.org/10.1177/0956797612439720>. [PubMed: 22700331]
 44. Jackson, JS., Williams, D., Torres, M. *Perceptions of Discrimination: The Stress Process and Physical and Psychological Health*. Washington, DC: National Institute of Mental Health; 1997.
 45. Sigelman, L., Welch, S. *Black Americans’ Views of Racial Inequality: The Dream Deferred*. New York: Cambridge University Press; 1991.
 46. Seaton EK, Caldwell CH, Sellers RM, Jackson JS. The prevalence of perceived discrimination among African American and Caribbean Black youth. *Dev Psychol*. 2008; 44(5):1288. <http://dx.doi.org/10.1037/a0012747>. [PubMed: 18793063]
 47. Gibbons FX, Gerrard M, Cleveland MJ, Wills TA, Brody G. Perceived discrimination and substance use in African American parents and their children: a panel study. *J Pers Soc Psychol*. 2004; 86(4):517. <http://dx.doi.org/10.1037/0022-3514.86.4.517>. [PubMed: 15053703]
 48. Pearlin LI, Schieman S, Fazio EM, Meersman SC. Stress, health, and the life course: some conceptual perspectives. *J Health Soc Behav*. 2005; 46(2):205–219. <http://dx.doi.org/10.1177/002214650504600206>. [PubMed: 16028458]
 49. Pearlin LI. The sociological study of stress. *J Health Soc Behav*. 1989; 30(3):241–256. <http://dx.doi.org/10.2307/2136956>. [PubMed: 2674272]
 50. Thoits PA. Stress and health: major findings and policy implications. *J Health Soc Behav*. 2010; 51(suppl):S41–S53. <http://dx.doi.org/10.1177/0022146510383499>. [PubMed: 20943582]
 51. Vraney EA, Berntson JM, Khambaty T, Stewart JC. Depressive symptoms clusters and insulin resistance: race/ethnicity as a moderator in 2005–2010 NHANES data. *Ann Behav Med*. 2016; 50(1):1–11. <http://dx.doi.org/10.1007/s12160-015-9725-0>. [PubMed: 26318593]
 52. Case SM, Stewart JC. Race/ethnicity moderates the relationship between depressive symptom severity and C-reactive protein: 2005–2010 NHANES data. *Brain Behav Immun*. 2014; 41:101–108. <http://dx.doi.org/10.1016/j.bbi.2014.04.004>. [PubMed: 24859042]
 53. Stewart JC. One effect size does not fit all: is the depression-inflammation link missing in racial/ethnic minority individuals? *JAMA Psychiatry*. 2016; 73(3):301–302. <http://dx.doi.org/10.1001/jamapsychiatry.2015.3205>. [PubMed: 26842009]

54. Deverts DJ, Cohen S, DiLillo VG, et al. Depressive symptoms, race, and circulating C-reactive protein: the Coronary Artery Risk Development in Young Adults (CARDIA) study. *Psychosom Med.* 2010; 72(8):734. <http://dx.doi.org/10.1097/PSY.0b013e3181ec4b98>. [PubMed: 20668285]
55. Murray KB, Vogel CM. Using a hierarchy-of-effects approach to gauge the effectiveness of corporate social responsibility to generate goodwill toward the firm: financial versus nonfinancial impacts. *J Bus Res.* 1997; 38(2):141–159. [http://dx.doi.org/10.1016/S0148-2963\(96\)00061-6](http://dx.doi.org/10.1016/S0148-2963(96)00061-6).
56. Ryff, CD., Seeman, T., Weinstein, M. National Survey of Midlife Development in the United States (MIDUS II): Biomarker Project, 2004–2009. ICPSR29282-v6. Ann Arbor, MI: Inter-university Consortium for Political and Social Research [distributor]; 2013. 05-02
57. Dienberg Love G, Seeman TE, Weinstein M, Ryff CD. Bioindicators in the MIDUS national study: protocol, measures, sample, and comparative context. *J Aging Health.* 2010; 22(8):1059–1080. <http://dx.doi.org/10.1177/0898264310374355>. [PubMed: 20876364]
58. Berg AH, Scherer PE. Adipose tissue, inflammation, and cardiovascular disease. *Circ Res.* 2005; 96(9):939–949. <http://dx.doi.org/10.1161/01.RES.0000163635.62927.34>. [PubMed: 15890981]
59. Toker S, Shirom A, Shapira I, Berliner S, Melamed S. The association between burnout, depression, anxiety, and inflammation biomarkers: C-reactive protein and fibrinogen in men and women. *J Occup Health Psychol.* 2005; 10(4):344. <http://dx.doi.org/10.1037/1076-8998.10.4.344>. [PubMed: 16248685]
60. Ross R. Atherosclerosis—an inflammatory disease. *N Engl J Med.* 1999; 340(2):115–126. <http://dx.doi.org/10.1056/NEJM199901143400207>. [PubMed: 9887164]
61. Williams D, Yu Y, Jackson JS, Anderson NB. Racial differences in physical and mental health: socioeconomic status, stress and discrimination. *J Health Psychol.* 1997; 2(3):335–351. <http://dx.doi.org/10.1177/135910539700200305>. [PubMed: 22013026]
62. Wang PS, Berglund P, Kessler RC. Recent care of common mental disorders in the United States: prevalence and conformance with evidence-based recommendations. *J Gen Intern Med.* 2000; 15(5):284–292. <http://dx.doi.org/10.1046/j.1525-1497.2000.9908044.x>. [PubMed: 10840263]
63. Patrick CJ, Curtin JJ, Tellegen A. Development and validation of a brief form of the Multidimensional Personality Questionnaire. *Psychol Assess.* 2002; 14(2):150–163. <http://dx.doi.org/10.1037/1040-3590.14.2.150>. [PubMed: 12056077]
64. Choi J, Joseph L, Pilote L. Obesity and C-reactive protein in various populations: a systematic review and meta-analysis. *Obes Rev.* 2013; 14(3):232–244. <http://dx.doi.org/10.1111/obr.12003>. [PubMed: 23171381]
65. Daviglius ML, Talavera GA, Avilés-Santa ML, et al. Prevalence of major cardiovascular risk factors and cardiovascular diseases among Hispanic/Latino individuals of diverse backgrounds in the United States. *JAMA.* 2012; 308(17):1775–1784. <http://dx.doi.org/10.1001/jama.2012.14517>. [PubMed: 23117778]
66. O’Connell J, Yi R, Wilson C, Manson SM, Acton KJ. Racial disparities in health status: a comparison of the morbidity among American Indian and U.S. adults with diabetes. *Diabetes Care.* 2010; 33(7):1463–1470. <http://dx.doi.org/10.2337/dc09-1652>. [PubMed: 20357367]




Figure 1.
Social determinants of systemic inflammation: conceptual model.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Characteristics of the Pooled Sample and by Racial Background

Variable ^a	All (n=1,054)	Blacks (n=170)	Whites (n=884)
Systemic inflammation markers			
Fibrinogen, mg/dL (range, 94.0–857.0)	348.36 (88.24)	388.70 (101.67)***	340.61 (83.27)
E-selectin, ng/mL (range, 0.1–161.9)	42.51 (22.01)	49.19 (25.75)***	41.22 (20.99)
CRP, ug/mL (range, 0.1–59.3)	3.11 (5.01)	4.78 (6.90)***	2.79 (4.49)
IL-6, pg/mL (range, 0.2–21.8)	2.97 (2.90)	3.91 (3.05)***	2.79 (2.83)
SES			
Education, years (range, 2–20)	14.75 (2.56)	13.58 (2.75)***	14.98 (2.46)
Income, log \$ (range, 0–30)	10.27 (3.77)	9.81 (3.45)***	10.36 (3.83)
Perceived discrimination			
Daily (range, 9–32)	12.87 (4.60)	14.65 (6.46)***	12.53 (4.07)
Lifetime (range, 0–11)	1.23 (1.90)	3.02 (2.82)***	0.88 (1.44)
Psychological factors			
Depression (range, 0–7)	0.72 (1.85)	0.88 (2.06)	0.69 (1.81)
Anxiety (range, 0–10)	0.15 (.94)	0.41 (1.64)***	0.10 (.73)
Stress reactivity (range, 3–12)	6.15 (2.31)	6.45 (2.61)	6.09 (2.24)
Aggression (range, 4–14)	5.31 (1.66)	5.36 (1.65)	5.30 (1.66)
Alienation (range, 3–12)	5.14 (1.89)	6.04 (2.39)***	4.96 (1.73)
Lifestyle factors			
Ever regular smoker	44.6	55.9**	42.4
Currently regular smoker	12.9	25.9***	10.4
Vigorous physical activity	30.5	22.4*	32.0
Moderate physical activity	44.1	30.6***	46.7
BMI (range, 14.23–161.10)	30.6 (13.99)	34.59 (17.79)***	29.84 (13.01)
Health factors			
Preventive aspirin	31.2	24.7*	32.5
Chronic conditions			
Hypertension	72.0	58.8***	75.0
Diabetes	10.0	18.8***	8.3
Joint/bone diseases	27.0	37.6**	25.0
Persistent skin trouble	9.1	7.1	9.5
Teeth problems	6.5	14.1***	5.1
Sleep problems	12.6	20.6**	11.1
Demographic factors			
Age, y (range, 35–82)	54.56 (11.62)	51.27 (10.48)***	55.19 (11.73)
Woman	56.6	67.6**	54.5

Source: Survey of Midlife in the U.S. (MIDUS II).

Note: Values are M (SD) for continuous variables and percentages for dichotomous variables. *T*-tests were used to compare blacks to whites on continuous variables. χ^2 tests were used to compare blacks to whites on categorical variables. Boldface indicates statistical significance of the differences between blacks and whites (* p <0.05, ** p <0.01, *** p <0.001; two-tailed tests).

^aRanges are given for continuous variables only.

CRP, C-reactive protein; IL-6, interleukin 6; y, years.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2
Coefficients from Linear Regression Models for Systemic Inflammation Markers: Pooled Sample (*n*=1,054)

Variable	Fibrinogen ^a			E-selectin ^b			CRP ^c			IL-6 ^d		
	Model 2	Model 3	Model 6	Model 2	Model 3	Model 6	Model 2	Model 3	Model 6	Model 2	Model 3	Model 6
Demographic factors												
Age, y	1.13*** (0.25)	1.17*** (0.25)	1.07*** (0.28)	-0.19** (0.05)	-0.17** (0.06)	-0.23** (0.07)	0.005 (0.003)	0.01 (0.003)	0.001 (0.003)	0.01*** (0.002)	0.01*** (0.002)	0.01*** (0.002)
Woman	29.51*** (5.15)	28.33*** (5.14)	30.18** (5.46)	-4.25** (1.35)	-4.54** (1.34)	-4.15** (1.40)	0.37*** (0.07)	0.36*** (0.07)	0.07*** (0.39)	0.06 (0.04)	0.05 (0.05)	0.07 (0.05)
Social determinants												
Race, black	45.49*** (8.34)	34.53*** (9.51)	29.25** (9.71)	7.25** (2.10)	4.52* (2.28)	2.52 (2.34)	0.45*** (0.10)	0.36** (0.11)	0.16 (0.12)	0.43*** (0.06)	0.36*** (0.07)	0.28*** (0.07)
SES												
Education, y	-2.59** (1.07)	-2.86** (1.09)	-2.34* (1.10)	-0.43 (0.027)	-0.46 (0.27)	-0.21 (0.28)	-0.04** (0.01)	-0.04** (0.02)	-0.03 (0.02)	-0.01 (0.01)	-0.02 (0.01)	-0.001 (0.01)
Income, log \$	0.82 (0.56)	0.91 (0.56)	0.82 (0.57)	0.11 (0.014)	0.13 (0.13)	0.08 (0.15)	0.01 (0.01)	0.01 (0.01)	0.01 (0.01)	0.0004 (0.01)	0.001 (0.01)	-0.003 (0.01)
Perceived discrimination												
Daily		-0.06 (0.69)	-0.42 (0.73)		0.18 (0.18)	0.07 (0.19)		0.01 (0.01)	0.003 (0.01)		0.002 (0.01)	-0.01 (0.01)
Lifetime		5.18** (1.95)	4.52* (1.90)		1.14* (0.47)	0.81 (0.48)		0.04 (0.02)	0.02 (0.02)		0.03* (0.01)	0.02 (0.01)
Psychological factors												
Depressed affect			0.46 (1.71)			0.10 (0.43)			-0.01 (0.02)			0.02 (0.02)
General anxiety			-2.33 (2.66)		-1.71** (0.62)				0.05 (0.04)			0.001 (0.03)
Stress reactivity			-0.88 (1.37)		-0.33 (0.38)				-0.01 (0.02)			-0.02 (0.01)
Aggression			0.14 (1.89)		0.32 (0.47)				-0.01 (0.02)			0.02 (0.01)
Alienation			1.31 (1.68)		0.53 (0.47)				0.02 (0.02)			0.03 (0.01)
Lifestyle factors												

Variable	Fibrinogen ^a			E-selectin ^b			CRP ^c			IL-6 ^d		
	Model 2	Model 3	Model 6	Model 2	Model 3	Model 6	Model 2	Model 3	Model 6	Model 2	Model 3	Model 6
Ever regular smoker	-3.30 (6.11)				1.51 (1.52)				0.04 (0.08)			0.07 (0.05)
Currently regular smoker	0.78 (9.05)				2.18 (2.63)				0.06 (0.21)			0.01 (0.07)
Vigorous physical activity	1.88 (6.63)				-1.98 (1.64)				-0.07 (0.09)			-0.12* (0.06)
Moderate physical activity	-9.65 (6.15)				-0.98 (1.55)				-0.16 (0.08)			-0.07 (0.05)
BMI	0.63* (0.24)				0.02 (0.04)				0.01*** (0.003)			0.01*** (0.0002)
Health factors												
Preventive aspirin	5.13 (6.51)				-1.21 (1.65)				0.13 (0.08)			0.02 (0.05)
Hypertension	-6.43 (6.72)				-2.34 (1.62)				-0.24** (0.08)			-0.18** (0.05)
Diabetes	16.14 (9.75)				8.75** (2.84)				0.31** (0.11)			0.06 (0.07)
Joint/bone diseases	-8.06 (6.72)				1.68 (1.65)				0.09 (0.08)			-0.01 (0.05)
Persistent skin trouble	14.01 (10.17)				0.94 (2.69)				-0.08 (0.12)			0.12 (0.08)
Teeth problems	17.93 (11.90)				4.21 (3.81)				0.26 (0.15)			0.04 (0.09)
Sleep problems	3.89 (9.00)				0.18 (2.33)				0.13 (0.12)			0.07 (0.07)
Intercept	292.40***	290.03***	284.89***	59.31***	55.32***	57.42***	0.38	0.24	0.46	0.15	0.09	0.12
Model F	17.94***	15.08***	5.69***	8.24***	7.82***	3.33***	14.10***	11.66***	7.82***	23.16***	17.30***	8.77***
R ² , %	9.45	10.44	13.23	3.83	5.06	8.74	6.36	6.86	14.10	8.85	9.46	15.68

Source: Survey of Midlife in the U.S. (MIDUS II).

Note: Values are coefficients unless otherwise noted; standard errors appear in parentheses. Robust estimators are used. Boldface indicates statistical significance (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; two-tailed tests).

^a mg/dL.

^b ng/mL.

^c log ug/mL.

CRP, C-reactive protein; IL-6, interleukin 6; Y, years.

\log_{10} pg/mL

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3

Coefficients from Linear Regression Models for Systemic Inflammation Markers by Racial Background

Variable	Whites (n=884)											
	Fibrinogen ^a		E-selectin ^b		CRP ^c		IL-6 ^d					
	Model 2	Model 3	Model 2	Model 3	Model 2	Model 3	Model 2	Model 3	Model 2	Model 3	Model 2	Model 3
Demographic factors												
Age, y	0.16*** (0.25)	1.20*** (0.25)	1.17*** (0.29)	-0.15** (0.06)	-0.12* (0.06)	-0.19** (0.07)	0.01* (0.003)	0.01* (0.004)	0.02*** (0.002)	0.01*** (0.002)	0.01*** (0.002)	0.01*** (0.002)
Woman	24.93*** (5.43)	23.11*** (5.37)	26.91*** (4.65)	-4.66** (1.43)	-5.21*** (1.43)	-4.56** (1.50)	0.34*** (0.08)	0.32*** (0.08)	0.37*** (0.08)	0.004 (0.05)	0.03 (0.05)	0.04 (0.05)
Social determinants												
SES												
Education, y	-2.87* (1.20)	-3.06* (1.23)	-2.33 (1.22)	-0.54 (0.29)	-0.48 (0.29)	-0.23 (0.29)	-0.04* (0.02)	-0.04* (0.02)	-0.02 (0.02)	-0.01 (0.01)	-0.01 (0.01)	0.0004 (0.01)
Income, log \$	0.71 (0.48)	0.76 (0.50)	0.62 (0.51)	0.31** (0.10)	0.33* (0.11)	0.31* (0.12)	0.02 (0.01)	0.02 (0.01)	0.01 (0.01)	-0.001 (0.01)	-0.0003 (0.01)	-0.003 (0.01)
Perceived discrimination												
Daily		-0.39 (0.78)	-0.59 (0.81)		0.37 (0.20)	0.23 (0.22)	0.01 (0.01)	0.01 (0.01)	0.002 (0.01)	0.01 (0.01)	0.01 (0.01)	0.001 (0.01)
Lifetime		5.39* (2.47)	4.47 (2.35)		1.21* (0.61)	0.70 (0.62)	0.05 (0.03)	0.05 (0.03)	0.03 (0.03)	0.06** (0.02)	0.04* (0.02)	
Psychological factors												
Depressed affect			-0.51 (1.82)			-0.27 (0.43)			-0.01 (0.02)			0.02 (0.02)
General anxiety			-1.17 (2.99)			-1.37 (0.96)			0.002 (0.05)			-0.02 (0.03)
Stress reactivity			-1.74 (1.46)			-0.66 (0.40)			-0.01 (0.02)			-0.03* (0.01)
Aggression			1.13 (2.01)			0.41 (0.49)			0.003 (0.03)			0.02 (0.02)
Alienation			1.79 (1.93)			1.14* (0.54)			0.01 (0.03)			0.02 (0.02)
Lifestyle factors												

Variable	Whites (n=884)											
	Fibrinogen ^a			E-selectin ^b			CRP ^c			IL-6 ^d		
	Model 2	Model 3	Model 6	Model 2	Model 3	Model 6	Model 2	Model 3	Model 6	Model 2	Model 3	Model 6
Ever regular smoker			-2.03 (6.31)			2.29 (1.61)			0.05 (0.09)			0.07 (0.05)
Currently regular smoker			2.58 (9.91)			-1.02 (2.83)			0.03 (0.14)			-0.04 (0.08)
Vigorous physical activity			0.07 (7.02)			-2.04 (1.70)			-0.09 (0.10)			-0.15* (0.06)
Moderate physical activity			-10.17 (6.44)			-0.71 (1.61)			-0.20* (0.09)			-0.08 (0.06)
BMI			0.48 (0.31)			0.09 (0.05)			0.01** (0.004)			0.01** (0.002)
Health factors												
Preventive aspirin			4.27 (6.95)			-1.45 (1.72)			0.13 (0.09)			0.03 (0.06)
Hypertension			-4.67 (7.48)			-1.18 (1.70)			-0.18 (0.10)			-0.19** (0.06)
Diabetes			14.77 (11.18)			6.76* (3.27)			0.37** (0.13)			0.04 (0.09)
Joint/bone diseases			14.02 (7.31)			2.92 (1.73)			0.04 (0.09)			-0.01 (0.06)
Persistent skin trouble			15.98 (10.65)			-0.54 (2.49)			-0.01 (0.12)			0.13 (0.09)
Teeth problems			20.87 (15.09)			7.70 (4.57)			0.19 (0.19)			0.09 (0.11)
Sleep problems			5.84 (10.29)			0.29 (2.44)			0.13 (0.14)			0.05 (0.09)
Intercept	298.34***	299.59***	286.24***	56.94***	48.50***	45.83***	0.18	0.00	0.09	0.11	0.06	0.15
Model F	12.96***	9.14***	3.74***	7.23***	6.39***	2.99***	8.31***	6.72***	4.40	14.40***	11.75***	5.65***
R ² , %	5.63	6.37	9.38	2.53	4.13	8.44	3.43	4.00	10.54	5.75	7.27	13.86

Source: Survey of Midlife in the U.S. (MIDUS II).

Note: Values are coefficients unless otherwise noted; standard errors appear in parentheses. Robust estimators are used. Boldface indicates statistical significance (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; two-tailed tests).

^a mg/dL.

^b ng/mL.

CRP, C-reactive protein; IL-6, interleukin 6; y, years.

d_{\log} pg/mL.

c_{\log} ug/mL.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 4

Coefficients from Linear Regression Models for Systemic Inflammation Markers by Racial Background

Variable	Blacks (n=170)											
	Fibrinogen ^a		E-selectin ^b		CRP ^c		IL-6 ^d					
	Model 2	Model 3	Model 2	Model 3	Model 2	Model 3	Model 2	Model 3				
Demographic factors												
Age, y	0.75 (0.91)	0.69 (0.94)	0.33 (1.08)	-0.42* (0.16)	-0.44* (0.17)	-0.34 (0.24)	-0.01 (0.01)	-0.03** (0.01)	0.01 (0.01)	0.01 (0.01)	0.01 (0.01)	
Woman	57.45*** (15.02)	62.13*** (14.51)	49.25** (16.82)	-2.39 (4.13)	-1.67 (4.24)	-2.42 (4.58)	0.64** (0.19)	0.67*** (0.19)	0.52** (0.19)	0.29* (0.12)	0.29* (0.12)	0.23 (0.13)
Social determinants												
SES												
Education, y	-1.96 (2.34)	-3.14 (2.40)	-2.85 (2.85)	0.05 (0.68)	-0.18 (0.72)	0.01 (0.75)	-0.06* (0.03)	-0.07* (0.03)	-0.05 (0.03)	-0.02 (0.02)	-0.02 (0.02)	-0.01 (0.02)
Income, log \$	1.98 (2.82)	2.36 (2.49)	1.75 (2.92)	-1.10* (0.54)	-0.99* (0.49)	-1.41** (0.45)	0.01 (0.01)	0.01 (0.01)	0.03 (0.02)	0.02 (0.01)	0.02 (0.01)	0.02 (0.01)
Perceived discrimination												
Daily		0.68 (1.50)	-0.11 (1.75)		-0.16 (0.40)	-0.08 (0.37)		0.01 (0.02)	-0.01 (0.02)		0.004 (0.01)	-0.01 (0.01)
Lifetime		5.57 (3.28)	4.91 (3.52)		1.26 (0.83)	1.04 (0.81)		0.03 (0.04)	0.02 (0.04)		0.01 (0.02)	-0.001 (0.03)
Psychological factors												
Depressed affect			5.58 (5.49)			2.39 (1.32)			0.06 (0.05)			0.04 (0.04)
General anxiety			-5.11 (5.83)		-3.20** (1.15)				0.10* (0.05)			0.03 (0.05)
Stress reactivity			2.47 (3.81)		1.29 (0.98)				-0.04 (0.04)			0.02 (0.03)
Aggression			-6.04 (5.43)		-0.99 (1.59)				-0.10 (0.07)			-0.02 (0.04)
Alienation			-1.19 (3.76)		-1.53 (0.91)				0.06 (0.04)			0.02 (0.03)
Lifestyle factors												

Variable	Blacks (n=170)											
	Fibrinogen ^d			E-selectin ^b			CRP ^c			IL-6 ^d		
	Model 2	Model 3	Model 6	Model 2	Model 3	Model 6	Model 2	Model 3	Model 6	Model 2	Model 3	Model 6
Ever regular smoker			-14.13 (18.83)			-3.80 (4.18)			-0.01 (0.20)			-0.02 (0.12)
Currently regular smoker			1.95 (21.96)			10.07 (6.51)			-0.02 (0.22)			0.16 (0.14)
Vigorous physical activity			7.39 (24.70)			-0.08 (5.25)			-0.13 (0.32)			-0.04 (0.18)
Moderate physical activity			4.65 (22.01)			-2.42 (4.68)			0.29 (0.29)			0.06 (0.17)
BMI			0.91 (0.50)			-0.11 (0.09)			0.01** (0.01)			0.01** (0.003)
Health factors												
Preventive aspirin			0.77 (21.91)			-3.76 (5.01)			0.08 (0.24)			-0.11 (0.14)
Hypertension			-13.98 (17.33)			-6.60 (5.36)			-0.49** (0.18)			-0.12 (0.11)
Diabetes			22.14 (19.16)			14.10* (5.84)			0.24 (0.19)			0.12 (0.14)
Joint/bone diseases			14.51 (16.54)			-5.42 (4.63)			0.27 (0.18)			-0.05 (0.11)
Persistent skin trouble			-18.29 (38.72)			4.16 (11.32)			-0.79 (0.43)			-0.01 (0.24)
Teeth problems			13.93 (20.99)			-2.25 (7.00)			0.40 (0.27)			-0.09 (0.16)
Sleep problems			3.53 (23.98)			-0.29 (6.09)			0.24 (0.24)			0.19 (0.14)
Intercept	318.50***	304.19***	351.47***	82.54***	83.52***	100.13***	1.84**	1.67*	2.96**	0.81*	0.86*	0.32
Model F	3.85**	5.00**	1.65*	3.26*	3.44**	2.26**	4.00**	3.46**	4.88	2.10	1.45	1.94*
R ² , %	7.90	11.07	17.87	4.81	6.16	22.65	8.52	9.80	29.92	5.34	5.48	17.50

Source: Survey of Midlife in the U.S. (MIDUS II).

Note: Values are coefficients unless otherwise noted; standard errors appear in parentheses. Robust estimators are used. Boldface indicates statistical significance (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; two-tailed tests).

^a mg/dL.

^b ng/mL.

CRP, C-reactive protein; IL-6, interleukin 6; y, years.

d_{\log} pg/mL.

c_{\log} ug/mL.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript