

Integrating insulin-like growth factor 1 and sex hormones into neuroprotection: Implications for diabetes

Jacob Huffman, Christina Hoffmann, George T Taylor

Jacob Huffman, Christina Hoffmann, George T Taylor, Behavioral Neuroscience Program, Department of Psychological Sciences, University of Missouri, St. Louis, MO 63121, United States

George T Taylor, Interfakultäre Biomedizinische Forschungseinrichtung der Universität Heidelberg, 69120 Heidelberg, Germany

Author contributions: Huffman J designed the figure and contributed to writing the manuscript; Hoffmann C and Taylor GT contributed to writing the manuscript and editing process.

Conflict-of-interest statement: The authors declare no conflict of interests for this article.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: George T Taylor, PhD, Professor, Behavioral Neuroscience Program, Department of Psychological Sciences, University of Missouri, 1 University Blvd, St. Louis, MO 63121, United States. geot@umsl.edu
Telephone: +1-314-5761709

Received: July 6, 2016

Peer-review started: July 12, 2016

First decision: September 12, 2016

Revised: September 24, 2016

Accepted: November 21, 2016

Article in press: November 22, 2016

Published online: February 15, 2017

Abstract

Brain integrity and cognitive aptitude are often impaired

in patients with diabetes mellitus, presumably a result of the metabolic complications inherent to the disease. However, an increasing body of evidence has demonstrated the central role of insulin-like growth factor 1 (IGF1) and its relation to sex hormones in many neuroprotective processes. Both male and female patients with diabetes display abnormal IGF1 and sex-hormone levels but the comparison of these fluctuations is seldom a topic of interest. It is interesting to note that both IGF1 and sex hormones have the ability to regulate phosphoinositide 3-kinase-Akt and mitogen-activated protein kinases-extracellular signal-related kinase signaling cascades in animal and cell culture models of neuroprotection. Additionally, there is considerable evidence demonstrating the neuroprotective coupling of IGF1 and estrogen. Androgens have also been implicated in many neuroprotective processes that operate on similar signaling cascades as the estrogen-IGF1 relation. Yet, androgens have not been directly linked to the brain IGF1 system and neuroprotection. Despite the sex-specific variations in brain integrity and hormone levels observed in diabetic patients, the IGF1-sex hormone relation in neuroprotection has yet to be fully substantiated in experimental models of diabetes. Taken together, there is a clear need for the comprehensive analysis of sex differences on brain integrity of diabetic patients and the relationship between IGF1 and sex hormones that may influence brain-health outcomes. As such, this review will briefly outline the basic relation of diabetes and IGF1 and its role in neuroprotection. We will also consider the findings on sex hormones and diabetes as a basis for separately analyzing males and females to identify possible hormone-induced brain abnormalities. Finally, we will introduce the neuroprotective interplay of IGF1 and estrogen and how androgen-derived neuroprotection operates through similar signaling cascades. Future research on both neuroprotection and diabetes should include androgens into the interplay of IGF1 and sex hormones.

Key words: Diabetes; Androgens; Estrogen; Insulin; Insulin-like growth factor 1; Neuroprotection; Brain

integrity; Cognition

© **The Author(s) 2017.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Insulin-like growth factor 1 (IGF1), estrogen, and androgens are known to have neuroprotective properties. Fluctuations in these hormones is observed in patients with diabetes, varies with sex, and may contribute to abnormalities in brain integrity and cognitive impairment typical of the disease. While the neuroprotective coupling of estrogen and IGF1 has been studied extensively, little research has focused similarly on androgens. Furthermore, research investigating the IGF1-sex hormones relation to diabetes and brain-health outcomes is minimal. One avenue of approach to extend this literature may be to examine sex differences by comparison of these hormone levels, brain integrity, and cognitive aptitude.

Huffman J, Hoffmann C, Taylor GT. Integrating insulin-like growth factor 1 and sex hormones into neuroprotection: Implications for diabetes. *World J Diabetes* 2017; 8(2): 45-55 Available from: URL: <http://www.wjgnet.com/1948-9358/full/v8/i2/45.htm> DOI: <http://dx.doi.org/10.4239/wjd.v8.i2.45>

INTRODUCTION

Diabetes mellitus is a metabolic syndrome known for impaired insulin production. This condition is associated with an abundance of sequelae including cardiovascular disease^[1,2], brain atrophy^[3,4], and more recently, Alzheimer's disease^[5-7]. Over the past thirty years, researchers have established strong evidence supporting a link between patients with diabetes and subsequent cognitive impairments and abnormalities in brain integrity.

While meta-analyses have found inconsistencies in the specifics of the literature^[8-10], general trends point to cognitive impairments and abnormalities in related structural and functional brain areas. For example, patients with type 1 diabetes (T1D) are frequently found to have decreased psychomotor speed, mental flexibility, and IQ scores^[8,11-13]. T1D patients also often show reductions in the volume of regional gray matter in areas such as the prefrontal cortex, hippocampus, and thalamus^[12,14,15]. On the other hand, affected skills in type 2 diabetes (T2D) are largely executive function, memory, and information processing^[16,17]. Neuroimaging studies done on T2D patients indicate global brain atrophy and microstructural changes^[4,7,9,18], while findings regarding white matter hyperintensities are mixed^[3].

In both T1D and T2D these decrements are considered mild across most age groups^[8,11,19]. The severity of cognitive impairments and brain abnormalities are correlated with age of onset in T1D^[11] and duration

of the disease in T2D^[20,21]. Age is also a risk factor as deficits in learning and memory have been reported to worsen considerably in T2D patients above 65 years of age^[22]. Findings suggest the decreased brain volume in patients with T2D is correlated with increased insulin resistance^[23], and both brain atrophy and microstructural changes are associated with impaired cognitive performance^[18,20].

These data lend support to the idea that brain integrity is compromised in patients with both T1D and T2D, but also emphasize the need to integrate peripheral biomarkers associated with neuroprotection into diabetes research in humans. Various hormones altered as a result of diabetes have been recognized as neuroprotective, including insulin-like growth factor 1 (IGF1) and sex hormones. Research has revealed differences in the serum levels of IGF1 and gonadal hormones in diabetic patients^[24-27], with clear sex differences in the effects of androgens and estrogens on the brain in animal models^[28].

There is currently a movement in biomedical research to incorporate analyses of sex differences into studies^[29-31]; however, studies on brain integrity of diabetic patients often fail to examine men and women separately. This is despite findings of sex-specific differences in regional brain volume between men and women^[32-34]. For instance, DTI scans have also reported white matter hyperintensities are different in men and women diabetics^[35]. Others have shown that, by combining the data of men and women, T2D patients had smaller gray matter volume with larger ventricular volume and white matter lesions compared to healthy controls. However, when the sexes were analyzed separately, the data for men failed to reach statistical significance^[36].

Because sex hormones can act on similar molecular pathways as IGF1, and IGF1 is functionally related to insulin and diabetes, there is a need to further investigate how these hormones interact in the brains of diabetic patients. The relationship between estrogen and IGF1 is the most extensively studied in the neuroprotection literature^[37-39], but it has yet to expand experimentally into diabetes research. Furthermore, little attention has been paid to androgen-IGF1 interactions, even in the animal literature, despite the similar mechanisms underlying estrogenic and androgenic neuroprotection.

DIABETES AND IGF1 RELATION

IGF1 has a hypoglycemic response similar to insulin and, in some circumstances, is capable of modulating insulin receptor (IR) activities. Research has demonstrated that low IGF1 is associated with T1D and T2D^[40-42]. Moreover, genetic studies suggest decreased IGF1, due to a genetic polymorphism in the promoter region of the IGF1 gene, increases the risk of glucose intolerance and T2D^[43].

On the other hand, T2D has also been correlated

with excessively high levels of IGF1. For example, people with acromegaly - a condition known for its overproduction of pituitary growth hormone - have both high levels of IGF1 and a greater risk of developing T2D^[44]. These findings were corroborated by two large studies from Denmark ($n = 3354$) and Germany ($n = 7777$) which found U-shaped associations between IGF1 levels and the likelihood of developing insulin resistance and T2D^[24,25]. Moreover, treatment with IGF1 can improve glycemic control in patients with T1D and T2D^[45,46], which may suggest an optimal range of IGF1 for normal glycemic control.

Although IGF1 is synthesized in the brain, peripheral values cannot be used to accurately infer brain levels of IGF1 in humans as local synthesis of IGF1 in the brain appears not to correlate with the quantity of IGF1 receptors (IGF1R)^[47-49]. Evidence from animal models suggest that brain atrophy and loss of DNA are prevented following injection of insulin and IGF1, but not insulin alone, into cerebrospinal fluid of mice^[50]. Thus, proper systemic levels of IGF1 and its transport from the periphery into the brain is likely necessary for the maintenance of various cognitive processes^[51].

Collectively, these data support the involvement of IGF1 in diabetes but also point to an "optimal range" of IGF1. Future research should examine the significance of an optimal peripheral range in the development and maintenance of diabetes and cognitive decline. Moreover, there is a need for data on the role of central vs peripheral IGF1 levels and the subsequent impact on cognitive impairment and brain atrophy.

THE IGF1 SYSTEM

Transportation

IGF1 is a polypeptide, structurally similar to insulin, that is released in response to growth hormones secreted by the anterior pituitary^[52]. While synthesized predominantly by hepatocytes in the liver and released into general circulation, both paracrine and autocrine functions contribute through local tissue synthesis of IGF1. The concentration of IGF1 is greatest during perinatal development and decreases markedly into adulthood. IGF1R are expressed in nearly all neural cells of the CNS, being most highly expressed in the cortex, hippocampus, cerebellum, brainstem, hypothalamus, and spinal cord^[53].

The blood brain barrier and blood-cerebrospinal fluid barrier are the two primary routes involved with transporting systemic IGF1 into the brain. Both barriers utilize lipoprotein receptor-related proteins along with IGF1R as transporters to enter the brain^[54,55]. However, the bioavailability of IGF1 is largely determined by the amount of hormone bound to IGF binding proteins (IGFBPs). Most circulating IGF is bound by IGFBPs, which are proteins that control the distribution and functional capabilities of IGF1 throughout the body. Six different IGFBPs modulate the activity of IGFs *via* binding affinities exceeding that of its respective receptor

and, thus, help regulate the amount of IGF1 that enters the brain^[56].

Signaling pathways

The role of IGF1 is dependent on its binding to insulin-like peptide receptors. The three most important include the IGF1R, IR, and a hybrid receptor formed from heterodimer α - β IR and IGF1R subunits^[53,57]. These receptors are important to the functional efficacy of IGF1 and have defined downstream molecular pathways. As part of the tyrosine kinase receptor family, activation of IGF1R leads to the signaling of either the mitogen-activated protein kinases-extracellular signal-related kinase (MAPK-ERK) or phosphoinositide 3-kinase (PI3K)-Akt pathways^[53,57]. These pathways are involved in several important cellular processes including the regulation of gene transcription, apoptosis, oxidative stress, and cellular proliferation and differentiation.

The affinity of IGF1 varies among the three receptors with the highest affinity for IGF1R. Activation of the IGF1R is capable of directly stimulating the RAS-ERK pathway, leading to the modulation of gene transcription by way of activating ETS-like transcription factor, ELK1^[57]. The capacity of insulin-like peptide receptors to initiate downstream molecular activity is modified in part by the recruitment of insulin receptor substrate (IRS) scaffolding proteins^[57-59]. This scaffolding helps adjust pathway choice following receptor phosphorylation. The result is activation of PI3K-Akt and subsequent expression of downstream effectors, including glycogen synthase 3 kinase (GSK3 β) and mammalian target of rapamycin^[53,57,60].

Relationship to the insulin system

IGF1 acts primarily through binding to the IGF1R, but also shares with insulin the capacity to bind the IR and hybrid receptor^[53,56,57]. Insulin is produced exclusively by β -cells of the pancreas and, hence, is strictly transported in the systemic circulation. The amount of insulin capable of entering the brain varies considerably^[54,55]. Unlike IGF1, insulin appears not to be locally synthesized in adult brain cells^[53,56]. Similar to IGF1, IR located on endothelial and epithelial cell membranes allow insulin to be transported into the brain from systemic circulation. IRs are concentrated mostly in the olfactory bulb, cerebral cortex, hypothalamus, hippocampus, and cerebellum^[55]. The movement of systemic insulin into the brain is not controlled by binding proteins.

Both insulin and IGF1 produced in the periphery contribute to varied physiological processes. Proper peripheral IGF1 activation is necessary for insulin secretion from the pancreas and, hence, is implicated in many facets of diabetes^[61]. However, their functions differ once entering the brain. IGF1R are expressed at notably higher rates in the brain than the rate IGF1 is synthesized. This differential suggests that active transport of IGF1 into the brain is required to furnish sufficient IGF1 for proper neuronal function^[47-49]. For example, peripheral IGF1 supplies the brain with

information regarding body mass, is related to neural plasticity and cognitive processes, and attenuates cognitive impairment induced by diabetes^[51,62,63]. Deficiency of IGF1 can also lead to hippocampal atrophy and impaired learning^[64]. Indeed, IGF1 in the brain is required for proper tissue growth in both the brain and periphery, as well as sufficient glucose regulation and insulin sensitivity^[65,66].

Insulin in the periphery is well-known for its role in glucose regulation and communication with the brain to maintain energy homeostasis. Similar to IGF1, insulin is involved in modifying BBB permeability in the brain^[55] with T2D patients showing greater permeability of the BBB^[67]. Insulin also acts on the PI3K and MAPK signaling cascades to enhance neuronal survival, plasticity, and subsequent cognitive processes^[55,68,69]. With that said, insulin does not necessarily regulate glucose activity in neuronal cells after entering the brain. Rather, insulin modulates energy homeostasis through its actions at the level of the hypothalamus^[70].

INTEGRATING SEX HORMONES INTO DIABETES AND IGF1

Diabetes is associated with imbalances in sex steroid hormone levels. This is not surprising as androgens and estrogens are known to play an important role in body composition^[71] while maintaining glucose and lipid homeostasis^[72,73]. Research into these imbalances suggests a complex relation between estradiol (E2) and insulin insensitivity. Several studies have reported that postmenopausal women with T2D have increased levels of circulating E2^[27,74,75]. Elevated E2 has been correlated with the development of insulin resistance and T2D in these women^[76,77]. Nevertheless, there are at least two studies that have shown inconsistencies between E2 levels and the development of diabetes in postmenopausal women^[78,79].

There is also a link between high levels of E2 and diabetes in men. Diabetic men have shown relatively high basal levels of E2^[27,78], while men with higher levels of circulating E2 have an increased risk of developing T2D^[80]. Although this may simply be a product of higher body fat content as adrenal androgens are readily converted to E2 in adipose tissue^[81-83], two studies reported E2 results in men were independent of obesity^[78,80].

Findings with animal models suggest an opposite conclusion for E2 and diabetes, at least during reproductive ages. Male mice with streptozotocin-induced insulin insensitivity are more likely to develop diabetes than their female cohorts. This increased risk of diabetes in the males can be attenuated with E2 supplements^[84]. Also, mice lacking the alpha subtype of estrogen receptor (ER α) have been reported to develop insulin insensitivity^[85]. In contrast, these data in animals mirror those from postmenopausal women in which glucose homeostasis was positively impacted with estrogen therapy in

the short term^[86].

Sex differences in androgen-diabetes relations have also been reported. Postmenopausal women with diabetes displayed elevated circulating testosterone (TS) levels^[27,75]. Reports suggest that premenopausal women with higher levels of TS^[76,79], as well as female mice administered the androgen^[84], had a greater risk of developing diabetes. Another example is the link between T2D development and hyperandrogenism experienced by patients with polycystic ovarian syndrome^[87]. Still, much like E2, there are also studies that dispute these reports, particularly in postmenopausal women^[77,78].

A clear sex difference is also indicated in that diabetic men tend to have either lower total, free, or bioavailable TS than healthy men^[27,88,89]. Indeed, men with the highest levels of TS were at the lowest risk and men with lowest levels of TS were at highest risk for developing T2D^[78,79,90]. Moreover, men undergoing androgen deprivation treatments for prostatic cancer had a greatly increased risk of developing T2D^[91]. Yet again, these reports are not without contradiction^[92] and some studies found this relationship to be dependent on obesity^[80,93].

Taken together, there are clear inconsistencies in the findings on sex hormones and diabetes. There is also an apparent lack of research focusing on sex hormones in premenopausal diabetic women that should be addressed^[26]. It is again important to note that many studies fail to acknowledge the possible relation of sex hormones to the IGF1 system. Findings with serum E2 data are consistent with findings from meta-analyses examining IGF1^[24,25]. Their proposed U-shaped association of IGF1 and T2D fits into the well-defined mechanistic relationship between E2 and IGF1, described in more detail below. The relation between sex hormones and IGF1 suggests that a delicate hormonal balance is likely an important facet of diabetes-induced brain and cognitive impairment.

NEUROPROTECTION: SEX HORMONES AND IGF1

Estrogen and IGF1

An intriguing feature of neuroactive hormones is their ability to protect the CNS from damage, especially in regards to estrogen. ER activation is implicated in the maintenance of various metabolic processes that are also associated with diabetes, including glucose homeostasis and obesity^[94,95]. Only recently has research with animal models focused on neuroprotection from IGF1-E2 interactions. Evidence suggests that neuroprotective properties of E2 are directly related to receptor activities of insulin-like peptide receptors, mainly IGF1R. E2 and IGF1 work in tandem to reciprocally modulate and facilitate ER and IGF1R activation of the PI3K-Akt and MAPK-ERK signaling cascades^[96-100].

IGF1 shows differential sensitivities to the two

estrogen receptor subtypes with ER α being more sensitive than ER β ^[97,101]. Selective inhibition of IGF1R, for instance, downregulates ER α expression in the hypothalamus, hippocampus, and cerebral cortex, with the only significant changes of ER β occurring in the cerebellum^[38]. Many glial and neuronal cells in the brain express IGF1R and both ER subtypes^[102]. In particular, ER α is uniquely capable of increasing IGF1R activity of downstream PI3K-Akt signaling in rodent models^[103,104]. ER α activation also increases the binding of p85 and IRS-1 regulatory subunits of PI3K and, thus, may be one mechanism assisting in Akt pro-survival signaling through the IGF1R^[39,97] (Figure 1).

Administration of E2 to mice increased IGF1R and ER α activity in the brain, enabling activation of IGF1R and downstream PI3K-Akt pathway signaling^[97]. Similarly, IGF1 and insulin modulated ER effects on gene transcription and the PI3K-Akt-GSK3 β signaling cascade^[38,98,103,105,106]. GSK3 β is a protein kinase known particularly well for its role in glycogen synthesis. However, as reviewed by Jacobs *et al.*^[60], recent attention has turned to the dual pro- and anti-apoptosis capabilities of GSK3 β regulated through multiple different pathways. Indeed, the neuroprotective effects of IGF1 may be consequent to Akt-derived inhibition of GSK3 β in a hypoxic state^[107] (Figure 1).

Activation of the MAPK pathway is another important signal transduction pathway involved with regulating gene transcription and cellular proliferation and differentiation, particularly in cancer^[108]. However, multiple studies have demonstrated that the neuroprotective properties of estrogen are also derived from its ability to regulate MAPK signaling in the brain^[38]. Both estrogen and IGF1 can facilitate MAPK signaling through the IGF1R, with IGF1 increasing ER α activities in the presence of E2^[104]. Akt inhibitors are capable of nullifying the neuroprotective effects of IGF1 and E2 regardless of MAPK signaling^[99,104], while ERK suppression increases PI3K-Akt activity *via* ER and IGF1R heterodimers^[39]. Thus, it appears the PI3K-Akt pro-survival signaling cascade is the most involved with the neuroprotective coupling of E2 and IGF1^[39].

It is important to note that IGF1 and E2 have a remarkable reciprocity. Inhibition of ER activity can downregulate IGF1R expression in the hippocampus^[109], a brain region known to atrophy in patients with diabetes and glucose intolerance^[110-112]. Similarly, IGF1 has the capacity to upregulate ER α in the hippocampus and is impaired following administration of IGF1R antagonists^[109]. Agonists or antagonists of either hormone can respectively facilitate or inhibit the neuroprotective and memory enhancing properties of the other^[96,109,113-116]. This has led some to suggest that cooperation between IGF1R and ER is required for many E2-induced neuroprotective processes. The present section does not, however, do justice to the complexity of the relation between estrogen and IGF1 receptors. A fuller explanation can be found in one of several reviews^[37-39,101,109,117].

Androgens and IGF1

Far less research has examined a functional link between IGF1 and androgens in the brain. This is an unfortunate but common trend in neuroendocrinology. Estrogens are the most intensely studied gonadal hormone, despite estrogens and androgens sharing metabolic pathways and functional properties. Much of the current literature on IGF1-androgen relations are directed at the periphery, particularly prostate cancer and motor systems, for which there are a number of recent reviews^[118,119]. Few studies have examined IGF1-androgen interactions in neuroprotection^[120,121] and none, to our knowledge, have empirically examined this interaction in diabetes. Therefore, we have relied on peripheral data, often from *in vitro* experiments, to extrapolate the androgen receptor (AR) brain discussion.

There is evidence that the two main androgens, TS and dihydrotestosterone (DHT), are capable of neuroprotection through binding the AR^[122-126]. Similar to ER α , androgen activation of the AR in mouse vas deferens epithelial cells can modulate the p85 regulatory subunits of PI3K and subsequently trigger Akt expression (Figure 1). Inhibiting the AR prevents these signaling effects^[127]. Phosphorylation of MAPK and Akt can also increase AR activation in low androgen and estrogen concentrations, as well as increase the neuroprotective activities of ER α and AR^[128]. Recent findings showed that DHT, which has a higher affinity than TS for the AR, prevents apoptosis in a C6 glial cell line through the PI3K-Akt signaling cascade^[129]. These effects were also impaired by inhibition of PI3K and suggest a functional relationship between apoptosis and AR activities.

Interestingly, studies have demonstrated that binding of DHT to the transmembrane AR impairs MAPK and PI3K signaling and subsequent neuroprotection from DHT or E2^[130-132]. This suggests that nuclear activation of the AR by DHT is likely one mechanism behind DHT's neuroprotective properties^[130]. DHT may also interact with effectors downstream of ER and IGF1R signaling. Both TS and DHT can activate the MAPK-ERK signaling cascade^[132] which has been shown to induce ribosomal S6 kinase (Rsk) expression. Rsk signaling can lead to the inhibition of the pro-apoptosis Bad protein and the activation of downstream effectors including the ER, GSK3 β and ELK1^[133] (Figure 1).

One possible explanation for the neuroprotective role of androgens is the conversion in the steroid metabolic cascade of TS into E2 by the enzyme aromatase. That is, TS may be involved in neuroprotection only to the extent that TS is a precursor for E2, which is capable of activating MAPK or PI3K signaling through the ER and IGF1R. The aromatization of TS into E2, as well as the aromatase enzyme, have been suggested to play an important role in neuroprotection^[134-139].

The ratio of endogenous TS to E2, and subsequent influences of aromatized TS, is indeed a topic of recent interest^[26]. Increased local synthesis of E2 from elevated aromatase expression is seen in models




Figure 1 Similar signaling cascades involved with neuroprotection for insulin-like peptides and sex hormones. The insulin receptor (IR), insulin-like growth factor 1 receptor (IGF1R), and insulin-IGF1 hybrid receptor enact their neuroprotection through the mitogen-activated protein kinases-extracellular signal-related kinase (MAPK-ERK) or phosphoinositide 3-kinase (PI3K)-Akt pathways signaling cascades. Although IGF1R can directly activate the RAS-ERK pathway, both the insulin-like peptide receptors and the estrogen receptor alpha (ER α) firstly interact with insulin receptor substrate 1 (IRS-1) scaffolding proteins. ER α and the androgen receptor (AR) can also directly modulate PI3K-Akt and MAPK-ERK signaling. Both IRS-1 and p85 binding of PI3K are increased with ER α activation, leading to downstream Akt-derived inhibition of glycogen synthase kinase 3 (GSK3) and mammalian target of rapamycin (mTOR). GSK3, specifically, is involved with glycogen synthesis, while both effectors are involved in apoptosis. A similar effect may occur with AR's ability to modulate p85 binding to PI3K. AR-induced MAPK-ERK signaling also results in ribosomal S6 kinase (Rsk) expression that can inhibit the pro-apoptosis bcl-2-associated death promoter protein, as well as effects on the ER, GSK3, and the ETS-like transcription factor, ELK1. Solid black arrows indicate downstream interaction. Dashed black arrows represent the influence of kinases or proteins on the cellular environment. Dashed blue arrows represent the binding capabilities of IGF1 and insulin across all three receptor types.

of neuroprotection from other brain disorders, *e.g.*, stroke^[140]. More pertinent to this review, streptozotocin-induced diabetes causes a considerable reduction in aromatase synthesis in female and male reproductive systems^[141]. Notably, inhibition of aromatase decreases E2 and impairs insulin sensitivity and peripheral glucose disposal in healthy males^[142], although the influence this may have on brain integrity and cognitive outcomes remains debated^[143].

Another explanation places greater emphasis on the other pathway in the steroid metabolic cascade leading to DHT. Metabolites of DHT, 3 α -Diol and 3 β -Diol, are also bioactive and may bind the ER or insulin-like peptide receptors to initiate MAPK or PI3K signaling cascades. Indeed, research shows that 3 α -Diol stimulated PI3K-Akt signaling enhances cell survival in the prostate^[144]. Similarly, DHT metabolites may influence transcriptional activities of nuclear ER by modulating ER-induced MAPK or PI3K signaling cascades.

Few *in vivo* studies examining these sex steroid metabolites have focused on MAPK or PI3K signal cascades in the brain. There is, however, evidence that 3 α -Diol inhibits protein kinase A expression in the rat hippocampus^[145]. Others have reported that strep-

tozotocin-induced diabetic mice had lower levels of TS and 3 α -Diol in the cerebral cortex, and lower levels of DHT and 3 α -Diol in the spinal cord^[146]. It is still unclear, though, whether 3 α -Diol and 3 β -Diol interact with or initiate the MAPK or PI3K signaling cascades following activation of the ER, AR, or, possibly, IGF1R.

None of these explanations clarify fully the ability of the AR to directly trigger these signaling cascades. We do not aim to discount the neuroprotective mechanisms of ER and AR, or the clear link between E2 and IGF1 processes in neuroprotection. Rather, we simply suggest that androgen-derived neuroprotection may be intertwined with IGF1, the activation of insulin-like peptide receptors, and/or the IGF1R and ER coupling. Given the common signaling pathways between these hormones, we suggest future research should aim to include androgens and AR activities into the ER-IGF1R neuroprotective coupling, as well as serum comparisons in brain-health outcomes of diabetic patients.

CONCLUSION

The reciprocity of IGF1 and estrogen in neuroprotective processes is well-established in cell cultures and

animal models^[38]. Interactions between androgens and IGF1 may also play an important role in the E2-IGF1 neuroprotective coupling. Both estrogens and androgens enact their neuroprotection through similar, but not identical, signal transduction pathways. Recognition of this has led us to consider the possibility that these sex hormones may work together with IGF1 and insulin-like peptide receptors to modulate MAPK and PI3K signaling and their neuroprotective properties.

Regulation of MAPK and PI3K activity may also be a driving force behind the structural changes, atrophy of brain regions, or functional changes, often observed in diabetic patients. Drawing conclusions from imaging data in humans to those found in animal models is indeed difficult. Nevertheless, there is a need for a clearer mechanistic explanation grounding the cognitive decline and brain abnormalities observed in diabetic patients.

Future studies in human research on diabetic brain integrity should integrate hormone titer measures to help substantiate sex differences in brain-health outcomes of diabetic patients. This approach may also assist in identifying region-specific brain abnormalities resulting from fluctuations in IGF1 and sex hormones between men and women. Moreover, animal models examining the E2-IGF1 coupling in neuroprotection should employ streptozotocin-induced diabetes, as well as the possible role of androgens and AR activities. These conclusions warrant further examination of the variability present in cognitive and brain-health outcomes for patients with diabetes as a result of sex hormone relations to IGF1, insulin, and the insulin-like peptide receptors.

REFERENCES

- 1 **Rydén L**, Grant PJ, Anker SD, Berne C, Cosentino F, Danchin N, Deaton C, Escaned J, Hammes HP, Huikuri H, Marre M, Marx N, Mellbin L, Ostergren J, Patrono C, Seferovic P, Uva MS, Taskinen MR, Tendera M, Tuomilehto J, Valensi P, Zamorano JL. ESC guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD - summary. *Diab Vasc Dis Res* 2014; **11**: 133-173 [PMID: 24800783 DOI: 10.1177/1479164114525548]
- 2 **Tillin T**, Hughes AD, Mayet J, Whincup P, Sattar N, Forouhi NG, McKeigue PM, Chaturvedi N. The relationship between metabolic risk factors and incident cardiovascular disease in Europeans, South Asians, and African Caribbeans: SABRE (Southall and Brent Revisited) -- a prospective population-based study. *J Am Coll Cardiol* 2013; **61**: 1777-1786 [PMID: 23500273 DOI: 10.1016/j.jacc.2012.12.046]
- 3 **van Elderen SG**, de Roos A, de Craen AJ, Westendorp RG, Blauw GJ, Jukema JW, Bollen EL, Middelkoop HA, van Buchem MA, van der Grond J. Progression of brain atrophy and cognitive decline in diabetes mellitus: a 3-year follow-up. *Neurology* 2010; **75**: 997-1002 [PMID: 20837967 DOI: 10.1212/WNL.0b013e3181f25f06]
- 4 **de Bresser J**, Tiehuis AM, van den Berg E, Reijmer YD, Jongen C, Kappelle LJ, Mali WP, Viergever MA, Biessels GJ. Progression of cerebral atrophy and white matter hyperintensities in patients with type 2 diabetes. *Diabetes Care* 2010; **33**: 1309-1314 [PMID: 20299484 DOI: 10.2337/dc09-1923]
- 5 **Peila R**, Rodriguez BL, Launer LJ. Type 2 diabetes, APOE gene, and the risk for dementia and related pathologies: The Honolulu-Asia Aging Study. *Diabetes* 2002; **51**: 1256-1262 [PMID: 11916953 DOI: 10.2337/diabetes.51.4.1256]
- 6 **Cheng G**, Huang C, Deng H, Wang H. Diabetes as a risk factor for dementia and mild cognitive impairment: a meta-analysis of longitudinal studies. *Intern Med J* 2012; **42**: 484-491 [PMID: 22372522 DOI: 10.1111/j.1445-5994.2012.02758.x]
- 7 **Moran C**, Phan TG, Chen J, Blizzard L, Beare R, Venn A, Münch G, Wood AG, Forbes J, Greenaway TM, Pearson S, Srikanth V. Brain atrophy in type 2 diabetes: regional distribution and influence on cognition. *Diabetes Care* 2013; **36**: 4036-4042 [PMID: 23939539 DOI: 10.2337/dc13-0143]
- 8 **Brands AM**, Biessels GJ, de Haan EH, Kappelle LJ, Kessels RP. The effects of type 1 diabetes on cognitive performance: a meta-analysis. *Diabetes Care* 2005; **28**: 726-735 [PMID: 15735218 DOI: 10.2337/diacare.28.3.726]
- 9 **van Harten B**, de Leeuw FE, Weinstein HC, Scheltens P, Biessels GJ. Brain imaging in patients with diabetes: a systematic review. *Diabetes Care* 2006; **29**: 2539-2548 [PMID: 17065699 DOI: 10.2337/dc06-1637]
- 10 **Sadanand S**, Balachandar R, Bharath S. Memory and executive functions in persons with type 2 diabetes: a meta-analysis. *Diabetes Metab Res Rev* 2016; **32**: 132-142 [PMID: 25963303 DOI: 10.1002/dmrr.2664]
- 11 **Ferguson SC**, Blane A, Wardlaw J, Frier BM, Perros P, McCrimmon RJ, Deary IJ. Influence of an early-onset age of type 1 diabetes on cerebral structure and cognitive function. *Diabetes Care* 2005; **28**: 1431-1437 [PMID: 15920064 DOI: 10.2337/diacare.28.6.1431]
- 12 **Northam EA**, Rankins D, Lin A, Wellard RM, Pell GS, Finch SJ, Werther GA, Cameron FJ. Central nervous system function in youth with type 1 diabetes 12 years after disease onset. *Diabetes Care* 2009; **32**: 445-450 [PMID: 19151204 DOI: 10.2337/dc08-1657]
- 13 **Biessels GJ**, Deary IJ, Ryan CM. Cognition and diabetes: a lifespan perspective. *Lancet Neurol* 2008; **7**: 184-190 [PMID: 18207116 DOI: 10.1016/S1474-4422(08)70021-8]
- 14 **Hughes TM**, Ryan CM, Aizenstein HJ, Nunley K, Gianaros PJ, Miller R, Costacou T, Strotmeyer ES, Orchard TJ, Rosano C. Frontal gray matter atrophy in middle aged adults with type 1 diabetes is independent of cardiovascular risk factors and diabetes complications. *J Diabetes Complications* 2013; **27**: 558-564 [PMID: 23994432 DOI: 10.1016/j.jdiacomp.2013.07.001]
- 15 **Foghi K**, Ahmadpour S. Role of neuronal apoptosis in volumetric change of hippocampus in diabetes mellitus type 1: a predictive model. *ISRN Anat* 2013; **2013**: 958461 [PMID: 25938109 DOI: 10.5402/2013/958461]
- 16 **van den Berg E**, Kloppenborg RP, Kessels RP, Kappelle LJ, Biessels GJ. Type 2 diabetes mellitus, hypertension, dyslipidemia and obesity: A systematic comparison of their impact on cognition. *Biochim Biophys Acta* 2009; **1792**: 470-481 [PMID: 18848880 DOI: 10.1016/j.bbdis.2008.09.004]
- 17 **Palta P**, Schneider AL, Biessels GJ, Touradji P, Hill-Briggs F. Magnitude of cognitive dysfunction in adults with type 2 diabetes: a meta-analysis of six cognitive domains and the most frequently reported neuropsychological tests within domains. *J Int Neuropsychol Soc* 2014; **20**: 278-291 [PMID: 24555960 DOI: 10.1017/S1355617713001483]
- 18 **Reijmer YD**, Brundel M, de Bresser J, Kappelle LJ, Leemans A, Biessels GJ. Microstructural white matter abnormalities and cognitive functioning in type 2 diabetes: a diffusion tensor imaging study. *Diabetes Care* 2013; **36**: 137-144 [PMID: 22961577 DOI: 10.2337/dc12-0493]
- 19 **Biessels GJ**, Strachan MW, Visseren FL, Kappelle LJ, Whitmer RA. Dementia and cognitive decline in type 2 diabetes and prediabetic stages: towards targeted interventions. *Lancet Diabetes Endocrinol* 2014; **2**: 246-255 [PMID: 24622755 DOI: 10.1016/S2213-8587(13)70088-3]
- 20 **Manschot SM**, Brands AM, van der Grond J, Kessels RP, Algra A, Kappelle LJ, Biessels GJ. Brain magnetic resonance imaging correlates of impaired cognition in patients with type 2 diabetes. *Diabetes* 2006; **55**: 1106-1113 [PMID: 16567535 DOI: 10.2337/diabetes.55.04.06.db05-1323]
- 21 **Saczynski JS**, Siggurdsson S, Jonsson PV, Eiriksdottir G, Olafsdottir E, Kjartansson O, Harris TB, van Buchem MA,

- Gudnason V, Launer LJ. Glycemic status and brain injury in older individuals: the age gene/environment susceptibility-Reykjavik study. *Diabetes Care* 2009; **32**: 1608-1613 [PMID: 19509008 DOI: 10.2337/dc08-2300]
- 22 **Ryan CM**, Geckle M. Why is learning and memory dysfunction in Type 2 diabetes limited to older adults? *Diabetes Metab Res Rev* 2000; **16**: 308-315 [PMID: 11025555 DOI: 10.1002/1520-7560(2000)9999]
- 23 **Tan ZS**, Beiser AS, Fox CS, Au R, Himali JJ, Debette S, Decarli C, Vasan RS, Wolf PA, Seshadri S. Association of metabolic dysregulation with volumetric brain magnetic resonance imaging and cognitive markers of subclinical brain aging in middle-aged adults: the Framingham Offspring Study. *Diabetes Care* 2011; **34**: 1766-1770 [PMID: 21680719 DOI: 10.2337/dc11-0308]
- 24 **Friedrich N**, Thuesen B, Jørgensen T, Juul A, Spielhagen C, Wallaschofski H, Linneberg A. The association between IGF-I and insulin resistance: a general population study in Danish adults. *Diabetes Care* 2012; **35**: 768-773 [PMID: 22374641 DOI: 10.2337/dc11-1833]
- 25 **Schneider HJ**, Friedrich N, Klotsche J, Schipf S, Nauck M, Völzke H, Sievers C, Pieper L, März W, Wittchen HU, Stalla GK, Wallaschofski H. Prediction of incident diabetes mellitus by baseline IGF1 levels. *Eur J Endocrinol* 2011; **164**: 223-229 [PMID: 21059863 DOI: 10.1530/EJE-10-0963]
- 26 **Kim C**, Halter JB. Endogenous sex hormones, metabolic syndrome, and diabetes in men and women. *Curr Cardiol Rep* 2014; **16**: 467 [PMID: 24585109 DOI: 10.1007/s11886-014-0467-6]
- 27 **Ding EL**, Song Y, Malik VS, Liu S. Sex differences of endogenous sex hormones and risk of type 2 diabetes: a systematic review and meta-analysis. *JAMA* 2006; **295**: 1288-1299 [PMID: 16537739 DOI: 10.1001/jama.295.11.1288]
- 28 **Pesaresi M**, Maschi O, Giatti S, Garcia-Segura LM, Caruso D, Melcangi RC. Sex differences in neuroactive steroid levels in the nervous system of diabetic and non-diabetic rats. *Horm Behav* 2010; **57**: 46-55 [PMID: 19422828 DOI: 10.1016/j.yhbeh.2009.04.008]
- 29 **Franconi F**, Seghieri G, Canu S, Straface E, Campesi I, Malorni W. Are the available experimental models of type 2 diabetes appropriate for a gender perspective? *Pharmacol Res* 2008; **57**: 6-18 [PMID: 18221886 DOI: 10.1016/j.phrs.2007.11.007]
- 30 **Legato MJ**, Gelzer A, Goland R, Ebner SA, Rajan S, Villagra V, Kosowski M. Gender-specific care of the patient with diabetes: review and recommendations. *Gen Med* 2006; **3**: 131-158 [PMID: 16860272 DOI: 10.1016/S1550-8579(06)80202-0]
- 31 **McCullum M**, Hansen LS, Lu L, Sullivan PW. Gender differences in diabetes mellitus and effects on self-care activity. *Gen Med* 2005; **2**: 246-254 [PMID: 16464736 DOI: 10.1016/S1550-8-579(05)80054-3]
- 32 **Goldstein JM**, Seidman LJ, Horton NJ, Makris N, Kennedy DN, Caviness VS, Faraone SV, Tsuang MT. Normal sexual dimorphism of the adult human brain assessed by in vivo magnetic resonance imaging. *Cereb Cortex* 2001; **11**: 490-497 [PMID: 11375910 DOI: 10.1093/cercor/11.6.490]
- 33 **Menzler K**, Belke M, Wehrmann E, Krakow K, Lengler U, Jansen A, Hamer HM, Oertel WH, Rosenow F, Knake S. Men and women are different: diffusion tensor imaging reveals sexual dimorphism in the microstructure of the thalamus, corpus callosum and cingulum. *Neuroimage* 2011; **54**: 2557-2562 [PMID: 21087671 DOI: 10.1016/j.neuroimage.2010.11.029]
- 34 **Nopoulos P**, Flaum M, O'Leary D, Andreasen NC. Sexual dimorphism in the human brain: evaluation of tissue volume, tissue composition and surface anatomy using magnetic resonance imaging. *Psychiatry Res* 2000; **98**: 1-13 [PMID: 10708922 DOI: 10.1016/S0925-4927(99)00044-X]
- 35 **Sachdev PS**, Parslow R, Wen W, Anstey KJ, Eastale S. Sex differences in the causes and consequences of white matter hyperintensities. *Neurobiol Aging* 2009; **30**: 946-956 [PMID: 17950492 DOI: 10.1016/j.neurobiolaging.2007.08.023]
- 36 **Jongen C**, van der Grond J, Kappelle LJ, Biessels GJ, Viergever MA, Pluim JP. Automated measurement of brain and white matter lesion volume in type 2 diabetes mellitus. *Diabetologia* 2007; **50**: 1509-1516 [PMID: 17492428 DOI: 10.1007/s00125-007-0688-y]
- 37 **Arevalo MA**, Azcoitia I, Garcia-Segura LM. The neuroprotective actions of oestradiol and oestrogen receptors. *Nat Rev Neurosci* 2015; **16**: 17-29 [PMID: 25423896 DOI: 10.1038/nrn3856]
- 38 **Garcia-Segura LM**, Arévalo MA, Azcoitia I. Interactions of estradiol and insulin-like growth factor-I signalling in the nervous system: new advances. *Prog Brain Res* 2010; **181**: 251-272 [PMID: 20478442 DOI: 10.1016/S0079-6123(08)81014-X]
- 39 **Sohrabji F**. Estrogen-IGF-1 interactions in neuroprotection: ischemic stroke as a case study. *Front Neuroendocrinol* 2015; **36**: 1-14 [PMID: 24882635 DOI: 10.1016/j.yfrne.2014.05.003]
- 40 **Ezzat VA**, Duncan ER, Wheatcroft SB, Kearney MT. The role of IGF-I and its binding proteins in the development of type 2 diabetes and cardiovascular disease. *Diabetes Obes Metab* 2008; **10**: 198-211 [PMID: 18269635 DOI: 10.1111/j.1463-1326.2007.00709.x]
- 41 **Janssen JA**, Jacobs ML, Derckx FH, Weber RF, van der Lely AJ, Lamberts SW. Free and total insulin-like growth factor I (IGF-I), IGF-binding protein-1 (IGFBP-1), and IGFBP-3 and their relationships to the presence of diabetic retinopathy and glomerular hyperfiltration in insulin-dependent diabetes mellitus. *J Clin Endocrinol Metab* 1997; **82**: 2809-2815 [PMID: 9284701 DOI: 10.1210/jcem.82.9.4180]
- 42 **Teppala S**, Shankar A, Sabanayagam C. Association between IGF-1 and chronic kidney disease among US adults. *Clin Exp Nephrol* 2010; **14**: 440-444 [PMID: 20567872 DOI: 10.1007/s10157-010-0307-y]
- 43 **Vaessen N**, Heutink P, Janssen JA, Witteman JC, Testers L, Hofman A, Lamberts SW, Oostra BA, Pols HA, van Duijn CM. A polymorphism in the gene for IGF-I: functional properties and risk for type 2 diabetes and myocardial infarction. *Diabetes* 2001; **50**: 637-642 [PMID: 11246885 DOI: 10.2337/diabetes.50.3.637]
- 44 **Melmed S**. Medical progress: Acromegaly. *N Engl J Med* 2006; **355**: 2558-2573 [PMID: 17167139 DOI: 10.1056/NEJMra062453]
- 45 **Carroll PV**, Christ ER, Umpleby AM, Gowrie I, Jackson N, Bowes SB, Hovorka R, Croos P, Sönksen PH, Russell-Jones DL. IGF-I treatment in adults with type 1 diabetes: effects on glucose and protein metabolism in the fasting state and during a hyperinsulinemic-euglycemic amino acid clamp. *Diabetes* 2000; **49**: 789-796 [PMID: 10905488 DOI: 10.2337/diabetes.49.5.789]
- 46 **Zenobi PD**, Jaeggi-Groisman SE, Riesen WF, Roder ME, Froesch ER. Insulin-like growth factor-I improves glucose and lipid metabolism in type 2 diabetes mellitus. *J Clin Invest* 1992; **90**: 2234-2241 [PMID: 1469083 DOI: 10.1172/JCI116109]
- 47 **Carro E**, Trejo JL, Núñez A, Torres-Aleman I. Brain repair and neuroprotection by serum insulin-like growth factor I. *Mol Neurobiol* 2003; **27**: 153-162 [PMID: 1277685]
- 48 **Carro E**, Torres-Aleman I. Serum insulin-like growth factor I in brain function. *Keio J Med* 2006; **55**: 59-63 [PMID: 16830417 DOI: 10.2302/kjm.55.59]
- 49 **Nishijima T**, Piriz J, Dufflot S, Fernandez AM, Gaitan G, Gomez-Pinedo U, Verdugo JM, Leroy F, Soya H, Nuñez A, Torres-Aleman I. Neuronal activity drives localized blood-brain-barrier transport of serum insulin-like growth factor-I into the CNS. *Neuron* 2010; **67**: 834-846 [PMID: 20826314 DOI: 10.1016/j.neuron.2010.08.007]
- 50 **Serbedzija P**, Madl JE, Ishii DN. Insulin and IGF-I prevent brain atrophy and DNA loss in diabetes. *Brain Res* 2009; **1303**: 179-194 [PMID: 19781531 DOI: 10.1016/j.brainres.2009.09.063]
- 51 **Lupien SB**, Bluhm EJ, Ishii DN. Systemic insulin-like growth factor-I administration prevents cognitive impairment in diabetic rats, and brain IGF regulates learning/memory in normal adult rats. *J Neurosci Res* 2003; **74**: 512-523 [PMID: 14598295 DOI: 10.1002/jnr.10791]
- 52 **Vardatsikos G**, Sahu A, Srivastava AK. The insulin-like growth factor family: molecular mechanisms, redox regulation, and clinical implications. *Antioxid Redox Signal* 2009; **11**: 1165-1190 [PMID: 19014342 DOI: 10.1089/ars.2008.2161]
- 53 **Fernandez AM**, Torres-Aleman I. The many faces of insulin-like peptide signalling in the brain. *Nat Rev Neurosci* 2012; **13**: 225-239 [PMID: 22430016 DOI: 10.1038/nrn3209]
- 54 **Banks WA**. The source of cerebral insulin. *Eur J Pharmacol* 2004;

- 490: 5-12 [PMID: 15094069 DOI: 10.1016/j.ejphar.2004.02.040]
- 55 **Banks WA**, Owen JB, Erickson MA. Insulin in the brain: there and back again. *Pharmacol Ther* 2012; **136**: 82-93 [PMID: 22820012 DOI: 10.1016/j.pharmthera.2012.07.006]
- 56 **Russo VC**, Gluckman PD, Feldman EL, Werther GA. The insulin-like growth factor system and its pleiotropic functions in brain. *Endocr Rev* 2005; **26**: 916-943 [PMID: 16131630 DOI: 10.1210/er.2004-0024]
- 57 **Benarroch EE**. Insulin-like growth factors in the brain and their potential clinical implications. *Neurology* 2012; **79**: 2148-2153 [PMID: 23170013 DOI: 10.1212/WNL.0b013e3182752eef]
- 58 **Ramalingam M**, Kim SJ. Mechanisms of action of brain insulin against neurodegenerative diseases. *J Neural Transm (Vienna)* 2014; **121**: 611-626 [PMID: 24398779 DOI: 10.1007/s00702-013-1147-1]
- 59 **Werner H**, LeRoith D. Insulin and insulin-like growth factor receptors in the brain: physiological and pathological aspects. *Eur Neuropsychopharmacol* 2014; **24**: 1947-1953 [PMID: 24529663 DOI: 10.1016/j.euroneuro.2014.01.020]
- 60 **Jacobs KM**, Bhawe SR, Ferraro DJ, Jaboin JJ, Hallahan DE, Thotala D. GSK-3 β : A Bifunctional Role in Cell Death Pathways. *Int J Cell Biol* 2012; **2012**: 930710 [PMID: 22675363 DOI: 10.1155/2012/930710]
- 61 **Withers DJ**, Burks DJ, Towery HH, Altamuro SL, Flint CL, White MF. Irs-2 coordinates Igf-1 receptor-mediated beta-cell development and peripheral insulin signalling. *Nat Genet* 1999; **23**: 32-40 [PMID: 10471495 DOI: 10.1038/12631]
- 62 **Aleman A**, Torres-Alemán I. Circulating insulin-like growth factor I and cognitive function: neuromodulation throughout the lifespan. *Prog Neurobiol* 2009; **89**: 256-265 [PMID: 19665513 DOI: 10.1016/j.pneurobio.2009.07.008]
- 63 **Torres-Aleman I**. Toward a comprehensive neurobiology of IGF-I. *Dev Neurobiol* 2010; **70**: 384-396 [PMID: 20186710 DOI: 10.1002/dneu.20778]
- 64 **Trejo JL**, Carro E, Garcia-Galloway E, Torres-Aleman I. Role of insulin-like growth factor I signaling in neurodegenerative diseases. *J Mol Med (Berl)* 2004; **82**: 156-162 [PMID: 14647921 DOI: 10.1007/s00109-003-0499-7]
- 65 **Huffman DM**, Farias Quipildor G, Mao K, Zhang X, Wan J, Apontes P, Cohen P, Barzilay N. Central insulin-like growth factor-1 (IGF-1) restores whole-body insulin action in a model of age-related insulin resistance and IGF-1 decline. *Aging Cell* 2016; **15**: 181-186 [PMID: 26534869 DOI: 10.1111/accel.12415]
- 66 **Kappeler L**, De Magalhães Filho C, Dupont J, Leneuve P, Cervera P, Périn L, Loudes C, Blaise A, Klein R, Epelbaum J, Le Bouc Y, Holzenberger M. Brain IGF-1 receptors control mammalian growth and lifespan through a neuroendocrine mechanism. *PLoS Biol* 2008; **6**: e254 [PMID: 18959478 DOI: 10.1371/journal.pbio.0060254]
- 67 **Starr JM**, Wardlaw J, Ferguson K, MacLulich A, Deary IJ, Marshall I. Increased blood-brain barrier permeability in type II diabetes demonstrated by gadolinium magnetic resonance imaging. *J Neurol Neurosurg Psychiatry* 2003; **74**: 70-76 [PMID: 12486269 DOI: 10.1136/jnnp.74.1.70]
- 68 **Duarte AI**, Moreira PI, Oliveira CR. Insulin in central nervous system: more than just a peripheral hormone. *J Aging Res* 2012; **2012**: 384017 [PMID: 22500228 DOI: 10.1155/2012/384017]
- 69 **Zeng Y**, Zhang L, Hu Z. Cerebral insulin, insulin signaling pathway, and brain angiogenesis. *Neurol Sci* 2016; **37**: 9-16 [PMID: 26442674 DOI: 10.1007/s10072-015-2386-8]
- 70 **Sánchez-Lasheras C**, Könnér AC, Brüning JC. Integrative neurobiology of energy homeostasis-neurocircuits, signals and mediators. *Front Neuroendocrinol* 2010; **31**: 4-15 [PMID: 19729032 DOI: 10.1016/j.yfne.2009.08.002]
- 71 **Power ML**, Schulkin J. Sex differences in fat storage, fat metabolism, and the health risks from obesity: possible evolutionary origins. *Br J Nutr* 2008; **99**: 931-940 [PMID: 17977473 DOI: 10.1017/S0007114507853347]
- 72 **Mauvais-Jarvis F**. Sex differences in metabolic homeostasis, diabetes, and obesity. *Biol Sex Differ* 2015; **6**: 14 [PMID: 26339468 DOI: 10.1186/s13293-015-0033-y]
- 73 **Varlamov O**, Bethea CL, Roberts CT. Sex-specific differences in lipid and glucose metabolism. *Front Endocrinol (Lausanne)* 2014; **5**: 241 [PMID: 25646091 DOI: 10.3389/fendo.2014.00241]
- 74 **Goodman-Gruen D**, Barrett-Connor E. Sex differences in the association of endogenous sex hormone levels and glucose tolerance status in older men and women. *Diabetes Care* 2000; **23**: 912-918 [PMID: 10895840 DOI: 10.2337/diacare.23.7.912]
- 75 **Phillips GB**, Tuck CH, Jing TY, Boden-Albala B, Lin IF, Dahodwala N, Sacco RL. Association of hyperandrogenemia and hyperestrogenemia with type 2 diabetes in Hispanic postmenopausal women. *Diabetes Care* 2000; **23**: 74-79 [PMID: 10857972 DOI: 10.2337/diacare.23.1.74]
- 76 **Ding EL**, Song Y, Manson JE, Rifai N, Buring JE, Liu S. Plasma sex steroid hormones and risk of developing type 2 diabetes in women: a prospective study. *Diabetologia* 2007; **50**: 2076-2084 [PMID: 17701157 DOI: 10.1007/s00125-007-0785-y]
- 77 **Kalyani RR**, Franco M, Dobs AS, Ouyang P, Vaidya D, Bertoni A, Gapstur SM, Golden SH. The association of endogenous sex hormones, adiposity, and insulin resistance with incident diabetes in postmenopausal women. *J Clin Endocrinol Metab* 2009; **94**: 4127-4135 [PMID: 19789205 DOI: 10.1210/jc.2009-0910]
- 78 **Hu J**, Zhang A, Yang S, Wang Y, Goswami R, Zhou H, Zhang Y, Wang Z, Li R, Cheng Q, Zhen Q, Li Q. Combined effects of sex hormone-binding globulin and sex hormones on risk of incident type 2 diabetes. *J Diabetes* 2016; **8**: 508-515 [PMID: 26119029 DOI: 10.1111/1753-0407.12322]
- 79 **Oh JY**, Barrett-Connor E, Wedick NM, Wingard DL. Endogenous sex hormones and the development of type 2 diabetes in older men and women: the Rancho Bernardo study. *Diabetes Care* 2002; **25**: 55-60 [PMID: 11772901 DOI: 10.2337/diacare.25.1.55]
- 80 **Vikan T**, Schirmer H, Njølstad I, Svartberg J. Low testosterone and sex hormone-binding globulin levels and high estradiol levels are independent predictors of type 2 diabetes in men. *Eur J Endocrinol* 2010; **162**: 747-754 [PMID: 20061333 DOI: 10.1530/EJE-09-0943]
- 81 **Cauley JA**, Gutai JP, Kuller LH, LeDonne D, Powell JG. The epidemiology of serum sex hormones in postmenopausal women. *Am J Epidemiol* 1989; **129**: 1120-1131 [PMID: 2729251 DOI: 10.1016/0378-5122(89)90030-3]
- 82 **Mayes JS**, Watson GH. Direct effects of sex steroid hormones on adipose tissues and obesity. *Obes Rev* 2004; **5**: 197-216 [PMID: 15458395 DOI: 10.1111/j.1467-789X.2004.00152.x]
- 83 **Turgeon JL**, Carr MC, Maki PM, Mendelsohn ME, Wise PM. Complex actions of sex steroids in adipose tissue, the cardiovascular system, and brain: Insights from basic science and clinical studies. *Endocr Rev* 2006; **27**: 575-605 [PMID: 16763155 DOI: 10.1210/er.2005-0020]
- 84 **Paik SG**, Michelis MA, Kim YT, Shin S. Induction of insulin-dependent diabetes by streptozotocin. Inhibition by estrogens and potentiation by androgens. *Diabetes* 1982; **31**: 724-729 [PMID: 6219020 DOI: 10.2337/diab.31.8.724]
- 85 **Heine PA**, Taylor JA, Iwamoto GA, Lubahn DB, Cooke PS. Increased adipose tissue in male and female estrogen receptor-alpha knockout mice. *Proc Natl Acad Sci USA* 2000; **97**: 12729-12734 [PMID: 11070086 DOI: 10.1073/pnas.97.23.12729]
- 86 **Andersson B**, Mattsson LA, Hahn L, Mårin P, Lapidus L, Holm G, Bengtsson BA, Björntorp P. Estrogen replacement therapy decreases hyperandrogenicity and improves glucose homeostasis and plasma lipids in postmenopausal women with noninsulin-dependent diabetes mellitus. *J Clin Endocrinol Metab* 1997; **82**: 638-643 [PMID: 9024268 DOI: 10.1210/jcem.82.2.3746]
- 87 **Moran LJ**, Misso ML, Wild RA, Norman RJ. Impaired glucose tolerance, type 2 diabetes and metabolic syndrome in polycystic ovary syndrome: a systematic review and meta-analysis. *Hum Reprod Update* 2010; **16**: 347-363 [PMID: 20159883 DOI: 10.1093/humupd/dmq001]
- 88 **Dhindsa S**, Prabhakar S, Sethi M, Bandyopadhyay A, Chaudhuri A, Dandona P. Frequent occurrence of hypogonadotropic hypogonadism in type 2 diabetes. *J Clin Endocrinol Metab* 2004; **89**: 5462-5468 [PMID: 15531498 DOI: 10.1210/jc.2004-0804]
- 89 **Rhoden EL**, Ribeiro EP, Teloken C, Souto CA. Diabetes mellitus is associated with subnormal serum levels of free testosterone in men.

- BJU Int* 2005; **96**: 867-870 [PMID: 16153219 DOI: 10.1111/j.1464-410X.2005.05728.x]
- 90 **Stellato RK**, Feldman HA, Hamdy O, Horton ES, McKinlay JB. Testosterone, sex hormone-binding globulin, and the development of type 2 diabetes in middle-aged men: prospective results from the Massachusetts male aging study. *Diabetes Care* 2000; **23**: 490-494 [PMID: 10857940 DOI: 10.2337/diacare.23.4.490]
- 91 **Keating NL**, O'Malley A, Freedland SJ, Smith MR. Diabetes and cardiovascular disease during androgen deprivation therapy: observational study of veterans with prostate cancer. *J Natl Cancer Inst* 2012; **104**: 1518-1523 [PMID: 23210129 DOI: 10.1200/JCO.2006.06.2497]
- 92 **Laaksonen DE**, Niskanen L, Punnonen K, Nyssönen K, Tuomainen TP, Valkonen VP, Salonen R, Salonen JT. Testosterone and sex hormone-binding globulin predict the metabolic syndrome and diabetes in middle-aged men. *Diabetes Care* 2004; **27**: 1036-1041 [PMID: 15111517 DOI: 10.2337/diacare.27.5.1036]
- 93 **Corona G**, Monami M, Rastrelli G, Aversa A, Sforza A, Lenzi A, Forti G, Mannucci E, Maggi M. Type 2 diabetes mellitus and testosterone: a meta-analysis study. *Int J Androl* 2011; **34**: 528-540 [PMID: 20969599 DOI: 10.1111/j.1365-2605.2010.01117.x]
- 94 **Foryst-Ludwig A**, Kintscher U. Metabolic impact of estrogen signalling through ERalpha and ERbeta. *J Steroid Biochem Mol Biol* 2010; **122**: 74-81 [PMID: 20599505 DOI: 10.1016/j.jsbmb.2010.06.012]
- 95 **Fuente-Martín E**, Garcia-Caceres C, Morselli E, Clegg DJ, Chowen JA, Finan B, Brinton RD, Tschöp MH. Estrogen, astrocytes and the neuroendocrine control of metabolism. *Rev Endocr Metab Disord* 2013; **14**: 331-338 [PMID: 24009071 DOI: 10.1007/s1154-013-9263-7]
- 96 **Azcoitia I**, Sierra A, Garcia-Segura LM. Neuroprotective effects of estradiol in the adult rat hippocampus: interaction with insulin-like growth factor-I signalling. *J Neurosci Res* 1999; **58**: 815-822 [PMID: 10583912 DOI: 10.1002/(SICI)1097-4547(19991215)58:]
- 97 **Mendez P**, Azcoitia I, Garcia-Segura LM. Estrogen receptor alpha forms estrogen-dependent multimolecular complexes with insulin-like growth factor receptor and phosphatidylinositol 3-kinase in the adult rat brain. *Brain Res Mol Brain Res* 2003; **112**: 170-176 [PMID: 12670715 DOI: 10.1016/S0169-328X(03)00088-3]
- 98 **D'Astous M**, Mendez P, Morissette M, Garcia-Segura LM, Di Paolo T. Implication of the phosphatidylinositol-3 kinase/protein kinase B signaling pathway in the neuroprotective effect of estradiol in the striatum of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine mice. *Mol Pharmacol* 2006; **69**: 1492-1498 [PMID: 16434614 DOI: 10.1124/mol.105.018671]
- 99 **Quesada A**, Lee BY, Micevych PE. PI3 kinase/Akt activation mediates estrogen and IGF-1 nigral DA neuronal neuroprotection against a unilateral rat model of Parkinson's disease. *Dev Neurobiol* 2008; **68**: 632-644 [PMID: 18278798 DOI: 10.1002/dneu.20609]
- 100 **Cardona-Gomez GP**, Mendez P, Garcia-Segura LM. Synergistic interaction of estradiol and insulin-like growth factor-I in the activation of PI3K/Akt signaling in the adult rat hypothalamus. *Brain Res Mol Brain Res* 2002; **107**: 80-88 [PMID: 12414126 DOI: 10.1016/S0169-328X(02)00449-7]
- 101 **Marin R**, Diaz M, Alonso R, Sanz A, Arévalo MA, Garcia-Segura LM. Role of estrogen receptor alpha in membrane-initiated signaling in neural cells: interaction with IGF-1 receptor. *J Steroid Biochem Mol Biol* 2009; **114**: 2-7 [PMID: 19167493 DOI: 10.1016/j.jsbmb.2008.12.014]
- 102 **Cardona-Gómez GP**, DonCarlos L, Garcia-Segura LM. Insulin-like growth factor I receptors and estrogen receptors colocalize in female rat brain. *Neuroscience* 2000; **99**: 751-760 [PMID: 10974438 DOI: 10.1016/S0306-4522(00)00228-1]
- 103 **Cardona-Gomez P**, Perez M, Avila J, Garcia-Segura LM, Wandosell F. Estradiol inhibits GSK3 and regulates interaction of estrogen receptors, GSK3, and beta-catenin in the hippocampus. *Mol Cell Neurosci* 2004; **25**: 363-373 [PMID: 15033165 DOI: 10.1016/j.mcn.2003.10.008]
- 104 **Mendez P**, Wandosell F, Garcia-Segura LM. Cross-talk between estrogen receptors and insulin-like growth factor-I receptor in the brain: cellular and molecular mechanisms. *Front Neuroendocrinol* 2006; **27**: 391-403 [PMID: 17049974 DOI: 10.1016/j.yfme.2006.09.001]
- 105 **Mendez P**, Azcoitia I, Garcia-Segura LM. Interdependence of oestrogen and insulin-like growth factor-I in the brain: potential for analysing neuroprotective mechanisms. *J Endocrinol* 2005; **185**: 11-17 [PMID: 15817823 DOI: 10.1677/joe.1.06058]
- 106 **Varea O**, Arevalo MA, Garrido JJ, Garcia-Segura LM, Wandosell F, Mendez P. Interaction of estrogen receptors with insulin-like growth factor-I and Wnt signaling in the nervous system. *Steroids* 2010; **75**: 565-569 [PMID: 19778547 DOI: 10.1016/j.steroids.2009.09.006]
- 107 **Brywe KG**, Mallard C, Gustavsson M, Hedtjörn M, Leverin AL, Wang X, Blomgren K, Isgaard J, Hagberg H. IGF-I neuroprotection in the immature brain after hypoxia-ischemia, involvement of Akt and GSK3beta? *Eur J Neurosci* 2005; **21**: 1489-1502 [PMID: 15845077 DOI: 10.1111/j.1460-9568.2005.03982.x]
- 108 **Dhillon AS**, Hagan S, Rath O, Kolch W. MAP kinase signalling pathways in cancer. *Oncogene* 2007; **26**: 3279-3290 [PMID: 17496922 DOI: 10.1038/sj.onc.1210421]
- 109 **Cardona-Gómez GP**, Mendez P, DonCarlos LL, Azcoitia I, Garcia-Segura LM. Interactions of estrogen and insulin-like growth factor-I in the brain: molecular mechanisms and functional implications. *J Steroid Biochem Mol Biol* 2002; **83**: 211-217 [PMID: 12650718 DOI: 10.1016/S0960-0760(02)00261-3]
- 110 **Convit A**, Wolf OT, Tarshish C, de Leon MJ. Reduced glucose tolerance is associated with poor memory performance and hippocampal atrophy among normal elderly. *Proc Natl Acad Sci USA* 2003; **100**: 2019-2022 [PMID: 12571363 DOI: 10.1073/pnas.0336073100]
- 111 **Gold SM**, Dziobek I, Sweat V, Tirsi A, Rogers K, Bruehl H, Tsui W, Richardson S, Javier E, Convit A. Hippocampal damage and memory impairments as possible early brain complications of type 2 diabetes. *Diabetologia* 2007; **50**: 711-719 [PMID: 17334649 DOI: 10.1007/s00125-007-0602-7]
- 112 **Kamiyama K**, Wada A, Sugihara M, Kurioka S, Hayashi K, Hayashi T, Yoshisako T, Yamamoto N, Tsuchie Y, Yamaguchi S, Sugimoto T, Kitagaki H. Potential hippocampal region atrophy in diabetes mellitus type 2: a voxel-based morphometry VSRAD study. *Jpn J Radiol* 2010; **28**: 266-272 [PMID: 20512543 DOI: 10.1007/s11604-009-0416-2]
- 113 **Nelson BS**, Springer RC, Daniel JM. Antagonism of brain insulin-like growth factor-1 receptors blocks estradiol effects on memory and levels of hippocampal synaptic proteins in ovariectomized rats. *Psychopharmacology (Berl)* 2014; **231**: 899-907 [PMID: 24146138 DOI: 10.1007/s00213-013-3310-7]
- 114 **Perez-Martín M**, Azcoitia I, Trejo JL, Sierra A, Garcia-Segura LM. An antagonist of estrogen receptors blocks the induction of adult neurogenesis by insulin-like growth factor-I in the dentate gyrus of adult female rat. *Eur J Neurosci* 2003; **18**: 923-930 [PMID: 12925018 DOI: 10.1046/j.1460-9568.2003.02830.x]
- 115 **Takeuchi K**, Yang Y, Takayasu Y, Gertner M, Hwang JY, Aromolaran K, Bennett MV, Zukin RS. Estradiol pretreatment ameliorates impaired synaptic plasticity at synapses of insulted CA1 neurons after transient global ischemia. *Brain Res* 2015; **1621**: 222-230 [PMID: 25463028 DOI: 10.1016/j.brainres.2014.11.016]
- 116 **Witty CF**, Gardella LP, Perez MC, Daniel JM. Short-term estradiol administration in aging ovariectomized rats provides lasting benefits for memory and the hippocampus: a role for insulin-like growth factor-I. *Endocrinology* 2013; **154**: 842-852 [PMID: 23264616 DOI: 10.1210/en.2012-1698]
- 117 **Cardona-Gómez GP**, Mendez P, DonCarlos LL, Azcoitia I, Garcia-Segura LM. Interactions of estrogens and insulin-like growth factor-I in the brain: implications for neuroprotection. *Brain Res Brain Res Rev* 2001; **37**: 320-334 [PMID: 11744097 DOI: 10.1016/S0165-0173(01)00137-0]
- 118 **Aggarwal RR**, Ryan CJ, Chan JM. Insulin-like growth factor pathway: a link between androgen deprivation therapy (ADT), insulin resistance, and disease progression in patients with prostate cancer? *Urol Oncol* 2013; **31**: 522-530 [PMID: 21658978 DOI: 10.1016/j.urolonc.2011.05.001]
- 119 **Oki K**, Law TD, Loucks AB, Clark BC. The effects of testosterone

- and insulin-like growth factor 1 on motor system form and function. *Exp Gerontol* 2015; **64**: 81-86 [PMID: 25681641 DOI: 10.1016/j.exger.2015.02.005]
- 120 **Castilla-Cortázar I**, García-Fernández M, Delgado G, Puche JE, Sierra I, Barhoum R, González-Barón S. Hepatoprotection and neuroprotection induced by low doses of IGF-II in aging rats. *J Transl Med* 2011; **9**: 103 [PMID: 21733157 DOI: 10.1186/1479-5876-9-103]
- 121 **Lin SY**, Cui H, Yusta B, Belsham DD. IGF-I signaling prevents dehydroepiandrosterone (DHEA)-induced apoptosis in hypothalamic neurons. *Mol Cell Endocrinol* 2004; **214**: 127-135 [PMID: 15062551 DOI: 10.1016/j.mce.2003.10.064]
- 122 **Bialek M**, Zaremba P, Borowicz KK, Czuczwar SJ. Neuroprotective role of testosterone in the nervous system. *Pol J Pharmacol* 2004; **56**: 509-518 [PMID: 15591638]
- 123 **Creta M**, Riccio R, Chiancone F, Fusco F. Androgens exert direct neuroprotective effects on the brain: a review of pre-clinical evidences. *J Androl Sci* 2010; **17**: 49-55
- 124 **Garcia-Segura LM**, Balthazart J. Steroids and neuroprotection: New advances. *Front Neuroendocrinol* 2009; **30**: v-ix [PMID: 19393683 DOI: 10.1016/j.yfrne.2009.04.006]
- 125 **Hammond J**, Le Q, Goodyer C, Gelfand M, Trifiro M, LeBlanc A. Testosterone-mediated neuroprotection through the androgen receptor in human primary neurons. *J Neurochem* 2001; **77**: 1319-1326 [PMID: 11389183 DOI: 10.1046/j.1471-4159.2001.00345.x]
- 126 **Nguyen TV**, Jayaraman A, Quaglino A, Pike CJ. Androgens selectively protect against apoptosis in hippocampal neurons. *J Neuroendocrinol* 2010; **22**: 1013-1022 [PMID: 20561156 DOI: 10.1111/j.1365-2826.2010.02044.x]
- 127 **Baron S**, Manin M, Beaudoine C, Leotoing L, Communal Y, Veysiere G, Morel L. Androgen receptor mediates non-genomic activation of phosphatidylinositol 3-OH kinase in androgen-sensitive epithelial cells. *J Biol Chem* 2004; **279**: 14579-14586 [PMID: 14668339 DOI: 10.1074/jbc.M306143200]
- 128 **Rochette-Egly C**. Nuclear receptors: integration of multiple signalling pathways through phosphorylation. *Cell Signal* 2003; **15**: 355-366 [PMID: 12618210 DOI: 10.1016/S0898-6568(02)00115-8]
- 129 **Bing L**, Wu J, Zhang J, Chen Y, Hong Z, Zu H. DHT inhibits the A β 25-35-induced apoptosis by regulation of seladin-1, survivin, XIAP, bax, and bcl-xl expression through a rapid PI3-K/Akt signaling in C6 glial cell lines. *Neurochem Res* 2015; **40**: 41-48 [PMID: 25347962 DOI: 10.1007/s11064-014-1463-3]
- 130 **Gatson JW**, Kaur P, Singh M. Dihydrotestosterone differentially modulates the mitogen-activated protein kinase and the phosphoinositide 3-kinase/Akt pathways through the nuclear and novel membrane androgen receptor in C6 cells. *Endocrinology* 2006; **147**: 2028-2034 [PMID: 16410299 DOI: 10.1210/en.2005-1395]
- 131 **Gatson JW**, Singh M. Activation of a membrane-associated androgen receptor promotes cell death in primary cortical astrocytes. *Endocrinology* 2007; **148**: 2458-2464 [PMID: 17303658 DOI: 10.1210/en.2006-1443]
- 132 **Nguyen TV**, Yao M, Pike CJ. Androgens activate mitogen-activated protein kinase signaling: role in neuroprotection. *J Neurochem* 2005; **94**: 1639-1651 [PMID: 16011741 DOI: 10.1111/j.1471-4159.2005.03318.x]
- 133 **Pike CJ**, Nguyen TV, Ramsden M, Yao M, Murphy MP, Rosario ER. Androgen cell signaling pathways involved in neuroprotective actions. *Horm Behav* 2008; **53**: 693-705 [PMID: 18222446 DOI: 10.1016/j.yhbeh.2007.11.006]
- 134 **Azcoitia I**, Sierra A, Veiga S, Honda S, Harada N, Garcia-Segura LM. Brain aromatase is neuroprotective. *J Neurobiol* 2001; **47**: 318-329 [PMID: 11351342 DOI: 10.1002/neu.1038]
- 135 **Azcoitia I**, Sierra A, Veiga S, Garcia-Segura LM. Aromatase expression by reactive astroglia is neuroprotective. *Ann N Y Acad Sci* 2003; **1007**: 298-305 [PMID: 14993062 DOI: 10.1196/annals.1286.028]
- 136 **Garcia-Segura LM**, Veiga S, Sierra A, Melcangi RC, Azcoitia I. Aromatase: a neuroprotective enzyme. *Prog Neurobiol* 2003; **71**: 31-41 [PMID: 14611865 DOI: 10.1016/j.pneurobio.2003.09.005]
- 137 **Garcia-Segura LM**. Aromatase in the brain: not just for reproduction anymore. *J Neuroendocrinol* 2008; **20**: 705-712 [PMID: 18601693 DOI: 10.1111/j.1365-2826.2008.01713.x]
- 138 **Roselli CF**. Brain aromatase: roles in reproduction and neuroprotection. *J Steroid Biochem Mol Biol* 2007; **106**: 143-150 [PMID: 17643294 DOI: 10.1016/j.jsmb.2007.05.014]
- 139 **Saldanha CJ**, Duncan KA, Walters BJ. Neuroprotective actions of brain aromatase. *Front Neuroendocrinol* 2009; **30**: 106-118 [PMID: 19450619 DOI: 10.1016/j.yfrne.2009.04.016]
- 140 **Carswell HV**, Dominiczak AF, Garcia-Segura LM, Harada N, Hutchison JB, Macrae IM. Brain aromatase expression after experimental stroke: topography and time course. *J Steroid Biochem Mol Biol* 2005; **96**: 89-91 [PMID: 15896953 DOI: 10.1016/j.jsmb.2005.02.016]
- 141 **Burul-Bozkurt N**, Pekiner C, Kelicen P. Diabetes alters aromatase enzyme levels in gonadal tissues of rats. *Naunyn Schmiedebergs Arch Pharmacol* 2010; **382**: 33-41 [PMID: 20428845 DOI: 10.1007/s00210-010-0518-5]
- 142 **Gibb FW**, Homer NZ, Faqehi AM, Upreti R, Livingstone DE, McInnes KJ, Andrew R, Walker BR. Aromatase Inhibition Reduces Insulin Sensitivity in Healthy Men. *J Clin Endocrinol Metab* 2016; **101**: 2040-2046 [PMID: 26967690 DOI: 10.1210/jc.2015-4146]
- 143 **Pintana H**, Chattipakorn N, Chattipakorn S. Testosterone deficiency, insulin-resistant obesity and cognitive function. *Metab Brain Dis* 2015; **30**: 853-876 [PMID: 25703239 DOI: 10.1007/s11011-015-9655-3]
- 144 **Dozmorov MG**, Yang Q, Matwalli A, Hurst RE, Culkin DJ, Kropp BP, Lin HK. 5 α -androstane-3 α ,17 β -diol selectively activates the canonical PI3K/AKT pathway: a bioinformatics-based evidence for androgen-activated cytoplasmic signaling. *Genomic Med* 2007; **1**: 139-146 [PMID: 18923939 DOI: 10.1007/s11568-008-9018-9]
- 145 **Narenji SA**, Naghdi N, Azadmanesh K, Edalat R. 3 α -diol administration decreases hippocampal PKA (II) mRNA expression and impairs Morris water maze performance in adult male rats. *Behav Brain Res* 2015; **280**: 149-159 [PMID: 25451551 DOI: 10.1016/j.bbr.2014.11.038]
- 146 **Caruso D**, Scurati S, Maschi O, De Angelis L, Roglio I, Giatti S, Garcia-Segura LM, Melcangi RC. Evaluation of neuroactive steroid levels by liquid chromatography-tandem mass spectrometry in central and peripheral nervous system: effect of diabetes. *Neurochem Int* 2008; **52**: 560-568 [PMID: 17686551 DOI: 10.1016/j.neuint.2007.06.004]

P- Reviewer: Comasco E, Egea J, Phillips J **S- Editor:** Qiu S
L- Editor: A **E- Editor:** Wu HL





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: bpgoffice@wjgnet.com

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>

