

RESEARCH ARTICLE

The costs of providing antiretroviral therapy services to HIV-infected individuals presenting with advanced HIV disease at public health centres in Dar es Salaam, Tanzania: Findings from a randomised trial evaluating different health care strategies

Godfather Dickson Kimaro^{1,2*}, Sayoki Mfinanga¹, Victoria Simms², Sokoine Kivuyo¹, Christian Bottomley², Neil Hawkins³, Thomas S. Harrison⁴, Shabbar Jaffar⁵, Lorna Guinness³, on behalf of the REMSTART trial team[¶]

1 Muhimbili Medical Research Centre, National Institute for Medical Research, Dar es Salaam, Tanzania, **2** Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, United Kingdom, **3** Department of Global Health and Development, London School of Hygiene and Tropical Medicine, London, United Kingdom, **4** Institute for Infection and Immunity, St Georges University of London, London, United Kingdom, **5** Department of International Public Health, Liverpool School of Tropical Medicine, Liverpool, United Kingdom

[¶] Membership of the REMSTART trial team is provided in the Acknowledgments.
* fatherdk@yahoo.com



OPEN ACCESS

Citation: Kimaro GD, Mfinanga S, Simms V, Kivuyo S, Bottomley C, Hawkins N, et al. (2017) The costs of providing antiretroviral therapy services to HIV-infected individuals presenting with advanced HIV disease at public health centres in Dar es Salaam, Tanzania: Findings from a randomised trial evaluating different health care strategies. PLoS ONE 12(2): e0171917. doi:10.1371/journal.pone.0171917

Editor: Giuseppe Vittorio De Socio, Azienda Ospedaliera Universitaria di Perugia, ITALY

Received: February 24, 2016

Accepted: January 28, 2017

Published: February 24, 2017

Copyright: © 2017 Kimaro et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This study was nested within the REMSTART trial which was funded by the European and Developing Countries Clinical Trials Partnership (IP.2009.33011.003). GDK received a scholarship from HIV Research Trust Scholarship

Abstract

Background

Understanding the costs associated with health care delivery strategies is essential for planning. There are few data on health service resources used by patients and their associated costs within antiretroviral (ART) programmes in Africa.

Material and methods

The study was nested within a large trial, which evaluated screening for cryptococcal meningitis and tuberculosis and a short initial period of home-based adherence support for patients initiating ART with advanced HIV disease in Tanzania and Zambia. The economic evaluation was done in Tanzania alone. We estimated costs of providing routine ART services from the health service provider's perspective using a micro-costing approach. Incremental costs for the different novel components of service delivery were also estimated. All costs were converted into US dollars (US\$) and based on 2012 prices.

Results

Of 870 individuals enrolled in Tanzania, 434 were enrolled in the intervention arm and 436 in the standard care/control arm. Overall, the median (IQR) age and CD4 cell count at enrolment were 38 [31, 44] years and 52 [20, 89] cells/mm³, respectively. The mean per patient

(HIVRT15-008) for his course on economic evaluation and attachment with experienced researchers at London School of Hygiene and Tropical Medicine. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: TTSH has been given antigen tests by IMMY for studies of cryptococcal meningitis treatment and prevention. However, this does not alter our adherence to PLOS ONE policies on sharing data and materials. All other authors declared that no competing interests exist.

costs over the first three months and over a one year period of follow up following ART initiation in the standard care arm were US\$ 107 (95%CI 101–112) and US\$ 265 (95%CI 254–275) respectively. ART drugs, clinic visits and hospital admission constituted 50%, 19%, and 19% of the total cost per patient year, while diagnostic tests and non-ART drugs (co-trimoxazole) accounted for 10% and 2% of total per patient year costs. The incremental costs of the intervention to the health service over the first three months was US\$ 59 ($p < 0.001$; 95%CI 52–67) and over a one year period was US\$ 67 ($p < 0.001$; 95%CI 50–83). This is equivalent to an increase of 55% (95%CI 51%–59%) in the mean cost of care over the first three months, and 25% (95%CI 20%–30%) increase over one year of follow up.

Background

Over 12 million HIV-infected people now have access to antiretroviral therapy (ART) in Africa [1] where there are severe constraints on health care resources, in particular, a severe shortage of health care workers. Understanding the costs associated with different approaches to health care delivery is essential to inform policy, and especially important in resource-limited settings [2]. Despite this, there are surprisingly few published studies describing the costs of ART programmes in low income countries in Africa [3].

Costing studies use either the “top-down” or “bottom-up” approach or a combination of these [4,5]. The former estimates costs by dividing the past expenditure by the number of services provided during the period that expenditure was incurred. These analyses are crude and it is not possible to assess how patient characteristics influence costs and or the extent to which costs vary between settings. The “bottom up” approach, also referred to as the ingredient-based approach, quantifies the inputs used to provide the service outputs. Micro-costing is a form of “bottom up” approach and is essential for evaluating new interventions [4,6,7] as it allows for the statistical analysis of the key cost drivers at the individual level. Since 2003, just four published HIV costing studies from low income sub-Saharan Africa have used an individual level micro-costing approach, but all were retrospective [8–11] and their findings are now outdated [12–16].

Materials and methods

In this study, we estimated the costs of ART delivery in the primary care setting in Tanzania using a micro-costing approach. Our study was nested within a trial, which aimed to reduce HIV-mortality among patients presenting in the very advanced stages of HIV-infection (REMSTART) [17]. Half of the participants received routine standard care. The other half received routine standard care combined with additional screening and adherence support. The participants were managed by health care staff according to national guidelines so that the trial provided an opportunity to estimate costs of real-life ART care in an urban setting. We analysed the resource use and costs of ART in both the routine health services and with the added intervention within REMSTART [17].

Study settings

The REMSTART trial was implemented in Dar es Salaam, Tanzania, and Lusaka, Zambia, but the cost study was restricted to Tanzania for logistical reasons. Dar es Salaam has an estimated population of 4.4 million people [18] and an HIV prevalence among 15–49 year old of 6.9%

[19]. The city is divided into three districts—Kinondoni, Ilala and Temeke. The trial was implemented at three government-run primary care health centres, one in each district (Tandale in Kinondoni, Buguruni in Ilala and Mbagala Rangi Tatu in Temeke).

In Tanzania, HIV is managed largely from primary care clinics. National guidelines at the time of the study were that patients presenting with a diagnosis of HIV-infection should have been offered ART if they had a CD4 count ≤ 350 cells/mm³ or if they were at WHO clinical stages 3 or 4 [20]. Follow-up clinic visits were initially every month and then two-monthly, when the patient was considered stable on therapy according to the clinician’s assessment. The REMSTART trial was restricted to patients who presented with advanced disease because they have very high rates of mortality [21–24]. Initially, patients with CD4 count < 100 cells/mm³ were enrolled, but this criteria was changed subsequently to enrol patients presenting with < 200 cells/mm³, in order to speed up recruitment [17].

Participants were individually randomised [17] to receive either routine care or routine care plus additional clinic and community-based services (the REMSTART intervention). All participants (in both arms) were screened for tuberculosis (TB), irrespective of symptoms and tested for TB using GeneXpert[®] MTB/RIF assay (Cepheid, Sunnyvale, USA) (hereon referred to as Xpert) and ART was initiated rapidly where possible. Fig 1 shows both the pathway for and services provided to individuals in intervention and standard care arms. At the time of the trial implementation, Xpert was not part of standard care but all participants were tested for TB using Xpert because it had been endorsed by the WHO as a screening tool for TB. Apart from screening for TB using Xpert test, study participants in the routine care arm were managed according to the standard of care. The clinic plus community support arm received the following additional services: i) screening for cryptococcal infection using serum cryptococcal antigen combined with pre-emptive fluconazole therapy for patients testing antigen positive; ii) weekly home visits for the first 4 weeks on ART by trained lay workers to provide adherence




Fig 1. Pathway of care for HIV-infected individuals presenting in with CD4 count < 200 cells/mm³ in intervention and standard care arms in Tanzania. *Under standard care, TB screening is done using smear microscopy examination or chest x-ray. For this study, Xpert screening was used in both arms at baseline.

doi:10.1371/journal.pone.0171917.g001

support; and (iii) re-screening for tuberculosis using the Xpert test at 6–8 weeks after ART initiation in participants in whom tuberculosis was not diagnosed at enrolment [17] (Fig 1).

Data collection

Participants were recruited from February 2012 to September 2013, and each participant was followed for up to 1 year. Resource use and costs were measured from a provider's perspective using a micro-costing approach. Health service resource use by the patient was tracked in trial records over a one year period for both patients under routine care and those in the clinic plus community plus support arms. This included number of clinic visits made by the patient, lay-worker home visits, diagnostic screening and treatment and inpatient costs. Cost data were collected as described in the supporting information (S1 Table) which lists the data collected and methods of collection at each of the sources. Costs were classified as recurrent or capital.

Allocation of costs. The time medical staff spent on different types of clinic visits was estimated using patient observations conducted over a one week period and personnel diaries self-completed by individual staff. Clinic visits were categorised as follows: i) a visit during which a clinical examination and CD4 count testing were done, ii) a visit during which the patient was assessed for ART eligibility, iii) a visit in which a patient self-referred to the clinic and was seen by clinician (sick visit), (iv) a visit during which a patient was clinically re-assessed following ART initiation and CD4 count testing was repeated, and v) a routine visit for the collection of drugs alone. Staff time spent on other activities was allocated to each visit in proportion to the daily patient contact time derived from personnel diaries. Such activities included morning report meetings, tea and lunch breaks, cleaning and preparing daily reports. As the clinic provides dedicated paediatric care on one day per week, 80% (4 out of 5 working days) of the shared costs were allocated to adult care. These costs were then divided by the average number of adult visits per month to obtain the cost per patient visit. Personnel time for laboratory work and lumbar punctures was derived from staff interviews. The laboratory equipment-related cost was calculated per test by dividing the annualised cost by the number of tests done. Research costs were excluded.

Costs incurred outside the clinic, other than laboratory costs, such as administrative costs, were not available for this study. Administrative costs reported elsewhere in resource limited settings ranged between 3% and 15.5% of the total costs of care [25,26]. Thus, we assumed an administrative cost of 10% of the clinic visit costs.

Cost valuation. All prices were converted to 2012 prices using the GDP deflator [27] and converted to US dollars (US\$) (Exchange rate: 1US\$ = 1,573.70 Tanzanian shillings -Tzshs) [28]. The annualization of capital costs, including training, was done using a discount rate of 7.8% equal to the 2-year Tanzanian government bond (at the time of calculation, June 2014) [29]. The annualization is the process of spreading the current purchase price of each item of equipment across its working life, taking into account its opportunity cost. The expected useful life for equipment varied from 3 to 15 years depending on the item [30]. Training of the lay-workers was estimated to have a life span of 3 years.

Analysis

Total and average resource use and costs per patient were estimated by month, quarter and year using time-series commands in Stata 13.0 (StataCorp). Costs were calculated separately for the routine care arm and then for the intervention arm. The denominator for the mean costs at all time points is all randomised patients rather than the those patients who were still alive at each time point.

Generalised linear regression analyses. We calculated a mean cost per patient (for 1 year of treatment) and used univariate and multivariate linear regression to explore the influence of study site [30], severity of illness (level of CD4 count and WHO clinical stage), marital status, level of education, gender and age [31–34] on the mean cost of HIV/AIDS care among those who received standard care.

Sensitivity analyses. To account for uncertainty in some of the key variables, univariate sensitivity analyses were done in which parameters were varied one at a time using values given in the supporting information (S2 Table). The discount rate was varied to ensure comparability with other studies [35]. Prices for ART, reagents and the Xpert cartridges were varied to explore the impact of any changes in the international market prices. Costs incurred above the clinic level were varied to reflect the uncertainty in the baseline estimate. In addition, the analysis was run replacing the cost of the Xpert with a cost for smear microscopy, given the limited use of the Xpert to date in sub-Saharan Africa. The transportation costs of Xpert sputum samples (from the clinics to the central tuberculosis reference laboratory where Xpert machines are located) are critical to cost-effectiveness of a programme [25], therefore this transport cost was varied to reflect any uncertainty in replicating the trial transport mechanism. Further, the number of home visits per day by a lay worker was varied from the base case scenario to a more likely work load scenario in a real life setting in which more home visits would be conducted. Detailed methods for the sensitivity analysis can be found in the supporting information (S2 Table). The uncertainty analyses were based on most likely, minimum and maximum values.

Ethics statement. The study was nested within a large trial (ISCRTN 20410413) which was approved by the ethics committee of the London School of Hygiene & Tropical Medicine, the Ethics and Research Science committee in Zambia, and the National Health Research Ethics Sub-Committee in Tanzania. All study participants were older than 18 years and each provided a written informed consent.

Results

Characteristics of the study participants in Tanzania

A total of 870 individuals, 436 in standard care arm and 434 in the intervention arm, were enrolled in the study. The median (IQR) age of patients was 38 [31,44] years, 62% were female, and the median (IQR) CD4 cell count at enrolment was 52 [20,89] cells/mm³ (Table 1). Overall, the median (IQR) length of follow up was 333 (347, 365) days. The majority of participants were initiated on TDF-containing (60%) or AZT-containing (36%) ARV regimens. Overall, after a year of follow up, 684 (78.6%) participants were alive and receiving care at the study clinics, 153 (17.6%) had died, 8 (0.9%) had withdrawn from the study and survival status for 25 (2.9%) participants was unknown. Sixty six of 153 (43.1%) participants who had died were from the intervention arm. Overall, 22 (5%) of the 434 participants screened for cryptococcal antigen in the intervention group tested positive for serum cryptococcal antigen.

Unit costs for different components of HIV care

Table 2 presents estimates of unit costs, at each clinic, for the different components of ART services. The mean clinic cost for outpatient visits ranged between US\$ 5.76 for a drug collection visit without clinical assessment to US\$ 6.74 for a visit that included a clinical assessment and a CD4 count test. The cost of a home visit conducted by a lay-worker was \$14.74. The cost of diagnostic tests ranged between US\$19.07 for an Xpert test and US\$ 0.31 for a creatinine test. The novel CrAg test was estimated to cost \$3.96 per participant.

Table 1. Characteristics, HIV disease progression and ART treatment regimen of for study participants (HIV patients presenting in the advanced stages of HIV-infection—CD4 <200cells/mm³), in Dar es Salaam, Tanzania.

Characteristics	Study sites			
	Buguruni	Tandale	Mbagala	All sites
Sample size	196	280	394	870
Standard (control) arm	99	140	197	436
Intervention arm	97	140	197	434
Sex				
Male n(%)	58 (29.6)	105 (37.5)	164 (41.6)	327 (37.6)
Age at enrolment mean (SD)	39.1 (10.4)	37.4 (8.6)	39.2(9.9)	38.6 (9.7)
CD4 cells/mm ³ , median (min, max)	52 (2, 189)	57.5 (1, 199)	47 (1, 199)	52. (1, 199)
ART^a regimen initiated at enrolment n(%)				
d4T-containing regimen	0 (0.00)	0 (0.00)	10(2.54)	10 (1.15)
AZT-containing regimen	68 (34.69)	146 (52.14)	100 (25.38)	314 (36.09)
TDF -containing regimen	125 (63.78)	127 (45.36)	269 (68.27)	521 (59.89)
ABC -containing regimen*	2 (1.02)	5 (1.79)	1 (0.25)	8 (0.92)
Did not initiate on ART	1 (0.51)	2 (0.71)	14 (3.55)	17 (1.95)

^aART—Antiretroviral Therapy; ABC—abacavir; TDF—tenofovir disoproxil fumarate AZT—zidovudine; d4T –stavudine

*ABC is the nucleoside reverse transcriptase inhibitors (NRTIs) used as the second line drugs for adult and adolescents in Tanzania.

doi:10.1371/journal.pone.0171917.t001

Costs for personnel and non-medical material plus supplies constituted more than 60% and 13% respectively of the unit costs for out-patient clinic visits, excluding diagnostic procedures. Costs of the reagents for CD4, CrAg and Xpert were 85%, 99% and 70% of the unit costs for these tests (see [S3 Table](#)). There was little variation between the clinics in the breakdown of costs ([Table 2](#)).

Resource utilization

[Table 3](#) summarizes the information on resources used by the study participants. Seventeen individuals (7 randomised to standard care and 10 to the intervention arm) of 870 (2%) did not start ART. Twelve of these individuals died soon after presentation, 1 was lost to follow up and 4 withdrew from the study before starting ART.

Of 436 patients in the standard care arm, 270 (62%) completed one year of follow up, which was defined as those individuals who were alive and receiving care at one of the study clinics, 66 (15%) had died, 60 (13.8%) did not return to a study site at 12 months, 37 (8.5%) withdrew from the study and information for 3 participants was missing. Overall, the study participants in the standard care arm spent 308.9 person years in ART care, equivalent to a mean follow up on ART of 0.71 year per patient. Of the total time on treatment, 62% and 37% was spent on ART regimens containing TDF and AZT, respectively; and 1% was spent on other regimens. Thirty of 429 (7%) patients who initiated on ART substituted at least one drug in their regimen during the one-year period of follow up. Patients in the standard care arm were on co-trimoxazole for a mean of 0.73 years during the study period.

In the intervention arm, 289/434 (66.6%) patients completed one year of follow up. Of those who continued to receive care from the study clinics, 51/434 (12%) had died, 36/434 (8%) did not return to a study clinic, 56/434 (13%) withdrew from the study and information for 2 participants was missing. Overall, the study participants in the intervention arm spent 316.4 years in ART care, equivalent to a mean follow up on ART of 0.73 year per patient. Of the total time on treatment, 64% and 34% was spent on ART regimens containing TDF and

Table 2. Unit costs (in 2012 US\$) of the different components of ART services for patients initiating treatment with CD4 count <200 cells/mm³ in Dar es Salaam, Tanzania.

Variable	Unit cost (2012 US\$) (per patient)			
	Buguruni	Mbagala	Tandale	Mean cost for all sites
Costs for clinic outpatient visits under ART programme				
Initial visit (assessed by clinician, blood for CD4 test taken)	6.61	6.13	7.48	6.74
ART eligibility assessment visit	6.04	5.10	6.96	6.04
Sick visit e.g. patient self-referred and seen by clinician*	6.04	5.10	6.96	6.04
6-monthly clinic (reassessed by clinician, blood sample for CD4 test taken)**	6.61	6.13	7.48	6.74
Routine follow up visit e.g. drug pick up and not seen by clinician	5.89	4.75	6.65	5.76
Costs of diagnostic tests in addition to outpatient visit costs				
CD4 count test	15.83	14.75	16.72	15.76
Alanine aminotransferase (ALAT) test	0.88	0.84	0.91	0.88
Creatinine test	0.32	0.28	0.35	0.31
Haemoglobin (Hb) test	0.83	0.86	0.91	0.87
Random blood glucose (RBG) test	0.85	0.81	0.89	0.85
Venereal disease reference laboratory (VDRL) test	1.88	1.88	1.89	1.89
Pregnancy test	0.53	0.53	0.53	0.53
Full Blood Count (FBC) test	2.06	1.79	2.29	2.04
Serum cryptococcal meningitis (CRAG) test	5.00	2.68	4.19	3.96
Costs of the service components performed outside the clinic				
Xpert test ^b				19.07
Collection and processing of cerebral spinal fluid ^c				16.29
Chest x-ray ^d				3.18
Lay worker visit to the patients home				14.74
Information obtained from the literature				
Smear microscopy test				1.81
Sputum culture (Lowenstein Jensen) for TB [25]				6.84
ART regimens[15] (costs for daily dose)				
	d4T + 3TC +NVP			0.16
	d4T + 3TC + EFV			0.28
	AZT + 3TC +NVP			0.31
	AZT + 3TC + EFV			0.61
	TDF + 3TC + NVP			0.32
	TDF + 3TC + EFV			0.60
	ABC +3TC + NVP			0.13
	ABC +3TC + EFV			0.76
	TDF + FTC + EFV			0.60
	TDF + FTC + NVP			0.48
	TDF + FTC + ABC			0.84
Fluconazole per a dose of 10 weeks				4.80
Cotrimoxazole per daily dose				0.02
District Hospital inpatient bed day[36]				26.45
Regional Hospital inpatient bed day[36]				24.72
Tertiary Hospital inpatient bed day ^e				24.01

(Continued)

Table 2. (Continued)

Variable	Unit cost (2012 US\$) (per patient)			
	Buguruni	Mbagala	Tandale	Mean cost for all sites
TB treatment (6 monthly dose)				225

* Involves similar activities and therefore unit costs were assumed to be the same as for the ART eligibility assessment visit

** Involves similar activities and therefore unit costs were assumed to be the same as for the initial visit.

^bPerformed at the Central TB laboratory

^cDone at Muhimbili National Hospital

^dperformed at the respective district hospital

^eCosts for tertiary hospital were not available; we therefore assumed these costs were the same as those of the regional hospital

doi:10.1371/journal.pone.0171917.t002

AZT, respectively, and 2% was spent on other regimens. Forty-four of 388 (11%) patients who were initiated on ART substituted at least one drug in their regimen during the one year period of follow up. Patients in the intervention arm were on co-trimoxazole for a mean of 0.74 years during the study period. Compared to patients in standard care, patients in the intervention

Table 3. Unit costs (in 2012 US\$), average quantity of resources utilized over 12 months of treatment and mean annual cost per patient, for patients initiating treatment with CD4 count <200 cells/mm³ in Dar es Salaam, Tanzania.

Item	Unit cost (US \$2012)	Mean? resource utilisation per person		Cost per patient (US\$2012)		Difference (Intervention—control)
		Intervention	Standard care	Intervention arm	Standard care	
Out patients visits						
Initial visits	6.74	1.05	1.03	7.08	6.94	0.13
ART eligibility assessment visits	6.04	1.00	1.00	6.04	6.04	0.00
Sick visits	6.04	0.17	0.10	1.03	0.60	0.42
6-monthly clinic review	6.74	0.89	0.86	6.00	5.80	0.20
Routine follow up visits	5.76	4.43	4.41	25.52	25.40	0.12
CD4 count test	15.72	1.60	1.56	25.15	24.52	0.63
ALT test	0.88	1.20	1.1	1.05	0.97	0.09
Creatinine test	0.31	0.77	0.74	0.24	0.23	0.01
Hb test	0.87	1.33	1.17	1.16	1.02	0.14
VDRL test	1.89	0.07	0.06	0.13	0.11	0.02
Pregnancy test	0.53	0.07	0.08	0.04	0.04	-0.01
Second Xpert test	19.07	0.88	0.00	16.78	0.00	16.78
Chest x-ray	3.18	0.06	0.05	0.17	0.16	0.02
CrAg test	8.43	1.00	0.00	8.43	0.00	8.43
Lay worker visit to the patients home	14.74	2.89	0.00	42.60	0.00	42.60
Collection and processing of cerebral spinal fluid	16.27	0.02	0.00	0.37	0.00	0.37
Days on ART drugs	0.42	323.12	213.60	135.71	131.71	4.00
Days on Co-trimoxazole	0.02	270.22	265.19	5.40	5.30	0.10
Days of hospital admissions	26.43	3.79	4.08	100.13	107.78	-7.66
Number of people put on fluconazole	4.8	0.09	0.00	0.43	0.00	0.43
Mean cost per patient during the first three months and one year in treatment (in 2012US\$)						
First three months in treatment, mean (95% CI)				166 (161–171)	107 (101–112)	59 (52–66)
One year in treatment, mean (95% CI)				331 (319–344)	265 (254–275)	67(50–83)

doi:10.1371/journal.pone.0171917.t003




Fig 2. The cost breakdown of ART services by month for patients initiating treatment with CD4 count <200 cells/mm³ in Tanzania for standard care.

doi:10.1371/journal.pone.0171917.g002

arm had significantly more sick clinic visits ($p = 0.001$), alanine aminotransferase tests ($p = 0.039$), and Hb tests ($p = 0.0013$).

Average costs per patient for routine care

The mean cost per patient year was US\$ 265 (95% CI 254–275) in the standard care arm (Table 3). ART drugs, clinic visits, hospital admission, diagnostic tests and non-ART drugs constituted 50%, 19%, 19%, 10% and 2% of the total cost per patient year respectively (Fig 2). Initial high costs of US\$71 per patient month in the first month fell to US\$ 30 per patient month at 12 months (see Fig 3). The cost per patient was US\$107 (101–112) in the first quarter, after which it fell and stabilised at between \$45 and \$59 per quarter.

The mean annual cost of ART per participant varied from \$34.89 for stavudine-based regimens to \$140.60 for tenofovir-based regimens. Costs of co-trimoxazole prophylaxis were just over \$5 per patient per year. Routine follow up visits were more common than other visits and consequently accounted for a higher cost than other visits. CD4 count and the Xpert testing were the most expensive laboratory tests, with serum CrAg testing costing just under \$4 per test per patient. Hospital admission costs exceed \$43 per admission and comprised a mean of 4.87 days of inpatient care per patient.

Table 3 also compares the costs in the standard and intervention arms. The cost of adding CrAg screening, home visits by lay-workers and a second Xpert test, as implemented in the clinic plus community support arm of the trial, were US\$ 8.43, US\$42.60 and US\$16.78 per patient, respectively (see Table 3). The incremental costs per patient of the full intervention to




Fig 3. Mean cost (in USD 2012) per patient month for patients initiating ART treatment with CD4 count <200 cells/mm³ in Tanzania for both standard care (control) and the REMSTART trial intervention arms.

doi:10.1371/journal.pone.0171917.g003

the health service over the first three months was US\$ 59 ($p < 0.001$; 95%CI 51.5–66.5) and over a one year period was US\$ 67 ($p < 0.001$; 95%CI 50.0–83.2). This is equivalent to a 55% (95%CI 50.7%–59.2%) increase in the mean cost of care over the first three months and a 25% (95%CI 19.7%–30.2%) increase over one year of follow up.

Sensitivity analyses

The cost of care was robust to changes in the variables explored in the sensitivity analysis, except to changes in the average number of home visits made by lay worker in a day, costs of clinic visits involving a CD4 test and the cost of antiretroviral drugs. A 25% decrease in the price of antiretroviral drugs led to a reduction in the mean costs per patient year of 10.6%, and an increase in the number of home visits from 1.4 to 4 per day resulted in a 22% reduction in the additional cost for the intervention component during the first three months, and a 12% over a one year period of follow up, whereas, an increase of costs for clinic visits involving a CD4 test, by 50%, during the first three month of ART led to an increase in the mean costs per patient year of 22.7%. A decrease in this by 50% resulted in a reduction of per patient cost by 11.4%, (Table 4).

Regression

Table 5 reports on the results of the univariate and multivariate regression analyses of factors that could influence health service costs in the standard care arm. None of the factors was significantly associated with health service costs.

Table 4. Results of the sensitivity analyses exploring the impact of uncertainty in key variables on the average cost estimates for patients initiating ART treatment with CD4 count <200 cells/mm³ in Dar es Salaam, Tanzania.

Arm	Parameters varied	Varied values		US\$ (% divergence from base case)				
		Minimum	Maximum	1 st three month		Per Patient Year		
				Minimum	Maximum	Minimum	Maximum	
Standard care	Base case*			106.9	106.9	264.7	264.7	
	Discount rate	1%	12%	106.3 (0.6)	107.3 (0.4)	263.4 (0.5)	265.8 (0.4)	
	Administrative costs	3%	15.5%	104.8 (2.0)	108.4 (1.4)	262.6 (0.8)	266.3 (0.6)	
	Costs for supporting staff	-50%	+50%	104.1 (2.6)	112.5 (5.2)	261.5 (1.2)	271.2 (2.4)	
	Costs for clinic visits involving a CD4 test	-50%	+50%	94.7 (11.4)	131.2 (22.7)	259.9 (1.8)	274.2 (3.6)	
	Costs for clinic visits involving seeing a clinician	-50%	+50%	103.7 (3.0)	113.2 (5.9)	262.3 (0.9)	269.5 (1.8)	
	Costs for routine follow up	-50%	+50%	106.8 (0.1)	124.1 (16.1)	255.7 (3.4)	268.9 (1.6)	
	Costs for Xpert	-5%	-25%	106.0 (0.8)	102.4 (4.2)	263.9 (0.3)	268.1 (1.3)	
	Price for Xpert cartridge	-5%	-25%	106.4 (0.5)	104.2 (2.5)	264.2 (0.2)	262.6 (0.8)	
	Costs for transport for Xpert sputum sample	0%	-50%	106.4 (0.5)	106.6 (0.3)	264.2 (0.2)	264.4 (0.1)	
	Price of ARV	-5%	-25%	104.8 (2.0)	96.3 (9.9)	259.1 (2.1)	236.6 (10.6)	
	Costs for sputum smear microscopy	0%	100%	106.9 (0.0)	105.5** (1.5)	264.7 (0.0)	262.3** (0.9)	
	Intervention	Base case*			165.9	165.9	331.4	331.4
		Price for CRAG reagents	-5%	-25%	165.9 (0.0)	165.9 (0.0)	331.4 (0.0)	331.4 (0.0)
Monthly salary for lay workers		US\$177	US\$383	156.6 (5.6)	165.9 (0.0)	325.1 (1.9)	334.4 (0.9)	
Monthly income (salary & benefits) for lab technician for processing the cerebral spinal fluid		US\$379	US\$397	165.7 (0.1)	166.1 (0.1)	331.4 (0.0)	331.5 (0.0)	
Number of home visits by lay-worker		1	4	174.1 (4.9)	129.2 (22.1)	335.4 (1.2)	290.5 (12.3)	

*Does not include costs for smear tests

**Does not include costs for Xpert test

doi:10.1371/journal.pone.0171917.t004

Discussion

The costs of antiretroviral therapy in Africa have fallen remarkably. Nonetheless, as this study demonstrates, the average cost of care remains high at nearly \$300 per patient per year and over half of this expenditure was on drugs. Using a micro costing approach, this study has calculated the average cost per patient year of ART as well as the incremental cost of an innovative intervention able to reduce mortality in ART patients. The analysis shows how costs fall over time and that ART drugs continue to take up the largest share of ART costs. We did not calculate the costs incurred by patients to access care, but previous studies have found these to be remarkably high [37]. It is evident, that HIV continues to be a substantial burden in Africa.

The costs of the clinic plus community support intervention, which mostly targeted the first month of ART, and which reduced all-cause mortality substantially [17], were about 25%

Table 5. Univariate and multivariate regression (Generalized Linear model) analyses of factors that could influence ART service costs (in 2012 US \$) for patients initiating treatment with CD4 count <200 cells/mm³ in Dar es Salaam, Tanzania.

Independent variable	Results of the Generalized Linear regression analyses (N = 870)			
	Univariate analysis	F-test	Multiple regression [‡]	F-test
	Coefficient (95% CI)	p-value	Coefficient (95% CI)	p-value
Age (reference is "Age less than 45years")		0.5415		0.2702
≥ 45 years	7.71 (-17.03–32.44)		14.67 (-11.38–40.66)	
Level of CD4 count at enrolment (reference is "CD4 <25cells/mm³")		0.3698		0.3984
CD4 25–49cells/mm ³	20.57 (-10.75–51.90)		20.83 (-10.94–52.59)	
CD4 ≥50cells/mm ³	13.64 (-10.37–37.65)		12.17 (-12.20–36.54)	
WHO clinical stage at enrolment (reference is "WHO stage 1 or 2")				0.9154
WHO stage 3 or 4	-4.94 (-32.50–22.62)	0.7254	-1.54 (-29.98–26.90)	
Sex				0.1168
Female	11.70 (-10.08–33.49)	0.2924	18.70 (-4.67–42.06)	
Level of education (reference is "no formal education")		0.4358		0.3954
Primary	19.45 (-10.26–49.16)		20.98 (-9.78–51.73)	
Secondary and above	17.79 (-20.78–56.37)		21.36 (-18.35–61.07)	
Clinic (reference is ("Buguruni clinic"))		0.1036		0.1284
Tandale clinic	18.46 (-10.53–47.45)		22.63 (-7.15–52.42)	
Mbagala clinic	-7.89 (-35.08–19.31)		-1.88 (-29.92–26.17)	
Marital status (reference is "married or living with someone")		0.9545		0.8574
Widowed/separated/divorced	-2.45 (-26.08–21.18)		-5.60 (-29.82–19.70)	
Never married	-4.33 (-34.44–25.78)		-7.91 (-39.33–23.52)	

[‡]All variables included simultaneously in a single model

*Mean difference in health service costs (in US\$) compared to the reference group

doi:10.1371/journal.pone.0171917.t005

higher than the standard treatment arm, when calculated over a 12-month period. The analysis by month and by quarter confirm that the increase in cost is only associated with the initial stage of treatment and as such do not have a major impact on the lifetime costs of ART, providing a strong argument for investment given the likely reduction in mortality that was demonstrated by the trial. Although this study was done as part of a trial, it was integrated into the health system, with patients managed by routine health care staff. Thus the costs presented are likely to be typical of urban clinics in Tanzania among HIV-infected individuals presenting with advanced disease. However, a formal economic evaluation is needed to determine if and to what extent this intervention is cost-effective compared to standard care.

The cost of a single home visit was nearly \$15, just over twice the cost incurred by health services when a patient visits the clinic for routine care and similar to the cost of a CD4 count test. The cost of home visits may be lower in real life settings if lay-workers can visit a greater number patients per day. In our trial, most patients (in both arms) were initiated on ART on their second visit to clinic, within a median of about two weeks [17], because we introduced expedited initiation of antiretroviral therapy, whereas in most African HIV programmes, patients are required to attend 3–4 times before ART is initiated. Our study demonstrates that these early pre initiation visits are probably unnecessary and that it would be better to provide additional support after the patient has started antiretroviral therapy.

Another striking finding was that the care delivered by the clinics differed substantially from guidelines. The mean number of clinic visits involving a clinician was less than 2 visits per patient during the first year on antiretroviral therapy, compared to the recommended 6 visits [20,32,38]; and CD4 count and safety blood testing were done much less frequently

than recommended. It is possible that this was because of shortages of staff and reagents, that patients were not returning for required visits or that clinics felt that these visits were not necessary.

The average cost per patient year of standard care in this study was marginally higher than that reported from Zambia [11] and Nigeria [12], although the Nigeria study excluded overhead costs. In accordance with our findings, in other studies conducted across Africa, antiretroviral drugs have been the most expensive component of care [9–14,16]. We found that reducing the price of antiretroviral drugs by 25% lowers the annual cost per patient by 11%; this highlights the importance of international efforts to negotiate reductions in antiretroviral drugs prices.

Initially, our study enrolled patients with CD4 count <100 cells/mm³, but this criteria was changed subsequently to enrol patients presenting with <200 cells/mm³, in order to speed up recruitment. This change affected both trial arms equally, and presumably was not felt to compromise the interpretation of the study from a clinical standpoint. The univariate and multivariate regression analyses of factors that could influence health service costs, the level of CD4 count was not significantly associated with health service costs, so it is unlikely this change had any effect on the cost estimate.

Our study was conducted among urban patients with advanced disease. Whether the findings are generalizable to rural patients, who are usually more geographically scattered and less mobile, or among patients initiating ART in the early stages of HIV-infection needs further research. We used university graduates as the lay workers and it is not clear if using lay-workers with fewer qualifications would have the same effect. This needs further research as it may be difficult to scale-up the programme using university graduates.

Conclusion

Micro costing approaches provide a way to obtain an in-depth understanding of cost structures and cost variation between individuals as well as over time. This study confirms that despite price reductions, antiretroviral drugs constitute half of the cost of care for people newly starting on ART and confirms that monthly costs of ART decline after the first month of treatment. The added costs of the clinic plus community support intervention, which mostly targets the first month, and which substantially reduced all-cause mortality, represented only a small increase in life time ART costs, suggesting this important innovation in ART could be an affordable approach to improving HIV care in resource constrained settings.

Supporting information

S1 Table. Information on resources use and prices by cost centre.

(DOCX)

S2 Table. Reasons for uncertainty.

(DOCX)

S3 Table. Recourse utilization by the components of ART.

(DOCX)

Acknowledgments

We thank the study participants and staff involved with the REMSTART trials and our funders, EDCTP (European and Developing Countries Clinical Trials Partnership) and HIV Research Trust. We also appreciate the following REMSTART team members for successfully

implementing this trial: Amos Kahwa^{1##}, Saidi Egwaga², Peter Mwaba³, Chanda Duncan³, Victor Chalwe³, Walter Shoo⁴, Paulina Chale⁵, Okeng'o Kigocha⁵, Raymond Phillips Shirima², Yacobo Lema¹, Said Mfaume⁶, Mabula Kasubi⁵, Milka, Mathania⁷, Magreth Makuchilo⁸, Flora Mziray⁹, Clement Kwayu⁷, Iadislau Rwezaura⁷, Joyce Mgohamwenda⁸, John Minde⁷, Theophilla Luhimbo⁹, Maijo Biseko⁴, Emmanuel Kapesa⁴, Ester Diarz⁴, Ramadhani Shemtandulo⁴ and Asheri Barankena⁴.

Affiliation of the REMSTART Trial team members

- ¹National Institute for Medical Research, Muhimbili Medical Research Centre, Tanzania
²National Tuberculosis and Leprosy Program, Tanzania
³Institute for Medical Research and Training, University Teaching Hospital, Lusaka, Zambia
⁴REMSTART Project, Muhimbili Medical Research Centre, National Institute for Medical Research, Dar es Salaam, Tanzania
⁵Department of Internal Medicine, Muhimbili National Hospital, Dar es Salaam, Tanzania
⁶Central Tuberculosis Reference Laboratory, Muhimbili Medical Research Centre, National Institute for Medical Research, Dar es Salaam, Tanzania
⁷Buguruni Health Centre, Ilala District, Dar es Salaam, Tanzania
⁸Mabagala Rangi Tatu. Health Centre, Temeke District, Dar es Salaam, Tanzania
⁹Tandala Health Centre, Kinondoni District, Dar es Salaam, Tanzania
^{##}Lead author for the REMSTART Trial team members
 E-mail: akahwa@hotmail.com

Author Contributions

- Conceptualization:** GDK SM TSH SJ LG.
Data curation: GDK SM VS SK CB TSH SJ LG.
Formal analysis: GDK SM VS SK CB NH SJ LG.
Funding acquisition: SM SJ LG.
Investigation: GDK SM VS SK CB TSH SJ LG.
Methodology: GDK SM TSH SJ LG.
Project administration: SM SJ LG.
Supervision: SM CB NH SJ LG.
Validation: GDK VS CB LG.
Visualization: GDK SM NH SJ LG.
Writing – original draft: GDK.
Writing – review & editing: GDK SM VS SK CB NH TSH SJ LG.

References

1. UNAIDS. Global report on the global AIDS epidemic. 2013.
2. Galárraga O, Wirtz VJ, Figueroa-Lara A, Santa-Ana-Tellez Y, Coulibaly I, Viisainen K, et al. Unit costs for delivery of antiretroviral treatment and prevention of mother-to-child transmission of HIV: a systematic review for low- and middle-income countries. *Pharmacoeconomics*. 2011; 29: 579–99. doi: [10.2165/11586120-000000000-00000](https://doi.org/10.2165/11586120-000000000-00000) PMID: [21671687](https://pubmed.ncbi.nlm.nih.gov/21671687/)

3. Beck EJ, Harling G, Gerbase S, DeLay P. The cost of treatment and care for people living with HIV infection: implications of published studies, 1999–2008. *Curr Opin HIV AIDS*. 2010; 5: 215–24. doi: [10.1097/COH.0b013e31819bc064](https://doi.org/10.1097/COH.0b013e31819bc064) PMID: [20539077](https://pubmed.ncbi.nlm.nih.gov/20539077/)
4. Gray AM, Clarke P.M., Wolstenhole Jane L., Wordsworth S. *Applied Methods of Cost-effectiveness Analysis in Health Care*. Gray AM, Briggs A, editor. New York, United States: Oxford University Press; 2011.
5. UNAIDS. *Workbook for the collection of cost information on HIV facilities and services*. 2011.
6. Frick KD. Microcosting quantity data collection methods. *Med Care*. 2009; 47: S76–81. doi: [10.1097/MLR.0b013e31819bc064](https://doi.org/10.1097/MLR.0b013e31819bc064) PMID: [19536026](https://pubmed.ncbi.nlm.nih.gov/19536026/)
7. Institute for Quality and Efficiency in Health Care. *Working Paper Cost Estimation* [Internet]. 2009. http://www.ispor.org/peguidelines/source/Germany_WorkPaperCostEst.pdf
8. Loubiere S, Sodqi M, Loundou A, Cleary S, Moatti J, Himmich H. When to initiate highly active antiretroviral therapy in low-resource settings: the Moroccan experience. *Antivir Ther*. 2008; 13: 241–251. PMID: [18505175](https://pubmed.ncbi.nlm.nih.gov/18505175/)
9. Marseille E, Giganti MJ, Mwango A, Chisembele-Taylor A, Mulenga L, Over M, et al. Taking ART to scale: determinants of the cost and cost-effectiveness of antiretroviral therapy in 45 clinical sites in Zambia. *PLoS One*. 2012; 7: e51993. doi: [10.1371/journal.pone.0051993](https://doi.org/10.1371/journal.pone.0051993) PMID: [23284843](https://pubmed.ncbi.nlm.nih.gov/23284843/)
10. Larson BA, Bii M, Henly-thomas S, McCoy K, Sawe F, Shaffer D, et al. ART treatment costs and retention in care in Kenya: a cohort study in three rural outpatient clinics. *J Int AIDS Soc*. 2013; 16: 1–5. Available: <http://www.jiasociety.org/index.php/jias/article/view/18026>
11. Scott C a, Iyer HS, McCoy K, Moyo C, Long L, Larson B a, et al. Retention in care, resource utilization, and costs for adults receiving antiretroviral therapy in Zambia: a retrospective cohort study. *BMC Public Health*. *BMC Public Health*; 2014; 14: 296. doi: [10.1186/1471-2458-14-296](https://doi.org/10.1186/1471-2458-14-296) PMID: [24684772](https://pubmed.ncbi.nlm.nih.gov/24684772/)
12. Aliyu HB, Chuku NN, Kola-Jebutu A, Abubakar Z, Torpey K, Chabikuli ON. What is the cost of providing outpatient HIV counseling and testing and antiretroviral therapy services in selected public health facilities in Nigeria? *J Acquir Immune Defic Syndr*. 2012; 61: 221–5. doi: [10.1097/QAI.0b013e3182683b04](https://doi.org/10.1097/QAI.0b013e3182683b04) PMID: [22820805](https://pubmed.ncbi.nlm.nih.gov/22820805/)
13. Cianci F, Sweeney S, Konate I, Nagot N, Low A, Mayaud P, et al. The cost of providing combined prevention and treatment services, including ART, to female sex workers in Burkina Faso. *PLoS One*. 2014; 9: e100107. doi: [10.1371/journal.pone.0100107](https://doi.org/10.1371/journal.pone.0100107) PMID: [24950185](https://pubmed.ncbi.nlm.nih.gov/24950185/)
14. Menzies N a, Berruti A a, Blandford JM. The determinants of HIV treatment costs in resource limited settings. *PLoS One*. 2012; 7: e48726. doi: [10.1371/journal.pone.0048726](https://doi.org/10.1371/journal.pone.0048726) PMID: [23144946](https://pubmed.ncbi.nlm.nih.gov/23144946/)
15. *Frontieres MS*. UNTANGLING THE WEB OF ANTIRETROVIRAL PRICE REDUCTIONS [Internet]. 2013. http://d2pd3b5abq75bb.cloudfront.net/2013/09/11/10/25/44/896/MSF_Access_UTW_16th_Edition_2013.pdf
16. Long L, Brennan A, Fox MP, Ndibongo B, Jaffray I, Sanne I, et al. Treatment outcomes and cost-effectiveness of shifting management of stable ART patients to nurses in South Africa: an observational cohort. *PLoS Med*. 2011; 8: e1001055. doi: [10.1371/journal.pmed.1001055](https://doi.org/10.1371/journal.pmed.1001055) PMID: [21811402](https://pubmed.ncbi.nlm.nih.gov/21811402/)
17. Mfinanga S, Chanda D, Kivuyo SL, Guinness L, Bottomley C, Simms V, et al. Cryptococcal meningitis screening and community-based early adherence support in people with advanced HIV infection starting antiretroviral therapy in Tanzania and Zambia: an open-label, randomised controlled trial. *Lancet*. 2015; 6736: 1–10.
18. National Bureau of Statistics. *The United Republic of Tanzania Population Distribution by Age and Sex* [Internet]. 2013. http://ihi.eprints.org/2169/1/Age_Sex_Distribution.pdf
19. National Bureau of Statistics. *The United Republic of Tanzania. HIV/AIDS and Malaria Indicator Survey 2011–12*. 2013.
20. NACP. *The United Republic of Tanzania. National Guidelines for the Management of HIV and AIDS*. 2012.
21. Lawn SD, Myer L, Orrell C, Bekker L-G, Wood R. Early mortality among adults accessing a community-based antiretroviral service in South Africa: implications for programme design. *AIDS*. 2005; 19: 2141–8. Available: <http://www.ncbi.nlm.nih.gov/pubmed/16284464> PMID: [16284464](https://pubmed.ncbi.nlm.nih.gov/16284464/)
22. Amuron B, Namara G, Birungi J, Nabiryo C, Levin J, Grosskurth H, et al. Mortality and loss-to-follow-up during the pre-treatment period in an antiretroviral therapy programme under normal health service conditions in Uganda. *BMC Public Health*. 2009; 9: 290. doi: [10.1186/1471-2458-9-290](https://doi.org/10.1186/1471-2458-9-290) PMID: [19671185](https://pubmed.ncbi.nlm.nih.gov/19671185/)
23. Rosen S, Long L, Sanne I. The outcomes and outpatient costs of different models of antiretroviral treatment delivery in South Africa. *Trop Med Int Health*. 2008; 13: 1005–15. doi: [10.1111/j.1365-3156.2008.02114.x](https://doi.org/10.1111/j.1365-3156.2008.02114.x) PMID: [18631314](https://pubmed.ncbi.nlm.nih.gov/18631314/)
24. Kranzer K, Govindasamy D, Ford N, Johnston V, Lawn SD. Quantifying and addressing losses along the continuum of care for people living with HIV infection in sub-Saharan Africa: a systematic review. *J*

- Int AIDS Soc. 2012; 15: 17383. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3503237&tool=pmcentrez&rendertype=abstract> doi: 10.7448/IAS.15.2.17383 PMID: 23199799
25. Vassall A, van Kampen S, Sohn H, Michael JS, John KR, den Boon S, et al. Rapid diagnosis of tuberculosis with the Xpert MTB/RIF assay in high burden countries: a cost-effectiveness analysis. *PLoS Med*. 2011; 8: e1001120. doi: 10.1371/journal.pmed.1001120 PMID: 22087078
 26. Bratt JH, Torpey K, Kabaso M, Gondwe Y. Costs of HIV/AIDS outpatient services delivered through Zambian public health facilities. *Trop Med Int Health*. 2011; 16: 110–8. doi: 10.1111/j.1365-3156.2010.02640.x PMID: 20958891
 27. World Bank. GDP deflator [Internet]. <http://data.worldbank.org/indicator/NY.GDP.DEFL.KD.ZG>
 28. Bank of Tanzania. Exchange rate [Internet]. 2012. http://www.bot-tz.org/publications/QuarterlyEconomicBulletins/QEB_Mar_2013.pdf#page=2&zoom=auto,-115,788
 29. Bank of Tanzania. The United Republic of Tanzania. The Treasury bond [Internet]. Dar es Salaam; 2012. <http://www.bot-tz.org/FinancialMarkets/TBonds/TBondCallforTender/2012-JUNE-13-TBOND.pdf>
 30. Siapka M, Remme M, Obure D, Maier CB, Dehne L, Vassall A. Is there scope for cost savings and efficiency gains in HIV services? A systematic review of the evidence from low- and middle-income countries. *WHO Bull*. 2014; 4: 499–511. <http://dx.doi.org/10.2471/BLT.13.127639>
 31. Losina E, Bassett IV, Giddy J, Chetty S, Regan S, Walensky RP, et al. The “ART” of linkage: pre-treatment loss to care after HIV diagnosis at two PEPFAR sites in Durban, South Africa. *PLoS One*. 2010; 5: e9538. doi: 10.1371/journal.pone.0009538 PMID: 20209059
 32. Bassett I V, Regan S, Chetty S, Giddy J, Uhler LM, Holst H, et al. Who starts antiretroviral therapy in Durban, South Africa? . . . not everyone who should. *AIDS*. 2010; 24 Suppl 1: S37–44.
 33. Ingle S, May M, Uebel K, Timmerman V, Kotze E, Sterne JAC, et al. Differences in access and patient outcomes across antiretroviral treatment clinics in the Free State province: prospective cohort study. *S Afr Med J*. 2011; 100: 675–681. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3059232/pdf/nihms247545.pdf>
 34. Ingle S, May M, Uebel K, Kotze E, Bachmann M, Sterne JAC, et al. Outcomes in Patients Waiting for Antiretroviral Treatment in the Free State Province, South Africa: Prospective Linkage Study. *AIDS*. 2011; 24: 2717–2725.
 35. Kirkwood Betty R, and Sterne JA. *Essential Medical statistics*. 2nd Editio. Massachusetts: Blackwell science Inc; 2003.
 36. James C, Bura M, Ensor T. *The Costs of Delivering Health Services in Tanzania: Findings from a comprehensive costing analysis*. 2013.
 37. Jaffar S, Amuron B, Foster S, Birungi J, Levin J, Namara G, et al. Rates of virological failure in patients treated in a home-based versus a facility-based HIV-care model in Jinja, southeast Uganda: a cluster-randomised equivalence trial. *Lancet*. 2009; 374: 2080–9. doi: 10.1016/S0140-6736(09)61674-3 PMID: 19939445
 38. WHO. *The WHO consolidated guidelines on the use of antiretroviral drugs for treating and preventing HIV infection; recommendations for public health approach—Implications for the African region*. 2013.