



Published in final edited form as:

*AIDS Behav.* 2017 May ; 21(5): 1452–1466. doi:10.1007/s10461-016-1510-y.

## Seroadaptive Strategies of Gay & Bisexual Men (GBM) With the Highest Quartile Number of Sexual Partners in Vancouver, Canada

Kiffer G. Card<sup>1,2</sup>, Nathan J Lachowsky<sup>1,3</sup>, Zishan Cui<sup>1</sup>, Paul Sereda<sup>1</sup>, Ashleigh Rich<sup>1</sup>, Jody Jollimore<sup>1,4</sup>, Terry Howard<sup>5</sup>, Robert Birch<sup>6</sup>, Allison Carter<sup>1,2</sup>, Julio Montaner<sup>1</sup>, David Moore<sup>1,3</sup>, Robert S. Hogg<sup>1,2</sup>, and Eric Abella Roth<sup>6,7</sup>

<sup>1</sup>BC Centre for Excellence in HIV/AIDS, Vancouver, British Columbia, Canada

<sup>2</sup>Faculty of Health Science, Simon Fraser University, Vancouver, British Columbia, Canada

<sup>3</sup>Faculty of Medicine, University of British Columbia, Vancouver, British Columbia, Canada

<sup>4</sup>Health Initiative for Men, Vancouver, British Columbia, Canada

<sup>5</sup>Positive Living Society of BC, Vancouver, British Columbia, Canada

<sup>6</sup>Department of Anthropology, University of Victoria, Victoria, British Columbia, Canada

<sup>7</sup>Centre for Addiction Research of British Columbia, University of Victoria, Victoria, British Columbia, Canada

### Abstract

Despite continued research among men with more sexual partners, little information exists on their seroadaptive behavior. Therefore, we examined seroadaptive anal sex strategies among 719 Vancouver gay and bisexual men (GBM) recruited using respondent driven sampling (RDS). Our

---

**Corresponding Author:** Nathan J. Lachowsky, 505-1200 Burrard Street, Vancouver, British Columbia, V6Z 2C7, Phone: 604-558-2017, nlachowsky@cfenet.ubc.ca.

#### Compliance with Ethical Standards

##### Ethical Approval:

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional research committees at Simon Fraser University, The University of British Columbia, and the University of Victoria and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent was obtained from all individual participants included in the study.

This article does not contain any studies with animals performed by any of the authors.

##### Conflict of Interest:

Kiffer G. Card declares that he has no conflict of interest.

Nathan J. Lachowsky declares that he has no conflict of interest.

Zishan Cui declares that she has no conflict of interest.

Paul Sereda declares that he has no conflict of interest.

Ashleigh Rich declares that she has no conflict of interest.

Jody Jollimore declares that he has no conflict of interest.

Terry Howard declares that he has no conflict of interest.

Robert Birch declares that he has no conflict of interest.

Allison Carter declares that she has no conflict of interest.

Julio Montaner declares that he has no conflict of interest.

David Moore declares that he has no conflict of interest.

Robert S. Hogg declares that he has no conflict of interest.

Eric A. Roth declares that he has no conflict of interest.

objectives were to (1) describe the distribution in frequency of male sexual partnering among Vancouver GBM, and (2) identify important covariates associated with the number of male sexual partners. To this aim, we provide descriptive, univariate, and multivariate adjusted statistics, stratified by HIV status, for the association between having  $\geq 7$  male anal sex partners in the past six months (Population Q3, versus  $<7$ ). Sensitivity Analysis were also performed to assess the robustness of this cut-off point. Results suggest that GBM with more sexual partners are more likely to employ seroadaptive strategies than men with fewer partners. These strategies may be used in hopes of offsetting risk, assessing needs for subsequent HIV testing, and balancing personal health with sexual intimacy. Further research is needed to determine the efficacy of these strategies, assess how GBM perceive their efficacy, and understand the social and health impacts of their widespread uptake.

### Keywords

Strength-based; Seroadaptive; HIV; Gay and Bisexual Men; Highly Sexually Active Men

---

## INTRODUCTION

Recent global reviews of HIV literature have demonstrated continued disparities in the sexual health and wellbeing of gay and bisexual men (GBM) (1,2). While many structural and social factors have been explored to explain the disproportionate burden of HIV/STI rates this population faces (3), researchers have often pointed to differences in the sexual partnering patterns of GBM compared with other men and women (4,5). Specifically, studies have shown that, on average, GBM are less likely to discriminate against sexual partners based on age, have greater numbers of lifetime and annual sexual partners, continue to form new partnerships later in life, and report higher prevalence of partner concurrency (6). While these characteristics may contribute to the broader social and cultural identity of GBM, the density and interconnectedness observed in these sexual networks may also play a role in HIV transmission (7,8). That is, men with greater numbers of male anal sex partners may be at greater risk for HIV transmission than men with fewer sexual partners (9–11).

Despite the continued attention that men with more sex partners receive from the academic and public health communities, there is little research on the seroadaptive behavioral strategies of this group. Therefore, GBM and their sexual practices continue to face stigma from what has colloquially been described as “slut-shaming” (12). Such attitudes towards these men and their relationship patterns inhibit constructive communication about sexual risk (13). In contrast, sex-positive community engagement approaches could promote risk reduction strategies which help men with greater numbers of male anal sex partners avoid seroconversion or transmission, while being supportive of the diversity of values and needs found within the GBM communities. Such approaches have focused on reducing the risks associated with anal sex, as it is the most common route of transmission for HIV among GBM (14). While some of these risk reduction strategies are empirically supported (e.g., condom use), other perceived harm reduction strategies may not be fully effective in preventing the transmission of HIV (e.g., withdrawal, negotiated safety agreements) (15). Seroadaptive behaviors, defined as “any modification of sexual behavior based on the

person's (perceived) serostatus, the (perceived) status of the partner and/or HIV transmission risk by type of sex act," (16), can be broadly divided into three categories: First, those seeking seroconcordant partnerships (i.e., serosorting, where both partners have the same HIV serostatus); Second, those aiming to reduce the probability of HIV transmission (e.g., anal sex abstinence, strategic positioning); and Third, those attempting to limit exposure to the virus (e.g., condom use, treatment/viral load sorting) (16). These seroadaptive practices are widely practiced among GBM (17,18) and have been cited as evidence of GBM adaptive resilience in preventing HIV seroconversion (19).

In Vancouver, expanded access to highly active antiretroviral therapy (HAART), and associated individual- and population-level viral load reduction (referred to as "Treatment as Prevention (TasP)") (20,21), may serve an important role in changing population patterns of these behaviors and practices (18). This closely relates to the theory of risk homeostasis which suggests decreased perceptions of risk corresponding to increased risk-taking behavior (22). This has raised concerns that the benefits associated with biological HIV prevention technologies may be offset by reduced condom use (23). If such is the case, the reinforcement of seroadaptive behaviors along with condom use will serve an important role in helping men with greater numbers of male anal sex partners to manage their sexual health.

When combined with condom use, serosorting and strategic positioning were estimated in one prospective cohort study to reduce HIV transmission among GBM by as much as 98% (24). However, more conservative estimates suggest that these strategies result in a 38% to 83% reduction of risk, depending on the strategy being employed (15). Of course, seroadaptive effectiveness requires low prevalence of undiagnosed STI/HIV infection coupled with high levels of partner disclosure and accurate HIV testing (17,25). Further, the effectiveness of seroadaptive strategies relies on an accurate understanding of the protective benefits – or lack thereof – associated with each strategy. For example, while withdrawal may be thought by some men to prevent transmission of HIV, this method is actually associated with a fivefold increase in seroconversions compared with no condomless anal sex (26) and is not considered an effective risk reduction strategy for those engaging in condomless anal sex because HIV is transmissible through pre-ejaculate (27,28).

Consistent with these observations and our understanding of GBM adaptation and resilience, we hypothesized that men with the highest quartile number of male anal sex partners were more likely to utilize seroadaptive behaviors to ameliorate the risks associated with having more sexual partners. We aimed to develop findings that would help shape sex-positive HIV-prevention messaging that will simultaneously reduce the stigma these men face and better encourage the use of more effective risk reduction strategies. To accomplish these goals, we examined reported seroadaptive strategies and individual characteristics relating to HIV-negative and HIV-positive men with the highest quartile number of male anal sex partners in Vancouver, British Columbia, Canada.

## METHODS

### Study Setting

The Momentum Health Study is a prospective biobehavioral cohort study investigating possible Treatment Optimism (29) and risk compensation associated with British Columbia's expanded Treatment as Prevention program (21). All data used in this analysis are drawn from the baseline study visit, which occurred between February 2012 and February 2014. Momentum uses respondent-driven sampling (30) to recruit HIV-positive and HIV-negative GBM in Metro Vancouver, British Columbia, Canada. Initially-recruited "seeds" distributed a maximum of 6 paper and/or electronic vouchers to other Vancouver GBM (31). Voucher recipients were screened for study eligibility criteria, which included being 16 years of age and older, identifying as male, having had sex with another man in the past six months, living in Metro Vancouver, and competency to understand a questionnaire written in English. Eligible participants completed a computer-assisted self-interview (CASI) questionnaire and biological tests including point-of-care HIV testing, venipuncture blood tests for hepatitis C and syphilis serology, and optional urine and swab tests for gonorrhoea and chlamydia. Study participants received a \$50 CAD honorarium and earned an additional \$10 CAD for each eligible recruit who completed the study protocol. All procedures received human ethics clearances from Simon Fraser University, the University of British Columbia, and the University of Victoria. Additional information about the Momentum study questionnaire and protocol can be found elsewhere (32,33).

### Dependent Variables

The study's dependent variable was a dichotomous categorical variable distinguishing men with high numbers of anal sex partners from other men in the sample. While other sources in the literature have described this group as "Highly Sexual Active Men" (9), we have attempted to describe sexual partnering and concurrency with greater sensitivity to the stigma that this group faces, as well as to more accurately describe their partnering pattern compared with frequency of sexual activity. To identify men with higher numbers of sexual partners, we asked this question: "During the past 6 months, with how many males have you had anal sex with (as top or bottom)?" From the resulting distribution, with the total number of anal sex partners in the past six months capped at 100, we used the global third quartile value to divide our sample first into two groups: 1) men with  $\geq 7$  anal sex partners ( $\geq Q3$ ,  $n=195$ ), and 2) men with  $<7$  recent anal sex partners ( $<Q3$ ,  $n=523$ ). These results are more conservative than those used by other researchers (9,11), which focused on men having 9 or more anal sex partners in the previous three months. To assess the use of our  $\geq Q3$  versus  $<Q3$  cut-off point, a sensitivity analysis was conducted at the univariate level with cut-off points at the global median ( $\geq 3$  versus  $<3$ ) as well as HIV status specific third quartile cut-offs (HIV-positive:  $\geq 15$  versus  $<15$ , HIV-negative:  $\geq 6$  versus  $<6$ ).

### Independent Variables

Seroadaptive strategies were represented in the study questionnaire (32) by a series of yes/no responses with necessary variation in the question wording to be applicable to HIV-negative versus HIV-positive participants. The question block was introduced by saying, "Some guys use strategies to prevent getting/transmitting HIV. Do you do any of the following to prevent

getting/transmitting HIV? (check ALL that apply).” Three strategies were asked the same for HIV-negative and HIV-positive men: condom use (“Always using condoms for anal sex”), anal sex avoidance (“Having sex which doesn’t include anal sex”), and serostatus inquiry (“Asking my sex partners about their HIV status before sex”). Four strategies were asked in a manner specific to participants’ serostatus: HIV-negative men were asked if they used strategic positioning (“Being a top for anal sex”), serosorting (“Having anal sex without condoms only with guys I know are HIV-negative”), viral load sorting (“Having anal sex without condoms with HIV-positive guys who have low viral loads or are on HIV treatment”), and withdrawal (“not letting my sex partners cum inside me”). Likewise, HIV-positive men were asked about strategic positioning (“Being the bottom for anal sex”), serosorting (“Having anal sex without condoms only with guys I know are HIV-positive”), viral load sorting (“Having anal sex without condoms if my viral load is low or I’m on HIV treatment”), and withdrawal (“Not cumming inside my sex partners”).

Additional independent variables pertained to sociodemographic factors, substance use, psychosocial traits, and sexual behavior in order to gain insights into the characteristics of men with the highest quartile number of male anal sex partners and to assess possible confounding variables. Socio-demographic variables included measures of age, education, annual income, race/ethnicity, residence, sexual orientation, and anal sex role preference (i.e., top, bottom or versatile). Substance use questions pertaining to alcohol use classified participants as “harmful drinkers” via the AUDIT Scale (34) and whether they used erectile dysfunction drugs (EDD), poppers, crystal methamphetamine, and/or Ecstasy in the past six months. Previously reported associations between EDD and crystal methamphetamine (35,36), led us to test the inclusion of an interaction term ( $EDD \times$  crystal methamphetamine).

Psychosocial measures included three validated scales. The first was the Escape Motivation Scale (12 questions, study Cronbach’s  $\alpha = 0.90$ ) (37), assessing if GBM used alcohol and illicit substances to diminish cognitive recognition of sexual risk, (e.g., “When I am high, I find it difficult to stay within my sexual limits”). Each item of the Escape Motivation Scale was scored on a 4-point Likert scale (Strongly disagree, Disagree, Agree, Strongly Agree), meaning the total score possible was 12–48 points with higher scores indicating greater escape motivations. The second was the Sexual Sensation Seeking Scale (revised, 11 questions, study  $\alpha = 0.73$ ) (38), which measured respondents’ attitudes towards sexual thrill-seeking (e.g., “I like wild, ‘uninhibited’ sexual encounters”). Each item on the Sexual Sensation Seeking Scale was scored on a 4-point Likert Scale (Not at all like me, Not like me, Like me, Very much like me.), meaning the total score possible was 11–44 points, with higher scores indicates greater sensation seeking. The last scale used was the Treatment Optimism Scale (12 questions, study  $\alpha = 0.85$ ) (39), examining possible changing sexual risk perceptions associated with (e.g., “HIV/AIDS is a less serious threat than it used to be because of new treatments”). Each item on the treatment optimism scale was scored on a 4-point Likert scale (Strongly disagree, Disagree, Agree, Strongly Agree), meaning the total score possible was 12–48 points with higher scores indicating greater treatment optimism.

## Analysis

All analyses were conducted using SAS version 9.3 (40) and stratified by HIV serostatus. The data for this analysis were adjusted by the respondent driven sampling program (RDSAT) version 7.1.46 to generate point estimates and 95% confidence intervals (41). Independent variables with probability values  $< 0.20$  were selected from initial univariable models for inclusion in subsequent multivariable models. Final multivariable models were determined using backward selection elimination procedure based on the optimisation (minimization) of two criteria at each step: Akaike Information Criterion (AIC) (42) and Type-III p-values (43). These two criteria balance model selection between finding an explanatory model with lower p-values (indicating greater significance) and lower AIC values (indicating goodness-of-fit). This model building procedure is described in greater detail elsewhere (44). A central premise of RDS is that respondents' social network size can be used to estimate sampling probabilities and generate population estimates (45).

## RESULTS

Between February 2012 and February 2014, we recruited a total of 719 GBM from 119 (16.6%) seeds or initial recruits. Table 1 provides descriptive socio-demographic and behavioral statistics of the cross-sectional study population, which was predominantly composed of white (68.0%), gay (80.3%) men with a median age of 33 years (Q1–Q3: 26–47). Approximately three-quarters (76.7%) of the sample were HIV-negative/unknown. Behavioral reports indicated that 65.1% had participated in insertive anal sex in the past six months, and 63.5% participated in receptive anal sex. One in three men reported a preference for receptive anal sex (35.9%) and 27.6% reported a versatile preference. The distribution of the number of sexual partners is shown in Figure 1. The median number of anal sex partners in the past 6 months reported by respondents was 3 (Q1–Q3: 1–7). Popper use in the past six months was reported by 34.3% of men. Other substance use reports included crystal methamphetamine (19.6%), EDD (17.3%), and Ecstasy (18.9%).

Descriptive statistics and univariable results for HIV-negative/unknown GBM are provided in Table 2. Among HIV-negative/unknown men with the highest quartile of male anal sex partners, 70.6% reported asking HIV status before sex (versus 56.3% for lower three quartiles), 62.8% reported always using condoms (versus 67.2%), 36.5% reported strategic positioning (versus 18.7%), 32.3% reported serosorting (versus 30.2%), 27.2% reported withdrawal (versus 22.2%), 25.2% reported abstaining from anal sex (versus 47.8%), and 17.6% reported treatment/viral load sorting (versus 5.4%). In univariable analysis, HIV-negative men with the highest quartile number of male anal sex partners were more likely to report these strategies: asking partners their status (OR=1.86, 95% CI [1.25, 2.87]), strategic positioning (OR= 2.55, 95% CI [1.61, 3.90]), and treatment/viral load sorting (OR=3.78, 95% CI [2.17, 7.69]), but were less likely to report abstention from anal sex (OR=0.37, 95% CI [0.23, 0.58]) compared with those in the lower three quartiles of male anal sex partners (*See Table 2*).

Table 3 provides the descriptive statistics and univariable results for HIV-positive GBM. Among HIV-positive men with the highest quartile number of male anal sex partners, 68.8% reported serosorting (versus 35.7% for the lower three quartiles), 49.9% reported asking

HIV status before sex (versus 31.1%), 48.8% reported treatment/viral load sorting (versus 22.2%), 47.6% reported strategic positioning (versus 24.4%), 38.7% reported avoiding anal sex (versus 38.0%), 34.2% reported withdrawal (versus 25.1%), and 26.6% reported always using condoms (versus 37.6%). In univariate analysis, HIV-positive men with the highest quartile number of male anal sex partners had an increased odds of reporting strategies such as asking partners their HIV status (OR=2.29, 95% CI [1.13,4.32]), strategic positioning (OR=2.82, 95% CI [1.41,5.62]), serosorting (OR=3.97, 95% CI [1.98,7.96]), and treatment/viral load sorting (OR=3.33, 95% CI [1.66,6.71]) compared with those in the lower three quartiles of male anal sex partners (See Table 3).

Multivariable models for HIV-negative/unknown GBM are presented in Table 4. Among HIV-negative/unknown men, being in the highest quartile of male anal sex partners was associated with being more likely to engage in strategic positioning (AOR=3.81, 95% CI [1.79,8.11]), ask HIV status prior to sex (AOR=2.15, 95% CI [1.18,3.93]), prefer a versatile versus bottom anal sex position (AOR=2.75, 95% CI [1.31,5.75]), identify as gay compared with bisexual or other (AOR=3.88, 95% CI [1.52,9.90]), live in downtown Vancouver compared with the metro Vancouver area (AOR=3.13, 95% CI [1.47,6.67]), achieve education greater than the high school level (AOR=3.13, 95% CI [1.20,8.13]), use ecstasy in the past six months (AOR=2.96, 95% CI [1.49, 5.89]), engage in “watersports” (46) (AOR=2.78, 95% CI [1.23,6.28]), attend a group sex event in the past six months (AOR=6.07, 95% CI [3.08,11.98]), and receive money in exchange for sex in the past six months (AOR=5.35, 95% CI [1.79,15.98]) compared with men in the lower three quartiles of male anal sex partners; they were also less likely to avoid anal sex (AOR=0.22, 95% CI [1.11,0.42]) (See Table 4).

Multivariable models for HIV-positive GBM are presented in Table 5. These results showed that the majority of seroadaptive strategies were not selected into the model. Being in the highest quartile of male anal sex partners for HIV-positive men was associated with a greater likelihood of asking about HIV status prior to sex (AOR=3.10, 95% CI [1.17,8.26]), having higher scores on the Sexual Sensation Seeking scale (AOR=1.17, 95% CI [1.05,1.31], per 1 unit increase), having used crystal methamphetamine in the past six months (AOR=3.06, 95% CI [1.16,8.10]), having attended a group sex event in the past six months (AOR=3.41, 95% CI [1.21,9.66]), having received money in exchange for sex in the past six months (AOR=5.07, 95% CI [1.09,23.61]), and having less preference for a versatile versus bottom anal sex position (AOR=0.16, 95% CI [0.05,0.53]) compared with men in the lower three quartiles of male anal sex partners (See Table 5).

The sensitivity analysis, which was conducted to assess how differing dichotomous cut-off points at the global median ( $\geq 3$  versus  $<3$ ) and the HIV status specific third quartile cut-offs (HIV-Positive:  $\geq 15$  versus  $<15$ , HIV-Negative:  $\geq 6$  versus  $<6$ ), showed that several variables acted in a dose response relationship to having more sexual partners. However, the overall results of the final model did not appear to be highly sensitive to changes in the cut-off point employed.

## DISCUSSION

We observed a higher prevalence of seroadaptive behaviors among GBM with more sexual partners, using a global data-driven definition of the highest quartile number of male anal sex partners (i.e., more than 7 in the past six months). Although both HIV-positive and HIV-negative men with 7 partners were more likely to engage in condomless anal sex on the univariable level, this factor was not retained in either of the multivariable models. However, in both groups, men in with the highest quartile number of male anal sex partners were more likely to employ other seroadaptive behaviors and prevention practices. HIV-negative men were significantly more likely to engage in strategic positioning, and both groups were more likely to engage in serostatus inquiry, which is a necessary precursor to effective use of other seroadaptive strategies. Contrary to this trend we observed that HIV-negative men were less likely to avoid anal sex as a means of seroadaptation.

We also note that, in general, HIV-positive men with the highest quartile number of male anal sex partners reported higher prevalence of seroadaptive behaviors than HIV-negative men with the highest quartile number of male anal sex partners. This latter finding may reflect a high level of either concern (e.g., HIV prevention altruism) and/or awareness (e.g., HIV risk perception) among HIV-positive men regarding HIV transmission (47,48). This supports other findings suggesting that HIV-positive men aware of their HIV-infection are less likely to infect sexual partners than those who are not aware (48,49).

Together these findings highlight seroadaptive behaviors as common harm reduction strategies for GBM in the highest quartile number of male anal sex partners, which parallels other documented harm reduction tactics in other populations. For example, in their analysis of injection drug users who remain HIV and HCV infection-free, Friedman and colleagues (50,51) adapted the concept of Positive Deviance, originating in infant/child nutrition studies to identify parental strategies associated with well-nourished children living in communities with high malnutrition rates (52) to explain how “subjects control their personal risk, even though they have engaged in high-risk activities for lengthy periods”. Seroadaptive strategies may therefore provide insights into how GBM with the highest quartile number of male anal sex partners manage their sexual health and prevent seroconversion despite having higher than average rates of partnering.

This interpretation of our findings raises the question of the effectiveness of various seroadaptive behaviors. For example, Vallabhaneni and colleagues (15) studied the risks associated with condomless anal sex, serosorting, strategic positioning, and partner concurrency and found that while condomless seroadaptive behaviors were more risky than sex with condoms, these behaviors did seem to offer some benefits compared with condomless sex without seroadaptation. Other research by Kurtz and colleagues (19) doubted that sero-sorting, while widely practiced, could be effective in major cities because: 1) HIV-negative men in their study averaged 10 partners and almost 20 episodes of condomless anal intercourse during a three-month period, 2) 45% of HIV-positive GBM were unaware of their status and, 3) 31% of HIV-negative GBM seroconverted within 5 years of moving to South Florida. In contrast, only 0.6% of Momentum Health Study participants did not know they were HIV-positive, 83% of HIV-negative men had an HIV



test within at least two years, and HIV incidence in our study was at least 1 per 100 person-years within 1.5 years of follow-up (53,54). Of course, some participants lost to follow-up may also have seroconverted, and we note that the number of new diagnoses in British Columbia remains stable (3). Under these conditions, some seroadaptive behaviors may have the potential to reduce HIV transmission risks, though further research is needed to establish the efficacy of each behavior with regards to preventing HIV in different epidemic settings. To accomplish this, we recommend a more purposive sexual health needs assessment to determine which seroadaptive behaviors or which combination of these behaviors might effectively reduce HIV-transmission and how condom use might be better utilized with consideration of these practices.

Finally, our results offer further insight into the behavioral strategies of men with the highest quartile number of male anal sex partners and may therefore provide public health leaders the information necessary to identify different overlapping sub-groups of men (e.g., those who participate in the sex industry as escorts, those who attend group sex events, those who engage in condomless anal sex, and those who use “sex drugs”) who might most benefit from more targeted risk reduction and HIV prevention programming.

Also important, our multivariable results indicated significant associations with substance use (i.e., crystal methamphetamine, EDD, and Ecstasy) for men with the highest quartile number of male anal sex partners. Yet despite this association, the Escape Motivation Scale – which has been used to link substance use and sexual risk (37) – was non-significant in the multivariable model for either HIV-negative or HIV-positive men, suggesting that substance use among these men is not explained by escape behavior but is likely attributable to other motivators, such as pleasure seeking. This association between sexual risk and substance use, especially EDD, is not entirely surprising considering the role sex-drugs play in some venues and cultures (35).

Further, the HAART Optimism scale did not appear to be significant in either the univariable or multivariable results, suggesting that HAART treatment optimism and risk compensation do not explicitly determine patterns of sexual partnering, and, conversely, having more sexual partners does not seem to greatly influence treatment optimism. Considering the risks associated with more densely connected social-sexual networks, this finding speaks positively to the expansion of HAART access. In contrast, the Sexual Sensation Seeking scale, which is a commonly used measure for pleasure seeking and sexual risk (38), was only significant for HIV-positive men with the highest quartile number of male anal sex partners. However, because items on this scale relate to the number of sex partners, it is difficult to assess whether higher scores among this group are attributable to increased pleasure-seeking or are simply a result of the natural correlation between scale items and our primary outcome.

## STRENGTHS & LIMITATIONS

Our study offered several important strengths with regard to past research focusing on sexual partnering and seroadaptation. This analysis included the use of RDS methodology to recruit a large number of gay and bisexual men not specifically from sexual health clinics or

community venues, and re-weighting their responses to provide more accurate population estimates of seroadaptation and sexual partnering (30). Further, by considering demographic, psychosocial, sexual behavior, and substance use together, our explanatory model provides useful analysis for understanding the diverse covariates associated with having higher numbers of male anal sex partners.

There are also limitations to this study. First, the strategies analyzed were self-reports, not actual events. Therefore, we lack a measurable association between what participants say and what they actually do (55). For example, even event-level sexual histories may not include instances when GBM decide not to have sex, perhaps because of serodiscordance. Similarly, anal sexual positioning may simply represent preferred sex roles, rather than seroadaptation. To determine seroadaptation intentionality requires a longitudinal study design (17). In addition, while data were corrected for RDS-bias, they pertain only to Metro Vancouver, and may not be representative of areas where HAART access/uptake and Treatment as Prevention programs are not as well developed. This is particularly relevant when reviewing the high reported prevalence of viral load sorting, which may not be found in areas with decreased HAART availability. In consideration of the psychosocial scales and other categorical measures used in this analysis it is difficult to assess whether differences between groups are clinically meaningful, despite being statistically significant. However, as this is an exploratory analysis, we feel that the direction of the association may serve as an important indicator for understanding the factors associated with having more sexual partners. Finally, speaking to our dichotomous comparison groups, the use of quartiles, while potentially arbitrary and difficult to apply clinically, better describes the natural distribution of sexual partnering frequency, offers simpler interpretations of effect measures resulting from statistical modeling, and allows us to avoid the linearity assumption implicit in modeling continuous covariates (56). Further, the use of data-driven cut-off points serve to provide a descriptive measure of sexual partner frequency.

## CONCLUSION

Our findings show that men with the highest quartile number of male anal sex partners are more likely to employ seroadaptive strategies (other than condom use and anal sex avoidance) than men with fewer sexual partners. This suggests that these men are taking steps that are at least perceived to improve their sexual health and reduce HIV transmission while balancing their needs for sexual intimacy and pleasure. Consistent with these findings we recommend further research to: (1) determine the efficacy of these strategies, (2) assess how GBM perceive the efficacy of these strategies, and (3) to determine the social and health impacts of widespread uptake of these strategies. Each of these suggested areas remain important because, even if these strategies were sufficiently effective to prevent seroconversion, they are by no means universally employed, and confusion regarding their efficacy is almost certain to continue. Further, though these strategies are more common among those with more sexual partners, they may not be used, nor understood, in sufficiently high rates to reduce population level HIV incidence. Yet still, we do not know what widespread uptake of these strategies might mean for condom use, which is still regarded as the most efficacious prevention strategy. Finally, we note that these strategies are

not necessarily effective in combating other STIs, which may put GBM at increased risk for HIV transmission or acquisition.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

The authors would like to thank the Momentum Study participants, office staff, and community advisory board, as well as our community partner agencies, Health Initiative for Men, YouthCoHIV and Hep C Society, and Positive Living Society of BC. Momentum is funded through the National Institute on Drug Abuse (R01DA031055-01A1) and the Canadian Institutes for Health Research (MOP-107544). NJL is supported by a CANFAR/CTN Postdoctoral Fellowship Award. DMM is supported by a Scholar Award from the Michael Smith Foundation for Health Research (#5209).


## REFERENCES

1. Beyrer C, Baral SD, van Griensven F, Goodreau SM, Chariyalertsak S, Wirtz AL, et al. Global epidemiology of HIV infection in men who have sex with men. *Lancet*. 2012 Jul 28; 380(9839): 367–377. [PubMed: 22819660]
2. Halkitis PN, Moeller RW, Siconolfi DE, Storholm ED, Solomon TM, Bub KL. Measurement model exploring a syndemic in emerging adult gay and bisexual men. *AIDS Behav*. 2013 Feb; 17(2):662–673. [PubMed: 22843250]
3. BC PHO. HIV, Stigma and Society: Tackling a Complex Epidemic and Renewing HIV Prevention for Gay and Bisexual Men in British Columbia [Internet]. Provincial Health Officer's 2010 Annual Report. 2014. Available from: <http://www2.gov.bc.ca/assets/gov/health/about-bc-s-health-care-system/office-of-the-provincial-health-officer/reports-publications/annual-reports/hiv-stigma-and-society.pdf>
4. Aral SO, Patel DA, Holmes KK, Foxman B. Temporal trends in sexual behaviors and sexually transmitted disease history among 18- to 39-year-old Seattle, Washington, residents: results of random digit-dial surveys. *Sex Transm Dis*. 2005 Nov; 32(11):710–717. [PubMed: 16254547]
5. Blair J. A Probability Sample of Gay Urban Males: The Use of Two-Phase Adaptive Sampling. *J Sex Res*. 1999 Feb 1; 36(1):39–44.
6. Glick SN, Morris M, Foxman B, Aral SO, Manhart LE, Holmes KK, et al. A comparison of sexual behavior patterns among men who have sex with men and heterosexual men and women. *J Acquir Immune Defic Syndr*. 2012 May 1; 60(1):83–90.
7. Koblin BA, Husnik MJ, Colfax G, Huang Y, Madison M, Mayer K, et al. Risk factors for HIV infection among men who have sex with men. *AIDS Lond Engl*. 2006 Mar 21; 20(5):731–739.
8. Morris M, Kretzschmar M. Concurrent partnerships and transmission dynamics in networks. *Soc Netw*. 1995 Jul; 17(3–4):299–318.
9. Grov C, Golub SA, Parsons JT. HIV status differences in venues where highly sexually active gay and bisexual men meet sex partners: results from a pilot study. *AIDS Educ Prev Off Publ Int Soc AIDS Educ*. 2010 Dec; 22(6):496–508.
10. Grov C, Whitfield THF, Rendina HJ, Ventuneac A, Parsons JT. Willingness to Take PrEP and Potential for Risk Compensation Among Highly Sexually Active Gay and Bisexual Men. *AIDS Behav*. 2015 Mar 4.
11. Parsons JT, Rendina HJ, Moody RL, Ventuneac A, Grov C. Syndemic Production and Sexual Compulsivity/Hypersexuality in Highly Sexually Active Gay and Bisexual Men: Further Evidence for a Three Group Conceptualization. *Arch Sex Behav*. 2015 Oct; 44(7):1903–1913. [PubMed: 26081246]
12. On Slut-shaming and Gay Men [Internet]. *Daily Kos*. [cited 2016 Jan 22] Available from: <http://www.dailykos.com/story/2012/2/11/1063712/-On-Slut-shaming-and-Gay-Men>.

13. McDavitt B, Mutchler MG. “Dude, You’re Such a Slut!” Barriers and Facilitators of Sexual Communication Among Young Gay Men and Their Best Friends. *J Adolesc Res.* 2014 Jul; 29(4): 464–498. [PubMed: 25419044]
14. Baggaley RF, White RG, Boily M-C. HIV transmission risk through anal intercourse: systematic review, meta-analysis and implications for HIV prevention. *Int J Epidemiol.* 2010 Aug; 39(4): 1048–1063. [PubMed: 20406794]
15. Vallabhaneni S, Li X, Vittinghoff E, Donnell D, Pilcher CD, Buchbinder SP. Seroadaptive practices: association with HIV acquisition among HIV-negative men who have sex with men. *PLoS One.* 2012; 7(10):e45718. [PubMed: 23056215]
16. Rönn M, White PJ, Hughes G, Ward H. Developing a Conceptual Framework of Seroadaptive Behaviors in HIV-Diagnosed Men Who Have Sex With Men. *J Infect Dis.* 2014 Dec 1; 210(Suppl 2):S586–S593. [PubMed: 25381379]
17. McFarland W, Chen Y-H, Nguyen B, Grasso M, Levine D, Stall R, et al. Behavior, intention or chance? A longitudinal study of HIV seroadaptive behaviors, abstinence and condom use. *AIDS Behav.* 2012 Jan; 16(1):121–131. [PubMed: 21644001]
18. Snowden JM, Wei C, McFarland W, Raymond HF. Prevalence, correlates and trends in seroadaptive behaviours among men who have sex with men from serial cross-sectional surveillance in San Francisco, 2004–2011. *Sex Transm Infect.* 2014 Sep; 90(6):498–504. [PubMed: 24687128]
19. Kurtz SP, Buttram ME, Surratt HL, Stall RD. Resilience, syndemic factors, and serosorting behaviors among HIV-positive and HIV-negative substance-using MSM. *AIDS Educ Prev Off Publ Int Soc AIDS Educ.* 2012 Jun; 24(3):193–205.
20. Cohen MS, Chen YQ, McCauley M, Gamble T, Hosseinipour MC, Kumarasamy N, et al. Prevention of HIV-1 Infection with Early Antiretroviral Therapy. *N Engl J Med.* 2011 Aug 11; 365(6):493–505. [PubMed: 21767103]
21. Montaner JSG, Lima VD, Harrigan PR, Lourenço L, Yip B, Nosyk B, et al. Expansion of HAART Coverage Is Associated with Sustained Decreases in HIV/AIDS Morbidity, Mortality and HIV Transmission: The “HIV Treatment as Prevention” Experience in a Canadian Setting. *PLoS ONE.* 2014 Feb 12; 9(2):e87872. [PubMed: 24533061]
22. Eaton LA, Kalichman SC. Risk Compensation in HIV Prevention: Implications for Vaccines, Microbicides, and Other Biomedical HIV Prevention Technologies. *Curr HIV/AIDS Rep.* 2007 Dec; 4(4):165–172. [PubMed: 18366947]
23. Blower SM, Gershengorn HB, Grant RM. A tale of two futures: HIV and antiretroviral therapy in San Francisco. *Science.* 2000 Jan 28; 287(5453):650–654. [PubMed: 10649998]
24. McConnell JJ, Bragg L, Shiboski S, Grant RM. Sexual Seroadaptation: Lessons for Prevention and Sex Research from a Cohort of HIV-Positive Men Who Have Sex with Men. *PLoS ONE.* 2010 Jan 21; 5(1):e8831. [PubMed: 20098616]
25. Parsons JT, Schrimshaw EW, Wolitski RJ, Halkitis PN, Purcell DW, Hoff CC, et al. Sexual harm reduction practices of HIV-seropositive gay and bisexual men: serosorting, strategic positioning, and withdrawal before ejaculation. *AIDS Lond Engl.* 2005 Apr; 19(Suppl 1):S13–S25.
26. Jin F, Crawford J, Prestage GP, Zablotska I, Imrie J, Kippax SC, et al. HIV risk reduction behaviours in gay men. *AIDS Lond Engl.* 2009 Jan 14; 23(2):243–252.
27. Ilaria G, Jacobs JL, Polsky B, Koll B, Baron P, MacLow C, et al. Detection of HIV-1 DNA sequences in pre-ejaculatory fluid. *Lancet Lond Engl.* 1992 Dec 12; 340(8833):1469.
28. Pudney J, Oneta M, Mayer K, Seage G, Anderson D. Pre-ejaculatory fluid as potential vector for sexual transmission of HIV-1. *Lancet Lond Engl.* 1992 Dec 12; 340(8833):1470.
29. Chen Y. Treatment-related optimistic beliefs and risk of HIV transmission: a review of recent findings (2009–2012) in an era of treatment as prevention. *Curr HIV/AIDS Rep.* 2013 Mar; 10(1): 79–88. [PubMed: 23239272]
30. Heckathorn D. Respondent-Driven Sampling: A New Approach to the Study of Hidden Populations\*. *Soc Study Soc Probl.* 1997 May; 44(2)
31. Forrest JI, Stevenson B, Rich A, Michelow W, Pai J, Jollimore J, et al. Community mapping and respondent-driven sampling of gay and bisexual men’s communities in Vancouver, Canada. *Cult Health Sex.* 2014 Feb 10.

32. Moore DM, Cui Z, Lachowsky N, Raymond HF, Roth E, Rich A, et al. HIV Community Viral Load and Factors Associated With Elevated Viremia Among a Community-Based Sample of Men Who Have Sex With Men in Vancouver, Canada: JAIDS. *J Acquir Immune Defic Syndr*. 2016 May; 72(1):87–95. [PubMed: 26825177]
33. Lachowsky NJ, Lal A, Forrest JI, Card KG, Cui Z, Sereda P, et al. Including Online-Recruited Seeds: A Respondent-Driven Sample of Men Who Have Sex With Men. *J Med Internet Res*. 2016; 18(3):e51. [PubMed: 26980147]
34. Saunders JB, Aasland OG, Babor TF, de la Fuente JR, Grant M. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption--II. *Addict Abingdon Engl*. 1993 Jun; 88(6):791–804.
35. Fisher DG, Reynolds GL, Napper LE. Use of Crystal Meth, Viagra and Sexual Behaviour. *Curr Opin Infect Dis*. 2010 Feb; 23(1):53–56. [PubMed: 19918176]
36. Rich A, Lachowsky N, Cui Z, Sereda P, Lal A, Moore D, et al. Anal Sex Roles Explain Substance Use Patterns Among a Sample of HIV-Negative and HIV-Positive Gay, Bisexual, and Other Men who have Sex with Men in Vancouver, British Columbia, Canada. *Arch Sex Behav*. 2015
37. McKirnan DJ, Vanable PA, Ostrow DG, Hope B. Expectancies of sexual “escape” and sexual risk among drug and alcohol-involved gay and bisexual men. *J Subst Abuse*. 2001; 13(1–2):137–154. [PubMed: 11547615]
38. Kalichman SC, Rompa D. Sexual sensation seeking and Sexual Compulsivity Scales: reliability, validity, and predicting HIV risk behavior. *J Pers Assess*. 1995 Dec; 65(3):586–601. [PubMed: 8609589]
39. Van de Ven P, Crawford J, Kippax S, Knox S, Prestage G. A scale of optimism-scepticism in the context of HIV treatments. *AIDS Care*. 2000 Apr; 12(2):171–176. [PubMed: 10827857]
40. SAS. Cary, NC, USA: SAS Institute Inc;
41. Voltz, E., Wejnert, C., Cameron, C., Spiller, M., Barash, V., Degani, I., et al. Respondent-Driven Sampling Analysis Tool (RDSAT). Ithaca, New York: Cornell University; 2012.
42. Akaike, H. A New Look at the Statistical Model Identification. In: Parzen, E. Tanabe, K., Kitagawa, G., editors. *Selected Papers of Hirotugu Akaike* [Internet]. New York: Springer; 1974. p. 215-222. (Springer Series in Statistics). Available from: [http://link.springer.com/chapter/10.1007/978-1-4612-1694-0\\_16](http://link.springer.com/chapter/10.1007/978-1-4612-1694-0_16) [cited 2016 May 26]
43. PROC GENMOD. Type 3 Analysis :: SAS/STAT(R) 9.2 User’s Guide, Second Edition [Internet]. [cited 2016 May 26] Available from: [https://support.sas.com/documentation/cdl/en/statug/63033/HTML/default/viewer.htm#statug\\_genmod\\_sect035.htm](https://support.sas.com/documentation/cdl/en/statug/63033/HTML/default/viewer.htm#statug_genmod_sect035.htm).
44. Lima VD, Geller J, Bangsberg DR, Patterson TL, Daniel M, Kerr T, et al. The effect of adherence on the association between depressive symptoms and mortality among HIV-infected individuals first initiating HAART. *AIDS Lond Engl*. 2007 May 31; 21(9):1175–1183.
45. Heckathorn DD, Semaan S, Broadhead RS, Hughes JJ. Extensions of Respondent-Driven Sampling: A New Approach to the Study of Injection Drug Users Aged 18–25. *AIDS Behav*. 2002 Mar; 6(1):55–67.
46. Rehor JE. Sensual, Erotic, and Sexual Behaviors of Women from the “Kink” Community. *Arch Sex Behav*. 2015; 44(4):825–836. [PubMed: 25795531]
47. Wolitski RJ, Bailey CJ, O’Leary A, Gómez CA, Parsons JT. Seropositive Urban Men’s Study (SUMS). Self-perceived responsibility of HIV-seropositive men who have sex with men for preventing HIV transmission. *AIDS Behav*. 2003 Dec; 7(4):363–372. [PubMed: 14707533]
48. O’Dell BL, Rosser BRS, Miner MH, Jacoby SM. HIV prevention altruism and sexual risk behavior in HIV-positive men who have sex with men. *AIDS Behav*. 2008 Sep; 12(5):713–720. [PubMed: 17985229]
49. Marks G, Crepaz N, Janssen RS. Estimating sexual transmission of HIV from persons aware and unaware that they are infected with the virus in the USA. *AIDS Lond Engl*. 2006 Jun 26; 20(10):1447–1450.
50. Friedman SR, Mateu-Gelabert P, Sandoval M, Hagan H, Des Jarlais DC. Positive deviance control-case life history: a method to develop grounded hypotheses about successful long-term avoidance of infection. *BMC Public Health*. 2008; 8:94. [PubMed: 18366699]

51. Vazan P, Mateu-Gelabert P, Cleland CM, Sandoval M, Friedman SR. Correlates of staying safe behaviors among long-term injection drug users: psychometric evaluation of the staying safe questionnaire. *AIDS Behav.* 2012 Aug; 16(6):1472–1481. [PubMed: 22038081]
52. Bisits Bullen PA. The positive deviance/hearth approach to reducing child malnutrition: systematic review. *Trop Med Int Health TM IH.* 2011 Nov; 16(11):1354–1366. [PubMed: 21749582]
53. Lachowsky N, Stephenson K, Rich A, Lal A, Cui Z, Colley G, et al. Characteristics and Rates of Newly Infected HIV-Positive Gay, Bisexual and other Men Who Have Sex With Men (MSM) in Vancouver, British Columbia: Preliminary Findings of the Momentum Health Study. *Can J Infect Dis Med Microbiol.* 2015 Spring;26(B):83b.
54. Moore, D., Cui, Z., Lachowsky, N., Fisher, RH., Roth, E., Rich, A., et al. HIV positive MSM with unsuppressed viral load are more likely to engage in risky sex. Conference on Retroviruses and Opportunistic Infections (CROI); 2015 Feb 23; Seattle, United States.
55. Cassels S, Katz DA. Seroadaptation among Men Who Have Sex with Men: Emerging Research Themes. *Curr HIV/AIDS Rep.* 2013 Dec; 10(4):305–313. [PubMed: 24234489]
56. Williams, BA., Mandrekar, JN., Mandrekar, SJ., Cha, SS., Furth, AF. Finding Optimal Cutpoints for Continuous Covariates with Binary and Time-to-Event Outcomes [Internet]. Rochester, Minnesota: Department of Health Sciences Research Mayo Clinic; 2006 Jun. (Technical Report Series). Report No.: 79. Available from: <http://www.mayo.edu/research/documents/biostat-79pdf/doc-10027230> [cited 2016 May 30]



**Figure 2.** Distribution of the number of reported male anal sex partners in the past 6 months for HIV-negative (top) and HIV-positive (bottom) men.

Table 1

## Selected Descriptive Statistics for Overall Sample

Categorical Variables	n	%	RDS <sup>a</sup> %	RDS <sup>a</sup> % 95% CI <sup>b</sup>
<b>Demographics</b>				
<b>Ethnicity</b>				
White	539	75.0	68.0	60.6 74.4
Asian	72	10.0	9.9	6.2 14.8
Aboriginal	50	7.0	10.3	5.6 16.2
Other	58	8.1	11.8	7.0 17.1
<b>Sexual Identity</b>				
Gay	612	85.1	80.3	75.6 85.2
Bisexual/Other	107	14.9	19.7	14.8 24.4
<b>Education</b>				
Less than high school	61	8.7	14.2	9.5 19.5
Completed/more than high school	644	91.4	85.8	80.5 90.5
<b>Neighborhood</b>				
Downtown	356	49.5	51.8	43.8 59.2
Vancouver	223	31.0	30.5	24.0 36.7
Outside Vancouver	140	19.5	17.7	13.0 24.1
<b>Annual Income</b>				
< 30K	457	63.6	74.5	69.8 79.9
30–60K	182	25.3	17.2	13.2 20.9
60K	80	11.1	8.3	5.2 11.8
<b>Self-reported HIV Status</b>				
Negative/Unknown	524	72.9	76.9	68.8 84.4
Positive	195	27.1	23.1	15.6 31.2
<b>Sexual Behaviors</b>				
<b>Unprotected Anal Sex in P6M<sup>d</sup></b>				
No	256	36.4	38.2	32.8 45.4



Categorical Variables	n	%	RDS <sup>a</sup> %	RDS <sup>a</sup> % 95% CI <sup>b</sup>
Yes	185	26.3	26.0	20.6 31.6
High Risk Sex	262	37.3	35.8	29.1 41.4
<b>Insertive Anal Sex in P6M<sup>d</sup></b>				
No	207	28.8	34.9	28.8 41.7
Yes	512	71.2	65.1	58.3 71.2
<b>Receptive Anal Sex in P6M<sup>d</sup></b>				
No	235	32.7	36.5	30.7 42.6
Yes	484	67.3	63.5	57.4 69.3
<b>Anal Sex Preference</b>				
Bottom	241	35.1	35.9	29.5 41.3
Versatile	193	28.1	27.6	22.3 33.4
Top	253	36.8	36.5	31.5 42.8
No anal				
<b>Attended Group Sex Event in P6M<sup>d</sup></b>				
No	539	75.0	78.6	73.8 83.7
Yes	180	25.0	21.4	16.3 26.2
<b>Received Money in Exchange for Sex in P6M<sup>d</sup></b>				
No	655	91.2	89.9	85.2 94.3
Yes	63	8.8	10.1	5.7 14.8
<b>Substance Use</b>				
<b>AUDIT<sup>c</sup> Harmful Drinker</b>				
Yes	243	34.0	31.2	25.2 37.0
<b>Used EDD<sup>f</sup> in P6M<sup>d</sup></b>				
Yes	162	22.5	17.3	12.2 21.6
<b>Used methamphetamine in P6M<sup>d</sup></b>				
Yes	136	18.9	19.6	13.7 25.4
<b>Used Poppers in P6M<sup>d</sup></b>				
Yes	266	37.0	34.3	28.7 40.3

Categorical Variables	n	%	RDS <sup>a</sup> %	RDS <sup>a</sup> % 95% CI <sup>b</sup>
<b>Used Ecstasy in P6M<sup>d</sup></b>				
Yes	176	24.5	18.9	14.2 - 24.0
<b>Continuous Variables</b>				
	Median	Q1	Q3 <sup>e</sup>	
<i>Age</i>	33	26.47		
<i>HAART<sup>c</sup> Treatment Optimism Scale (n=716)</i>	25	21.28		
<i>Sexual Sensations Scale (n=698)</i>	31	28.34		
<i>Cognitive Escape Scale (n=705)</i>	29	25.33		

<sup>a</sup> Respondent Driven Sampling

<sup>b</sup> Confidence Interval

<sup>c</sup> Highly Active Antiretroviral Therapy

<sup>d</sup> Past Six Months

<sup>e</sup> Alcohol Use Disorder Identification Test

<sup>f</sup> Erectile Dysfunction Drugs

<sup>g</sup> Quartile 1, Quartile 3

**Table II**  
Univariable associations with the highest quartile of male anal sex partners among HIV-negative or unknown status men

Categorical Variables	<7 Anal Sex Partners (n=398)			7 Anal Sex Partners (n=126)			OR <sup>b</sup>	95% CI <sup>c</sup>	p-value
	n	%	RDS <sup>a</sup> %	n	%	RDS <sup>a</sup> %			
<b>Demographics</b>									
<b>Ethnicity</b>									
White	295	74.1	69.6	95	75.4	70.5	1.00		0.482
Asian	51	12.8	11.8	9	7.1	8.5	0.71	0.35 1.45	
Aboriginal	20	5.0	6.3	9	7.1	9.6	1.51	0.73 3.98	
Other	32	8.0	12.3	13	10.3	11.4	0.92	0.48 1.73	
<b>Sexual Identity</b>									
Gay	331	83.2	79.0	114	90.5	92.9	1.00		<0.001
Bisexual/Other	67	16.8	21.0	12	9.5	7.1	<b>0.29</b>	<b>0.14 0.60</b>	
<b>Education</b>									
Greater than high school	23	5.9	9.0	15	12.3	21.5	1.00		<0.001
Some/completed high school	369	94.1	91.0	107	87.7	78.5	<b>0.36</b>	<b>0.29 0.63</b>	
<b>Neighborhood</b>									
Downtown	155	38.9	42.4	67	53.2	54.1	1.00		<b>0.010</b>
Vancouver	153	38.4	34.5	34	27.0	20.4	<b>0.46</b>	<b>0.28 0.77</b>	
Outside Vancouver	90	22.6	23.1	25	19.8	25.5	0.86	0.53 1.49	
<b>Annual Income</b>									
<30K	246	61.8	71.6	67	53.2	60.7	1.00		<b>0.051</b>
30-60K	102	25.6	17.5	40	31.8	26.6	<b>1.79</b>	<b>1.16 2.95</b>	
60K	50	12.6	10.9	19	15.1	12.7	1.37	0.73 2.57	
<b>Seroadaptive Sexual Behaviors</b>									
<b>Strategy: Always Using Condoms</b>									
No	130	32.9	32.8	55	44.0	37.2	1.00		0.358
Yes	265	67.1	67.2	70	56.0	62.8	0.82	0.54 1.25	
<b>Strategy: Strategic Positioning</b>									

Categorical Variables	<7 Anal Sex Partners (n=398)			7 Anal Sex Partners (n=126)			p-value
	n	%	RDS <sup>a</sup> %	n	%	RDS <sup>a</sup> %	
Strategy: Anal Sex Avoidance	No	311	78.7	71	56.8	63.5	1.00
	Yes	84	21.3	54	43.2	36.5	<b>2.55</b>
Strategy: Serosorting	No	183	46.3	79	63.2	74.8	1.00
	Yes	212	53.7	46	36.8	25.2	<b>0.37</b>
Strategy: Viral-load Sorting	No	262	66.3	76	60.8	67.7	1.00
	Yes	133	33.7	49	39.2	32.3	1.15
Strategy: Withdrawal	No	372	94.2	97	77.6	82.4	1.00
	Yes	23	5.8	28	22.4	17.6	<b>3.78</b>
Strategy: Ask HIV Status	No	295	74.7	79	63.2	72.8	1.00
	Yes	100	25.3	46	36.8	27.2	1.34
Other Sexual Behaviors	No	159	40.3	35	28.0	29.4	1.00
	Yes	236	59.8	90	72.0	70.6	<b>1.86</b>
Anal Sex Preference	Bottom	136	36.3	37	29.6	28.8	1.00
	Versatile	94	25.1	38	30.4	36.7	<b>2.12</b>
	Top	145	38.7	50	40.0	34.4	0.97
Condomless Anal Sex in P6M <sup>c</sup>	No	174	44.7	28	22.8	25.6	1.00
	Yes, but only with seroconcordant partners	117	30.1	16	13.0	15.1	0.92
Watersports	Yes, with serodiscordant/unknown partners	98	25.2	79	64.2	59.3	<b>4.76</b>
	No	363	91.2	94	74.6	80.3	1.00
95% CI <sup>c</sup>							<b>1.61</b>
							<b>3.90</b>
95% CI <sup>c</sup>							<b>0.23</b>
							<b>0.58</b>
95% CI <sup>c</sup>							<b>2.17</b>
							<b>7.69</b>
95% CI <sup>c</sup>							<b>1.25</b>
							<b>2.87</b>

Categorical Variables	<7 Anal Sex Partners (n=398)			7 Anal Sex Partners (n=126)			p-value
	n	%	RDS <sup>a</sup> %	n	%	RDS <sup>a</sup> %	
Yes	35	8.8	6.3	32	25.4	19.7	
<b>Insertive Anal Sex in P6M<sup>e</sup></b>							
No	133	33.4	35.9	13	10.3	17.5	<0.001
Yes	265	66.6	64.1	113	89.7	82.5	
<b>Receptive Anal Sex in P6M<sup>e</sup></b>							
No	153	38.4	42.6	23	18.3	16.3	<0.001
Yes	245	61.6	57.4	103	81.8	83.7	
<b>Attended Sex Party/Darkroom/Blackout in P6M<sup>e</sup></b>							
No	342	85.9	87.8	68	54.0	50.7	<0.001
Yes	56	14.1	12.2	58	46.0	49.3	
<b>Received Money in Exchange for Sex in P6M<sup>e</sup></b>							
No	380	95.7	95.3	103	81.8	73.3	<0.001
Yes	17	4.3	4.7	23	18.3	26.7	
<b>Substance Use</b>							
<b>AUDIT<sup>b</sup>Harmful Drinker</b>							
No	246	61.8	65.3	72	57.6	57.7	0.125
Yes	152	38.2	34.7	53	42.4	42.3	
<b>Used EDD<sup>c</sup>in P6M<sup>e</sup></b>							
No	357	89.7	91.2	88	69.8	70.2	<0.001
Yes	41	10.3	8.8	38	30.2	29.8	
<b>Used methamphetamine in P6M<sup>e</sup></b>							
No	364	91.5	89.7	101	80.2	80.5	0.006
Yes	34	8.5	10.3	25	19.8	19.5	
<b>Used Poppers in P6M<sup>e</sup></b>							
No	296	74.4	77.6	59	46.8	51.8	<0.001
Yes	102	25.6	22.4	67	53.2	48.2	
<b>Used Ecstasy in P6M<sup>e</sup></b>							
No	296	74.4	77.6	59	46.8	51.8	<0.001
Yes	102	25.6	22.4	67	53.2	48.2	

Categorical Variables	<7 Anal Sex Partners (n=398)			7 Anal Sex Partners (n=126)			p-value	
	n	%	RDS <sup>a</sup> %	n	%	RDS <sup>a</sup> %		
No	312	78.4	81.2	77	61.1	60.4	<0.001	
Yes	86	21.6	18.8	49	38.9	39.6		
<b>Continuous Variables</b>								
	Median	Q1, Q3 <sup>h</sup>	Median	Median	Q1, Q3 <sup>h</sup>	OR <sup>b</sup>	95% CI <sup>c</sup>	p-value
Age	30	24, 39	29	29	25, 38	0.98	0.96, 1.00	0.929
HAART <sup>d</sup> Treatment Optimism Scale (n=716)	24	20, 26	24	24	20, 28	1.03	0.99, 1.08	0.167
Sexual Sensations Scale (n=698)	30	27, 33	32	32	30, 35	1.13	1.08, 1.19	<0.001
Cognitive Escape Scale (n=705)	28	24, 31	30	30	25, 33	1.03	0.99, 1.06	0.014

<sup>a</sup> Respondent Driven Sampling

<sup>b</sup> Odds Ratio

<sup>c</sup> Confidence Interval

<sup>d</sup> Highly Active Antiretroviral Therapy

<sup>e</sup> Past Six Months

<sup>f</sup> Alcohol Use Disorder Identification Test

<sup>g</sup> Erectile Dysfunction Drugs

<sup>h</sup> Quartile 1, Quartile 3

**Table III**

Univariable associations with the highest quartile of male anal sex partners among HIV-positive men.

Categorical Variables	<7 Anal Sex Partners (n=125)			7 Anal Sex Partners (n=69)			OR <sup>b</sup>	95% CI <sup>c</sup>	p-value
	n	%	RDS <sup>a</sup> %	n	%	RDS <sup>a</sup> %			
<b>Demographics</b>									
<b>Ethnicity</b>									
White	96	76.8	72.6	53	76.8	59.4	1.00		0.349
Asian	7	5.6	6.5	5	7.3	6.8	1.28	0.34	4.84
Aboriginal	15	12.0	13.8	6	8.7	22.2	1.97	0.83	4.68
Other	7	5.6	7.2	5	7.3	11.6	1.98	0.64	6.13
<b>Sexual Identity</b>									
Gay	103	82.4	80.2	63	91.3	87.7	1.00		0.231
Bisexual/Other	22	17.6	19.8	6	8.7	12.3	0.57	0.22	1.45
<b>Education</b>									
Greater than high school	19	15.6	14.1	4	5.9	11.1	1.00		0.591
Some/completed high school	103	84.4	85.9	64	94.1	88.9	1.32	0.48	3.64
<b>Neighborhood</b>									
Downtown	87	69.6	66.3	46	66.7	74.1	1.00		0.550
Vancouver	23	18.4	21.1	13	18.8	14.5	0.62	0.25	1.51
Outside Vancouver	15	12.0	12.6	10	14.5	11.4	0.87	0.29	2.25
<b>Annual Income</b>									
< 30K	101	80.8	81.6	42	60.9	74.8	1.00		0.270
30-60K	19	15.2	16.4	21	30.4	18.4	1.22	0.52	2.89
60K	5	4.0	2.0	6	8.7	6.8	3.76	0.66	21.28
<b>Seroadaptive Sexual Behaviors</b>									
<b>Strategy: Always Using Condoms</b>									
No	72	57.6	62.4	54	78.3	73.4	1.00		0.162
Yes	53	42.4	37.6	15	21.7	26.6	0.61	0.29	1.23
<b>Strategy: Strategic Positioning</b>									

Categorical Variables	<7 Anal Sex Partners (n=125)				7 Anal Sex Partners (n=69)				p-value	
	n	%	RDS <sup>a</sup> %	n	%	RDS <sup>a</sup> %	OR <sup>b</sup>	95% CI <sup>c</sup>		
Strategy: Anal Sex Avoidance	No	84	67.2	75.6	40	58.0	52.4	1.00		0.003
	Yes	41	32.8	24.4	29	42.0	47.6	2.82	1.41	5.62
Strategy: Serosorting	No	84	67.2	62.0	43	62.3	61.3	1.00		0.932
	Yes	41	32.8	38.0	26	37.7	38.7	1.30	0.53	2.14
Strategy: Viral-load Sorting	No	69	55.2	64.3	24	34.8	31.2	1.00		<0.001
	Yes	56	44.8	35.7	45	65.2	68.8	3.97	1.98	7.96
Strategy: Withdrawal	No	92	73.6	77.8	34	49.3	51.2	1.00		0.001
	Yes	33	26.4	22.2	35	50.7	48.8	3.33	1.66	6.71
Strategy: Ask HIV Status	No	94	75.2	74.9	45	65.2	65.8	1.00		0.229
	Yes	31	24.8	25.1	24	34.8	34.2	1.54	0.76	3.14
<b>Other Sexual Behaviors</b>										
<b>Anal Sex Preference</b>										
Bottom	40	34.2	29.2	28	40.6	63.9	1.00			<0.001
Versatile	39	33.3	41.2	21	30.4	13.7	0.15	0.59	0.39	
Top	38	32.5	29.6	20	29.0	22.4	0.35	0.15	0.85	
<b>Condomless Anal Sex in P6M<sup>e</sup></b>										
No	49	39.8	39.5	5	7.4	5.7	1.00			<0.001
Yes, but only with seroconcordant partners	34	27.6	30.4	18	26.5	19.3	4.44	1.09	18.16	
Yes, with serodiscordant/unknown partners	40	32.5	30.1	45	66.2	75.0	17.36	4.74	63.59	
<b>Watersports</b>										
No	116	92.8	96.7	46	66.7	77.6	1.00			<0.001



Categorical Variables	<7 Anal Sex Partners (n=125)						7 Anal Sex Partners (n=69)						p-value
	n	%	RDS <sup>a</sup> %	n	%	RDS <sup>a</sup> %	n	%	RDS <sup>a</sup> %	OR <sup>b</sup>	95% CI <sup>c</sup>		
<b>Insertive Anal Sex in P6M<sup>e</sup></b>													
Yes	9	7.2	3.3	23	33.3	22.4	8.60	2.49	29.72				
No	51	40.8	47.0	10	14.5	18.4	1.00						<0.001
Yes	74	59.2	53.0	59	85.5	81.6	3.93	1.80	8.62				
<b>Receptive Anal Sex in P6M<sup>e</sup></b>													
No	50	40.0	45.6	9	13.0	7.4	1.00						<0.001
Yes	75	60.0	54.4	60	87.0	92.6	10.55	3.56	31.31				
<b>Attended Group Sex Event, P6M<sup>e</sup></b>													
No	101	80.8	84.4	27	39.1	52.9	1.00						<0.001
Yes	24	19.2	15.6	42	60.9	47.1	4.80	2.28	10.11				
<b>Received Money in Exchange for Sex in P6M<sup>e</sup></b>													
No	116	92.8	93.3	56	81.2	85.8	1.00						0.121
Yes	9	7.2	6.7	13	18.8	14.2	2.29	0.79	6.69				
<b>Substance Use</b>													
<b>AUDIT<sup>f</sup> Harmful Drinker</b>													
No	97	78.9	76.9	56	82.4	86.2	1.00						0.164
Yes	26	21.1	23.1	12	17.7	13.8	0.53	0.22	1.34				
<b>Used EDD<sup>g</sup> in P6M<sup>e</sup></b>													
No	87	69.6	78.1	24	34.8	47.0	1.00						<0.001
Yes	38	30.4	21.9	45	65.2	53.0	4.25	2.00	8.12				
<b>Used methamphetamine in P6M<sup>e</sup></b>													
No	92	73.6	77.7	25	36.2	46.3	1.00						<0.001
Yes	33	26.4	22.3	44	63.8	53.7	4.43	2.70	8.14				
<b>Used Poppers in P6M<sup>e</sup></b>													
No	75	60.0	59.2	22	31.9	36.0	1.00						0.005
Yes	50	40.0	40.8	47	68.1	64.0	2.59	1.32	5.79				
<b>Used Ecstasy in P6M<sup>e</sup></b>													

Categorical Variables	<7 Anal Sex Partners (n=125)				7 Anal Sex Partners (n=69)				p-value
	n	%	RDS <sup>a</sup> %	n	%	RDS <sup>a</sup> %	OR <sup>b</sup>	95% CI <sup>c</sup>	
No	110	88.0	91.9	43	62.3	73.4	1.00		0.002
Yes	15	12.0	8.1	26	37.7	26.6	<b>4.13</b>	<b>1.65</b>	<b>1.32</b>

Continuous Variables	<7 Anal Sex Partners (n=125)		7 Anal Sex Partners (n=69)		p-value
	Median	Q1, Q3 <sup>h</sup>	Median	Q1, Q3 <sup>h</sup>	
Age (Median Q1, Q3)	47	41, 51	47	42, 52	0.564
HAAART <sup>d</sup> Treatment Optimism Scale (n=716)	28	25, 31	29	26, 32	0.044
Sexual Sensations Scale (n=698)	30	27, 33	34	31, 37	<b>&lt;0.001</b>
Cognitive Escape Scale (n=705)	30	26, 35	33	28, 36	<b>0.028</b>

<sup>a</sup> Respondent Driven Sampling

<sup>b</sup> Odds Ratio

<sup>c</sup> Confidence Interval

<sup>d</sup> Highly Active Antiretroviral Therapy

<sup>e</sup> Past Six Months

<sup>f</sup> Alcohol Use Disorder Identification Test

<sup>g</sup> Erectile Dysfunction Drugs

<sup>h</sup> Quartile 1, Quartile 3

Table IV

Multivariable results with the highest quartile of male anal sex partners among HIV-negative or unknown men.

	AOR <sup>a</sup>	95% CI <sup>b</sup>	p-value
<b>Sexual Identity</b>			
Gay	1.00		
Bisexual/Other	<b>0.26</b>	<b>0.10</b>	<b>0.66</b>
<b>0.005</b>			
<b>Education</b>			
Greater than high school	1.00		
Some/completed high school	<b>0.32</b>	<b>0.12</b>	<b>0.83</b>
<b>0.019</b>			
<b>Neighborhood</b>			
Downtown	1.00		
Vancouver	<b>0.32</b>	<b>0.15</b>	<b>0.68</b>
<b>0.003</b>			
Outside Vancouver	0.69	0.35	1.37
0.290			
<b>Strategy: Strategic Positioning</b>			
No	1.00		
Yes	<b>3.81</b>	<b>1.79</b>	<b>8.11</b>
<b>&lt;.001</b>			
<b>Strategy: Anal Sex Avoidance</b>			
No	1.00		
Yes	<b>0.22</b>	<b>0.11</b>	<b>0.42</b>
<b>&lt;.001</b>			
<b>Strategy: Ask HIV Status</b>			
No	1.00		
Yes	<b>2.15</b>	<b>1.18</b>	<b>3.93</b>
<b>0.013</b>			
<i>Sexual Sensations Scale (n=698)</i>	<i>1.07</i>	<i>0.99</i>	<i>1.16</i>
<i>0.087</i>			
<b>Anal Sex Preference</b>			
Bottom	1.00		
Versatile	<b>2.75</b>	<b>1.31</b>	<b>5.75</b>
<b>0.007</b>			
Top	<b>0.62</b>	<b>0.28</b>	<b>1.36</b>
<b>0.229</b>			
<b>Unprotected Anal Sex, P6Mc</b>			
No	1.00		
Yes, but only with seroconcordant partners	0.72	0.32	1.60
0.417			
Yes, with serodiscordant/unknown partners	1.78	0.87	3.65
0.115			

	AOR <sup>a</sup>	95% CI <sup>b</sup>	p-value
<b>Watersports</b>			
No	1.00		
Yes	<b>2.78</b>	<b>1.23 6.28</b>	<b>0.014</b>
<b>Attended Group Sex Event, P6M<sup>c</sup></b>			
No	1.00		
Yes	<b>6.07</b>	<b>3.08 11.98</b>	<b>&lt;.001</b>
<b>Received Money for Sex, P6M<sup>c</sup></b>			
No	1.00		
Yes	<b>5.35</b>	<b>1.79 15.98</b>	<b>0.003</b>
<b>Used Ecstasy, P6M<sup>c</sup></b>			
No	1.00		
Yes	<b>2.96</b>	<b>1.49 5.89</b>	<b>0.002</b>

<sup>a</sup> Adjusted Odds Ratio

<sup>b</sup> CI - Confidence Interval

<sup>c</sup> P6M - Past Six Months.

**Table V**

Multivariable associations with the highest quartile of male anal sex partners among HIV-positive men.

	AOR <sup>a</sup>	95% CI <sup>b</sup>	p-value
<b>Strategy: Ask HIV status</b>			
No	1.00		
Yes	<b>3.10</b>	<b>1.17 8.26</b>	<b>0.024</b>
<b>Sexual Sensations Scale (n=698)</b>			
	<b>1.17</b>	<b>1.05 1.31</b>	<b>0.004</b>
<b>Anal Sex Preference</b>			
Bottom	1.00		
Versatile	0.16	0.05 0.53	0.003
Top	0.44	0.15 1.25	0.122
<b>Attended Group Sex Event, P6M<sup>c</sup></b>			
No	1.00		
Yes	<b>3.41</b>	<b>1.21 9.66</b>	<b>0.021</b>
<b>Received Money for Sex, P6M<sup>c</sup></b>			
No	1.00		
Yes	<b>5.07</b>	<b>1.09 23.61</b>	<b>0.039</b>
<b>Used Methamphetamine, P6M<sup>c</sup></b>			
No	1.00		
Yes	<b>3.06</b>	<b>1.16 8.10</b>	<b>0.024</b>
<b>Used EDD, P6M<sup>c</sup></b>			
No	1.00		
Yes	2.52	0.93 6.83	0.068

<sup>a</sup> Adjusted Odds Ratio

<sup>b</sup> Confidence Interval

<sup>c</sup> Past Six Months.