



# Pathogenesis of Inflammatory Bowel Disease and Recent Advances in Biologic Therapies

Duk Hwan Kim<sup>1</sup> and Jae Hee Cheon<sup>2,3,4\*</sup>

<sup>1</sup>Digestive Disease Center, CHA Bundang Hospital, CHA University, Seongnam 13496, <sup>2</sup>Department of Internal Medicine and Institute of Gastroenterology, Yonsei University College of Medicine, <sup>3</sup>Severance Biomedical Science Institute, Yonsei University College of Medicine, <sup>4</sup>Brain Korea 21 PLUS Project for Medical Science, Yonsei University College of Medicine, Seoul 03722, Korea

Inflammatory bowel disease (IBD) is a chronic intestinal inflammatory disorder with an unknown etiology. IBD is composed of two different disease entities: Crohn's disease (CD) and ulcerative colitis (UC). IBD has been thought to be idiopathic but has two main attributable causes that include genetic and environmental factors. The gastrointestinal tract in which this disease occurs is central to the immune system, and the innate and the adaptive immune systems are balanced in complex interactions with intestinal microbes under homeostatic conditions. However, in IBD, this homeostasis is disrupted and uncontrolled intestinal inflammation is perpetuated. Recently, the pathogenesis of IBD has become better understood owing to advances in genetic and immunologic technology. Moreover, new therapeutic strategies are now being implemented that accurately target the pathogenesis of IBD. Beyond conventional immune-suppressive therapy, the development of biological agents that target specific disease mechanisms has resulted in more frequent and deeper remission in IBD patients, with mucosal healing as a treatment goal of therapy. Future novel biologics should overcome the limitations of current therapies and ensure that individual patients can be treated with optimal drugs that are safe and precisely target IBD.

[Immune Network 2017;17(1):25-40]

**Keywords:** Inflammatory bowel diseases, Crohn disease, Colitis, Ulcerative, Physiopathology

## INTRODUCTION

The entire surface of the human intestine reaches 200~400 m<sup>2</sup> (1). Moreover, it occupies a central position as the frontier of the innate immune system. The inner cell lining of the intestine works not only as a barrier to protect the host from harmful pathogens but also as a place where interactions with commensal microorganisms occur. These interactions are delicately modulated by the intestinal immune system and contribute to immune homeostasis. For various reasons, idiopathic intestinal

inflammations such as inflammatory bowel disease (IBD) can occur when this homeostasis is disrupted (2,3).

IBD is a multifactorial immune disorder characterized by chronic relapsing inflammation of the intestine (4). It is classified into two different disorders: Crohn's disease (CD) and ulcerative colitis (UC). Clinically, CD and UC share similar symptoms, including diarrhea, hematochezia, and abdominal pain, whereas the location and depth of inflammation, as well as complications and prevalence can differ. Currently, the exact etiology of IBD is unclear. However, it is believed that disturbance

Received on December 1, 2016. Revised on January 24, 2017. Accepted on February 7, 2017.

© This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

\*Corresponding Author. Jae Hee Cheon, Department of Internal Medicine and Institute of Gastroenterology, Yonsei University College of Medicine, 50-1 Yonsei-ro, Seodaemun-gu, Seoul 03722, Korea. Tel: 82-2-2228-1990; Fax: 82-2-393-6884; E-mail: [geniushee@yuhs.ac](mailto:geniushee@yuhs.ac)

of the immune system and/or imbalanced interactions with microbes leads to development of chronic intestinal inflammation when certain environmental factors trigger genetically susceptible hosts. Traditionally, Th1 cells have been thought to play an important role in pathogenesis related to the chronicity of intestinal inflammation, especially in CD, whereas Th2 cells have been thought to play an important role in UC (5). Recently, however, activation of Th17 cells and imbalance of Th17/regulatory T (Treg) cells are recognized to be an important component in the development of intestinal inflammation (6). Since tumor necrosis factor (TNF)- $\alpha$  has been identified as a key cytokine in IBD pathogenesis, the introduction of anti-TNF- $\alpha$  treatment has led to the development of disease-modifying drugs (7-9). Compared to conventional therapies, anti-TNF agents have higher rates of remission induction and maintenance. Moreover, these drugs have been able to obtain mucosal healing through targeted immune suppression. However, about a third of patients with IBD still do not show an appropriate response to existing therapies. This high rate of treatment failure suggests that there are still unknown aspects regarding the mechanism of IBD. Encouragingly, however, dozens of novel agents based on recent advances in our understanding of the mucosal immune system for IBD pathogenesis have been developed and are now in clinical trials worldwide. In this review, we will describe our current knowledge of the mucosal immune system in terms of IBD pathogenesis and discuss its therapeutic implications.

## THE INTESTINAL EPITHELIUM AND MICROBIOTA

The intestinal epithelial cell (IEC) layer consists of several different cells, including enterocytes, goblet cells, neuroendocrine cells, Paneth cells, M cells, and epithelial resident intestinal stem cells. These cells structurally constitute crypts and villi, with a single columnar cell lining with a tight junction, and secrete mucus containing anti-microbial peptides; these cells separate intra-luminal pathogens from the subepithelial lamina propria (3,10,11).

### Mucus layer

To protect mucosa, a mucus layer covers the outer epithelial surface. The mucus layer is composed of glycosylated mucin from goblet cells as well as defensins from Paneth cells and IECs. A major component of mucin is encoded by *Muc2*, and spontaneous colitis develops upon deletion of *Muc2* in mice (12). A study showed that aberrant mucin production was accompanied by endoplasmic

reticulum (ER) stress (13). Goblet cell depletion and a reduced mucus layer are characteristic findings in patients with UC (14). In addition to mucin, Paneth cells secrete  $\alpha$ -defensin, whereas most IECs produce  $\beta$ -defensin. Paneth cells are known to play an important role in the homeostasis of the intestinal epithelium. Genetic alterations or ER stress causing Paneth cell dysfunction or depletion result in dysbiosis of commensal flora and increased susceptibility to intestinal inflammation (15). It is known that IBD patients often have this Paneth cell dysfunction (16). Paneth cell abnormalities are thought to be a very early event in IBD development, particularly in CD. Therefore, there are studies examining the effects of applying ER-stress-reducing methods to IBD treatment. A study showed that the chemical chaperones tauroursodeoxycholate (TUDCA) and 4-phenylbutyrate (PBA)—small molecules that can reduce ER stress by facilitating protein folding—prevented the induction of intestinal inflammation in mice (17).

### Integrity of the intestinal epithelium

Epithelial integrity is maintained by tight junctions between IECs. When the permeability of the intestinal epithelium is increased, external pathogens are easily introduced, which is known to affect the pathogenesis of IBD (18). Several lines of evidence showed that single-nucleotide polymorphisms in the organic cation transporter (OCTN), which mediates the transport of organic cations across the cell membrane, were associated with CD susceptibility (19,20). IECs also play a role as communicator between pathogens and lamina propria. Only small amounts of bacteria are generally capable of moving into the intestinal epithelium. This translocation is a method of antigen sampling and immune surveillance for the intestinal mucosal immune system that is essential for the host's immune homeostasis (21). However, when the integrity of the intestinal epithelial layer is broken, a high influx of intestinal contents and/or a high burden of microorganisms is thought to initiate and maintain a sustained inflammatory response, which is considered to be one of the mechanisms underlying IBD (22). For example, in an animal model in which the barrier function of the intestinal epithelial layer is reduced, such as in mice with a dominant negative N-cadherin mutation (23) or mice lacking NOD1 and NOD2 (24), the mice develop IBD-like enteritis. Moreover, several genetic studies have identified several candidate genes in patients with UC (25,26), such as *CDH1* and *LAMB1*, which are involved in regulation of the epithelial barrier. Therapeutic attempts to restore mucosal barrier function have also been attempt-

ted. Phosphatidylcholine (lecithin) is abundant in the mucus of healthy colons, whereas reduced lecithin levels were observed in UC patients (27). A phase IIA, double-blind, randomized, placebo-controlled study showed that oral administration of lecithin was effective for achieving clinical remission in patients with chronic active UC (28).

### Intestinal microbiota

Generally, there are approximately  $10^{11}$ – $10^{14}$  enteric commensal microorganisms from 300–500 bacterial species (29,30). Under normal circumstances, most commensal bacteria play an essential role in protecting intestinal homeostasis. They affect crucial nutrient provision, development of the immune system, and modulation of energy metabolism (5,31). The majority of commensal bacteria consist of gram-negative bacteria, such as *Bacteroidetes*, and gram-positive bacteria, such as *Firmicutes* (32). Other minor divisions are comprised of *Proteobacteria*, *Actinobacteria*, *Fusobacteria*, and *Verrucomicrobia* (32). Those mucosa-associated phyla are reduced in diversity and amount in patients with IBD compared to that in healthy humans (5,33,34). However, commensal microorganisms can be noxious for intestinal inflammation under certain circumstances (35). There are some clues that commensal bacteria play an important role in the development of IBD. First, empiric antibiotic treatment has been effective in some IBD patients (36). Second, IBD patients have increased titers against indigenous bacteria (37). Third, genetic variants that are associated with bacterial detection, such as *NOD2* (38), and T cell immunity, such as *IL23R* (39), are implicated in IBD. Fourth, most animal models of colitis require commensal bacteria for the initiation of intestinal inflammation (40). In addition, recent studies have focused on the contribution of other enteric microorganisms, such as viruses or fungi, for IBD development. For example, a study (41) revealed that altered amounts and compositions of enteric virus were related to experimental colitis. In particular, mice without Toll-like receptor (TLR) 3 and TLR7 were more susceptible to the induction of colitis. Likewise, Iliev *et al.* reported that mice deficient for Dectin-1, which is an innate immune receptor responsible for interacting with commensal fungi, showed increased susceptibility to colitis (42).

To date, several pathogens have been proposed as causative microorganisms for IBD development. Recent studies showed *Proteobacteria*, especially adherent-invasive *Escherichia coli* (AIEC), as one of the candidates. AIEC has been detected more frequently in patients with CD than in healthy subjects (43–45). AIEC is known to

be able to invade epithelium and replicate within macrophages (46). Some investigators isolated AIEC from the ileum of patients with CD (47,48). However, AIEC was rarely found in the colon tissues of CD patients and was not identified in UC patients (49), suggesting that AIEC performs an important role in the occurrence of small bowel inflammation (18).

In contrast, *Clostridium* cluster *XIVa* and *IV* are thought to be a crucial part of gut homeostasis through Treg cell accumulation (50). Foxp3<sup>+</sup>CD4<sup>+</sup> Tregs are known to be abundant in the colonic lamina propria and are the most important immune-regulating cells (51). Several studies showed that Treg cells were strongly affected by intestinal microbiota (52). In particular, Treg cells stimulated by CBir1, a microbiota flagellin, induce IgA<sup>+</sup>B cells in the intestine. As a result, decreased pathogenic loading by IgA leads to down-regulation of systemic T cell activation (53). An experimental murine model with an increased *Clostridium XIVa/IV* population was resistant to allergy and intestinal inflammation (50). Conversely, patients with IBD showed a reduced *Clostridium XIVa/IV* compared to that in controls (34,54,55).

Observations of dysbiosis in IBD patients led to efforts to restore microbiota to a normal composition. Fecal microbial transplantation (FMT) has emerged as a novel treatment in patients with IBD. One randomized control trial involving 75 UC patients showed a significantly higher remission rate (24%) in patients receiving FMT from unrelated donor enemas than that in the placebo group (5%) (56). However, a second randomized control trial with 48 UC patients reported a negative result (57). Currently, there are no randomized control trials comparing FMT with placebo treatment in CD patients. A meta-analysis using four case series data in 38 CD patients revealed a 60.5% pooled response rate (58). However, their outcome was not that of mucosal remission but of clinical response. Therefore, the effectiveness of FMT as a therapeutic application for IBD remains unclear. Furthermore, optimal donor selection, delivery methods, and donor feces processing have not yet been standardized. Probiotics are nutritional supplements that contain microorganisms that benefit the host's health when administered in the proper amount. Attempts have also been made to treat IBD by improving intestinal microbial balance through probiotics. In an experimental colitis model, probiotics showed an anti-inflammatory effect through TLR9 signaling (59). A recent meta-analysis using 23 randomized controlled trials showed that administration of probiotics was associated with benefits regarding induction and maintenance of remission in

patients with UC but not in CD (60). Further studies are warranted to draw a concrete conclusion in terms of the therapeutic effects of probiotics in IBD.

## INNATE AND ADAPTIVE IMMUNITY IN IBD TREATMENT

### Innate immune recognition

The innate immune system is at the forefront of defending against external pathogens in the human immune system. The innate immune system provides rapid and non-specific protection to the host through pattern-recognition of pathogens, whereas the adaptive immune system mediates highly selective and long-lasting immunity. The innate immune system of the intestine is composed of intestine epithelia, macrophages, monocytes, neutrophils, eosinophils, basophils, dendritic cells (DCs), and natural killer cells. Intraluminal pathogens continuously communicate with innate immune cells through diverse innate immune receptors such as TLRs, NOD, leucine-rich repeat receptors (NLRs), C-type lectin receptors (CLRs), and retinoic acid-inducible gene 1-like receptors (RLRs) (61). When intestinal macrophages and DCs sense pathogen-associated molecular patterns (PAMPs) of microbes, activated signal pathways, such as NF- $\kappa$ B, produce proinflammatory cytokines, chemokines, and anti-microbial peptides (62). Activation of macrophages by these cytokines and chemokines plays a role in the direct elimination of pathogens through free radicals and proteases and also results in antigen presentation to the adaptive immune system. Antigen presenting cells (APCs), such as DCs and macrophages, have a key role in connecting the innate and adaptive immune system. Comparing that macrophages perform antigen presentation and have a phagocytic function, activated DCs present intraluminal pathogens to naïve CD4<sup>+</sup> T cells at secondary lymphoid organs of the gut and modulate the polarization of naïve CD4<sup>+</sup> T cells to Treg cells and helper T cells, including Th1, Th2, and Th17 cells.

Under non-inflammatory conditions, TLR signaling leads to tolerance towards luminal pathogens through down-regulation of pattern-recognition receptors and promotes mucosal wound healing. In IBD patients, impaired TLR signaling often leads to increased intestinal permeability and inappropriate mucosal healing. For example, TLR2-deficient mice showed an increased mortality rate after damage was induced to the colon mucosa via chemicals. TLR2 signaling stimulates the production of trefoil factor (TFF) 3 and restores damaged

mucosa. In mice lacking TLR2, mortality was reduced when TFF3 was administered (63). Likewise, genetic studies showed the association between CD and the nucleotide oligomerization domain (*NOD*) 2 gene. *NOD2* polymorphisms lead to an impaired response to bacterial peptidoglycan sensing. While the exact mechanism between CD and impaired *NOD2* function is still unclear, *NOD2*-mediated chronic stimulation is thought to be one of the factors controlling proinflammatory cytokine production. Recent genome-wide association studies (GWAS) showed relationships between a variety of single-nucleotide polymorphisms (SNPs) and IBD risks: microbial sensing (*NOD2*, *IRF5*, *NFKB1*, *RELA*, *REL*, *RIPK2*, *CARD9*, and *PTPN22*), microbial elimination (*ATG16L1*, *IRGM*, and *NCF4*), and integration of antimicrobial adaptive immune responses (*IL23R*, *IL10*, *IL12*, *IL18RAP/IL1R1*, *IFNGR/IFNAR1*, *JAK2*, *STAT3*, and *TYK2*) (64-66). Recent studies have shown that autophagy plays an important role in the innate immune system. Autophagy is a biological process that activates cellular autodigestion of the cell's own cytosolic materials, including intracellular microbes. Additionally, autophagy enables antigen presentation by major histocompatibility complex class II. Repeated GWAS studies consistently showed the association between CD and autophagy-related genetic polymorphisms, such as *ATG16L1* and *IRGM* (67-71). The accumulation of both macrophages and DCs is observed in the lamina propria of IBD patients and in experimental colitis models (3). If interactions between DCs and T cells are interrupted, experimental T cell-mediated colitis is prevented (72). Moreover, DCs and macrophages also play a role in maintaining gut homeostasis against the inflammatory conditions of the intestine. A study showed that a pro-resolution mediator, prostaglandin D (2), was specifically up-regulated in UC patients with long-term remission (73). Likewise, another study revealed that a SNP associated with low levels of CD39/ENTPD1, which hydrolyzes proinflammatory nucleotides and generates adenosine, was related to an increased risk of developing CD (74). Taken together, an impaired innate immune response might promote IBD development via inappropriate stimulation of adaptive immunity through failure to control microorganisms (66). Therefore, researchers have focused on enhancing innate immunity as a therapeutic target of IBD. For example, certain growth factors, such as granulocyte-macrophage colony-stimulating factor (GM-CSF) and granulocyte colony-stimulating factor (G-CSF), which are critical for modulation of cellular proliferation, differentiation, angiogenesis, and inflammation, have been evaluated for



treatment of intestinal inflammation (75).

### Adaptive immunity

Chronic inappropriate activation of the adaptive immune system against commensal microorganism has been thought to be the main pathogenesis of IBD. Increased production of IFN- $\gamma$  from Th1 cells and cytokines related with Th17 cell, such as IL-17A/F, IL-21, IL-22, and CXCL8, are observed in the intestine of CD patients, while T cells from the lamina propria of UC patients highly produce Th2 cell-related cytokines, such as IL-5 and IL-13 (3,76,77).

Classically, immune-modulating treatments of IBD have focused on adaptive immunity. Corticosteroids have been widely used to treat acute flares of IBD since Truelove and Witts reported the effectiveness of oral corticosteroids in patients with UC in 1955 (78). Suppression of pro-inflammatory cytokines, such as TNF- $\alpha$  and IL-1 $\beta$ , is known to be the primary mechanism underlying how corticosteroids control IBD (79). In addition, recent studies highlighted that corticosteroids play an important role in regulation of T helper cell differentiation and type I interferon (IFN) production (79). Clinically, corticosteroids are effective for remission induction of IBD. A study observed that the first course of oral corticosteroid treatment achieved 89.5% of therapeutic response after 1 month, 69.5% after 4 months, and 56.6% after 1 year in patients with CD (80). Likewise, various immunomodulators that down-regulate the proinflammatory cytokines of T cells have been a well-established treatment for IBD. For example, cyclosporine A and tacrolimus are used for remission induction of active UC, and methotrexate is used for chronically active CD (81-83). Of these treatments, the most widely used agent is thiopurines, such as azathioprine and 6-mercaptopurine. Thiopurine inhibits purine nucleotide synthesis and breaking of DNA in leukocytes via 6-thioguanine nucleotides (6-TGNs), which is the effector product of thiopurine metabolism (84). Moreover, thiopurine suppresses CD4<sup>+</sup> T cell activity and promotes T cell apoptosis by inhibiting GTPase Rac1 in inflamed intestine (85). Recently, a study showed that autophagy-related genetic variant, *ATG16L1*, was associated with the good clinical towards thiopurine treatment in patients with CD but not in UC (86). In addition, a study showed that local administration of thioguanine improved murine colitis by promoting autophagy and killing translocated bacteria at the site of the inflamed intestine independently of systemic myelosuppression (87). Therefore, thiopurine probably works by multiple mechanisms to

improve IBD. Meanwhile, thiopurine may cause life-threatening leukopenia. The *TPMT* mutation is known to be associated with this complication (88). More specifically, a study involving immunochip genotyping of Asians revealed that *NUDT15* SNP was strongly related to thiopurine-induced leukopenia, and another study showed an association between *FTO* variant and leukopenia by GWAS analysis (89,90). Although non-specific immunosuppression using immunomodulators is generally safe and effective for disease control to an extent, advances in the understanding of the specific mechanisms of IBD led to the development of targeted treatment, i.e., biologics.

### Modulation of anti-inflammatory cytokines

The era of biologic therapy began with an anti-TNF agent, infliximab, in patients with CD (91). TNF- $\alpha$  is a proinflammatory cytokine that is produced by activated macrophages, monocytes, and T lymphocytes (7). Intestinal specimens of CD patients were shown to have increased levels of TNF- $\alpha$  protein and mRNA expression (92). Excessive production of TNF- $\alpha$  using experimental deletion of the adenosine-uracil (AU)-rich elements (ARE) from the 38-untranslated region (38-UTR) of the TNF- $\alpha$  gene in mice resulted in development of chronic inflammatory arthritis and CD-like IBD phenotype (93). Another experimental study revealed that inhibiting TNF was able to improve dextran sulfate sodium (DSS)-induced colitis in a mice model (94). Therefore, TNF- $\alpha$  has been thought to play a pivotal role in the development of IBD.

TNF- $\alpha$  has two forms in the human intestine: transmembrane TNF (mTNF) and soluble TNF (sTNF). mTNF is generally expressed on the surface of CD14<sup>+</sup> macrophages and targets TNF-R2 of T cells, whereas sTNF is secreted by several immune cells as a signaling molecule and targets TNF-R1 of effector cells (95,96). In IBD, increased levels of both mTNF and sTNF play various pro-inflammatory functions in the inflamed gut, such as angiogenesis, Paneth cell death, matrix metalloproteinase production from myofibroblasts, and the undermining of the barrier function of IECs (77). Recent studies have shown that interaction between mTNF and TNF-R2 is more important for IBD pathogenesis than that of sTNF and TNF-R1 (97,98).

Several anti-TNF- $\alpha$  monoclonal antibodies have been developed since infliximab (chimeric antibody with a murine sequence) and adalimumab (fully humanized antibody) showed effectiveness for induction and maintenance of remission, as well as mucosal healing of

IBD (99-103). A humanized, pegylated anti-TNF Fab fragment, certolizumab pegol, also showed benefits and was approved for CD treatment (104). Moreover, golimumab, a transgenic fully human monoclonal immunoglobulin G1 antibody, was recently launched for the treatment of UC (105). However, another anti-TNF agent that targets sTNF, etanercept, showed no benefits regarding treatment of IBD (106,107). Although there are several limitations of anti-TNF treatment, such as safety issues, relatively high cost, and loss of effectiveness, the potential benefits of anti-TNF agent may outweigh these drawbacks, because blocking the TNF signal in IBD works through various mechanisms, including T cell apoptosis, inhibiting T cell differentiation, induction of Treg cells and macrophages, and barrier improvement (96,108-110). Therefore, efforts to overcome the drawbacks of anti-TNF agents, such as oral formulations, bacteria-producing nanobodies, and therapeutic vaccines against TNF are still in development (111-113).

Other important cytokines in the treatment of IBD are related to Th17 cells (IL-17A, IL21, IL-22, and IL-23) (114). Th17 cells are differentiated from naïve CD4<sup>+</sup> T cells that are stimulated with transforming growth factor (TGF)- $\beta$  and IL-6 in mice models. The inflamed intestinal tissue of IBD patients was shown to contain higher levels of Th17 cells and its cytokines (115). Additionally, a chemoattractant of Th17 cells, CCL20, is also elevated in the intestinal mucosal of IBD patients (116). While the exact role of Th17 cells and their cytokines in regards to intestinal inflammation has not yet been fully understood, the balance between Th17 and Treg cells are thought to be an important aspect in development of IBD (117). Based on those viewpoints, a humanized IFN- $\gamma$  antibody, fontolizumab, was developed. However, it did not show a satisfactory result in patients with moderately to severely active CD (118). Likewise, an attempt to target IL-17A through a monoclonal antibody, secukinumab, in patients with CD also failed (119). In experimental colitis models, Th17 cells and their related cytokines are thought to play both inflammatory and anti-inflammatory roles in the intestine (114). Anti-inflammatory cytokines such as IL-22 are also produced by Th17 cells. These cytokines are known to promote epithelial proliferation, mucosal healing, and anti-microbial peptides in the mucus (120). Moreover, plasticity reflecting the environmental conditions during the inflammatory process between Th1/Treg and Th17 cells is observed, and these reciprocal alterations are thought to be important for maintaining intestinal homeostasis (121). The contribution of Th17 cells for intestinal inflammation might be controlled by

more detailed interactions between immune cells.

Considering the complex interactions between various cytokines that contribute IBD, the targeting of multiple cytokines is thought to be a reasonable approach in the treatment of IBD. Ustekinumab, a human monoclonal antibody against the p40 subunit that is a component of both IL-12 and IL-23, is theoretically relevant for the treatment of CD involving both Th1 and Th17 aspects of CD. IL-12 induces Th1 polarization of naïve CD4<sup>+</sup> T cells and IL-23 promotes Th17 cell differentiation (122). In moderate-to-severe CD patients, ustekinumab showed a significant clinical benefit in both remission induction and maintenance (123). Similarly, other biologics targeting the IL-12/23 pathway, such as ABT-874 (124) and apilimod mesylate (125), are under evaluation. IL-10-deficient mice can develop spontaneous T cell-dependent colitis and colitic cancer (126,127). IL-10 is known as an anti-inflammatory cytokine and the UC-related *IL10* gene variation was also noted from a GWAS study. Given this, a study used an interesting approach to treat experimental murine colitis involving the genetically engineered IL-10-secreting bacteria *Lactococcus lactis*. Intra-gastric administration of this bacteria resulted in a 50% reduction of DSS-induced colitis in IL-10- knockout mice (128).

#### Targeting inter-/intra-cellular signaling pathways

Interactions between proinflammatory cytokines and their receptors lead to activation of intracellular signal transduction and production of inflammatory proteins. Janus kinase (JAK)-signaling transducers and activator of transcription (STAT) cytokine signaling pathways are recently thought to be a potential therapeutic target of IBD. Because various key cytokines such as IFN- $\gamma$ , IL-2, IL-4, IL-7, IL-9, IL-15, IL-12, IL-21, IL-22, and IL-23 depend on the JAK-signaling pathway, inhibiting JAK might result in the downregulation of multiple inflammatory cytokines (129). The JAK family consists of four intracellular proteins, JAK1, JAK2, JAK3, and tyrosine kinase (TYK) 2 (130). A JAK1/JAK3 inhibitory small molecule, tofacitinib, showed promising results for the treatment of UC in a phase II study. In 194 moderately to severely active UC patients, 78% of patients who received 15 mg of oral tofacitinib twice a day showed a favorable clinical response (131). However, these clinical responses were not repeated in patients with moderate-to-severe CD in another phase II trial (132). Currently, various kinds of JAK inhibitors are now in development and awaiting clinical results.

In IBD patients, defective tumor necrosis factor (TGF)- $\beta$ 1 activity is related to up-regulation of SMAD7.


Inhibition of TGF- $\beta$ 1 in healthy human intestines results in increased production of proinflammatory cytokines (133). In a double-blind, placebo-controlled, phase 2 trial, an oral SMAD7 anti-sense oligonucleotide, mongersen, showed a better clinical remission rate in patients with CD compared to that in patients who received a placebo (134).

Several approaches involving the targeting of differentiation and activation of T cells have been attempted. However, a cytotoxic T-lymphocyte antigen 4 (CTLA4) agonist, abatacept, which blocks APC and T cell interaction, showed no clinical benefits in regards to either remission induction or maintenance in patients with IBD (135). Similarly, a humanized monoclonal antibody to CD3 on the activated T cell, visilizumab, was shown to have no clinical benefits for treatment of IBD in phase III

study (91).

### Inhibition of lymphocyte trafficking

Effector lymphocytes must travel from the periphery to the intestine for development of IBD. In this process, various adhesion molecules act on specific lymphocytes. Different lymphocytes express specific cell surface adhesion molecules targeting specific organs. Therefore, selective inhibition of those adhesion molecules potentially has a therapeutic role for IBD. Natalizumab is a humanized monoclonal antibody that binds the  $\alpha$ 4 subunit of integrin on T cells. Theoretically, gut homing of T cell results from the interaction between  $\alpha$ 4 $\beta$ 7 integrin and mucosal vascular addressing cell adhesion molecule 1 (MAdCAM-1). Natalizumab showed clinical benefits for remission maintenance (136). However, natalizumab also




**Figure 1.** Intestinal immune system. IL, interleukin; IFN, interferon; TNF, tumor necrosis factor; TGF, transforming growth factor; Th, helper T cell; Treg, regulatory T cell; TCR, T cell receptor; NF- $\kappa$ B, nuclear factor kappa-light-chain-enhancer of activated B cell; TLR, toll-like receptor; NOD, nucleotide oligomerization domain.

blocks  $\alpha 4\beta 1$  integrin, which is important for T cell homing to the central nervous system, and fatal complications such as progressive multifocal leukoencephalopathy have emerged as a result (137). Therefore, using natalizumab for IBD treatment is restricted in some countries such as the USA. A more recently developed intestine-specific, anti-adhesion molecule, vedolizumab (monoclonal antibody against  $\alpha 4\beta 7$ ), showed promising results for inducing and maintaining remission in UC, and clinical remission in CD, with a relatively good safety profile (138,139). Etrolizumab, a monoclonal antibody against the  $\beta 7$  subunit of integrin, is under phase III studies for CD and UC. Etrolizumab acts as dual inhibitor of the  $\alpha 4\beta 7$ -MAdCAM-1 and  $\alpha E\beta 7$ -E-cadherin interactions. Therefore, etrolizumab prevents both gut homing of lymphocytes and intraepithelial leukocyte retention of intestinal mucosa. A recent phase II trial in UC patients treated with etrolizumab showed a significantly higher clinical remission rate than that in the placebo group (140). Direct inhibition of MAdCAM-1 by its monoclonal antibody, PF-00547659, was also developed

and attempted in clinical trials. Recently, mixed results were obtained, with one study showing significantly higher clinical remission rate in UC (TURANDOT study) compared to that in the placebo group, whereas another study yielded negative results for CD (OPERA study) (141,142).

**CONCLUSION**

As IBD-related research progresses, understanding of these diseases is deepening (Fig. 1). However, it is not believed that only one obvious mechanism of disease will be readily apparent. With the opening of the era of biologics, it has become possible to expect deep remission in IBD patients, unlike in the past; however, about one-third of patients still do not show clinical improvement to the biological agents. A variety of new biologics specific to IBD pathogenesis are now emerging and under clinical investigation (Fig. 2). With this development, more and more patients will benefit from these novel agents. More-



**Figure 2.** Biologics regarding therapeutic targets (black: showed benefits; violet: no benefits). APC, antigen presenting cell; IEC, intestinal epithelial cell; TNF, tumor necrosis factor; MHC, major histocompatibility complex; TCR, T cell receptor; JAK, Janus kinase; TGF, transforming growth factor; IL, interleukin; MAdCAM, mucosal vascular addressing cell adhesion molecule.



over, future IBD therapy should be approached in terms of “patient-customized treatment,” and it is anticipated that it will be a great help in clinical practice to have a drug repertoire targeting various mechanisms of the disease.

## ACKNOWLEDGEMENTS

This study was funded by the grant 15182MFDS507 from the Ministry of Food and Drug Safety, Korea.

## CONFLICTS OF INTEREST

All authors have no conflicts of interest to declare.

## REFERENCES

- Schulenburg, H., C. L. Kurz, and J. J. Ewbank. 2004. Evolution of the innate immune system: the worm perspective. *Immunol. Rev.* 198: 36-58.
- Maloy, K. J., and F. Powrie. 2011. Intestinal homeostasis and its breakdown in inflammatory bowel disease. *Nature* 474: 298-306.
- Cader, M. Z., and A. Kaser. 2013. Recent advances in inflammatory bowel disease: mucosal immune cells in intestinal inflammation. *Gut* 62: 1653-1664.
- Podolsky, D. K. 1991. Inflammatory bowel disease. *N. Engl. J. Med.* 325: 928-937.
- Abraham, C., and J. H. Cho. 2009. Inflammatory bowel disease. *N. Engl. J. Med.* 361: 2066-2078.
- Cheon, J. H. 2013. Genetics of inflammatory bowel diseases: a comparison between Western and Eastern perspectives. *J. Gastroenterol. Hepatol.* 28: 220-226.
- Papadakis, K. A., and S. R. Targan. 2000. Role of cytokines in the pathogenesis of inflammatory bowel disease. *Annu. Rev. Med.* 51: 289-298.
- Cheon, J. H. 2016. Understanding the complications of anti-TNF therapy in East Asian patients with inflammatory bowel disease. *J. Gastroenterol. Hepatol.* doi: 10.1111/jgh.13612.
- Choi, C. H., I. D. Song, Y. H. Kim, J. S. Koo, Y. S. Kim, J. S. Kim, N. Kim, E. S. Kim, J. H. Kim, J. W. Kim, T. O. Kim, H. S. Kim, H. J. Kim, Y. S. Park, D. I. Park, S. J. Park, H. J. Song, S. J. Shin, S. K. Yang, B. D. Ye, K. M. Lee, B. I. Lee, S. Y. Lee, C. K. Lee, J. P. Im, B. I. Jang, T. J. Jeon, Y. K. Cho, S. K. Chang, S. R. Jeon, S. A. Jung, Y. T. Jeon, J. M. Cha, D. S. Han, and W. H. Kim. 2016. Efficacy and safety of infliximab therapy and predictors of response in Korean patients with Crohn's disease: A nationwide, multicenter study. *Yonsei Med. J.* 57: 1376-1385.
- van der Flier, L. G., and H. Clevers. 2009. Stem cells, self-renewal, and differentiation in the intestinal epithelium. *Annu. Rev. Physiol* 71: 241-260.
- Geremia, A., P. Biancheri, P. Allan, G. R. Corazza, and S. A. Di. 2014. Innate and adaptive immunity in inflammatory bowel disease. *Autoimmun. Rev.* 13: 3-10.
- Van der, S. M., B. A. De Koning, A. C. De Bruijn, A. Velcich, J. P. Meijerink, J. B. Van Goudoever, H. A. Buller, J. Dekker, S. Van, I. B. Renes, and A. W. Einerhand. 2006. Muc2-deficient mice spontaneously develop colitis, indicating that MUC2 is critical for colonic protection. *Gastroenterology* 131: 117-129.
- Heazlewood, C. K., M. C. Cook, R. Eri, G. R. Price, S. B. Tauro, D. Taupin, D. J. Thornton, C. W. Png, T. L. Crockford, R. J. Cornall, R. Adams, M. Kato, K. A. Nelms, N. A. Hong, T. H. Florin, C. C. Goodnow, and M. A. McGuckin. 2008. Aberrant mucin assembly in mice causes endoplasmic reticulum stress and spontaneous inflammation resembling ulcerative colitis. *PLoS Med.* 5: e54.
- Jass, J. R., and M. D. Walsh. 2001. Altered mucin expression in the gastrointestinal tract: a review. *J. Cell. Mol. Med.* 5: 327-351.
- Kaser, A., S. Zeissig, and R. S. Blumberg. 2010. Inflammatory bowel disease. *Annu. Rev. Immunol.* 28: 573-621.
- Kaser, A., and R. S. Blumberg. 2010. Endoplasmic reticulum stress and intestinal inflammation. *Mucosal Immunol.* 3: 11-16.
- Cao, S. S., E. M. Zimmermann, B. M. Chuang, B. Song, A. Nwokoye, J. E. Wilkinson, K. A. Eaton, and R. J. Kaufman. 2013. The unfolded protein response and chemical chaperones reduce protein misfolding and colitis in mice. *Gastroenterology* 144: 989-1000.
- Wallace, K. L., L. B. Zheng, Y. Kanazawa, and D. Q. Shih. 2014. Immunopathology of inflammatory bowel disease. *World J. Gastroenterol.* 20: 6-21.
- Pelteková, V. D., R. F. Wintle, L. A. Rubin, C. I. Amos, Q. Huang, X. Gu, B. Newman, O. M. Van, D. Cescon, G. Greenberg, A. M. Griffiths, P. H. St George-Hyslop, and K. A. Siminovitch. 2004. Functional variants of OCTN cation transporter genes are associated with Crohn disease. *Nat. Genet.* 36: 471-475.
- Park, H. J., E. S. Jung, K. A. Kong, E. M. Park, J. H. Cheon, and J. H. Choi. 2016. Identification of OCTN2 variants and their association with phenotypes of Crohn's disease in a Korean population. *Sci. Rep.* 6: 22887.
- Slack, E., S. Hapfelmeier, B. Stecher, Y. Velykoredko, M. Stoel, M. A. Lawson, M. B. Geuking, B. Beutler, T. F. Tedder, W. D. Hardt, P. Bercik, E. F. Verdu, K. D. McCoy, and A. J.

- Macpherson. 2009. Innate and adaptive immunity cooperate flexibly to maintain host-microbiota mutualism. *Science* 325: 617-620.
22. Sartor, R. B. 2006. Mechanisms of disease: pathogenesis of Crohn's disease and ulcerative colitis. *Nat. Clin. Pract. Gastroenterol. Hepatol.* 3: 390-407.
  23. Hermiston, M. L., and J. I. Gordon. 1995. Inflammatory bowel disease and adenomas in mice expressing a dominant negative N-cadherin. *Science* 270: 1203-1207.
  24. Natividad, J. M., V. Petit, X. Huang, P. G. de, J. Jury, Y. Sanz, D. Philpott, C. L. Garcia Rodenas, K. D. McCoy, and E. F. Verdu. 2012. Commensal and probiotic bacteria influence intestinal barrier function and susceptibility to colitis in Nod1<sup>-/-</sup>; Nod2<sup>-/-</sup> mice. *Inflamm. Bowel Dis.* 18: 1434-1446.
  25. Barrett, J. C., J. C. Lee, C. W. Lees, N. J. Prescott, C. A. Anderson, A. Phillips, E. Wesley, K. Parnell, H. Zhang, H. Drummond, E. R. Nimmo, D. Massey, K. Blaszczyk, T. Elliott, L. Cotterill, H. Dallal, A. J. Lobo, C. Mowat, J. D. Sanderson, D. P. Jewell, W. G. Newman, C. Edwards, T. Ahmad, J. C. Mansfield, J. Satsangi, M. Parkes, C. G. Mathew, P. Donnelly, L. Peltonen, J. M. Blackwell, E. Bramon, M. A. Brown, J. P. Casas, A. Corvin, N. Craddock, P. Deloukas, A. Duncanson, J. Jankowski, H. S. Markus, C. G. Mathew, M. I. McCarthy, C. N. Palmer, R. Plomin, A. Rautanen, S. J. Sawcer, N. Samani, R. C. Trembath, A. C. Viswanathan, N. Wood, C. C. Spencer, J. C. Barrett, C. Bellenguez, D. Davison, C. Freeman, A. Strange, P. Donnelly, C. Langford, S. E. Hunt, S. Edkins, R. Gwilliam, H. Blackburn, S. J. Bumpstead, S. Dronov, M. Gillman, E. Gray, N. Hammond, A. Jayakumar, O. T. McCann, J. Liddle, M. L. Perez, S. C. Potter, R. Ravindrarajah, M. Ricketts, M. Waller, P. Weston, S. Widaa, P. Whittaker, P. Deloukas, L. Peltonen, C. G. Mathew, J. M. Blackwell, M. A. Brown, A. Corvin, M. I. McCarthy, C. C. Spencer, A. P. Attwood, J. Stephens, J. Sambrook, W. H. Ouwehand, W. L. McArdle, S. M. Ring, and D. P. Strachan. 2009. Genome-wide association study of ulcerative colitis identifies three new susceptibility loci, including the HNF4A region. *Nat. Genet.* 41: 1330-1334.
  26. Silverberg, M. S., J. H. Cho, J. D. Rioux, D. P. McGovern, J. Wu, V. Annese, J. P. Achkar, P. Goyette, R. Scott, W. Xu, M. M. Barmada, L. Klei, M. J. Daly, C. Abraham, T. M. Bayless, F. Bossa, A. M. Griffiths, A. F. Ippoliti, R. G. Lahaie, A. Latiano, P. Pare, D. D. Proctor, M. D. Regueiro, A. H. Steinhardt, S. R. Targan, L. P. Schumm, E. O. Kistner, A. T. Lee, P. K. Gregersen, J. I. Rotter, S. R. Brant, K. D. Taylor, K. Roeder, and R. H. Duerr. 2009. Ulcerative colitis-risk loci on chromosomes 1p36 and 12q15 found by genome-wide association study. *Nat. Genet.* 41: 216-220.
  27. Stremmel, W., and A. Gauss. 2013. Lecithin as a therapeutic agent in ulcerative colitis. *Dig. Dis.* 31: 388-390.
  28. Stremmel, W., U. Merle, A. Zahn, F. Autschbach, U. Hinz, and R. Ehehalt. 2005. Retarded release phosphatidylcholine benefits patients with chronic active ulcerative colitis. *Gut* 54: 966-971.
  29. Quigley, E. M., and R. Quera. 2006. Small intestinal bacterial overgrowth: roles of antibiotics, prebiotics, and probiotics. *Gastroenterology* 130: S78-S90.
  30. Pickard, K. M., A. R. Bremner, J. N. Gordon, and T. T. MacDonald. 2004. Microbial-gut interactions in health and disease. Immune responses. *Best Pract. Res. Clin. Gastroenterol.* 18: 271-285.
  31. Backhed, F., R. E. Ley, J. L. Sonnenburg, D. A. Peterson, and J. I. Gordon. 2005. Host-bacterial mutualism in the human intestine. *Science* 307: 1915-1920.
  32. Eckburg, P. B., E. M. Bik, C. N. Bernstein, E. Purdom, L. Dethlefsen, M. Sargent, S. R. Gill, K. E. Nelson, and D. A. Relman. 2005. Diversity of the human intestinal microbial flora. *Science* 308: 1635-1638.
  33. Eckburg, P. B., and D. A. Relman. 2007. The role of microbes in Crohn's disease. *Clin. Infect. Dis.* 44: 256-262.
  34. Frank, D. N., A. L. St Amand, R. A. Feldman, E. C. Boedeker, N. Harpaz, and N. R. Pace. 2007. Molecular-phylogenetic characterization of microbial community imbalances in human inflammatory bowel diseases. *Proc. Natl. Acad. Sci. U. S. A.* 104: 13780-13785.
  35. Nagao-Kitamoto, H., S. Kitamoto, P. Kuffa, and N. Kamada. 2016. Pathogenic role of the gut microbiota in gastrointestinal diseases. *Intest. Res.* 14: 127-138.
  36. Khan, K. J., T. A. Ullman, A. C. Ford, M. T. Abreu, A. Abadir, J. K. Marshall, N. J. Talley, and P. Moayyedi. 2011. Antibiotic therapy in inflammatory bowel disease: a systematic review and meta-analysis. *Am. J. Gastroenterol.* 106:661-673.
  37. Mitsuyama, K., M. Niwa, H. Takedatsu, H. Yamasaki, K. Kuwaki, S. Yoshioka, R. Yamauchi, S. Fukunaga, and T. Torimura. 2016. Antibody markers in the diagnosis of inflammatory bowel disease. *World J. Gastroenterol.* 22: 1304-1310.
  38. Hugot, J. P., M. Chamaillard, H. Zouali, S. Lesage, J. P. Cezard, J. Belaiche, S. Almer, C. Tysk, C. A. O'Morain, M. Gassull, V. Binder, Y. Finkel, A. Cortot, R. Modigliani, P. Laurent-Puig, C. Gower-Rousseau, J. Macry, J. F. Colombel, M. Sahbatou, and G. Thomas. 2001. Association of NOD2 leucine-rich repeat variants with susceptibility to Crohn's disease. *Nature* 411: 599-603.
  39. Duerr, R. H., K. D. Taylor, S. R. Brant, J. D. Rioux, M. S. Silverberg, M. J. Daly, A. H. Steinhardt, C. Abraham, M. Regueiro, A. Griffiths, T. Dassopoulos, A. Bitton, H. Yang, S. Targan, L. W. Datta, E. O. Kistner, L. P. Schumm, A. T. Lee, P. K. Gregersen, M. M. Barmada, J. I. Rotter, D. L. Nicolae, and J. H. Cho. 2006. A genome-wide association study identifies

- IL23R as an inflammatory bowel disease gene. *Science* 314: 1461-1463.
40. Elson, C. O., Y. Cong, V. J. McCracken, R. A. Dimmitt, R. G. Lorenz, and C. T. Weaver. 2005. Experimental models of inflammatory bowel disease reveal innate, adaptive, and regulatory mechanisms of host dialogue with the microbiota. *Immunol. Rev.* 206: 260-276.
  41. Yang, J. Y., M. S. Kim, E. Kim, J. H. Cheon, Y. S. Lee, Y. Kim, S. H. Lee, S. U. Seo, S. H. Shin, S. S. Choi, B. Kim, S. Y. Chang, H. J. Ko, J. W. Bae, and M. N. Kweon. 2016. Enteric viruses ameliorate gut inflammation via Toll-like receptor 3 and Toll-like receptor 7-mediated interferon-beta production. *Immunity* 44: 889-900.
  42. Iliiev, I. D., V. A. Funari, K. D. Taylor, Q. Nguyen, C. N. Reyes, S. P. Strom, J. Brown, C. A. Becker, P. R. Fleshner, M. Dubinsky, J. I. Rotter, H. L. Wang, D. P. McGovern, G. D. Brown, and D. M. Underhill. 2012. Interactions between commensal fungi and the C-type lectin receptor Dectin-1 influence colitis. *Science* 336: 1314-1317.
  43. Baumgart, M., B. Dogan, M. Rishniw, G. Weitzman, B. Bosworth, R. Yantiss, R. H. Orsi, M. Wiedmann, P. McDonough, S. G. Kim, D. Berg, Y. Schukken, E. Scherl, and K. W. Simpson. 2007. Culture independent analysis of ileal mucosa reveals a selective increase in invasive *Escherichia coli* of novel phylogeny relative to depletion of Clostridiales in Crohn's disease involving the ileum. *ISME J.* 1: 403-418.
  44. Martin, H. M., B. J. Campbell, C. A. Hart, C. Mpofo, M. Nayar, R. Singh, H. Englyst, H. F. Williams, and J. M. Rhodes. 2004. Enhanced *Escherichia coli* adherence and invasion in Crohn's disease and colon cancer. *Gastroenterology* 127: 80-93.
  45. Sasaki, M., S. V. Sitaraman, B. A. Babbin, P. Gerner-Smidt, E. M. Ribot, N. Garrett, J. A. Alpern, A. Akyildiz, A. L. Theiss, A. Nusrat, and J. M. Klaproth. 2007. Invasive *Escherichia coli* are a feature of Crohn's disease. *Lab. Invest.* 87: 1042-1054.
  46. Darfeuille-Michaud, A. 2002. Adherent-invasive *Escherichia coli*: a putative new *E. coli* pathotype associated with Crohn's disease. *Int. J. Med. Microbiol.* 292:185-193.
  47. Martinez-Medina, M., X. Aldeguer, M. Lopez-Siles, F. Gonzalez-Huix, C. Lopez-Oliu, G. Dahbi, J. E. Blanco, J. Blanco, L. J. Garcia-Gil, and A. rfeuille-Michaud. 2009. Molecular diversity of *Escherichia coli* in the human gut: new ecological evidence supporting the role of adherent-invasive *E. coli* (AIEC) in Crohn's disease. *Inflamm. Bowel Dis.* 15: 872-882.
  48. Manichanh, C., L. Rigottier-Gois, E. Bonnaud, K. Gloux, E. Pelletier, L. Frangeul, R. Nalin, C. Jarrin, P. Chardon, P. Marteau, J. Roca, and J. Dore. 2006. Reduced diversity of faecal microbiota in Crohn's disease revealed by a metagenomic approach. *Gut* 55: 205-211.
  49. Darfeuille-Michaud, A., J. Boudeau, P. Bulois, C. Neut, A. L. Glasser, N. Barnich, M. A. Bringer, A. Swidsinski, L. Beaugerie, and J. F. Colombel. 2004. High prevalence of adherent-invasive *Escherichia coli* associated with ileal mucosa in Crohn's disease. *Gastroenterology* 127: 412-421.
  50. Atarashi, K., T. Tanoue, T. Shima, A. Imaoka, T. Kuwahara, Y. Momose, G. Cheng, S. Yamasaki, T. Saito, Y. Ohba, T. Taniguchi, K. Takeda, S. Hori, I. I. Ivanov, Y. Umesaki, K. Itoh, and K. Honda. 2011. Induction of colonic regulatory T cells by indigenous *Clostridium* species. *Science* 331: 337-341.
  51. Honda, K., and D. R. Littman. 2012. The microbiome in infectious disease and inflammation. *Annu. Rev. Immunol.* 30: 759-795.
  52. Feuerer, M., J. A. Hill, K. Kretschmer, B. H. von, D. Mathis, and C. Benoist. 2010. Genomic definition of multiple ex vivo regulatory T cell subphenotypes. *Proc. Natl. Acad. Sci. U. S. A.* 107: 5919-5924.
  53. Cong, Y., T. Feng, K. Fujihashi, T. R. Schoeb, and C. O. Elson. 2009. A dominant, coordinated T regulatory cell-IgA response to the intestinal microbiota. *Proc. Natl. Acad. Sci. U. S. A.* 106: 19256-19261.
  54. Sokol, H., B. Pigneur, L. Watterlot, O. Lakhdari, L. G. Bermudez-Humaran, J. J. Gratadoux, S. Blugeon, C. Bridonneau, J. P. Furet, G. Corthier, C. Grangette, N. Vasquez, P. Pochart, G. Trugnan, G. Thomas, H. M. Blottiere, J. Dore, P. Marteau, P. Seksik, and P. Langella. 2008. Faecalibacterium *prausnitzii* is an anti-inflammatory commensal bacterium identified by gut microbiota analysis of Crohn disease patients. *Proc. Natl. Acad. Sci. U. S. A.* 105: 16731-16736.
  55. Willing, B., J. Halfvarson, J. Dicksved, M. Rosenquist, G. Jarnerot, L. Engstrand, C. Tysk, and J. K. Jansson. 2009. Twin studies reveal specific imbalances in the mucosa-associated microbiota of patients with ileal Crohn's disease. *Inflamm. Bowel Dis.* 15: 653-660.
  56. Moayyedi, P., M. G. Surette, P. T. Kim, J. Libertucci, M. Wolfe, C. Onischi, D. Armstrong, J. K. Marshall, Z. Kassam, W. Reinisch, and C. H. Lee. 2015. Fecal microbiota transplantation induces remission in patients with active ulcerative colitis in a randomized controlled trial. *Gastroenterology* 149: 102-109.
  57. Rossen, N. G., S. Fuentes, M. J. van der Spek, J. G. Tijssen, J. H. Hartman, A. Duflo, M. Lowenberg, G. R. van den Brink, E. M. Mathus-Vliegen, W. M. de Vos, E. G. Zoetendal, G. R. D'Haens, and C. Y. Ponsioen. 2015. Findings from a randomized controlled trial of fecal transplantation for patients with ulcerative colitis. *Gastroenterology* 149: 110-118.
  58. Colman, R. J., and D. T. Rubin. 2014. Fecal microbiota

- transplantation as therapy for inflammatory bowel disease: a systematic review and meta-analysis. *J. Crohns. Colitis.* 8: 1569-1581.
59. Rachmilewitz, D., K. Katakura, F. Karmeli, T. Hayashi, C. Reinus, B. Rudensky, S. Akira, K. Takeda, J. Lee, K. Takabayashi, and E. Raz. 2004. Toll-like receptor 9 signaling mediates the anti-inflammatory effects of probiotics in murine experimental colitis. *Gastroenterology* 126:520-528.
  60. Shen, J., Z. X. Zuo, and A. P. Mao. 2014. Effect of probiotics on inducing remission and maintaining therapy in ulcerative colitis, Crohn's disease, and pouchitis: meta-analysis of randomized controlled trials. *Inflamm. Bowel Dis.* 20: 21-35.
  61. Kawai, T., and S. Akira. 2011. Toll-like receptors and their crosstalk with other innate receptors in infection and immunity. *Immunity* 34: 637-650.
  62. Akira, S., S. Uematsu, and O. Takeuchi. 2006. Pathogen recognition and innate immunity. *Cell* 124: 783-801.
  63. Podolsky, D. K., G. Gerken, A. Eyking, and E. Cario. 2009. Colitis-associated variant of TLR2 causes impaired mucosal repair because of TFF3 deficiency. *Gastroenterology* 137: 209-220.
  64. Cho, J. H., and S. R. Brant. 2011. Recent insights into the genetics of inflammatory bowel disease. *Gastroenterology* 140: 1704-1712.
  65. Jostins, L., S. Ripke, R. K. Weersma, R. H. Duerr, D. P. McGovern, K. Y. Hui, J. C. Lee, L. P. Schumm, Y. Sharma, C. A. Anderson, J. Essers, M. Mitrovic, K. Ning, I. Cleynen, E. Theate, S. L. Spain, S. Raychaudhuri, P. Goyette, Z. Wei, C. Abraham, J. P. Achkar, T. Ahmad, L. Amininejad, A. N. Ananthakrishnan, V. Andersen, J. M. Andrews, L. Baidoo, T. Balschun, P. A. Bampton, A. Bitton, G. Boucher, S. Brand, C. Buning, A. Cohain, S. Cichon, M. D'Amato, J. D. De, K. L. Devaney, M. Dubinsky, C. Edwards, D. Ellinghaus, L. R. Ferguson, D. Franchimont, K. Fransen, R. Gearry, M. Georges, C. Gieger, J. Glas, T. Haritunians, A. Hart, C. Hawkey, M. Hedl, X. Hu, T. H. Karlsen, L. Kupcinskas, S. Kugathasan, A. Latiano, D. Laukens, I. C. Lawrance, C. W. Lees, E. Louis, G. Mahy, J. Mansfield, A. R. Morgan, C. Mowat, W. Newman, O. Palmieri, C. Y. Ponsioen, U. Potocnik, N. J. Prescott, M. Regueiro, J. I. Rotter, R. K. Russell, J. D. Sanderson, M. Sans, J. Satsangi, S. Schreiber, L. A. Simms, J. Sventoraityte, S. R. Targan, K. D. Taylor, M. Tremelling, H. W. Verspaget, V. M. De, C. Wijmenga, D. C. Wilson, J. Winkelmann, R. J. Xavier, S. Zeissig, B. Zhang, C. K. Zhang, H. Zhao, M. S. Silverberg, V. Annesse, H. Hakonarson, S. R. Brant, G. Radford-Smith, C. G. Mathew, J. D. Rioux, E. E. Schadt, M. J. Daly, A. Franke, M. Parkes, S. Vermeire, J. C. Barrett, and J. H. Cho. 2012. Host-microbe interactions have shaped the genetic architecture of inflammatory bowel disease. *Nature* 491:119-124.
  66. Steinbach, E. C., and S. E. Plevy. 2014. The role of macrophages and dendritic cells in the initiation of inflammation in IBD. *Inflamm. Bowel Dis.* 20: 166-175.
  67. Hampe, J., A. Franke, P. Rosenstiel, A. Till, M. Teuber, K. Huse, M. Albrecht, G. Mayr, L. De, V. J. Briggs, S. Gunther, N. J. Prescott, C. M. Onnie, R. Hasler, B. Sipos, U. R. Folsch, T. Lengauer, M. Platzer, C. G. Mathew, M. Krawczak, and S. Schreiber. 2007. A genome-wide association scan of nonsynonymous SNPs identifies a susceptibility variant for Crohn disease in ATG16L1. *Nat. Genet.* 39: 207-211.
  68. Rioux, J. D., R. J. Xavier, K. D. Taylor, M. S. Silverberg, P. Goyette, A. Huett, T. Green, P. Kuballa, M. M. Barnada, L. W. Datta, Y. Y. Shugart, A. M. Griffiths, S. R. Targan, A. F. Ippoliti, E. J. Bernard, L. Mei, D. L. Nicolae, M. Regueiro, L. P. Schumm, A. H. Steinhart, J. I. Rotter, R. H. Duerr, J. H. Cho, M. J. Daly, and S. R. Brant. 2007. Genome-wide association study identifies new susceptibility loci for Crohn disease and implicates autophagy in disease pathogenesis. *Nat. Genet.* 39: 596-604.
  69. Libioulle, C., E. Louis, S. Hansoul, C. Sandor, F. Farnir, D. Franchimont, S. Vermeire, O. Dewit, V. M. de, A. Dixon, B. Demarche, I. Gut, S. Heath, M. Foglio, L. Liang, D. Laukens, M. Mni, D. Zelenika, G. A. Van, P. Rutgeerts, J. Belaiche, M. Lathrop, and M. Georges. 2007. Novel Crohn disease locus identified by genome-wide association maps to a gene desert on 5p13.1 and modulates expression of PTGER4. *PLoS Genet.* 3: e58.
  70. Parkes, M., J. C. Barrett, N. J. Prescott, M. Tremelling, C. A. Anderson, S. A. Fisher, R. G. Roberts, E. R. Nimmo, F. R. Cummings, D. Soars, H. Drummond, C. W. Lees, S. A. Khawaja, R. Bagnall, D. A. Burke, C. E. Todhunter, T. Ahmad, C. M. Onnie, W. McArdle, D. Strachan, G. Bethel, C. Bryan, C. M. Lewis, P. Deloukas, A. Forbes, J. Sanderson, D. P. Jewell, J. Satsangi, J. C. Mansfield, L. Cardon, and C. G. Mathew. 2007. Sequence variants in the autophagy gene IRGM and multiple other replicating loci contribute to Crohn's disease susceptibility. *Nat. Genet.* 39: 830-832.
  71. Moon, C. M., D. J. Shin, S. W. Kim, N. H. Son, A. Park, B. Park, E. S. Jung, E. S. Kim, S. P. Hong, T. I. Kim, W. H. Kim, and J. H. Cheon. 2013. Associations between genetic variants in the IRGM gene and inflammatory bowel diseases in the Korean population. *Inflamm. Bowel Dis.* 19: 106-114.
  72. Uhlig, H. H., B. S. McKenzie, S. Hue, C. Thompson, B. Joyce-Shaikh, R. Stepankova, N. Robinson, S. Buonocore, H. Tlaskalova-Hogenova, D. J. Cua, and F. Powrie. 2006. Differential activity of IL-12 and IL-23 in mucosal and systemic innate immune pathology. *Immunity* 25: 309-318.
  73. Vong, L., J. G. Ferraz, R. Panaccione, P. L. Beck, and J. L. Wallace. 2010. A pro-resolution mediator, prostaglandin D(2), is specifically up-regulated in individuals in long-term



- remission from ulcerative colitis. *Proc. Natl. Acad. Sci. U. S. A.* 107: 12023-12027.
74. Friedman, D. J., B. M. Kunzli, Y. I. Rahim, J. Sevigny, P. O. Berberat, K. Enjoji, E. Csizmadia, H. Friess, and S. C. Robson. 2009. From the Cover: CD39 deletion exacerbates experimental murine colitis and human polymorphisms increase susceptibility to inflammatory bowel disease. *Proc. Natl. Acad. Sci. U. S. A.* 106: 16788-16793.
  75. Dieckgraefe, B. K., J. R. Korzenik, and S. Anant. 2006. Growth factors as treatment options for intestinal inflammation. *Ann. N. Y. Acad. Sci.* 1072: 300-306.
  76. Weaver, C. T., C. O. Elson, L. A. Fouser, and J. K. Kolls. 2013. The Th17 pathway and inflammatory diseases of the intestines, lungs, and skin. *Annu. Rev. Pathol.* 8: 477-512.
  77. Neurath, M. F. 2014. Cytokines in inflammatory bowel disease. *Nat. Rev. Immunol.* 14: 329-342.
  78. Truelove, S. 1954. Cortisone in ulcerative colitis; preliminary report on a therapeutic trial. *Gastroenterologia.* 81: 86-90.
  79. Flammer, J. R., and I. Rogatsky. 2011. Minireview: Glucocorticoids in autoimmunity: unexpected targets and mechanisms. *Mol. Endocrinol.* 25:1075-1086.
  80. Kim, D. H., J. H. Cheon, J. J. Park, J. Y. Yoon, C. M. Moon, S. P. Hong, T. I. Kim, and W. H. Kim. 2013. Clinical outcomes and predictive factors for response after the first course of corticosteroid therapy in patients with Crohn's disease. *Gut Liver* 7: 58-65.
  81. Kornbluth, A., D. H. Present, S. Lichtiger, and S. Hanauer. 1997. Cyclosporin for severe ulcerative colitis: a user's guide. *Am. J. Gastroenterol.* 92: 1424-1428.
  82. Feagan, B. G., R. N. Fedorak, E. J. Irvine, G. Wild, L. Sutherland, A. H. Steinhardt, G. R. Greenberg, J. Koval, C. J. Wong, M. Hopkins, S. B. Hanauer, and J. W. McDonald. 2000. A comparison of methotrexate with placebo for the maintenance of remission in Crohn's disease. *N. Engl. J. Med.* 342: 1627-1632.
  83. Feagan, B. G., J. Rochon, R. N. Fedorak, E. J. Irvine, G. Wild, L. Sutherland, A. H. Steinhardt, G. R. Greenberg, R. Gillies, M. Hopkins, and . 1995. Methotrexate for the treatment of Crohn's disease. The North American Crohn's Study Group Investigators. *N. Engl. J. Med.* 332: 292-297.
  84. Lennard, L. 1992. The clinical pharmacology of 6-mercaptopurine. *Eur. J. Clin. Pharmacol.* 43: 329-339.
  85. Tiede, I., G. Fritz, S. Strand, D. Poppe, R. Dvorsky, D. Strand, H. A. Lehr, S. Wirtz, C. Becker, R. Atreya, J. Mudter, K. Hildner, B. Bartsch, M. Holtmann, R. Blumberg, H. Walczak, H. Iven, P. R. Galle, M. R. Ahmadian, and M. F. Neurath. 2003. CD28-dependent Rac1 activation is the molecular target of azathioprine in primary human CD4<sup>+</sup> T lymphocytes. *J. Clin. Invest* 111: 1133-1145.
  86. Wildenberg, M. E., P. J. Koelink, K. Diederens, A. A. Te Velde, S. C. Wolfkamp, V. J. Nuij, M. P. Peppelenbosch, M. Nobis, O. J. Sansom, K. I. Anderson, C. J. van der Woude, G. R. D'Haens, and G. R. van den Brink. 2016. The ATG16L1 risk allele associated with Crohn's disease results in a Rac1-dependent defect in dendritic cell migration that is corrected by thiopurines. *Mucosal. Immunol.* doi: 10.1038/mi.2016.65.
  87. Oancea, I., R. Movva, I. Das, C. D. Guirre de, V. Schreiber, Y. Yang, A. Purdon, B. Harrington, M. Proctor, R. Wang, Y. Sheng, M. Lobb, R. Lourie, O. Cuiv, J. A. Duley, J. Begun, and T. H. Florin. 2017. Colonic microbiota can promote rapid local improvement of murine colitis by thioguanine independently of T lymphocytes and host metabolism. *Gut* 66: 59-69.
  88. Dewit, O., T. Moreels, F. Baert, H. Peeters, C. Reenaers, V. M. de, H. P. Van, V. Muls, G. Veereman, F. Mana, O. M. Van, J. Holvoet, S. Naegels, H. Piessevaux, Y. Horsmans, and J. L. Gala. 2011. Limitations of extensive TPMT genotyping in the management of azathioprine-induced myelosuppression in IBD patients. *Clin. Biochem.* 44: 1062-1066.
  89. Yang, S. K., M. Hong, J. Baek, H. Choi, W. Zhao, Y. Jung, T. Haritunians, B. D. Ye, K. J. Kim, S. H. Park, S. K. Park, D. H. Yang, M. Dubinsky, I. Lee, D. P. McGovern, J. Liu, and K. Song. 2014. A common missense variant in NUDT15 confers susceptibility to thiopurine-induced leukopenia. *Nat. Genet.* 46: 1017-1020.
  90. Kim, H. S., J. H. Cheon, E. S. Jung, J. Park, S. Aum, S. J. Park, S. Eun, J. Lee, U. Ruther, G. S. Yeo, M. Ma, K. S. Park, T. Naito, Y. Kakuta, J. H. Lee, W. H. Kim, and M. G. Lee. 2016. A coding variant in FTO confers susceptibility to thiopurine-induced leukopenia in East Asian patients with IBD. *Gut.* doi: 10.1136/gutjnl-2016-311921.
  91. Melmed, G. Y., and S. R. Targan. 2010. Future biologic targets for IBD: potentials and pitfalls. *Nat. Rev. Gastroenterol. Hepatol.* 7: 110-117.
  92. Present, D. H., P. Rutgeerts, S. Targan, S. B. Hanauer, L. Mayer, R. A. van Hogezaand, D. K. Podolsky, B. E. Sands, T. Braakman, K. L. DeWoody, T. F. Schaible, and S. J. van Deventer. 1999. Infliximab for the treatment of fistulas in patients with Crohn's disease. *N. Engl. J. Med.* 340: 1398-1405.
  93. Kontoyiannis, D., M. Pasparakis, T. T. Pizarro, F. Cominelli, and G. Kollias. 1999. Impaired on/off regulation of TNF biosynthesis in mice lacking TNF AU-rich elements: implications for joint and gut-associated immunopathologies. *Immunity* 10:387-398.
  94. Kojouharoff, G., W. Hans, F. Obermeier, D. N. Mannel, T. Andus, J. Scholmerich, V. Gross, and W. Falk. 1997. Neutralization of tumour necrosis factor (TNF) but not of IL-1 reduces inflammation in chronic dextran sulphate sodium-induced colitis in mice. *Clin. Exp. Immunol.* 107:

- 353-358.
95. Wajant, H., K. Pfizenmaier, and P. Scheurich. 2003. Tumor necrosis factor signaling. *Cell Death Differ.* 10: 45-65.
96. Neurath, M. F. 2014. New targets for mucosal healing and therapy in inflammatory bowel diseases. *Mucosal Immunol.* 7: 6-19.
97. Holtmann, M. H., E. Douni, M. Schutz, G. Zeller, J. Mudter, H. A. Lehr, J. Gerspach, P. Scheurich, P. R. Galle, G. Kollias, and M. F. Neurath. 2002. Tumor necrosis factor-receptor 2 is up-regulated on lamina propria T cells in Crohn's disease and promotes experimental colitis *in vivo*. *Eur. J. Immunol.* 32: 3142-3151.
98. Perrier, C., H. G. de, J. Cremer, S. Vermeire, P. Rutgeerts, A. G. Van, D. E. Szymkowski, and J. L. Ceuppens. 2013. Neutralization of membrane TNF, but not soluble TNF, is crucial for the treatment of experimental colitis. *Inflamm. Bowel Dis.* 19: 246-253.
99. Colombel, J. F., W. J. Sandborn, W. Reinisch, G. J. Mantzaris, A. Kornbluth, D. Rachmilewitz, S. Lichtiger, G. D'Haens, R. H. Diamond, D. L. Broussard, K. L. Tang, C. J. van der Woude, and P. Rutgeerts. 2010. Infliximab, azathioprine, or combination therapy for Crohn's disease. *N. Engl. J. Med.* 362: 1383-1395.
100. Colombel, J. F., W. J. Sandborn, P. Rutgeerts, R. Enns, S. B. Hanauer, R. Panaccione, S. Schreiber, D. Byczkowski, J. Li, J. D. Kent, and P. F. Pollack. 2007. Adalimumab for maintenance of clinical response and remission in patients with Crohn's disease: the CHARM trial. *Gastroenterology* 132: 52-65.
101. Hanauer, S. B., W. J. Sandborn, P. Rutgeerts, R. N. Fedorak, M. Lukas, D. MacIntosh, R. Panaccione, D. Wolf, and P. Pollack. 2006. Human anti-tumor necrosis factor monoclonal antibody (adalimumab) in Crohn's disease: the CLASSIC-I trial. *Gastroenterology* 130: 323-333.
102. Hanauer, S. B., B. G. Feagan, G. R. Lichtenstein, L. F. Mayer, S. Schreiber, J. F. Colombel, D. Rachmilewitz, D. C. Wolf, A. Olson, W. Bao, and P. Rutgeerts. 2002. Maintenance infliximab for Crohn's disease: the ACCENT I randomised trial. *Lancet* 359: 1541-1549.
103. Colombel, J. F., P. Rutgeerts, W. Reinisch, D. Esser, Y. Wang, Y. Lang, C. W. Marano, R. Strauss, B. J. Oddens, B. G. Feagan, S. B. Hanauer, G. R. Lichtenstein, D. Present, B. E. Sands, and W. J. Sandborn. 2011. Early mucosal healing with infliximab is associated with improved long-term clinical outcomes in ulcerative colitis. *Gastroenterology* 141: 1194-1201.
104. Evans, A. T., and S. D. Lee. 2012. A review and expert opinion of the use of certolizumab for Crohn's disease. *Expert Opin. Biol. Ther.* 12: 363-370.
105. Lowenberg, M., N. K. de Boer, and F. Hoentjen. 2014. Golimumab for the treatment of ulcerative colitis. *Clin. Exp. Gastroenterol.* 7: 53-59.
106. Van den Brande, J. M., H. Braat, G. R. van den Brink, H. H. Versteeg, C. A. Bauer, I. Hoedemaeker, M. C. van, D. W. Hommes, M. P. Peppelenbosch, and S. J. van Deventer. 2003. Infliximab but not etanercept induces apoptosis in lamina propria T-lymphocytes from patients with Crohn's disease. *Gastroenterology* 124: 1774-1785.
107. Sandborn, W. J., S. B. Hanauer, S. Katz, M. Safdi, D. G. Wolf, R. D. Baerg, W. J. Tremaine, T. Johnson, N. N. Diehl, and A. R. Zinsmeister. 2001. Etanercept for active Crohn's disease: a randomized, double-blind, placebo-controlled trial. *Gastroenterology* 121: 1088-1094.
108. Zeissig, S., C. Bojarski, N. Buerge, J. Mankertz, M. Zeitz, M. Fromm, and J. D. Schulzke. 2004. Downregulation of epithelial apoptosis and barrier repair in active Crohn's disease by tumour necrosis factor alpha antibody treatment. *Gut* 53: 1295-1302.
109. Vos, A. C., M. E. Wildenberg, M. Duijvestein, A. P. Verhaar, G. R. van den Brink, and D. W. Hommes. 2011. Anti-tumor necrosis factor-alpha antibodies induce regulatory macrophages in an Fc region-dependent manner. *Gastroenterology* 140: 221-230.
110. Vos, A. C., M. E. Wildenberg, I. Arijis, M. Duijvestein, A. P. Verhaar, H. G. de, S. Vermeire, P. Rutgeerts, G. R. van den Brink, and D. W. Hommes. 2012. Regulatory macrophages induced by infliximab are involved in healing *in vivo* and *in vitro*. *Inflamm. Bowel Dis.* 18: 401-408.
111. Bhol, K. C., D. E. Tracey, B. R. Lemos, G. D. Lyng, E. C. Erlich, D. M. Keane, M. S. Quesenberry, A. D. Holdorf, L. D. Schlehuber, S. A. Clark, and B. S. Fox. 2013. AVX-470: a novel oral anti-TNF antibody with therapeutic potential in inflammatory bowel disease. *Inflamm. Bowel Dis.* 19: 2273-2281.
112. Vandembroucke, K., H. H. de, E. Beirnaert, T. Dreier, M. Lauwereys, L. Huyck, H. J. Van, P. Demetter, L. Steidler, E. Remaut, C. Cuvelier, and P. Rottiers. 2010. Orally administered *L. lactis* secreting an anti-TNF Nanobody demonstrate efficacy in chronic colitis. *Mucosal Immunol.* 3:49-56.
113. Assier, E., L. Semerano, E. Duvallet, L. Delavallee, E. Bernier, M. Laborie, G. Grouard-Vogel, P. Larcier, N. Bessis, and M. C. Boissier. 2012. Modulation of anti-tumor necrosis factor alpha (TNF-alpha) antibody secretion in mice immunized with TNF-alpha kinoid. *Clin. Vaccine Immunol.* 19:699-703.
114. Monteleone, I., F. Pallone, and G. Monteleone. 2011. Th17-related cytokines: new players in the control of chronic intestinal inflammation. *BMC Med.* 9: 122.
115. Zenewicz, L. A., A. Antov, and R. A. Flavell. 2009. CD4

- T-cell differentiation and inflammatory bowel disease. *Trends Mol. Med.* 15: 199-207.
116. Caruso, R., D. Fina, I. Peluso, C. Stolfi, M. C. Fantini, V. Gioia, F. Caprioli, B. G. Del Vecchio, O. A. Paoluzi, T. T. Macdonald, F. Pallone, and G. Monteleone. 2007. A functional role for interleukin-21 in promoting the synthesis of the T-cell chemoattractant, MIP-3alpha, by gut epithelial cells. *Gastroenterology* 132: 166-175.
  117. Huang, Y., and Z. Chen. 2016. Inflammatory bowel disease related innate immunity and adaptive immunity. *Am. J. Transl. Res.* 8: 2490-2497.
  118. Reinisch, W., V. W. de, L. Bene, L. Simon, I. Racz, S. Katz, I. Altorjay, B. Feagan, D. Riff, C. N. Bernstein, D. Hommes, P. Rutgeerts, A. Cortot, M. Gaspari, M. Cheng, T. Pearce, and B. E. Sands. 2010. Fontolizumab in moderate to severe Crohn's disease: a phase 2, randomized, double-blind, placebo-controlled, multiple-dose study. *Inflamm. Bowel Dis.* 16: 233-242.
  119. Hueber, W., B. E. Sands, S. Lewitzky, M. Vandemeulebroecke, W. Reinisch, P. D. Higgins, J. Wehkamp, B. G. Feagan, M. D. Yao, M. Karczewski, J. Karczewski, N. Pezous, S. Bek, G. Bruin, B. Mellgard, C. Berger, M. Londei, A. P. Bertolino, G. Tougas, and S. P. Travis. 2012. Secukinumab, a human anti-IL-17A monoclonal antibody, for moderate to severe Crohn's disease: unexpected results of a randomised, double-blind placebo-controlled trial. *Gut* 61: 1693-1700.
  120. Pickert, G., C. Neufert, M. Leppkes, Y. Zheng, N. Wittkopf, M. Warntjen, H. A. Lehr, S. Hirth, B. Weigmann, S. Wirtz, W. Ouyang, M. F. Neurath, and C. Becker. 2009. STAT3 links IL-22 signaling in intestinal epithelial cells to mucosal wound healing. *J. Exp. Med.* 206: 1465-1472.
  121. Ueno, A., A. Ghosh, D. Hung, J. Li, and H. Jijon. 2015. Th17 plasticity and its changes associated with inflammatory bowel disease. *World J. Gastroenterol.* 21: 12283-12295.
  122. Shih, D. Q., S. R. Targan, and D. McGovern. 2008. Recent advances in IBD pathogenesis: genetics and immunobiology. *Curr. Gastroenterol. Rep.* 10: 568-575.
  123. Feagan, B. G., W. J. Sandborn, C. Gasink, D. Jacobstein, Y. Lang, J. R. Friedman, M. A. Blank, J. Johanns, L. L. Gao, Y. Miao, O. J. Adedokun, B. E. Sands, S. B. Hanauer, S. Vermeire, S. Targan, S. Ghosh, W. J. de Villiers, J. F. Colombel, Z. Tulassay, U. Seidler, B. A. Salzberg, P. Desreumaux, S. D. Lee, E. V. Loftus, Jr., L. A. Dieleman, S. Katz, and P. Rutgeerts. 2016. Ustekinumab as Induction and Maintenance Therapy for Crohn's Disease. *N. Engl. J. Med.* 375: 1946-1960.
  124. Mannon, P. J., I. J. Fuss, L. Mayer, C. O. Elson, W. J. Sandborn, D. Present, B. Dolin, N. Goodman, C. Groden, R. L. Hornung, M. Quezado, Z. Yang, M. F. Neurath, J. Salfeld, G. M. Veldman, U. Schwertschlag, and W. Strober. 2004. Anti-interleukin-12 antibody for active Crohn's disease. *N. Engl. J. Med.* 351:2069-2079.
  125. Sands, B. E., E. W. Jacobson, T. Sylwestrowicz, Z. Younes, G. Dryden, R. Fedorak, and S. Greenbloom. 2010. Randomized, double-blind, placebo-controlled trial of the oral interleukin-12/23 inhibitor apilimod mesylate for treatment of active Crohn's disease. *Inflamm. Bowel Dis.* 16:1209-1218.
  126. Davidson, N. J., M. W. Leach, M. M. Fort, L. Thompson-Snipes, R. Kuhn, W. Muller, D. J. Berg, and D. M. Rennick. 1996. T helper cell 1-type CD4<sup>+</sup> T cells, but not B cells, mediate colitis in interleukin 10-deficient mice. *J. Exp. Med.* 184:241-251.
  127. Erdman, S. E., V. P. Rao, T. Poutahidis, M. M. Ihrig, Z. Ge, Y. Feng, M. Tomczak, A. B. Rogers, B. H. Horwitz, and J. G. Fox. 2003. CD4(+)CD25(+) regulatory lymphocytes require interleukin 10 to interrupt colon carcinogenesis in mice. *Cancer Res.* 63: 6042-6050.
  128. Steidler, L., W. Hans, L. Schotte, S. Neiryck, F. Obermeier, W. Falk, W. Fiers, and E. Remaut. 2000. Treatment of murine colitis by *Lactococcus lactis* secreting interleukin-10. *Science* 289:1352-1355.
  129. Danese, S., M. Grisham, J. Hodge, and J. B. Telliez. 2016. JAK inhibition using tofacitinib for inflammatory bowel disease treatment: a hub for multiple inflammatory cytokines. *Am. J. Physiol. Gastrointest. Liver Physiol* 310: G155-G162.
  130. Ghoreschi, K., A. Laurence, and J. J. O'Shea. 2009. Janus kinases in immune cell signaling. *Immunol. Rev.* 228: 273-287.
  131. Sandborn, W. J., S. Ghosh, J. Panes, I. Vranic, C. Su, S. Rousell, and W. Niezychowski. 2012. Tofacitinib, an oral Janus kinase inhibitor, in active ulcerative colitis. *N. Engl. J. Med.* 367: 616-624.
  132. Sandborn, W. J., S. Ghosh, J. Panes, I. Vranic, W. Wang, and W. Niezychowski. 2014. A phase 2 study of tofacitinib, an oral Janus kinase inhibitor, in patients with Crohn's disease. *Clin. Gastroenterol. Hepatol.* 12: 1485-1493.
  133. Monteleone, G., M. Boirivant, F. Pallone, and T. T. MacDonald. 2008. TGF-beta1 and Smad7 in the regulation of IBD. *Mucosal Immunol.* 1 Suppl 1: S50-S53.
  134. Monteleone, G., M. F. Neurath, S. Ardizzone, S. A. Di, M. C. Fantini, F. Castiglione, M. L. Scribano, A. Armuzzi, F. Caprioli, G. C. Sturniolo, F. Rogai, M. Vecchi, R. Atreya, F. Bossa, S. Onali, M. Fichera, G. R. Corazza, L. Biancone, V. Savarino, R. Pica, A. Orlando, and F. Pallone. 2015. Mongersen, an oral SMAD7 antisense oligonucleotide, and Crohn's disease. *N. Engl. J. Med.* 372: 1104-1113.
  135. Sandborn, W. J., J. F. Colombel, B. E. Sands, P. Rutgeerts, S. R. Targan, R. Panaccione, B. Bressler, K. Geboes, S. Schreiber, R. Aranda, S. Gujrathi, A. Luo, Y. Peng, L. Salter-Cid, and S. B. Hanauer. 2012. Abatacept for Crohn's disease and

- ulcerative colitis. *Gastroenterology* 143: 62-69.
136. Sandborn, W. J., J. F. Colombel, R. Enns, B. G. Feagan, S. B. Hanauer, I. C. Lawrance, R. Panaccione, M. Sanders, S. Schreiber, S. Targan, D. S. van, R. Goldblum, D. Despaigne, G. S. Hogge, and P. Rutgeerts. 2005. Natalizumab induction and maintenance therapy for Crohn's disease. *N. Engl. J. Med.* 353: 1912-1925.
137. Maas, R. P., A. H. Muller-Hansma, R. A. Esselink, J. L. Murk, C. Warnke, J. Killestein, and M. P. Wattjes. 2016. Drug-associated progressive multifocal leukoencephalopathy: a clinical, radiological, and cerebrospinal fluid analysis of 326 cases. *J. Neurol.* 263: 2004-2021.
138. Sandborn, W. J., B. G. Feagan, P. Rutgeerts, S. Hanauer, J. F. Colombel, B. E. Sands, M. Lukas, R. N. Fedorak, S. Lee, B. Bressler, I. Fox, M. Rosario, S. Sankoh, J. Xu, K. Stephens, C. Milch, and A. Parikh. 2013. Vedolizumab as induction and maintenance therapy for Crohn's disease. *N. Engl. J. Med.* 369: 711-721.
139. Feagan, B. G., P. Rutgeerts, B. E. Sands, S. Hanauer, J. F. Colombel, W. J. Sandborn, A. G. Van, J. Axler, H. J. Kim, S. Danese, I. Fox, C. Milch, S. Sankoh, T. Wyant, J. Xu, and A. Parikh. 2013. Vedolizumab as induction and maintenance therapy for ulcerative colitis. *N. Engl. J. Med.* 369: 699-710.
140. Vermeire, S., S. O'Byrne, M. Keir, M. Williams, T. T. Lu, J. C. Mansfield, C. A. Lamb, B. G. Feagan, J. Panes, A. Salas, D. C. Baumgart, S. Schreiber, I. Dotan, W. J. Sandborn, G. W. Tew, D. Luca, M. T. Tang, L. Diehl, J. Eastham-Anderson, H. G. De, C. Perrier, J. G. Egen, J. A. Kirby, A. G. van, and P. Rutgeerts. 2014. Etrolizumab as induction therapy for ulcerative colitis: a randomised, controlled, phase 2 trial. *Lancet* 384: 309-318.
141. Sandborn, W., S. D. Lee, D. Tarabar, E. Louis, M. Klopocka, J. Klaus, W. Reinisch, X. Hebuterne, D. I. Park, and S. Schreiber. 2015. 825 Anti-MAdCAM-1 antibody (PF-00547659) for active refractory Crohn's disease: Results of the OPERA study. *Gastroenterology* 148: S-162.
142. Reinisch, W., W. Sandborn, S. Danese, F. Cataldi, X. Hebuterne, B. Salzberg, M. Klopocka, D. Tarabar, T. Vanasek, and M. Gregus. 2015. 901a A randomized, multicenter double-blind, placebo-controlled study of the safety and efficacy of anti-MAdCAM antibody PF-00547659 (PF) in patients with moderate to severe ulcerative colitis: Results of the TURANDOT study. *Gastroenterology* 148: S-1193.