



Draft Genome Sequence of *Ochrobactrum intermedium* Strain SA148, a Plant Growth-Promoting Desert Rhizobacterium

Feras F. Lafi,^{a,b} Intikhab Alam,^b Rene Geurts,^c Ton Bisseling,^c Vladimir B. Bajic,^b Heribert Hirt,^a  Maged M. Saad^a

King Abdullah University of Science and Technology (KAUST), Biological and Environmental Sciences and Engineering Division (BESE), Thuwal, Kingdom of Saudi Arabia^a; Computational Bioscience Research Center (CBRC), King Abdullah University of Science and Technology (KAUST), Thuwal, Kingdom of Saudi Arabia^b; Department of Plant Sciences, Laboratory of Molecular Biology, Wageningen University, Wageningen, The Netherlands^c

ABSTRACT *Ochrobactrum intermedium* strain SA148 is a plant growth-promoting bacterium isolated from sandy soil in the Jizan area of Saudi Arabia. Here, we report the 4.9-Mb draft genome sequence of this strain, highlighting different pathways characteristic of plant growth promotion activity and environmental adaptation of SA148.

The DARWIN21 project (<http://www.darwin21.net/>) is an initiative established with the aim of expanding our knowledge on microbial life in deserts. A diverse number of bacterial species were isolated from both soil and plants and were characterized for their potential application in modern agriculture. Within the DARWIN21 collection, diverse bacterial species show enhanced growth of the model plant *Arabidopsis thaliana* under stress conditions (1, 2). A number of bacterial *Ochrobactrum* species were isolated on the basis of their plant growth-promoting trait. *Ochrobactrum* species are widespread in the environment and can colonize a wide variety of habitats, including soil, plants, animals, and humans (3). In an attempt to decipher the genetic basis of the plant growth-promoting activity of a member of the *Ochrobactrum* species, the genome of strain SA148 was sequenced.

SA148 strain was isolated from the soil samples collected in the Jizan region (16°56.475'N, 42°36.694'E) in the Kingdom of Saudi Arabia. Based on the 16S rRNA gene sequences, the SA148 strain is closely related (99% gene similarity) to *Ochrobactrum intermedium* strain LMG3301 (accession no. NR_026039.1) (1). The genomic DNA of the SA148 strain was extracted using the Qiagen DNeasy blood and tissue kit, according to the manufacturer's protocol. The DNA was then sequenced using paired-end Illumina MiSeq, and the library was constructed as described previously (4). Contig assembly was done with SPAdes assembler version 3.6 (5), with a 1-kb contig cutoff size. *De novo* assembly of MiSeq reads for strain SA148 resulted in 38 contigs with a total length of 4,918,899 bp, a mean contig size of 129,445 bp, an N_{50} of 419,572 bp, L_{50} reached with five contigs, and a genome G+C content of 57.5%. MegaBLAST (6) searches of strain SA148 concatenated genomes against the NCBI reference genome database (<http://www.ncbi.nlm.nih.gov/genome/>) revealed that the closest relative genomes to strain SA148 was *O. intermedium* strain LMG3301, with sequence coverages of 50% and 35% of chromosomes I and II (accession no. ACQA010000.1) and similarities of 98% and 99%, respectively. The annotation of strain SA148 was carried out using the default INDIGO pipeline (7), with the exception of open reading frame (ORF) prediction by FragGeneScan (8). SA148 has 3,714 ORFs, three rRNA, 51 tRNA, and 64 noncoding RNA

Received 30 December 2016 Accepted 5 January 2017 Published 2 March 2017

Citation Lafi FF, Alam I, Geurts R, Bisseling T, Bajic VB, Hirt H, Saad MM. 2017. Draft genome sequence of *Ochrobactrum intermedium* strain SA148, a plant growth-promoting desert rhizobacterium. *Genome Announc* 5:e01707-16. <https://doi.org/10.1128/genomeA.01707-16>.

Copyright © 2017 Lafi et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Heribert Hirt, heribert.hirt@kaust.edu.sa.

(ncRNA). The annotation predicted a number of enzymes, such as glyphosate resistance enzyme 3-phosphoshikimate 1-carboxyvinyltransferase (EC 2.5.1.19) first identified in *Ochrobactrum anthropi* (9), which is targeted by the herbicide glyphosate. We also detected mannitol-1-phosphate 5-dehydrogenase (EC 1.1.1.17) (gene *mtID*) as a part of an osmoprotectant synthesis pathway that can be utilized to improve stress tolerance against temperature fluctuation (10). Moreover, a number of genes related to growth-promoting activity, including gene-coding clusters for phosphate solubilization with six genes for pyrroloquinoline quinone synthesis (PQQ) (EC 1.1.5.2) and glucose 1-dehydrogenase (GCD) (EC 1.1.5.2) were found. The genome analysis highlights the presence of 3-phytase (EC 3.1.3.8) (11) and acid phosphatase (EC 3.1.3.2) (12), two enzymes that aid plants during phosphate starvation.

Accession number(s). The genome of *Ochrobactrum intermedium* SA148 was deposited at DDBJ/EMBL/GenBank under the accession no. [LWEA00000000](https://www.ncbi.nlm.nih.gov/nuclink/LWEA00000000). The version described in this paper is the first version, LWEA01000000.

ACKNOWLEDGMENTS

Genome sequencing was performed at the Bioscience Core Laboratory of the King Abdullah University of Science and Technology (KAUST), Thuwal, Saudi Arabia. We are grateful for the use of Dragon and Snapdragon compute clusters at the Computational Bioscience Research Center (CBRC) of KAUST.

This work was supported by a base fund research grant to H.H. from the King Abdullah University of Science and Technology (KAUST). Computational aspects of this work have been supported by the KAUST Office of Sponsored Research (OSR) under awards URF/1/1976-02 and FCS/1/2448-01 to V.B.B.

REFERENCES

1. Lafi FF, Bokhari A, Alam I, Bajic VB, Hirt H, Saad MM. 2016. Draft genome sequence of the plant growth-promoting *Cupriavidus gilardii* strain JZ4 isolated from the desert plant *Tribulus terrestris*. *Genome Announc* 4(4): e00678-16. <https://doi.org/10.1128/genomeA.00678-16>.
2. Lafi FF, Ramirez-Prado JS, Alam I, Bajic VB, Hirt H, Saad MM. 2016. Draft genome sequence of *Halomonas elongata* strain K4, an endophytic growth-promoting bacterium enhancing salinity tolerance in planta. *Genome Announc* 4(6):e01214-16. <https://doi.org/10.1128/genomeA.01214-16>.
3. Bathe S, Achouak W, Hartmann A, Heulin T, Schlöter M, Leubner M. 2006. Genetic and phenotypic microdiversity of *Ochrobactrum* spp. *FEMS Microbiol Ecol* 56:272–280. <https://doi.org/10.1111/j.1574-6941.2005.00029.x>.
4. Velasco J, Romero C, López-Goñi I, Leiva J, Díaz R, Moriyón I. 1998. Evaluation of the relatedness of *Brucella* spp. and *Ochrobactrum anthropi* and description of *Ochrobactrum intermedium* sp. nov., a new species with a closer relationship to *Brucella* spp. *Int J Syst Bacteriol* 48:759–768. <https://doi.org/10.1099/00207713-48-3-759>.
5. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshtkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
6. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST plus: architecture and applications. *BMC Bioinformatics* 10:421. <https://doi.org/10.1186/1471-2105-10-421>.
7. Alam I, Antunes A, Kamau AA, Ba Alawi WB, Kalkatawi M, Stingl U, Bajic VB. 2013. INDIGO—INtegrated data warehouse of Microbial Genomes with examples from the Red Sea extremophiles. *PLoS One* 8:e82210. <https://doi.org/10.1371/journal.pone.0082210>.
8. Rho MN, Tang HX, Ye YZ. 2010. FragGeneScan: predicting genes in short and error-prone reads. *Nucleic Acids Res* 38:e191. <https://doi.org/10.1093/nar/gkq747>.
9. Tian YS, Xu J, Xiong AS, Zhao W, Fu XY, Peng RH, Yao QH. 2011. Improvement of glyphosate resistance through concurrent mutations in three amino acids of the *Ochrobactrum* 5-enopyruvylshikimate-3-phosphate synthase. *Appl Environ Microbiol* 77:8409–8414. <https://doi.org/10.1128/AEM.05271-11>.
10. Chiang YJ, Stushnoff C, McSay AE. 2005. Overexpression of mannitol-1-phosphate dehydrogenase increases mannitol accumulation and adds protection against chilling injury in petunia. *J Am Soc Hort Sci* 130:605–610.
11. Vohra A, Satyanarayana T. 2003. Phytases: microbial sources, production, purification, and potential biotechnological applications. *Crit Rev Biotechnol* 23:29–60. <https://doi.org/10.1080/713609297>.
12. Hong YF, Liu CY, Cheng KJ, Hour AL, Chan MT, Tseng TH, Chen KY, Shaw JF, Yu SM. 2008. The sweet potato sporamin promoter confers high-level phytase expression and improves organic phosphorus acquisition and tuber yield of transgenic potato. *Plant Mol Biol* 67:347–361. <https://doi.org/10.1007/s11103-008-9324-6>.