

CORRECTION


Correction: An increase in visceral fat is associated with a decrease in the taste and olfactory capacity

Jose Carlos Fernandez-Garcia, Juan Alcaide, Concepcion Santiago-Fernandez, MM. Roca-Rodriguez, Zaida Aguera, Rosa Baños, Cristina Botella, Rafael de la Torre, Jose M. Fernandez-Real, Gema Fruhbeck, Javier Gomez-Ambrosi, Susana Jimenez-Murcia, Jose M. Menchon, Felipe F. Casanueva, Fernando Fernandez-Aranda, Francisco J. Tinahones, Lourdes Garrido-Sanchez

The following information is missing from the Funding section: This study was supported by the project Instituto de Salud Carlos III (FIS PI15/01114).

Reference

1. Fernandez-Garcia JC, Alcaide J, Santiago-Fernandez C, Roca-Rodriguez M, Aguera Z, Baños R, et al. (2017) An increase in visceral fat is associated with a decrease in the taste and olfactory capacity. PLoS ONE 12(2): e0171204. doi: [10.1371/journal.pone.0171204](https://doi.org/10.1371/journal.pone.0171204) PMID: [28158237](#)



OPEN ACCESS

Citation: Fernandez-Garcia JC, Alcaide J, Santiago-Fernandez C, Roca-Rodriguez M, Aguera Z, Baños R, et al. (2017) Correction: An increase in visceral fat is associated with a decrease in the taste and olfactory capacity. PLoS ONE 12(3): e0173588. doi:10.1371/journal.pone.0173588

Published: March 3, 2017

Copyright: © 2017 Fernandez-Garcia et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.