



Published in final edited form as:

Cancer Epidemiol Biomarkers Prev. 2017 March ; 26(3): 321–327. doi:10.1158/1055-9965.EPI-16-0601.

History of gestational diabetes mellitus and risk of incident invasive breast cancer among parous women in the Nurses' Health Study II prospective cohort

Camille E. Powe^{*,1}, Deirdre K. Tobias^{*,2,3}, Karin B. Michels^{4,5,6}, Wendy Y. Chen^{4,9}, A. Heather Eliassen^{4,6}, JoAnn E. Manson^{2,6,8}, Bernard Rosner⁴, Walter C. Willett^{3,4,6}, Frank B. Hu^{3,4,6}, Cuilin Zhang⁷, Janet W. Rich-Edwards^{4,6,8}, and Kathryn M. Rexrode²

¹Diabetes Unit, Massachusetts General Hospital

²Division of Preventive Medicine, Department of Medicine, Brigham and Women's Hospital

³Department of and Nutrition, Harvard T.H. Chan School of Public Health

⁴Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital

⁵Obstetrics and Gynecology Epidemiology Center, Department of Obstetrics, Gynecology and Reproductive Biology, Brigham and Women's Hospital

⁶Department of Epidemiology, Harvard T.H. Chan School of Public Health

⁷Epidemiology Branch, Division of Intramural Population Health Research, Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health, Rockville, MD

⁸Connors Center for Women's Health and Gender Biology, Brigham and Women's Hospital

⁹Department of Medical Oncology, Dana Farber Cancer Institute

Abstract

Background—Type 2 diabetes is associated with breast cancer in epidemiologic studies. Pregnancy also modifies breast cancer risk. We hypothesized that women with a history of gestational diabetes mellitus (GDM), which shares pathogenesis and risk factors with type 2 diabetes, would have greater invasive breast cancer risk than parous women without a history of GDM.

Methods—We conducted a prospective analysis among parous women in the Nurses' Health Study II, with mean age 35 years in 1989. Multivariate Cox proportional hazard models were used to compare risks of incident invasive breast cancer in women with and without a history of GDM.

Results—Among 86,972 women studied, 5,188 women reported a history of GDM and 2,377 developed invasive breast cancer (100 with history of GDM, 2277 without GDM) over 22 years of

Corresponding author: Kathryn M. Rexrode, 900 Commonwealth Avenue Boston, MA 02215, Krexrode@Partners.ORG, phone: 617-278-0834, fax: 617-731-3843.

*These authors contributed equally (co-first authors)

The authors declare no potential conflicts of interest.

prospective follow-up. History of GDM was inversely associated with incident invasive breast cancer (HR 0.68, 95% CI 0.55-0.84, $p=0.0004$), compared with no history of GDM, after adjustment for body mass index, reproductive history, and other breast cancer risk factors. Findings were similar by menopausal status, although observed person-time was predominantly premenopausal (premenopausal: HR 0.73, 95% CI 0.56-0.96, $p=0.03$; postmenopausal: HR 0.63, 95% CI 0.43-0.92, $p=0.02$). Restricting to women undergoing mammography screening modestly attenuated the relationship (HR 0.74, 95% CI 0.57-0.96, $p=0.02$).

Conclusion—Among a large cohort of US women, history of GDM was not associated with an elevated risk of subsequent invasive breast cancer.

Impact—Our findings highlight the need to further investigate GDM's role in breast cancer development.

Keywords

gestational diabetes; breast cancer; epidemiology; cohort study; diabetes

Introduction

The risk of breast cancer has long been linked to reproductive risk factors including age at menarche, age at first birth, and age at menopause.(1) Pregnancy has bidirectional effects on breast cancer risk: it increases risk over the short-term, but in women with a young age at first birth, it is protective over the long-term.(1, 2) Hormonal changes in pregnancy are thought to induce proliferation and differentiation of the breast epithelial cells, modifying the risk of future malignancy.(3-5) Identifying pregnancy-specific characteristics that modulate breast cancer risk could inform screening and prevention practices.

Type 2 diabetes was associated with a 27% greater overall risk of breast cancer in a meta-analysis of epidemiologic studies, though significant between-study heterogeneity suggested potential differences by menopausal status, adjustment for body mass index (BMI), or study design (prospective vs. retrospective).(6) For example, no positive relationship was observed for diabetes and breast cancer among premenopausal women. Gestational diabetes mellitus (GDM) occurs in ~6% of pregnancies in the United States and confers a high risk of future type 2 diabetes.(7-9) During pregnancy, women with GDM have higher glucose and C-reactive protein levels, lower sex hormone binding globulin levels, and may be hyperinsulinemic, as compared to pregnant women without GDM.(10, 11) These findings suggest an altered metabolic milieu which may have biologic effects on breast tissue during a critical period of differentiation.(3-5) It is unclear whether GDM is associated with increased breast cancer risk, with mixed results in past studies.(12-16)

We hypothesized that a history of GDM would be associated with a greater invasive breast cancer risk among parous participants in the Nurses' Health Study II, a large prospective cohort of U.S. female nurses.

Materials and Methods

Study Population

This analysis was conducted in the Nurses' Health Study II (NHS II) longitudinal prospective cohort. NHS II began in 1989 with the enrollment of 116,430 female nurses, aged 25 to 42. Questionnaires are distributed every 2 years to update information on a variety of lifestyle and health-related characteristics, with follow-up >90% of total potential person-years. This study has been approved by the Brigham and Women's Hospital Institutional Review Board (Boston, MA), with participants' consent implied by the return of the questionnaires.

This analysis included participants reporting a previous birth on the baseline questionnaire (1989) or an incident first birth during follow-up, through 2001. Nulliparous women were excluded. The 2001 questionnaire cycle is the last in which pregnancy history was prospectively ascertained, as the majority of NHS II participants had passed reproductive age. Exclusion criteria were any history of cancer at baseline or prior to first birth, a prior multiple gestation pregnancy (i.e., twins or higher), and age at first birth <18 years of age. Women were censored during follow-up upon death, cancer diagnosis, reporting of a multiple gestation pregnancy, or being lost-to-follow-up. Follow-up continued through diagnosis of breast cancer or May 31, 2013, whichever occurred first.

Ascertainment of GDM History

History of GDM was captured on the baseline questionnaire and updated every 2 years through 2001 by self-report of a physician's diagnosis. Self-reported GDM was previously validated against medical records in a subgroup of NHS II participants, with 94% of cases confirmed.(17)

Ascertainment of Incident Breast Cancer Cases

Invasive breast cancer cases were identified on biennial follow-up questionnaires from baseline through the 2011 questionnaire. Medical records were obtained from willing participants to confirm the diagnosis (with >98% accuracy) as well as to capture estrogen receptor (ER) and progesterone receptor (PR) status.(18) ER and PR status (positive/negative) was determined from medical records for 92% and 91% of our confirmed cases, respectively. Reasons for undetermined ER/PR status included missing information (5%), test not performed (~2%), and borderline status (~1%). Deaths were identified by the postal service, next of kin, or National Death Index, with medical records or death certificates used for additional documentation of breast cancer.

Covariate Assessment

Detailed health, reproductive, and lifestyle histories were captured at baseline and updated every 2 years. Participants reporting a diagnosis of type 2 diabetes received a previously validated supplemental questionnaire,(19) from which confirmation of diagnosis was made in accordance with the National Diabetes Data Group criteria through 1998,(20) and the American Diabetes Association criteria for cases thereafter.(21) A family history of breast cancer in a mother or sister, and family history of diabetes in a parent or sibling was updated

approximately every 4 years. Participants indicated the number of years since the most recent mammogram at baseline and on follow-up questionnaires answered whether they had a mammogram in the past 2 years. Current body weight was self-reported and was highly correlated with technician-measured weight in a previous validation study.(22) A validated physical activity questionnaire was distributed approximately every 4 years,(23) asking participants the frequency in which they usually engaged in common recreational activities, from which we derived total metabolic equivalent task hours per week (MET-hours/week). Usual diet was ascertained every 4 years via a validated semi-quantitative food frequency questionnaire (FFQ), from which we estimated intake of fruits and vegetables (servings/day) and alcohol (grams/day). The birth index, which takes into account parity and the age at each birth, is inversely associated with breast cancer risk and was calculated based on participants' reproductive histories.(1)

Statistical Analysis

Participants' person-time was calculated from age at first birth through date of breast cancer diagnosis, lost-to-follow-up, death, or return of the 2011 questionnaire, whichever occurred first. A history of GDM was determined at baseline and updated every 2 years. A participant was considered exposed from her first report of GDM through the remainder of follow-up. Women reporting a diagnosis of type 2 diabetes prior to a GDM pregnancy were considered not exposed to GDM.

We used age- and multivariable-adjusted Cox proportional hazards models, stratified by time since first birth and age, to estimate the hazard ratios (HRs) and 95% confidence intervals (95% CI) for the association between a history of GDM (ever vs. never) and risk of incident invasive breast cancer. Time-varying covariates were updated every questionnaire cycle. The proportional hazards assumption was met with $p>0.05$ for the interaction of follow-up time (years) with exposure on breast cancer risk.

Our primary endpoint was invasive breast cancer. We also conducted analyses by breast cancer subgroup, including menopausal status at diagnosis and hormone receptor status. Cox proportional hazards competing risk analysis, as described by Lunn, et al, was used to assess heterogeneity by molecular subtype (any positive receptors ER+ or PR+ vs. both negative receptors ER-/PR-).(24) Briefly, data duplication methods were used, allowing the exposure variable and covariates with heterogeneity across subtypes to vary, keeping other covariate estimates constant. A likelihood ratio test is used to compare models with and without this approach to evaluate heterogeneity across subtype. We considered potential confounders in the multivariable model including body mass index (BMI, kg/m²) at age 18 (continuous), weight gain since age 18 (continuous), height (continuous), total physical activity (MET-hours/week, quintiles), age at menarche (10 years old, 11-12, 13-14, 15), birth index (1) (continuous), total lifetime breastfeeding (none, <6 months, 6 months), menopausal status (premenopausal, postmenopausal, unknown), hormone therapy use (never, ever use of estrogen plus progesterone, past – estrogen only or other, current – estrogen only or other), family history of breast cancer in mother or sister (yes/no), personal history of benign breast disease (yes/no), white race/ethnicity (yes/no), alcohol intake (none, 1-14 grams/day, 15 grams/day), and mammography within the past 2 years (<40 years old, 40 and no

mammography, 40 and mammography for screening, 40 and mammography for abnormality/symptoms). Additional supplemental models adjusted for self-reported pregnancy-associated hypertension and use of diabetes therapies (with separate terms for insulin and oral therapies). Missing covariate data for an individual was carried forward from a previous questionnaire, if available. A missing indicator category was created for categorical covariates if necessary.

Additionally, we evaluated a secondary exposure definition, taking into account progression from GDM to type 2 diabetes, classifying women as (1) no diabetes, (2) GDM only, without subsequent type 2 diabetes, (3) type 2 diabetes only, or (4) both GDM and subsequent type 2 diabetes. We updated participants' status prospectively over follow-up; women without either GDM or type 2 diabetes served as the reference group.

We next conducted stratified analyses to evaluate whether the relationship between GDM and breast cancer differed by established diabetes and breast cancer risk factors, including menopausal status (pre- vs. postmenopausal), current BMI classification (normal <25.0, overweight 25.0-29.0, and obese ≥30.0 kg/m²), age at first birth (<30 vs. ≥30 years old), family history of breast cancer in a sister or mother (yes vs. no), family history of diabetes in a parent or sibling (yes vs. no), physical activity (quintile 1-3 vs. quintile 4-5), pre-pregnancy menstrual cycle regularity at ages 18-22 (regular to mostly regular vs. usual or always irregular) and total lifetime breastfeeding history (<6 months vs. ≥6 months). We hypothesized that differences in underlying risk factors, such as insulin resistance and sex steroid hormones may modify the relationship between GDM and subsequent breast cancer risk. We calculated p-values for interaction from a multiplicative interaction term between the exposure and modifier in the main effects model.

We assessed bias due to differences in screening practices between women with and without a history of GDM by restricting to person-time among women ≥40 years of age with mammography screening in the past 2 years. We also modeled BMI at various time points, including current BMI, BMI at age 18, or both in the multivariable model. All analyses were run on SAS Version 9.3, with alpha <0.05 as the level of significance.

Results

The study population consisted of 81,784 parous women without a history of GDM and 5,188 with a history of GDM. The mean (standard deviation) age at GDM diagnosis was 33.1 (5.5) years. Overall, 63% of the 982,078 person-years was accrued prior to menopause. Baseline characteristics are shown in Table 1. Women with a history of GDM were less physically active, less likely to be white, more likely to be overweight or obese, and to have a family history of diabetes, compared to women without a history of GDM. Women with a history of GDM also had an older age at first birth, were more likely to have a history of pregnancy-induced hypertension, and were less likely to have breastfed for ≥6 months total in their lifetime. Over follow-up, women ≥40 years old with a history of GDM were slightly less likely than those without a history of GDM to report mammography screening in the past 2 years (GDM 65.5% vs. no GDM 68.1% person-years).

Over 22 years of prospective follow-up, there were 2,377 incident cases of invasive breast cancer, with 100 occurring among women with a history of GDM. The mean (standard deviation) age at breast cancer diagnosis was 49.5 (6.6) years. Women with a history of GDM were significantly less likely to develop invasive breast cancer (Table 2) in both the age-adjusted (HR 0.68, 95% CI 0.55-0.84, $p=0.0003$) and multivariable-adjusted models (HR 0.68, 95% CI 0.55-0.84, $p=0.0004$). Adjusting for oral contraceptive use, history of pregnancy-induced hypertension, and smoking status did not impact results. Similarly, inclusion of women who reported multiple gestation pregnancies in the analysis did not affect the results. Results were similar by menopausal status (premenopausal, $n=1377$ events: HR 0.73, 95% CI 0.56-0.96, $p=0.03$; postmenopausal, $n=916$ events: HR 0.63, 95% CI 0.43-0.92, $p=0.02$). There was an inverse association between history of GDM and hormone receptor positive breast cancer (either ER+ or PR+, $n=71$ GDM events, $n=1690$ non-GDM events; HR 0.65, 95% CI 0.50-0.84, $p=0.0009$), but not for hormone receptor negative cancers (both ER- and PR-, $n=23$ GDM events, $n=393$ non-GDM events; HR 0.96, 95% CI 0.60-1.52, $p=0.9$); however, test for heterogeneity was not statistically significant (p -heterogeneity=0.16).

Table 3 gives results stratified by established diabetes or breast cancer risk factors. Significant effect modification was observed for BMI (p -interaction=0.04) and total physical activity (p -interaction=0.04), indicating an inverse relationship between GDM history and breast cancer among overweight, obese and less physically active participants (below median MET-hours/week), with no association among normal weight or more physically active women (Table 3). There was a trend toward stronger inverse relationship between history of GDM and invasive breast cancer among women with <6 months total lifetime breastfeeding, as compared to their longer breastfeeding counterparts (p -interaction=0.08). Findings did not differ by age at first birth, family history of breast cancer, regularity of menstrual cycles, or family history of diabetes (all p -interactions >0.05).

Table 4 shows the risk of GDM with invasive breast cancer accounting for type 2 diabetes. There was a significant inverse relationship between GDM and breast cancer vs. no diabetes, regardless of progression to type 2 diabetes (GDM only: HR 0.72, 95% CI 0.58-0.89, $p=0.003$; GDM and subsequent type 2 diabetes: HR 0.26, 95% CI 0.10-0.68, $p=0.006$). Type 2 diabetes only without a prior history of GDM, compared with no diabetes, was not significantly associated with breast cancer (HR 0.69, 95% CI 0.40-1.18, $p=0.2$). Additional adjustment for use of diabetes therapies (insulin and oral therapies) did not modify the estimate (HR 0.68, 95% CI 0.0.55-0.85, $p=0.0005$).

When we restricted to follow-up among participants ≥ 40 years of age reporting mammography screening in the past 2 years, results were similar to the overall cohort (HR 0.74, 95% CI 0.57-0.96, $p=0.02$) (Supplemental Table 1). Results did not change when we adjusted for current BMI instead of BMI at age 18.

Discussion

In this large prospective study of parous women in the United States with >20 years of follow-up, we did not observe a greater invasive breast cancer risk among women with a

history of GDM. In fact, women with history of GDM experienced a lower incident breast cancer risk. This inverse relationship was independent of intermediate progression to type 2 diabetes and was stronger among women who were overweight, obese, or less physically active. Our analyses also suggested that the inverse relationship may be particularly strong for hormone receptor positive breast cancers (ER+ or PR+).

Results of previous studies examining GDM and breast cancer are mixed, and study populations, design, and control for potential confounders varied widely.(12-15, 25, 26) Two prior studies observed inverse relationships between GDM and breast cancer risk.(15, 25) One retrospective population-based study with a median follow-up of 8 years postpartum, found a lower risk of premenopausal breast cancer among women with a history of GDM (OR=0.86, 95% CI=0.75, 0.98).(16) Similarly, a retrospective US-based case-control study indicated an inverse relationship between GDM and postmenopausal breast cancer (OR=0.54, 95% CI=0.37, 0.79).(15) In contrast, there was a positive relationship between a history of GDM and breast cancer risk in a retrospective cohort of live births in 1964-1976, limited to women \leq 50 years of age (OR=1.7, 95% CI=1.1, 2.5).(12) A positive association was also observed between measures of glucose intolerance in a pregnancy cohort with long-term breast cancer,(13) Three additional studies observed null associations for GDM with breast cancer risk.(14, 26, 27) In the present study, we did not observe a positive association between GDM and postmenopausal breast cancer. We were able to address several limitations of prior studies with the prospective design, large sample size, and adjustment for many possible sources of bias including lifestyle and mammography screening practices.

There are several potential explanations for our unexpected observation of lower breast cancer risk among women with prior GDM. It is possible that the altered hormonal milieu of a GDM complicated pregnancy confers protection on breast tissue. Pregnancy is known to induce proliferation and functional differentiation of breast lobules and ducts, thought to mediate the epidemiologic association between pregnancy history and breast cancer risk. (3-5) It follows that differences in circulating growth factors or hormones in GDM-affected pregnancies may impact these processes and the future risk for breast cancer. Preeclampsia is a pregnancy-associated condition linked with GDM that may be associated with lower breast cancer risk.(28, 29) However, adjusting for a history of pregnancy-induced hypertension did not impact our findings.

Beyond pregnancy, GDM may be a manifestation of chronic subclinical abnormal glucose metabolism. We hypothesized, based on epidemiologic literature and rodent models, that a history of abnormal glucose metabolism in pregnancy would be associated with a higher risk of breast cancer. However, obesity (itself strongly associated with type 2 diabetes and abnormal glucose metabolism) is known to be inversely associated with the risk of premenopausal breast cancer.(30, 31) Consistent with this, a trend toward an inverse association between midlife fasting insulin levels (a marker of insulin resistance) and premenopausal breast cancer risk was observed in the NHS II,(32) in contrast to a previously observed positive relationship between fasting insulin levels and postmenopausal breast cancer risk.(30, 31) It is possible that GDM reflects an underlying metabolic state that specifically impacts premenopausal breast cancer risk, which accounted for the majority of the cases in this study. Notably, the magnitude of the reduction in breast cancer risk

associated with a history of GDM appeared to be greatest among women with additional diabetes risk factors: low physical activity, overweight/obesity, and less lifetime breastfeeding.(33) Previous studies suggest that the physiology underlying GDM differs in obese versus lean women; thus different subtypes of GDM may have different associations with breast cancer (34, 35).

Alternate explanations for our observations include residual or unmeasured confounding and screening bias. Unmeasured factors underlying the inverse relationship between obesity and premenopausal breast cancer risk may partially explain our findings. Some oral diabetes therapies have been associated with a lower cancer risk,(36) but adjusting for this factor did not account for our findings. A recent analysis identified three type 2 diabetes genetic risk variants as inversely related to breast cancer risk, although the overall diabetes genetic risk score was null; this suggests there may be some potential for shared pathways between diabetes and lower breast cancer risk that were uncontrolled for in our analysis.(37) We explored the potential for screening bias in sensitivity analyses. Women with GDM were slightly less likely to have been recently screened for breast cancer, which might lead to fewer diagnoses. However, mammography screening was included in our multivariate model. Further, when restricting to women 40 years old reporting mammography screening in the past 2 years, breast cancer risk remained lower among women with a history of GDM, suggesting that differences in screening did not account for our findings.

Strengths of the present study include its large sample size, prospective nature, and rigorous ascertainment of exposures and outcomes. We also captured information on numerous potential confounders. Analyses of breast cancer subtypes and subgroup stratification were limited in statistical power by the number of events; thus, we must interpret these with caution. Another limitation to note is the potential for misclassification of GDM exposure status, given that of the women deemed “probable GDM”, 21% had medical records indicating a physician's diagnosis, but no oral glucose tolerance test values to verify the diagnosis. Finally, because <5% of the participants in this study were from ethnic/racial minority groups, generalizability to non-white populations may be limited.

In summary, in contrast to our hypothesis, we found that a history of GDM was not associated with greater breast cancer risk among a large cohort of U.S. women, followed prospectively for >20 years. In fact, GDM appeared inversely associated with breast cancer risk, even after adjustment for the many known breast cancer risk factors. It is possible that the hormonal milieu of a GDM pregnancy protects the breast from future malignancy or that GDM is representative of an underlying metabolic state (including obesity) which is protective against premenopausal breast cancer. Further research is needed to validate our unexpected findings in other large prospective cohorts.

Acknowledgments

We would like to thank the participants and staff of the Nurses' Health Study II for their valuable contributions as well as the following state cancer registries for their help: AL, AZ, AR, CA, CO, CT, DE, FL, GA, ID, IL, IN, IA, KY, LA, ME, MD, MA, MI, NE, NH, NJ, NY, NC, ND, OH, OK, OR, PA, RI, SC, TN, TX, VA, WA, WY.

Financial support: W.C. Willett received grants UM1CA176726 and CA50385, D.K. Tobias received grant DK103720, C.E. Powe was supported by grant DK007028-41, F.B. Hu received grants DK58845, P30DK46200 and HHSN275201000020C.

References

1. Rosner B, Colditz GA, Willett WC. Reproductive risk factors in a prospective study of breast cancer: the Nurses' Health Study. *Am J Epidemiol*. 1994 Apr 15; 139(8):819–35. [PubMed: 8178795]
2. Lambe M, Hsieh C, Trichopoulos D, Ekblom A, Pavia M, Adami HO. Transient increase in the risk of breast cancer after giving birth. *N Engl J Med*. 1994 Jul 7; 331(1):5–9. [PubMed: 8202106]
3. Russo IH, Russo J. Pregnancy-induced changes in breast cancer risk. *J Mammary Gland Biol Neoplasia*. 2011 Sep; 16(3):221–33. [PubMed: 21805333]
4. Meier-Abt F, Bentires-Alj M. How pregnancy at early age protects against breast cancer. *Trends Mol Med*. 2014 Mar; 20(3):143–53. [PubMed: 24355762]
5. Schedin P. Pregnancy-associated breast cancer and metastasis. *Nat Rev Cancer*. 2006 Apr; 6(4):281–91. [PubMed: 16557280]
6. Boyle P, Boniol M, Koechlin A, Robertson C, Valentini F, Coppens K, et al. Diabetes and breast cancer risk: a meta-analysis. *British journal of cancer*. 2012 Oct 23; 107(9):1608–17. [PubMed: 22996614]
7. Kim C, Newton KM, Knopp RH. Gestational diabetes and the incidence of type 2 diabetes: a systematic review. *Diabetes Care*. 2002 Oct; 25(10):1862–8. [PubMed: 12351492]
8. Bellamy L, Casas JP, Hingorani AD, Williams D. Type 2 diabetes mellitus after gestational diabetes: a systematic review and meta-analysis. *Lancet*. 2009 May 23; 373(9677):1773–9. [PubMed: 19465232]
9. Lavery JA, Friedman AM, Keyes KM, Wright JD, Ananth CV. Gestational diabetes in the United States: temporal changes in prevalence rates between 1979 and 2010. *BJOG : an international journal of obstetrics and gynaecology*. 2016 Aug 11.
10. Wolf M, Sauk J, Shah A, Vossen Smirnakis K, Jimenez-Kimble R, Ecker JL, et al. Inflammation and glucose intolerance: a prospective study of gestational diabetes mellitus. *Diabetes care*. 2004 Jan; 27(1):21–7. [PubMed: 14693961]
11. Thadhani R, Wolf M, Hsu-Blatman K, Sandler L, Nathan D, Ecker JL. First-trimester sex hormone binding globulin and subsequent gestational diabetes mellitus. *Am J Obstet Gynecol*. 2003 Jul; 189(1):171–6. [PubMed: 12861158]
12. Perrin MC, Terry MB, Kleinhaus K, Deutsch L, Yanetz R, Tiram E, et al. Gestational diabetes and the risk of breast cancer among women in the Jerusalem Perinatal Study. *Breast cancer research and treatment*. 2008 Mar; 108(1):129–35. [PubMed: 17476589]
13. Dawson SI. Long-term risk of malignant neoplasm associated with gestational glucose intolerance. *Cancer*. 2004 Jan 1; 100(1):149–55. [PubMed: 14692035]
14. Troisi R, Weiss HA, Hoover RN, Potischman N, Swanson CA, Brogan DR, et al. Pregnancy characteristics and maternal risk of breast cancer. *Epidemiology*. 1998 Nov; 9(6):641–7. [PubMed: 9799175]
15. Rollison DE, Giuliano AR, Sellers TA, Laronga C, Sweeney C, Risendal B, et al. Population-based case-control study of diabetes and breast cancer risk in Hispanic and non-Hispanic White women living in US southwestern states. *Am J Epidemiol*. 2008 Feb 15; 167(4):447–56. [PubMed: 18033764]
16. Bejaimal SA, Wu CF, Lowe J, Feig DS, Shah BR, Lipscombe LL. Short-term risk of cancer among women with previous gestational diabetes: a population-based study. *Diabet Med*. 2015 May 13.
17. Solomon CG, Willett WC, Rich-Edwards J, Hunter DJ, Stampfer MJ, Colditz GA, et al. Variability in diagnostic evaluation and criteria for gestational diabetes. *Diabetes Care*. 1996 Jan; 19(1):12–6. [PubMed: 8720526]
18. Fung TT, Chiuve SE, Willett WC, Hankinson SE, Hu FB, Holmes MD. Intake of specific fruits and vegetables in relation to risk of estrogen receptor-negative breast cancer among postmenopausal women. *Breast Cancer Res Treat*. 2013 Apr; 138(3):925–30. [PubMed: 23532538]

19. Manson JE, Rimm EB, Stampfer MJ, Colditz GA, Willett WC, Krolewski AS, et al. Physical activity and incidence of non-insulin-dependent diabetes mellitus in women. *Lancet*. 1991 Sep 28; 338(8770):774–8. [PubMed: 1681160]
20. National Diabetes Data Group. Classification and diagnosis of diabetes mellitus and other categories of glucose intolerance. *Diabetes*. 1979 Dec; 28(12):1039–57. [PubMed: 510803]
21. Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care*. 1997 Jul; 20(7):1183–97. [PubMed: 9203460]
22. Rimm EB, Stampfer MJ, Colditz GA, Chute CG, Litin LB, Willett WC. Validity of self-reported waist and hip circumferences in men and women. *Epidemiology*. 1990 Nov; 1(6):466–73. [PubMed: 2090285]
23. Wolf AM, Hunter DJ, Colditz GA, Manson JE, Stampfer MJ, Corsano KA, et al. Reproducibility and validity of a self-administered physical activity questionnaire. *International Journal of Epidemiology*. 1994 Oct; 23(5):991–9. [PubMed: 7860180]
24. Lunn M, McNeil D. Applying Cox regression to competing risks. *Biometrics*. 1995 Jun; 51(2): 524–32. [PubMed: 7662841]
25. Bejaimal SA, Wu CF, Lowe J, Feig DS, Shah BR, Lipscombe LL. Short-term risk of cancer among women with previous gestational diabetes: a population-based study. *Diabetic Medicine : A Journal of the British Diabetic Association*. 2016 Jan; 33(1):39–46. [PubMed: 25970380]
26. Sella T, Chodick G, Barchana M, Heymann AD, Porath A, Kokia E, et al. Gestational diabetes and risk of incident primary cancer: a large historical cohort study in Israel. *Cancer Causes & Control*. 2011 Nov; 22(11):1513–20. [PubMed: 21847538]
27. Brasky TM, Li Y, Jaworowicz DJ Jr, Potischman N, Ambrosone CB, Hutson AD, et al. Pregnancy-related characteristics and breast cancer risk. *Cancer Causes & Control*. 2013 Sep; 24(9):1675–85. [PubMed: 23737027]
28. Pacheco NL, Andersen AN, Kamper-Jorgensen M. Preeclampsia and breast cancer: The influence of birth characteristics. *Breast*. 2015 Jul 2.
29. Joffe GM, Esterlitz JR, Levine RJ, Clemens JD, Ewell MG, Sibai BM, et al. The relationship between abnormal glucose tolerance and hypertensive disorders of pregnancy in healthy nulliparous women. Calcium for Preeclampsia Prevention (CPEP) Study Group. *Am J Obstet Gynecol*. 1998 Oct; 179(4):1032–7. [PubMed: 9790393]
30. Huang Z, Hankinson SE, Colditz GA, Stampfer MJ, Hunter DJ, Manson JE, et al. Dual effects of weight and weight gain on breast cancer risk. *JAMA*. 1997 Nov 5; 278(17):1407–11. [PubMed: 9355998]
31. Michels KB, Terry KL, Willett WC. Longitudinal study on the role of body size in premenopausal breast cancer. *Arch Intern Med*. 2006 Nov 27; 166(21):2395–402. [PubMed: 17130395]
32. Eliassen AH, Tworoger SS, Mantzoros CS, Pollak MN, Hankinson SE. Circulating insulin and c-peptide levels and risk of breast cancer among predominately premenopausal women. *Cancer Epidemiol Biomarkers Prev*. 2007 Jan; 16(1):161–4. [PubMed: 17220346]
33. Jager S, Jacobs S, Kroger J, Fritsche A, Schienkiewitz A, Rubin D, et al. Breast-feeding and maternal risk of type 2 diabetes: a prospective study and meta-analysis. *Diabetologia*. 2014 Jul; 57(7):1355–65. [PubMed: 24789344]
34. Catalano PM. Trying to understand gestational diabetes. *Diabetic Medicine : A Journal of the British Diabetic Association*. 2014 Mar; 31(3):273–81. [PubMed: 24341419]
35. Powe CE, Allard C, Battista MC, Doyon M, Bouchard L, Ecker JL, et al. Heterogeneous Contribution of Insulin Sensitivity and Secretion Defects to Gestational Diabetes Mellitus. *Diabetes Care*. 2016 Jun; 39(6):1052–5. [PubMed: 27208340]
36. Franciosi M, Lucisano G, Lapice E, Strippoli GF, Pellegrini F, Nicolucci A. Metformin therapy and risk of cancer in patients with type 2 diabetes: systematic review. *PLoS One*. 2013; 8(8):e71583. [PubMed: 23936520]
37. Zhao Z, Wen W, Michailidou K, Bolla MK, Wang Q, Zhang B, et al. Association of genetic susceptibility variants for type 2 diabetes with breast cancer risk in women of European ancestry. *Cancer Causes & Control*. 2016 May; 27(5):679–93. [PubMed: 27053251]

Table 1
Characteristics of 86,972 parous US women in the Nurses' Health Study II cohort in 1989,
by history of gestational diabetes mellitus (GDM)

	No History of GDM n=81,784	History of GDM n=5,188
Age*	35.0 +/- 4.7	33.8 +/- 4.4
Lifestyle Factors		
BMI	23.9 +/- 4.7	25.8 +/- 5.9
Normal weight (BMI<25.0) - %	71	55
Overweight (BMI 25.0-29.9) - %	18	25
Obese (BMI ≥30) - %	10	20
BMI at age 18 [‡]	21.0 +/- 3.0	21.5 +/- 3.6
Normal weight at 18 (BMI<25.0) - %	91	87
Overweight/obese at 18 (BMI ≥25.0) - %	8	12
Height (inches)	64.9 +/- 2.6	64.5 +/- 2.6
Total physical activity (MET-hrs/week)	23.7 +/- 35.2	21.8 +/- 33.0
Alternative Healthy Eating Index Score [‡]	47.9 (10.7)	47.2 (10.5)
Fruit and vegetable intake (servings/day) [‡]	4.2 +/- 2.3	4.3 +/- 2.5
Alcohol intake (grams/day) [‡]	2.9 +/- 5.6	2.2 +/- 4.8
Smoking total pack years	7.4 +/- 60.7	6.9 +/- 55.4
Smoking status - %		
Never	66	66
Past	22	21
Current smoker	12	13
Current multivitamin use - %	46	47
Other Risk Factors		
Age at menarche - %		
10 years	7	10
11-12 years	46	48
13-14 years	38	35
15 years	8	7
Regularity of menses at ages 18-22 - %		
Regular or very regular	74	72
Usually or always irregular	22	25
Age at first birth (years)	26.5 +/- 4.5	27.6 +/- 5.0
Parity (Pregnancies ≥6 months)	1.8 +/- 1.1	1.8 +/- 1.2
Birth Index [‡]	16.1 (14.3)	14.3 (14.8)
Total breastfeeding - % [‡]		
None	15	13

	No History of GDM n=81,784	History of GDM n=5,188
<6 months	13	13
6 months	46	43
Oral contraceptive use - %		
Never	15	16
Past	73	72
Current	12	12
History of pregnancy-induced hypertension	10	20
Mammography screening among women >40 years old - %		
<40 years old	80	80
40 years old, with screening	11	11
40 years old, no screening	5	5
40 years old, mammography for symptoms	4	4
Hysterectomy	5	5
Bilateral oophorectomy	1	1
Family history of diabetes - %	14	28
Family history of breast cancer - %	6	5
Personal history of benign breast disease - % [‡]	29	30
Race/ethnicity - %		
White	96	93
Black	2	3
Asian	2	3

* Value is not age adjusted

[‡] Missing data on BMI at age 18 in 1% of participants, food frequency and alcohol intake in 16.3% of participants, breastfeeding in 14.6% of parous participants, and history of benign breast disease in 3.6% of participants. All other data was missing in less than 1% of participants.

[‡] Birth index was calculated by summing total years from each birth to current age (or age at menopause for postmenopausal women) over all births.

Values represent means \pm SD unless otherwise indicated and are standardized to the age distribution of the study population

Values of categorical variables may not sum to 100% due to rounding

Table 2
Association between history of GDM and incident invasive breast cancer risk

	No GDM HR (95% CI)	GDM HR (95% CI)	p-value
Cases – no.	2,277	100	
Person-years *	923,694	58,384	
Age-adjusted	1.00 (reference)	0.68 (0.55, 0.84)	0.0003
Multivariable adjusted	1.00 (reference)	0.68 (0.55, 0.84)	0.0004

HR=hazard ratio, CI=confidence interval

* Person-years are calculated as the time from age at first birth or first GDM pregnancy, through end of follow-up (the date of incident breast cancer diagnosis, death, or last questionnaire return through May 31, 2013)

Multivariable model additionally adjusts for body mass index at age 18 (continuous), weight gain since age 18 (continuous), height (continuous), total physical activity (MET-hours/week, quintiles), alcohol intake (none, 1-14 grams/day, 15 grams/day), age at menarche (10 years old, 11-12, 13-14, 15), birth index (continuous), total breastfeeding (none, <6 months, 6 months), menopausal status (premenopausal, postmenopausal, unknown), hormone therapy use (never, ever use of estrogen + progesterone, past – estrogen only or other, current – estrogen only or other), family history of breast cancer in mother or sister (yes/no), personal history of benign breast disease (yes/no), White race/ethnicity (yes/no), and mammography within the past 2 years (<40 years old, 40 and no mammography, 40 and mammography for screening, 40 and mammography for abnormality/symptoms).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3
Association between a history of GDM and incident invasive breast cancer risk, stratified by risk factors

Stratified by Risk Factors	Cases– No. No GDM / GDM	No GDM HR (95% CI)	GDM HR (95% CI)	P for interaction
Current BMI category (kg/m ²)*				
Normal (<25.0)	1116 / 49	1.00 (reference)	0.97 (0.72-1.32)	0.04
Overweight (25.0-29.9)	646 / 25	1.00 (reference)	0.50 (0.32-0.78)	
Obese (30.0)	512 / 26	1.00 (reference)	0.50 (0.32-0.77)	
Age at first birth				
<30 years	1707 / 59	1.00 (reference)	0.57 (0.39-0.83)	0.85
30 years	570 / 41	1.00 (reference)	0.89 (0.64-1.24)	
Family history of breast cancer				
No	1900 / 83	1.00 (reference)	0.65 (0.52-0.83)	0.96
Yes	377 / 17	1.00 (reference)	0.76 (0.44-1.31)	
Family history of diabetes				
No	1669/62	1.00 (reference)	0.75 (0.57-0.98)	0.33
Yes	608/38	1.00 (reference)	0.59 (0.41-0.85)	
Physical Activity				
Low (Q1-Q3)	1457 / 58	1.00 (reference)	0.56 (0.42-0.74)	0.04
High (Q4-Q5)	820 / 42	1.00 (reference)	0.98 (0.70-1.37)	
Total lifetime breastfeeding				
None to <6 months	933 / 29	1.00 (reference)	0.55 (0.37-0.81)	0.08
6 months	1344 / 71	1.00 (reference)	0.77 (0.59-0.99)	

HR=hazard ratio, CI=confidence interval

Multivariable model adjusts for age, body mass index at age 18 (continuous), weight gain since age 18 (continuous), height (continuous), total physical activity (MET-hours/week, quintiles), alcohol intake (none, 1-14 grams/day, 15 grams/day), age at menarche (10 years old, 11-12, 13-14, 15), birth index (continuous), total breastfeeding (none, <6 months, 6 months), menopausal status (premenopausal, postmenopausal, unknown), hormone therapy use (never, ever use of estrogen + progesterone, past – estrogen only or other, current – estrogen only or other), family history of breast cancer in mother or sister (yes/no), personal history of benign breast disease (yes/no), White race/ethnicity (yes/no), and mammography within the past 2 years (<40 years old, 40 and no mammography, 40 and mammography for screening, 40 and mammography for abnormality/symptoms).

* Multivariable model is adjusted for current BMI (continuous).

Table 4
Association between a history of GDM and incident invasive breast cancer risk, by intermediate type 2 diabetes status

	No GDM or Type 2 Diabetes HR (95% CI)	GDM Only HR (95% CI)	Type 2 Diabetes Only HR (95% CI)	Both GDM and Type 2 Diabetes HR (95% CI)
All Women				
Cases – no.	2224	95	53	5
Person-years	907,823	53,635	20,576	5982
Age-adjusted	1.00 (reference)	0.72 (0.58-0.90)	0.63 (0.37-1.07)	0.23 (0.09-0.61)
Multivariable model	1.00 (reference)	0.72 (0.58-0.89)	0.69 (0.40-1.18)	0.26 (0.10-0.68)

HR=hazard ratio, CI=confidence interval

Multivariable model adjusts for age, body mass index at age 18 (continuous), weight gain since age 18 (continuous), height (continuous), total physical activity (MET-hours/week, quintiles), alcohol intake (none, 1-14 grams/day, 15 grams/day), age at menarche (10 years old, 11-12, 13-14, 15), birth index (continuous), total breastfeeding (none, <6 months, 6 months), menopausal status (premenopausal, postmenopausal, unknown), hormone therapy use (never, ever use of estrogen + progesterone, past – estrogen only or other, current – estrogen only or other), family history of breast cancer in mother or sister (yes/no), personal history of benign breast disease (yes/no), White race/ethnicity (yes/no), and mammography within the past 2 years (<40 years old, 40 and no mammography, 40 and mammography for screening, 40 and mammography for abnormality/symptoms).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript