

*THE COVALENT STRUCTURE OF  
AN ENTIRE  $\gamma$ G IMMUNOGLOBULIN MOLECULE\**

BY GERALD M. EDELMAN, BRUCE A. CUNNINGHAM, W. EINAR GALL,  
PAUL D. GOTTLIEB, URS RUTISHAUSER, AND MYRON J. WAXDAL

THE ROCKEFELLER UNIVERSITY

*Communicated by Theodore Shedlovsky, March 21, 1969*

*Abstract.*—The complete amino acid sequence of a human  $\gamma$ G1 immunoglobulin (Eu) has been determined and the arrangement of all of the disulfide bonds has been established. Comparison of the sequence with that of another myeloma protein (He) suggests that the variable regions of heavy and light chains are homologous and similar in length. The constant portion of the heavy chain contains three homology regions each of which is similar in size and homologous to the constant region of the light chain. Each variable region and each constant homology region contains one intrachain disulfide bond. The half-cystines participating in the interchain bonds are all clustered within a stretch of ten residues at the middle of the heavy chains.

These data support the hypothesis that immunoglobulins evolved by gene duplication after early divergence of V genes, which specified antigen-binding functions, and C genes, which specified other functions of antibody molecules. Each polypeptide chain may therefore be specified by two genes, V and C, which are fused to form a single gene (translocation hypothesis). The internal homologies and symmetry of the molecule suggest that homology regions may have similar three-dimensional structures each consisting of a compact domain which contributes to at least one active site (domain hypothesis). Both hypotheses are in accord with the linear regional differentiation of function in antibody molecules.

---

Antibodies or immunoglobulins can interact with a wide range of different antigenic determinants and, after specific binding to an antigen, they play a fundamental part in physiological functions of the immune response. The specificity of antigen binding depends ultimately upon amino acid sequences of the variable or V regions of antibody molecules. It is the diversity of these sequences which results in the range of specificities required for a selective immune response. In contrast, other regions of the antibody molecule have relatively constant sequences and are responsible for physiological functions. Like enzymes, these C regions appear to have evolved for a restricted set of interactions. This unusual picture of intramolecular differentiation has emerged from studies of the structure of immunoglobulins from different animal species.<sup>1</sup> To date, only portions of immunoglobulin molecules have been subjected to amino acid sequence determination.

We now report the amino acid sequence of an entire human  $\gamma$ G1 immunoglobulin (molecular weight 150,000), the location of all disulfide bonds, the arrangement of light and heavy chains, and the length of the heavy chain V region.

**Materials and Methods.**—The isolation of the myeloma protein Eu<sup>2</sup> and the preparation of its CNBr fragments<sup>3, 4</sup> have been described. Similar methods were used for the isolation of the  $\gamma$ G1 myeloma protein He and for the preparation of its CNBr fragments.

We have previously described the methods used for enzymatic digestion with trypsin, chymotrypsin, and pepsin, gel filtration, ion exchange chromatography, high voltage paper electrophoresis, determination of NH<sub>2</sub>-terminal and COOH-terminal residues, amino acid analysis, and determination of amino acid sequences by the dansyl-Edman procedure.<sup>4-8</sup>

The positions of glutamine and asparagine were assigned<sup>9</sup> by determining the electrophoretic mobility of peptides and by amino acid analysis of the peptides after enzymatic hydrolysis. The half-cystinyl residues contributing to each intrachain disulfide bond were determined<sup>10</sup> using the diagonal electrophoresis method.<sup>11</sup>

**Results.**—The organization of the whole molecule is shown in Figure 1; an unequivocal proof of the arrangement of the two identical light chains and two identical heavy chains has already been given.<sup>4</sup> Each light chain is linked to its neighboring heavy chain by a disulfide bond between corresponding half-




FIG. 1.—Over-all arrangement of chains and disulfide bonds of  $\gamma$ G1 immunoglobulin Eu. Half-cystinyl residues are numbered I–XI; numbers I–V designate corresponding residues in light and heavy chains. PCA: pyrrolidonecarboxylic acid. CHO: carbohydrate. “Fab(t)” and “Fc(t)” refer to fragments produced by trypsin, which cleaves the heavy chain as indicated by dashed lines above half-cystinyl residues VI. V<sub>H</sub>, V<sub>L</sub>: variable regions of heavy and light chains, C<sub>L</sub>: constant region of light chain. C<sub>H</sub>1, C<sub>H</sub>2, C<sub>H</sub>3: homology regions comprising C<sub>H</sub> or constant region of heavy chain.

cystines V. Half-cystines VI and VII form bonds linking the half-molecules via the heavy chains. Trypsin cleaves the molecule at lysyl residue 222 to form two Fab(t) and one Fc(t) fragments.<sup>2, 5</sup>

There are several strikingly linear arrangements in the primary structure. From their amino termini to half-cystines V, the light and heavy chains can be aligned or put in register. The intrachain disulfide bonds are linearly and periodically disposed.<sup>12, 13</sup> In accord with the alignment of light and heavy chains, corresponding intrachain disulfide bonds are in similar positions and the disulfide loops are of approximately the same size.

Previous studies<sup>7</sup> have suggested that V regions of light and heavy chains have similar lengths and begin at the NH<sub>2</sub>-termini; this will be confirmed below. The C<sub>L</sub> region of the light chain has the same length as V<sub>L</sub>, but the C<sub>H</sub> region of the heavy chain is about three times as long. C<sub>H</sub> may be divided into three

homologous regions of approximately equal length:  $C_{H1}$ ,  $C_{H2}$ , and  $C_{H3}$  (Fig. 1).

We have already reported the amino acid sequence of the first 87<sup>7</sup> and the last 224 residues<sup>8</sup> of the heavy chain as well as the partial sequence of the entire light chain.<sup>6</sup> The complete amino acid sequence of the light chain (214 residues) is shown in Figure 2. Positions of the half-cystinyl residues may be compared with Figure 1 and the methionyl residues may be correlated with previous studies on the CNBr fragments of Eu.<sup>3, 4</sup> The variable region extends through residue 108. In accord with other studies,<sup>1</sup> valine 191 is related to the Inv specificity.<sup>2</sup>

The complete sequence of the heavy chain (446 residues) is presented in Figure 3 which may be compared with Figure 2 for alignment with the light chain sequence. Isolation of a single glycopeptide<sup>8</sup> indicated that the polysaccharide




FIG. 2.—Complete amino acid sequence of the Eu light chain. Half-cystinyl residues are in boxes and methionyl residues are underlined.

portion of the molecule is attached at Asx residue 297.<sup>14</sup> In a previous study<sup>8</sup> we have suggested that glutamyl residue 356 and methionyl residue 358 may be associated with Gm 1 specificities. The sequence of Eu (Gm 4+) between residues 211–252 can be compared with the partial sequence of immunoglobulin Daw<sup>17</sup> (Gm 4–). The presence of arginine in position 214 of Eu and lysine in a comparable position of Daw may be associated with their Gm 4 specificities.<sup>18</sup>

Of particular significance is the determination of the point at which  $V_H$  ends and  $C_H$  begins. A CNBr fragment comparable to fragment  $H_4$  was isolated from myeloma protein He which has the same Gm specificity as protein Eu. The sequence of the amino terminal portion of the CNBr fragment from He differed from that of the  $H_4$  fragment from Eu.<sup>21</sup>

1  
 PCA-VAL-GLN-LEU-VAL-GLN-SER-GLY-ALA-GLU-VAL-LYS-LYS-PRO-GLY-SER-SER-VAL-LYS-VAL-20  
 30  
 SER-CYS-LYS-ALA-SER-GLY-GLY-THR-PHE-SER-ARG-SER-ALA-ILE-ILE-TRP-VAL-ARG-GLN-ALA-40  
 50  
 PRO-GLY-GLN-GLY-LEU-GLU-TRP-MET-GLY-GLY-ILE-VAL-PRO-MET-PHE-GLY-PRO-PRO-ASN-TYR-60  
 70  
 ALA-GLN-LYS-PHE-GLN-GLY-ARG-VAL-THR-ILE-THR-ALA-ASP-GLU-SER-THR-ASN-THR-ALA-TYR-80  
 90  
MET-GLU-LEU-SER-SER-LEU-ARG-SER-GLU-ASP-THR-ALA-PHE-TYR-PHE-CYS-ALA-GLY-GLY-TYR-100  
 110  
 GLY-ILE-TYR-SER-PRO-GLU-GLU-TYR-ASN-GLY-GLY-LEU-VAL-THR-VAL-SER-SER-ALA-SER-THR-120  
 130  
 LYS-GLY-PRO-SER-VAL-PHE-PRO-LEU-ALA-PRO-SER-SER-LYS-SER-THR-SER-GLY-GLY-THR-ALA-140  
 150  
 ALA-LEU-GLY-CYS-LEU-VAL-LYS-ASP-TYR-PHE-PRO-GLU-PRO-VAL-THR-VAL-SER-TRP-ASN-SER-160  
 170  
 GLY-ALA-LEU-THR-SER-GLY-VAL-HIS-THR-PHE-PRO-ALA-VAL-LEU-GLN-SER-SER-GLY-LEU-TYR-180  
 190  
 SER-LEU-SER-SER-VAL-VAL-THR-VAL-PRO-SER-SER-SER-LEU-GLY-THR-GLN-THR-TYR-ILE-CYS-200  
 210  
 ASN-VAL-ASN-HIS-LYS-PRO-SER-ASN-THR-LYS-VAL-ASP-LYS-ARG-VAL-GLU-PRO-LYS-SER-CYS-220  
 230  
 ASP-LYS-THR-HIS-THR-CYS-PRO-PRO-CYS-PRO-ALA-PRO-GLU-LEU-LEU-GLY-GLY-PRO-SER-VAL-240  
 250  
 PHE-LEU-PHE-PRO-PRO-LYS-PRO-LYS-ASP-THR-LEU-MET-ILE-SER-ARG-THR-PRO-GLU-VAL-THR-260  
 270  
CYS-VAL-VAL-VAL-ASP-VAL-SER-HIS-GLU-ASP-PRO-GLN-VAL-LYS-PHE-ASN-TRP-TYR-VAL-ASP-280  
 290  
 GLY-VAL-GLN-VAL-HIS-ASN-ALA-LYS-THR-LYS-PRO-ARG-GLU-GLN-GLN-TYR-ASX-SER-THR-TYR-300  
 310  
 ARG-VAL-VAL-SER-VAL-LEU-THR-VAL-LEU-HIS-GLN-ASN-TRP-LEU-ASP-GLY-LYS-GLU-TYR-LYS-320  
 330  
CYS-LYS-VAL-SER-ASN-LYS-ALA-LEU-PRO-ALA-PRO-ILE-GLU-LYS-THR-ILE-SER-LYS-ALA-LYS-340  
 350  
 GLY-GLN-PRO-ARG-GLU-PRO-GLN-VAL-TYR-THR-LEU-PRO-PRO-SER-ARG-GLU-GLU-MET-THR-LYS-360  
 370  
 ASN-GLN-VAL-SER-LEU-THR-CYS-LEU-VAL-LYS-GLY-PHE-TYR-PRO-SER-ASP-ILE-ALA-VAL-GLU-380  
 390  
 TRP-GLU-SER-ASN-ASP-GLY-GLU-PRO-GLU-ASN-TYR-LYS-THR-THR-PRO-PRO-VAL-LEU-ASP-SER-400  
 410  
 ASP-GLY-SER-PHE-PHE-LEU-TYR-SER-LYS-LEU-THR-VAL-ASP-LYS-SER-ARG-TRP-GLN-GLU-GLY-420  
 430  
 ASN-VAL-PHE-SER-CYS-SER-VAL-MET-HIS-GLU-ALA-LEU-HIS-ASN-HIS-TYR-THR-GLN-LYS-SER-440  
 446  
 LEU-SER-LEU-SER-PRO-GLY

FIG. 3.—Complete amino acid sequence of the Eu heavy chain. Half-cystinyl residues are in boxes and methionyl residues are underlined.

A comparison of the sequence of the two fragments from residues 101 to 121 is given in Figure 4. The sequences become identical at residue 115 (Eu numbering). Further studies<sup>21</sup> confirmed that from residue 115 to residue 252 the sequence of the He fragment was identical to that of Eu H<sub>4</sub>. In addition, tryptic fingerprints of the Fc fragments from Eu and He were identical. These

	101	105	110	115	120
EU	- GLY - ILE - TYR - SER -	PRO - GLU - GLU - TYR - ASN -	GLY - GLY - LEU - VAL -	THR - VAL - SER - SER -	ALA - SER - THR - LYS -
HE	- THR - LEU - ALA - PHE -	ASX - VAL - TRP - GLY -	GLX - GLY - THR - LYS -	VAL - ALA - VAL - SER -	SER - ALA - SER - THR - LYS -

FIG. 4.—Comparison of the amino acid sequence of the Eu heavy chain from residue 101–121 with the corresponding sequence of the heavy chain of myeloma protein He.

data suggest that the transition between  $V_H$  and  $C_H$  is located in the vicinity of residue 114 (Eu numbering). Studies on a number of additional proteins and a search for  $V_H$  region subgroups<sup>7</sup> will be required to locate this point definitively.

*Discussion.*—The present studies provide proof of the covalent structure and arrangement of chains in  $\gamma$ G1 immunoglobulin. The half-molecule of Eu is the largest protein unit (446 + 214 residues) for which a complete amino acid sequence has been determined. In a protein of this size, one cannot neglect the possibility of small errors in sequence assignment; for this reason we are carrying out a number of further checks using various methods of peptide cleavage and fractionation.

One of the most striking features of the immunoglobulin molecule that emerges from the completed sequence is the sharp demarcation of its polypeptide chains into linearly connected regions that are associated with different functions. Variations in the sequences of paired  $V_H$  and  $V_L$  regions for the function of antigen binding in the selective immune response, and at the same time, conservation of sequence in  $C_H$  and  $C_L$  regions for other immunological functions appear to require special genetic and evolutionary mechanisms.<sup>22</sup>

The amino acid sequence of Eu provides convincing evidence that the immunoglobulin molecule evolved by successive duplication of precursor genes.<sup>23, 24</sup> Our analysis of a complete heavy chain has revealed an additional homology region ( $C_{H1}$ ), the structure of which was not previously known. Earlier comparisons<sup>7, 8</sup> of polynucleotide sequences corresponding to both chains of Eu showed evidence of homology between  $V_H$  and  $V_L$  and homologies among  $C_L$ ,  $C_{H2}$ , and  $C_{H3}$ . A complete comparison of the amino acid sequences of  $C_L$ ,  $C_{H1}$ ,  $C_{H2}$ , and  $C_{H3}$  is given in Figure 5. In a stretch of 100 residues, any two regions are identical in 29 to 34 positions. It is noteworthy that the stretch in the heavy chain from residue 221 to 233 which contains the interchain disulfide bonds<sup>5</sup> has no homologous counterpart in other portions of heavy or light chains.

In the data accumulated so far, little or no homology has been found between V and C regions. This prompts the speculation that V genes and C genes diverged early in the evolution of antibodies to serve two major groups of functions: antigen recognition functions (ARF) and effector functions (EF) such as interaction with cells and complement. The order of emergence of  $C_L$  genes or  $C_H$  genes from a precursor gene is not apparent from the data. A comparison with sequences of  $\mu$ ,  $\alpha$ ,  $\delta$ , or  $\epsilon$  chains may indicate similarities in their  $V_H$  regions<sup>25</sup> and may show whether any of the homology regions are conserved in constant regions of heavy chains of these classes.<sup>26</sup>

Early evolutionary divergence of V and C genes is consistent with the evidence<sup>27, 28</sup> that *each* chain is specified by two genes, V and C, and the hypothesis<sup>7, 22, 29</sup> that V gene episomes are translocated to C genes to form a single VC

EU C <sub>L</sub> (RESIDUES 109-214)						110														120					
						THR	VAL	ALA	ALA	PRO	SER	VAL	PHE	ILE	PHE	PRO	PRO	SER							
EU C <sub>H</sub> 1 (RESIDUES 119-220)						SER	THR	LYS	GLY	PRO	SER	VAL	PHE	PRO	LEU	ALA	PRO	SER							
EU C <sub>H</sub> 2 (RESIDUES 234-341)						LEU	LEU	GLY	GLY	PRO	SER	VAL	PHE	LEU	PHE	PRO	PRO	LYS							
EU C <sub>H</sub> 3 (RESIDUES 342-446)						GLN	PRO	ARG	GLU	PRO	GLN	VAL	TYR	THR	LEU	PRO	PRO	SER							
130																									
ASP	GLU	GLN	-	-	LEU	LYS	SER	GLY	THR	ALA	SER	VAL	VAL	CYS	LEU	LEU	ASN	ASN	PHE						
SER	LYS	SER	-	-	THR	SER	GLY	GLY	THR	ALA	ALA	LEU	GLY	CYS	LEU	VAL	LYS	ASP	TYR						
PRO	LYS	ASP	THR	LEU	MET	ILE	SER	ARG	THR	PRO	GLU	VAL	THR	CYS	VAL	VAL	VAL	ASP	VAL						
ARG	GLU	GLU	-	-	MET	THR	LYS	ASN	GLN	VAL	SER	LEU	THR	CYS	LEU	VAL	LYS	GLY	PHE						
140																									
TYR	PRO	ARG	GLU	ALA	LYS	VAL	-	-	GLN	TRP	LYS	VAL	ASP	ASN	ALA	LEU	GLN	SER	GLY						
PHE	PRO	GLU	PRO	VAL	THR	VAL	-	-	SER	TRP	ASN	SER	-	GLY	ALA	LEU	THR	SER	GLY						
SER	HIS	GLU	ASP	PRO	GLN	VAL	LYS	PHE	ASN	TRP	TYR	VAL	ASP	GLY	-	VAL	GLN	VAL	HIS						
TYR	PRO	SER	ASP	ILE	ALA	VAL	-	-	GLU	TRP	GLU	SER	ASN	ASP	-	GLY	GLU	PRO	GLU						
150																									
160																									
ASN	SER	GLN	GLU	SER	VAL	THR	GLU	GLN	ASP	SER	LYS	ASP	SER	THR	TYR	SER	LEU	SER	SER						
-	VAL	HIS	THR	PHE	PRO	ALA	VAL	LEU	GLN	SER	-	SER	GLY	LEU	TYR	SER	LEU	SER	SER						
ASN	ALA	LYS	THR	LYS	PRO	ARG	GLU	GLN	GLN	TYR	-	ASP	SER	THR	TYR	ARG	VAL	VAL	SER						
ASN	TYR	LYS	THR	THR	PRO	PRO	VAL	LEU	ASP	SER	-	ASP	GLY	SER	PHE	PHE	LEU	TYR	SER						
170																									
180																									
THR	LEU	THR	LEU	SER	LYS	ALA	ASP	TYR	GLU	LYS	HIS	LYS	VAL	TYR	ALA	CYS	GLU	VAL	THR						
VAL	VAL	THR	VAL	PRO	SER	SER	SER	LEU	GLY	THR	GLN	-	THR	TYR	ILE	CYS	ASN	VAL	ASN						
VAL	LEU	THR	VAL	LEU	HIS	GLN	ASN	TRP	LEU	ASP	GLY	LYS	GLU	TYR	LYS	CYS	LYS	VAL	SER						
LYS	LEU	THR	VAL	ASP	LYS	SER	ARG	TRP	GLN	GLU	GLY	ASN	VAL	PHE	SER	CYS	SER	VAL	MET						
190																									
200																									
HIS	GLN	GLY	LEU	SER	SER	PRO	VAL	THR	-	LYS	SER	PHE	-	-	ASN	ARG	GLY	GLU	CYS						
HIS	LYS	PRO	SER	ASN	THR	LYS	VAL	-	ASP	LYS	ARG	VAL	-	-	GLU	PRO	LYS	SER	CYS						
ASN	LYS	ALA	LEU	PRO	ALA	PRO	ILE	-	GLU	LYS	THR	ILE	SER	LYS	ALA	LYS	GLY								
HIS	GLU	ALA	LEU	HIS	ASN	HIS	TYR	THR	GLN	LYS	SER	LEU	SER	LEU	SER	PRO	GLY								
210																									

FIG. 5.—Comparison of the amino acid sequence of C<sub>L</sub>, C<sub>H</sub>1, C<sub>H</sub>2, and C<sub>H</sub>3. Deletions indicated by dashes have been introduced to maximize the homology. Identical residues are darkly shaded; both dark and light shading are used to indicate identities which occur in pairs in the same positions.

gene in lymphoid cell precursors. Dreyer and Bennett<sup>30</sup> have previously suggested a translocation of C genes and, more recently, this has been abandoned in favor of a detailed "copy-splice" mechanism.<sup>31</sup> Translocation of genes may be the basis of the phenomena of clonal expression and allelic exclusion in antibody production. Irreversible differentiation and commitment of a lymphoid precursor cell may thus occur at the time of gene translocation.

The alignment of disulfide bonds, the arrangement of symmetry axes, and the fact that proteolytic enzymes cleave the molecule to produce Fab, Fc and Fc' fragments<sup>32</sup> suggest that each homology region may be folded in a compact domain<sup>29</sup> stabilized by a single intrachain disulfide bond and linked to neighboring regions by less tightly folded stretches of polypeptide chain. Such domains would have similar but not identical tertiary structures, and each domain would

contribute to at least one active site mediating a function of that class of immunoglobulin. This domain hypothesis is consistent with the hypothesis that the molecule evolved by gene duplication as well as with the translocation hypothesis. As mentioned above, comparison of the structure and function of  $C_H$  regions in different immunoglobulin classes should reveal whether addition or deletion of homology regions and corresponding domains is a major mechanism in the evolution of these classes.

Support for the domain hypothesis would come from finding that limited proteolysis of Fab fragments yields fragments containing halves of the Fd fragments. Similar treatment of Bence-Jones proteins may produce  $V_L$  and  $C_L$  fragments.<sup>33</sup> Additional evidence may come from location in  $C_H$  of sites for complement fixation and skin fixation. Final proof or disproof of this hypothesis obviously rests on the results of X-ray crystallographic analysis. It is clear that a rotation axis passes through the disulfide bonds linking the heavy chains and there may be an axis of pseudosymmetry between the light and heavy chains. The locations of the interchain and intrachain disulfide bonds, the extensive homologies, and the alignment of the light and heavy chains with each other suggest that the overall relationships described above will be conserved in the three-dimensional structure regardless of the details of folding.

The exact contribution of the variable regions to the antigen combining site must also await analysis of the three-dimensional structure. It is known that the Fab fragment contains both  $V_L$  and  $V_H$  regions, and affinity-labeling experiments<sup>24</sup> indicate that tyrosyl residues<sup>34, 35</sup> in these regions are directly involved in antigen binding. The constancy of the disulfide bonds in  $V_H$  and  $V_L$  regions and their coordinate location suggest the possibility that the site is fixed by these bonds and that variations in the branches of a chain connected by each bond may be sterically arranged around the bond as a center. The closely homologous  $C_L$  and  $C_{H1}$  regions may serve additionally to stabilize the structure in the face of the variation, so that both  $V_L$  and  $V_H$  can participate in the site.

In contrast to the diversity of V regions, the origin of which is still unknown,<sup>22</sup> the C regions appear to be quite stable in various animal species. Recent studies<sup>8</sup> show striking resemblances in the Fc portions of rabbit<sup>23</sup> and human  $\gamma G$  immunoglobulin. In this respect, C regions, like enzymes, may have evolved to interact with specific molecules, e.g., those of the complement system. The presence of genetic differences in C regions has not so far been related to their function, but there is no reason to expect that the origin of variations in C regions will differ from other genetic polymorphisms.

We are deeply indebted to Dr. Jack Brook for supplying Eu plasma. We are grateful to Miss Joan Low, Miss Catherine Volin, and Mrs. Helvi Hjelt for their expert technical assistance.

\* Supported by grant GB 6546 from the National Science Foundation and by grant AM 04256 from the National Institutes of Health.

<sup>1</sup> For a general review, see *Cold Spring Harbor Symposia on Quantitative Biology*, vol. 32 (1967), and *Nobel Symposium 3: Gamma Globulins, Structure and Control of Biosynthesis*, ed. J. Killander (Stockholm: Almqvist and Wiksell, 1967).

<sup>2</sup> Edelman, G. M., W. E. Gall, M. J. Waxdal, and W. H. Konigsberg, *Biochemistry*, **7**, 1950 (1968).

- <sup>3</sup> Waxdal, M. J., W. H. Konigsberg, W. L. Henley, and G. M. Edelman, *Biochemistry*, **7**, 1959 (1968).
- <sup>4</sup> Waxdal, M. J., W. H. Konigsberg, and G. M. Edelman, *Biochemistry*, **7**, 1967 (1968).
- <sup>5</sup> Gall, W. E., B. A. Cunningham, M. J. Waxdal, W. H. Konigsberg, and G. M. Edelman, *Biochemistry*, **7**, 1973 (1968).
- <sup>6</sup> Cunningham, B. A., P. D. Gottlieb, W. H. Konigsberg, and G. M. Edelman, *Biochemistry*, **7**, 1983 (1968).
- <sup>7</sup> Gottlieb, P. D., B. A. Cunningham, M. J. Waxdal, W. H. Konigsberg, and G. M. Edelman, these PROCEEDINGS, **61**, 168 (1968).
- <sup>8</sup> Rutishauser, U., B. A. Cunningham, C. Bennett, W. H. Konigsberg, and G. M. Edelman, these PROCEEDINGS, **61**, 1414 (1969).
- <sup>9</sup> Bennett, C., W. H. Konigsberg, and G. M. Edelman, manuscript in preparation.
- <sup>10</sup> Gall, W. E., and G. M. Edelman, manuscript in preparation.
- <sup>11</sup> Brown, J. R., and B. S. Hartley, *Biochem. J.*, **101**, 214 (1966).
- <sup>12</sup> Waxdal, M. J., W. H. Konigsberg, and G. M. Edelman, in *Cold Spring Harbor Symposia on Quantitative Biology*, vol. 32, 53 (1967).
- <sup>13</sup> Pink, J. R. L., and C. Milstein, *Nature*, **216**, 941 (1967).
- <sup>14</sup> Inasmuch as the mechanism of attachment of the carbohydrate is unknown, we have not specified whether residue 297 is Asp or Asn. This residue begins the tripeptide sequence Asx-Ser-Thr (residues 297-299), which is the same as the Asx-X-Thr sequence found at the point of attachment of the carbohydrate portion of several other glycoproteins.<sup>15, 16</sup>
- <sup>15</sup> Eylar, E. H., *J. Theoret. Biol.*, **10**, 89 (1966).
- <sup>16</sup> Catley, B. J., S. Moore, and W. H. Stein, *J. Biol. Chem.*, **244**, 933 (1969).
- <sup>17</sup> Steiner, L. A., and R. R. Porter, *Biochemistry*, **6**, 3957 (1967).
- <sup>18</sup> The association with Gm 1 specificities is based on definitive studies.<sup>19, 20</sup> Additional evidence is required for the correlation of sequence variation with Gm 4 specificities.
- <sup>19</sup> Thorpe, N. O., and H. F. Deutsch, *Immunochemistry*, **3**, 329 (1966).
- <sup>20</sup> Frangione, B., C. Milstein, and J. R. L. Pink, *Nature*, **221**, 145 (1969).
- <sup>21</sup> Cunningham, B. A., M. Pflumm, U. Rutishauser, and G. M. Edelman, manuscript in preparation.
- <sup>22</sup> Edelman, G. M., and J. A. Gally, in *Brookhaven Symposia in Biology*, No. 21, 328 (1968).
- <sup>23</sup> Hill, R. L., R. Delaney, R. E. Fellows, Jr., and H. E. Lebovitz, these PROCEEDINGS, **56**, 1762 (1966).
- <sup>24</sup> Singer, S. J., and R. F. Doolittle, *Science*, **153**, 13 (1966).
- <sup>25</sup> Wikler, M., H. Köhler, T. Shinoda, and F. W. Putnam, *Science*, **163**, 75 (1969).
- <sup>26</sup> For generality, we have used C<sub>L</sub> and C<sub>H1</sub>, C<sub>H2</sub>, etc., to designate homology regions. If such regions are found in other classes, a more specific nomenclature, e.g. C<sub>x</sub>, C<sub>γ1</sub>, C<sub>γ2</sub>, etc., and perhaps C<sub>μ1</sub>, etc., may be required. The same obviously holds for V regions.
- <sup>27</sup> Hood, L., and D. Ein, *Nature*, **220**, 764 (1968).
- <sup>28</sup> Milstein, C., C. P. Milstein, and A. Feinstein, *Nature*, **221**, 151 (1969).
- <sup>29</sup> Edelman, G. M., and W. E. Gall, *Ann. Rev. Biochem.*, **38**, (1969), in press.
- <sup>30</sup> Dreyer, W. J., and J. C. Bennett, these PROCEEDINGS, **54**, 864 (1965).
- <sup>31</sup> Dreyer, W. J., W. R. Gray, and L. Hood, in *Cold Spring Harbor Symposia on Quantitative Biology*, vol. 32, p. 353 (1967).
- <sup>32</sup> Turner, M. W., and H. Bennich, *Biochem. J.*, **107**, 171 (1968).
- <sup>33</sup> Berggård, I., and P. Peterson, in *Nobel Symposium 3: Gamma Globulins, Structure and Control of Biosynthesis*, ed. J. Killander (Stockholm: Almqvist and Wiksell 1967), p. 71.
- <sup>34</sup> Singer, S. J., and N. O. Thorpe, these PROCEEDINGS, **60**, 1371 (1968).
- <sup>35</sup> Although there are no direct data on affinity labeling of human antibodies, by analogy (see ref. 24), we would suggest that tyrosyl residues 86 of the light chain and 94 of the heavy chain may have a function in the active site.