





Complete Genome Sequence of Bifidobacterium longum W11 (LMG P-21586), Used as a Probiotic Strain

Rosanna Inturri,^a Marco Ventura,^b ® Patricia Ruas-Madiedo,^c Gabriele Andrea Lugli,^b Giovanna Blandino^a

Department of Biomedical and Biotechnological Sciences, University of Catania, Catania, Italya; Laboratory of Probiogenomics, Department of Life Sciences, University of Parma, Parma, Italyb; Department of Microbiology and Biochemistry of Dairy Products, Instituto de Productos Lácteos de Asturias, Consejo Superior de Investigaciones Científicas (IPLA-CSIC), Villaviciosa, Asturias, Spainc

ABSTRACT We report the complete genome sequence of *Bifidobacterium longum* W11 (LMG P-21586) isolated from the intestinal microbiota of a healthy man. The analysis of the sequence may provide insights into the microbiological characteristics and the functional activity of this probiotic strain.

members of the genus *Bifidobacterium* are nonmotile non-spore-forming Grampositive polymorphic anaerobic rods, with a high G+C DNA content. *Bifidobacterium* spp. ferment carbohydrates to produce acetic acid and lactic acid but no carbon dioxide, and they metabolize glucose via the fructose-6-phosphate phosphoketolase shunt (1–3). This genus represents an important commensal group of the human intestinal microbiota, particularly during breastfeeding (4). *Bifidobacterium longum* subsp. *longum* W11 (LMG P-21586) has been isolated from the intestinal microbiota of a healthy man and has been studied for some microbial characteristics (5–8).

Genomic DNA of *B. longum* W11 was subject to whole-genome sequencing using MiSeq (Illumina, United Kingdom) at GenProbio s.r.l. (Parma, Italy). Paired-end reads obtained from targeted genome sequencing were used as input for the genome assemblies through the MEGAnnotator pipeline (9). The MIRA program (version 4.0) was used for the *de novo* assembly, while proteins encoded by open reading frames (ORFs) were predicted using Prodigal. The circular chromosome contains 2,329,981 bp with 1,884 ORFs, with these total numbers being higher than those in the representative *B. longum* NCC2705. The average of G+C content of the genome is 59.9%.

The complete genome sequence of *B. longum* W11 may be useful to better understand the health-promoting characteristics of this probiotic strain and to clarify the mechanisms involved in its interaction with the host.

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number MRBG00000000. The version described in this paper is version MRBG01000000.

ACKNOWLEDGMENT

No support was received in order to perform this research.

REFERENCES

- Biavati B, Mattarelli P. 2006. The family Bifidobacteriaceae, p 322–382. In Dworkin M, Falkow S, Rosenberg E, Schleifer K-H, Stackebrandt E (ed), The prokaryotes, vol 3, 3rd ed. Springer Verlag, New York, NY.
- Shkoporov AN, Efimov BA, Khokhlova EV, Chaplin AV, Kafarskaya LI, Durkin AS, McCorrison J, Torralba M, Gillis M, Sutton G, Weibel DB, Nelson

Received 8 December 2016 Accepted 11
January 2017 Published 9 March 2017
Citation Inturri R, Ventura M, Ruas-Madiedo P,

Citation Inturn IP, Ventura M, Ruas-Madiedo P, Lugli GA, Blandino G. 2017. Complete genome sequence of *Bifidobacterium longum* W11 (LMG P-21586), used as a probiotic strain. Genome Announc 5:e01659-16. https://doi.org/10.1128/ qenomeA.01659-16

Copyright © 2017 Inturri et al. This is an openaccess article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Giovanna Blandino, blandino@unict.it.

KE, Smeianov VV. 2013. Draft genome sequences of two pairs of human intestinal *Bifidobacterium longum* subsp. *longum* strains, 44B and 1-6B and 35B and 2-2B, consecutively isolated from two children after a 5-year time period. Genome Announc 1(3):e00234-13. https://doi.org/10.1128/genomeA.00234-13.

Inturri et al. genærieAnnouncements"

 Ventura M, Turroni F, Motherway MO, MacSharry J, van Sinderen D. 2012. Host-microbe interactions that facilitate gut colonization by commensal bifidobacteria. Trends Microbiol 20:467–476. https://doi.org/10.1016/ j.tim.2012.07.002.

- 4. Valdés-Varela L, Ruas-Madiedo P, Gueimonde M. 2017. *In vitro* fermentation of different fructo-oligosaccharides by *Bifidobacterium* strains for the selection of synbiotic combinations. Int J Food Microbiol 242:19–23. https://doi.org/10.1016/j.ijfoodmicro.2016.11.011.
- Medina M, Izquierdo E, Ennahar S, Sanz Y. 2007. Differential immunomodulatory properties of *Bifidobacterium longum* strains: relevance to probiotic selection and clinical applications. Clin Exp Immunol 150: 531–538. https://doi.org/10.1111/j.1365-2249.2007.03522.x.
- EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA). 2012.
 Scientific opinion on the substantiation of health claims related to non-characterised micro-organisms (ID 2936, 2937, 2938, 2941, 2944, 2965, 2968, 2969, 3035, 3047, 3056, 3059, further assessment) pursuant to Article 13 (1) Of regulation (EC) no 1924/2006. EFSA J 10:2854. https://doi.org/10.2903/j.efsa.2012.2854.
- Inturri R, Stivala A, Sinatra F, Morrone R, Blandino G. 2014. Scanning electron microscopy observation of adhesion properties of *Bifidobacte-rium longum* W11 and chromatographic analysis of its exopolysaccharide. Food Nutr Sci 2014:1787–1792.
- Graziano T, Amoruso A, Nicola S, Deidda F, Allesina S, Pane M, Piffanelli P, Strozzi F, Mogna L, Del Piano M. 2016. The possible innovative use of Bifidobacterium longum W11 in association with rifaximin: a new horizon for combined approach? J Clin Gastroenterol 50(Suppl 2):S153–S156 (Proceedings from the 8th Probiotics, Prebiotics & New Foods for Microbiota and Human Health meeting held in Rome, Italy, 13 to 15 September 2016). https://doi.org/10.1097/MCG.000000000000683.
- Lugli GA, Milani C, Mancabelli L, van Sinderen D, Ventura M. 2016. MEGAnnotator: a user-friendly pipeline for microbial genomes assembly and annotation. FEMS Microbiol Lett 363. https://doi.org/10.1093/femsle/ fnw049.

Volume 5 lssue 10 e01659-16 genomea.asm.org **2**