

## Addendum

## A novel eukaryotic factor for cytosolic Fe-S cluster assembly

## Amit Roy, Natalia Solodovnikova, Tracy Nicholson, William Antholine and William E Walden

The EMBO Journal (2004) 23, 4878. doi:10.1038/sj.emboj.7600518

Addendum to: The EMBO Journal (2003) 22, 4826-4835. doi:10.1093/emboj/cdg455

In the above article, the authors twice made the statement that Cfd1p was the first iron-sulfur cluster assembly factor to be found in the cytoplasm. Other recent publications have suggested that proteins that have been shown to function in Fe-S cluster assembly in mitochondria may exist at low levels in the cytoplasm (Land and Rouault, 1998; Tong and Rouault, 2000; Nakai et al, 2001). Since this article went to press, two additional papers have added to these findings (Tong et al, 2003; Nakai et al, 2004).

While some investigators have questioned whether Cfd1p is indeed the first reported cytosolic Fe-S cluster assembly factor, we emphasize that the significance of our finding is in identifying a new class of cluster assembly factor whose localization is exclusively cytoplasmic and whose function is demonstrably cytoplasmic as well.

## References

Land T, Rouault TA (1998) Targeting of a human iron-sulfur cluster assembly enzyme, nifs, to different subcellular compartments is regulated through alternative AUG utilization. Mol Cell 2: 807-815 Nakai Y et al (2001) Nuclear localization of yeast Nfslp is required for cell survival. J Biol Chem 276: 8314-8320

Nakai Y et al (2004) Yeast Nfslp is involved in thio-modification of both mitochondrial and cytoplasmic tRNAs. J Biol Chem 279: 12363-12368

Tong WH, Rouault T (2000) Distinct iron-sulfur cluster assembly complexes exist in the cytosol and mitochondria of human cells. EMBO J 19: 5692-5700

Tong WH et al (2003) Subcellular compartmentalization of human Nfu, an iron-sulfur cluster scaffold protein, and its ability to assemble a [4Fe-4S] cluster. Proc Natl Acad Sci USA 100: 9762-