

REVIEW

Combinatorial strategies for combating invasive fungal infections

Michaela Spitzer[#], Nicole Robbins[#], and Gerard D. Wright

Michael G. DeGroot Institute for Infectious Disease Research and the Department of Biochemistry and Biomedical Sciences, McMaster University, Hamilton, ON, Canada

ABSTRACT

Invasive fungal infections are an important cause of human mortality and morbidity, particularly for immunocompromised populations. However, there remains a paucity of antifungal drug treatments available to combat these fungal pathogens. Further, antifungal compounds are plagued with problems such as host toxicity, fungistatic activity, and the emergence of drug resistance in pathogen populations. A promising therapeutic strategy to increase drug effectiveness and mitigate the emergence of drug resistance is through the use of combination drug therapy. In this review we describe the current arsenal of antifungals in medicine and elaborate on the benefits of combination therapy to expand our current antifungal drug repertoire. We examine those antifungal combinations that have shown potential against fungal pathogens and discuss strategies being employed to discover novel combination therapeutics, in particular combining antifungal agents with non-antifungal bioactive compounds. The findings summarized in this review highlight the promise of combinatorial strategies in combatting invasive mycoses.

ARTICLE HISTORY

Received 4 April 2016
Revised 24 May 2016
Accepted 25 May 2016

KEYWORDS

antifungal; *Aspergillus*; *Candida*; combination therapy; *Cryptococcus*; drug synergy

Introduction

The discovery of antimicrobial drugs to treat infections caused by bacterial and fungal pathogens is a challenging endeavor. The golden era of antibiotics, from the 1940s to 1970s, witnessed the discovery of novel chemical scaffolds produced by bacteria and fungi that revolutionized modern medicine. However in the past 30 y only 2 mechanistically and structurally new classes of antibacterial antibiotics have reached the clinic: linezolid and daptomycin.¹ Similarly, the only novel antifungal drug class to reach the clinic in ~30 y are the echinocandins.² This paucity of new antimicrobial drugs is further complicated given the widespread and ubiquitous evolution of resistance.³ Surveys of environmental microorganisms reveal that collections of microbial resistance genes (termed the antibiotic resistome) are genetically diverse,^{3–5} widespread across multiple environmental niches,⁶ and pre-date the modern antibiotic era by millennia.⁴ To support these conclusions, experience has shown time and again that the success of any new antimicrobial is ultimately undermined by the evolution of resistance that follows. Most recently the emergence and worldwide dissemination of resistance determinants NDM-1 and MCR-1 have breached the efficacy of

carbapenems and colistin respectively, “last-resort” antibiotics in our arsenal to combat gram-negative bacterial pathogens.^{7–9} Antimicrobial resistance is a serious threat to public health and as we enter a potential post-antibiotic age there is a dire need for novel treatment strategies to combat microbial infections.

Pathogenic fungi have emerged over the past decades as an increasing global threat to human health. Fungi are generally opportunistic in nature, preying on hosts with compromised immune systems due to infection with HIV or as a consequence of modern medical breakthroughs like chemotherapy and organ transplantation. They are the causative agents of a spectrum of diseases ranging from superficial infections affecting 1.7 billion individuals worldwide to invasive infections that kill 1.5 million humans per year.¹⁰ Notably, epidemiological data for fungal infections are unacceptably poor due to a lack of standards for reporting fungal disease and problems with misdiagnosis. Thus, fungal pathogens are likely an even greater contributor to human morbidity and mortality than current estimates. Species of *Candida*, *Cryptococcus*, *Aspergillus*, and *Pneumocystis* are at the forefront of fungal infections, accounting for an

CONTACT Gerard D. Wright  wrightge@mcmaster.ca  McMaster University, 1200 Main Street West, Hamilton, ON L8N 3Z5, Canada.

Color versions of one or more of the figures in the article can be found online at www.tandfonline.com/kvir.

[#]These authors contributed equally to this work.

© 2017 Taylor & Francis

estimated 90% of human mortality cases.¹⁰ Problems of antifungal drug resistance, both intrinsic and acquired, are common among all of these species across multiple antifungal drug classes. For example, the newest class of antifungal drug, the echinocandins, is ineffective against *Cryptococcus* species leaving the treatment of choice for cryptococcal meningitis reliant on medications developed in the 1950s that are plagued with problems of host toxicity.¹¹ Further, the increased deployment of the azole antifungals, both prophylactically and to treat active infections, has increased the prevalence of fluconazole-resistant *Candida* infections to ~3,400 annually in the United States alone, warranting the Center for Disease Control and Prevention to rank fluconazole-resistant *Candida* as a serious threat, the same threat level as methicillin-resistant *Staphylococcus aureus* (MRSA).¹²

A promising strategy for combatting resistant microbes is to extend the lifespan and efficacy of our currently employed drugs by using combination therapy. Combining drugs has the potential to confer enhanced efficacy and specificity compared to individual drug treatments and can slow the evolution of resistance.^{13,14} Further, by carefully selecting specific drug combinations, microbial drug resistance may not only be neutralized but also reversed through a process called selection inversion.¹⁵ Combination therapy is already the treatment of choice for many infectious diseases including HIV,¹⁶ tuberculosis¹⁷ and malaria.¹⁸ Consequently, the use of drug combinations to treat fungal pathogens has garnered considerable interest over the past several years. This review will describe the advantages of combination therapies and will highlight those antifungal drug combinations that show considerable promise in treating invasive mycoses.

Major classes of antifungal drugs

The therapeutic options for treating invasive fungal infections remain limited to only 3 structurally distinct classes of compounds. In contrast, there are over 2 dozen classes of antibacterials and 6 distinct classes of antiretroviral drugs available for clinical use.¹⁹ This is in large part due to the close evolutionary relatedness between humans and fungi, limiting the number of unique fungal cellular targets that can be exploited for drug development.

The polyenes (Fig. 1) are broad-spectrum natural product antifungals originally discovered in the 1950s. For decades it was accepted that the amphipathic nature of the polyenes allowed them to directly interact with the membrane lipid ergosterol, resulting in pore formation, membrane permeabilization, cell leakage and ultimately cell death.²⁰ Recently, elegant structure activity studies concluded channel formation is not the primary killing mechanism and that polyenes instead form large

extramembranous aggregates that extract the essential membrane-lipid ergosterol from the plasma membrane (Fig. 1).²¹ Since ergosterol serves as a central molecular node for multiple essential cellular processes, the ultimate cause of death is likely multifactorial. Given the structural similarities between ergosterol and the functional mammalian analog cholesterol, polyenes cause serious problems of systemic toxicity and nephrotoxicity in human patients when used in the clinic.²⁰ Consequently, several lipid formulations of amphotericin B have been developed with improved safety profiles and these continue to be widely deployed to treat life-threatening disseminated and invasive mycoses.^{22,23} Despite their long-term use in the clinic, resistance to the polyenes remains extremely rare.^{2,24}

The azoles are synthetic compounds first introduced as antifungals in the 1980s. They function by disrupting ergosterol biosynthesis through inhibition of the cytochrome P-450-dependent enzyme lanosterol 14- α -demethylase (Fig 1).² Subsequent ergosterol depletion and accumulation of toxic ergosterol precursors result in induction of severe membrane stress on the cell, thus inhibiting growth.² Azoles are chemically classified as either imidazoles if they possess 2 nitrogen atoms in the azole ring, or triazoles if they have three. Imidazoles are generally employed to treat superficial infections and only the triazoles fluconazole, itraconazole, voriconazole, and posaconazole are approved as drugs for systemic infections due to their favorable pharmacokinetic and safety profiles.²⁰ Given their low toxicity, they are widely used in the clinic as an initial therapy for most fungal infections, and as a prophylactic treatment for high-risk patients.^{22,25,26} Unfortunately, given their widespread use, resistance to the azoles is widespread particularly in *Candida* species.¹²

The echinocandins are the newest class of antifungal to reach the clinic, first entering the market in 2001. These molecules are large semi-synthetic lipopeptides that inhibit the cell wall enzyme complex β -1,3-D-glucan synthase, disrupting cell wall integrity and resulting in fungal cell death (Fig. 1).²⁷ Currently, there are 3 echinocandin drugs available: caspofungin, anidulafungin, and micafungin. Although echinocandins are well tolerated with little to no side effects, they are poorly absorbed when administered orally and completely ineffective against *Cryptococcus* or *Fusarium* species.²⁷ Echinocandin resistance has been reported in both laboratory and clinical settings, and the incidence of echinocandin resistant infections continues to rise.^{28,29}

Finally, although the rapid emergence of resistance to the fluorinated pyrimidine flucytosine precludes its use as a single agent, it is often combined with amphotericin B. This antifungal drug combination is the standard treatment for infections caused by *C.*




Figure 1. Structures and mechanisms of action of clinically relevant antifungal drugs. Fungal cells consist of a phospholipid membrane bilayer containing the sterol ergosterol. Ergosterol is synthesized within the cell in the endoplasmic reticulum. Fungi also possess a complex cell wall consisting of (1,3)- β -D-glucans covalently linked to (1,6)- β -D-glucans as well as chitin, mannans, and cell wall proteins (center panel). (A) Polyenes such as amphotericin B primarily exist in the form of large extramembranous aggregates that extract ergosterol from lipid bilayers. Structure of amphotericin B is shown. (B) Pyrimidines such as flucytosine becomes rapidly deaminated in the cytosol to generate 5-fluorouracil (5-FU) by fungal specific cytosine deaminases. 5-FU acts as a potent antimetabolite that causes RNA miscoding and inhibits DNA synthesis. Structure of flucytosine is shown. (C) The azoles function by targeting the ergosterol biosynthetic enzyme lanosterol demethylase, encoded by *ERG11* (*C. albicans* and *C. neoformans*) or *cyp51A* and *cyp51B* (*A. fumigatus*), causing a block in the production of ergosterol and the accumulation of a toxic sterol (yellow sunburst) produced by Erg3. This toxic sterol exerts a severe membrane stress on the cell. Structures of the clinically used azoles fluconazole and itraconazole are shown as examples. (D) The echinocandins act as noncompetitive inhibitors of (1,3)- β -D-glucan synthase (encoded by *FKS1* in *C. albicans*, *C. neoformans*, and *A. fumigatus* and by *FKS1* and *FKS2* in *C. glabrata* and *S. cerevisiae*) and thereby cause a loss of cell wall integrity and severe cell wall stress. Structures of the clinically used echinocandins caspofungin and anidulafungin are shown as examples.

neoformans, *C. gattii*, and other less common fungal species.^{10,11} Upon entry into the cytosol, flucytosine (also referred to as 5-fluorocytosine) becomes rapidly deaminated to generate 5-fluorouracil (5-FU) by

fungal-specific cytosine deaminases.³⁰ 5-FU acts as a potent antimetabolite that inhibits DNA synthesis and causes RNA miscoding, which ultimately impairs protein synthesis (Fig. 1).

Drug combinations to expand the therapeutic repertoire

Advantages of drug combinations

Combining antifungal drugs for treatment of invasive fungal infections has numerous advantages over monotherapy. In particular, drug combinations can delay or even prevent the development of resistance by rapidly reducing the pathogen population. Furthermore, it generally requires several mutations to accumulate to confer resistance to 2 drugs rather than one. Another combination strategy is the administration of an antibiotic in combination with an inhibitor of resistance enzymes. β -lactam antibiotics are routinely combined with inhibitors of serine β -lactamases, for example.³¹ Various studies have highlighted the potential of reversal of antibiotic resistance through combination treatment.¹⁵ Suppressive drug combinations have been shown to invert selective pressure to disfavor drug resistant mutants.³² The type of drug interactions can change in response to mutations³³ and drug combinations can become synergistic in resistant mutants.³⁴ Selection inversion can also occur when resistance to one drug also confers sensitivity to the second drug, a phenomenon called collateral sensitivity.³⁵ Most of these studies were done *in vitro* and need to be validated *in vivo*. Further, synergistic drug combinations allow for lower doses of each drug and shorter treatment duration reducing host-toxicity.^{13,14} Finally, compound combinations might improve fungicidal efficacy through synergy and result in greater therapeutic effect and broader activity than can be achieved with either drug alone.³⁶

The concept of synthetic lethality and its application to drug combinations

An additional advantage of combination therapy is that it has the potential to unveil a plethora of additional antifungal targets given that eukaryotic genomes are proven to be highly interconnected and functionally redundant. This is especially true for combinations of antifungal drugs with non-antifungal bioactive compounds. The complexity of cellular networks and the multifactorial nature of many diseases suggest that multi-component therapies might be more effective than single agents. Biological systems contain many features such as crosstalk, feedback and feed-forward loops which systems biology is only beginning to unravel. The functional redundancy and extensive buffering in biological networks is evident from the fact that only ~1,000 of the ~6,000 genes in budding yeast are essential.³⁷ Systematic screens have explored this robustness by mapping genetic interactions in the model organism *Saccharomyces cerevisiae*.³⁸ A genetic interaction between 2 genes is

observed when a phenotype caused by a mutation in one gene is exacerbated by a mutation in another gene such that the combined effect exceeds the sum of the individual effects.³⁹ Synthetic lethality represents the extreme case of a negative genetic interaction.^{1,40} Genome-wide analyses uncovered that most single deletion strains are sensitive to additional perturbations, such as a second genetic perturbation³⁸ or environmental and chemical stresses.⁴¹ The mapping of over 200,000 genetic interactions suggests that effective antifungal therapies may require the inhibition of multiple cellular targets simultaneously.⁴²⁻⁴⁶

For example, in *S. cerevisiae* the echinocandin target *FKS1* and its paralog *FKS2*, encode the biosynthetic enzyme for (1,3)- β -D-glucan synthesis. Genetic interaction networks have highlighted that *FKS1* is synthetic lethal with *CHS3*, a chitin synthase required for the synthesis of chitin,³⁸ and pharmacological inhibition of chitin synthases with nikkomycin strongly potentiates caspofungin against numerous fungal pathogens.^{47,48} Further, *FKS1* is synthetic lethal with *CNBI*, the regulatory subunit of calcineurin and calcineurin inhibitors exert potent synergy with echinocandins against diverse fungi.⁴⁹ While these examples highlight the power of genome-wide genetic interaction maps at predicting small molecule interactions, targeting 2 genes that genetically interact does not always result in successful pharmacological synergy.^{50,51} However, computational approaches have been developed that predict small molecule synergies based on gene expression data⁵² and chemical-genetic interactions⁵¹ that can increasingly be deployed to guide combination drug development.

Mechanisms of drug interactions

Various efforts have been made to classify mechanisms of drug interactions.⁵³⁻⁵⁵ However, many drug combinations have multiple mechanisms and do not fall cleanly into a single category. The combination of 2 drugs might be synergistic because compound one is effective against a disease on its own and the second drug increases its effective concentration at the target site (Fig. 2).⁵⁴ In this case the second drug does not have a direct effect of its own. It could affect pharmacokinetics of the first drug by increasing the rate or extent of absorption or the distribution. It could also target the first drug to the intended site of action or slow down metabolism or elimination of the first drug. This type of interaction has also been described as the bioavailability model.⁵⁰ Targeting of the fungal cell membrane by azoles and amphotericin B may explain synergism of these drugs with flucytosine.^{56,57} Alternatively, 2 drugs that target different stages of the same biological pathway or even the same protein (i.e., the same disease component) can also




Figure 2. Mechanisms of drug potentiation. (A) Drug potentiation through the bioavailability model suggests 2 drugs will be synergistic if one drug's action (Drug 2) helps another drug's availability (Drug 1) in the target cell. This could be achieved by increasing a drug's entry into the cell, by decreasing a drug's degradation or by affecting the pharmacokinetics of a drug within a host. (B) Alternatively, 2 drugs that target different stages of the same biological pathway can enhance pharmaceutical efficacy when combined, often in an additive manner. These combinations allow greater effects and/or reduced toxicities due to lower single doses. (C) A final model for drug synergy, the parallel pathway inhibition model, suggests 2 drugs will be synergistic if they target 2 proteins on parallel pathways that converge on an essential process.

increase pharmaceutical efficacy when combined, generally in an additive manner (Fig. 2). These combinations allow greater effects or reduced toxicities due to lower single doses. Both, terbinafine and azoles, target fungal ergosterol biosynthesis and impair the function of the cell membrane and when combined can improve efficacy.^{58,59} Finally, combination treatment with drugs that target separate cellular pathways converging on the same essential biological function can result in synergistic drug interactions, analogous to synthetic lethal interactions between 2 genes as mentioned above (Fig. 2). The drug interaction network of 21 antibiotics revealed synergistic interactions between specific classes of antibiotics that target different cellular functions.⁶⁰ Combinations of echinocandins with azoles and polyenes are synergistic and target the fungal cell wall and membrane simultaneously.^{61,62}

Concepts and terminology for combination therapy

Various models and concepts have been developed to study and assess antimicrobial compound combinations *in vitro* and *in vivo*.^{53,63} These tools have been reviewed extensively in the context of the debate about the definition of synergy.⁶³ In this section we will discuss the 2 main mathematical models that underlie most approaches developed to evaluate the effect of compound combinations. Both approaches consist of a conceptual model used to predict the expected result for a combination and the phrases used to categorize the results.

The first model was defined by Siegfried Walter Loewe and is based on the assumption of additive interactions.⁶⁴ Loewe assumed that an agent cannot interact with itself, and consequently, Loewe additivity is observed when an

agent is combined with itself and is the term used for indifferent combinations (Fig. 3B). In contrast, Chester Ittner Bliss assumed that the relative effect of a drug at a specific concentration is independent of the presence of another drug and Bliss independence is used to describe indifferent combinations (Fig. 3C).⁶⁵ For both, the additive and the multiplicative model, if the effect of a combination is better than expected the combination is said to be synergistic, and results worse than expected indicate antagonistic combinations. Many sophisticated approaches have been developed to deal with the complex nature of compound interactions that are based on the additive or multiplicative models just described.⁶³

Quantification of compound interactions in the laboratory is traditionally done by calculation of fractional inhibitory concentration (FIC) index (FICI) based on data obtained with the checkerboard method, a simple 2-dimensional array of serial dilutions of 2 test compounds (Fig. 3A).⁶⁶ Other methods to assess compound interactions include E-test, time-kill and disk diffusion assays.^{67,68} Due to its simplicity, the checkerboard method is the most commonly used approach to assess compound interactions *in vitro*. Checkerboard results are usually interpreted by constructing an isobologram based on dose pairs that give a specified effect (Fig. 3B). The axes of this graph are the doses of the 2 drugs, and




Figure 3. Loewe additivity and Bliss independence to define drug interactions. (A) Checkerboards that are representative of synergistic, indifferent and antagonistic drug interactions. Concentrations of drugs A and B increase along the x and y-axis, respectively. (B) Representative isoboles for synergistic, additive and antagonistic drug interactions based on the Loewe model. The x and y-axis represent the concentration of drugs A and B necessary to achieve a defined growth inhibition level X . The dashed line indicates additivity and is constructed by connecting the IC_X values for drugs A and B. Concentration pairs for drugs A and B used in combination to achieve the same growth inhibition X are added to this plot (red dots). The position of these points below, on or above the line result in concave, linear or convex isoboles and indicate synergistic, indifferent or antagonistic drug interactions, respectively. (C) Examples of drug interactions that are synergistic, indifferent and antagonistic according to the Bliss independence model. Growth in the presence of drugs A and B (red bars) is compared to the expected growth in the presence of the drug combination (E_{combo}) based on treatment with drugs A and B alone: $E_{\text{combo}} = G_{\text{Drug A}} * G_{\text{Drug B}}$ (indicated by dashed line). If the observed growth is less than expected, a drug combination is classified as synergistic. Conversely, more growth than expected indicates an antagonistic drug interaction.

the axial intercepts represent the dose of each agent that when used alone cause growth inhibition. If the selected effect is half the maximum effect (E_{max}), then the intercepts are the ED_{50} values. A linear isobole represents the expected concentration pairs when the 2 compounds do not interact ('indifference'). Deviation below the linear isobole indicates synergism and deviation in the other direction is indicative of antagonism (Fig. 3B). The deviation from indifference can be quantified by calculating FICI values. The FICI is defined as the sum of the FICs of each individual drug tested:

$$FICI = FIC_{Drug A} + FIC_{Drug B} \\ = \frac{MIC_{Drug A \text{ in combination}}}{MIC_{Drug A \text{ alone}}} + \frac{MIC_{Drug B \text{ in combination}}}{MIC_{Drug B \text{ alone}}}$$

In terms of the Loewe additivity model, we see indifference when $FICI = 1$, which is the same as to say that combining equal amounts of 2 indifferent compound yields the same effect as doubling the amount of a single compound. In terms of the isobole, this represents a point on the linear isobole.

When reproducible variations from an FICI of 1 are observed, the inherent inaccuracy of MIC methodologies and the question of biological relevance must be considered. Thus, conservative interpretation of results has been proposed by restricting interpretations of "synergy" to $FICI \leq 0.5$ and "antagonism" to $FICI > 4$ while $0.5 < FICI \leq 4$ indicates "indifference."⁶⁹ These interpretative categories are based on the assumption that checkerboard assays are carried out with 2-fold dilutions of compounds and that the range of experimental error for MIC determination is within one dilution step. For a result to be synergistic and achieve a $FICI \leq 0.5$, the MIC of both drugs has to decrease at least 4-fold. Importantly, a FICI of 0.5001 does not indicate synergy, meaning that if the MIC of one drug only decreases 2-fold, even if the FIC of the other drug decreases significantly, the interaction is reported as indifferent since the FIC of the first drug will be 0.5. Conversely, antagonism is observed when the MIC of at least one drug increases 4-fold. Since the condition for antagonism is $FICI > 4$, the situation where one drug dramatically increases the MIC of the other drug is included in the definition. As mentioned above, it is hard to assess drug interactions based on a single number (like the FICI) because interactions between compounds are not usually smooth and linear.

Combination therapy with antifungal drugs

The efficacy of various antifungal drug combinations against different fungal pathogens has been assessed

in vitro, *in vivo*, and in clinical settings.^{53,70-72} An advantage of *in vitro* studies is the ease of the methodology and analysis, making it feasible to test a variety of drug combinations against multiple fungal pathogens. However, checkerboard assays only assess the susceptibility of the fungus itself to the drug combination precluding host factors and pharmacokinetic parameters from being taken into consideration.⁷³ Factors that contribute to clinical efficacy of a drug combination include fungal virulence and resistance, the host immune condition and their interaction with the therapeutic agents. In addition, *in vitro* results for activity of specific drug combinations against specific fungi often differ between laboratories.⁵³ The checkerboard dilution method is most commonly used to assess drug interactions *in vitro*.⁶⁹ However, this method has demonstrated poor inter-laboratory reproducibility due to the fact that it is not standardized. Animal models of invasive fungal infections allow for pharmacokinetics of the administered drugs, tissue burden, and rate of clearance can be assessed. However, animal models generally do not account for a suppressed immune system and pharmacokinetics in animals differ from this in humans.⁷³ Despite their limitations, *in vitro* and *in vivo* studies are important for framing new hypotheses. Carefully designed clinical trials constitute the ultimate reference for treatment guidelines. The downsides of clinical data include the heterogeneous nature of subjects, the variety of fungal pathogen isolates, as well as comorbidity and dose verification across different centers. Since clinical data are the most informative with respect to treating systemic fungal infections, the following sections will focus on clinical studies that assess antifungal-antifungal drug combinations.

Cryptococcus

Current recommendations for the treatment of cryptococcal meningitis consists of 3 phases: an initial induction therapy for 2 weeks, followed by consolidation therapy for 8 weeks, and subsequent maintenance therapy for 6–12 months or until restoration of host immunity.⁷⁴ Treatment options for cryptococcosis are limited to amphotericin B, flucytosine, and fluconazole, alone or in combination. The induction therapy is an efficient fungicidal regimen, typically consisting of amphotericin B with flucytosine, and consolidation and maintenance therapies are usually monotherapies of fluconazole.⁷⁴ The other triazoles show activity against cryptococcal isolates *in vitro*,⁷⁵ but robust studies with these agents are not available yet. The combination of amphotericin B and flucytosine is the 'gold standard' for treatment of cryptococcal meningitis in the Infectious Disease Society of America and the World Health Organization treatment guidelines.^{74,76} Several

clinical studies have demonstrated superior efficacy of this combination using different mycological endpoints and observed faster cerebral spinal fluid sterilization, fewer relapses and lower mortality.^{11,77-81} Patients receiving the amphotericin B and flucytosine combination need to be closely monitored for renal dysfunction, however studies on cryptococcal meningitis had previously shown that this combination reduces toxicity of the polyene component because clinicians can give lower amphotericin B doses for a shorter time.^{77,82} Further, there is evidence that lower doses of flucytosine are just as fungicidal.⁸³ Flucytosine availability is limited in settings where disease burdens and mortality are highest and the inferior combination of amphotericin B and fluconazole is often an alternative in settings with limited resources.^{23,75}

Higher order combinations of antifungal agents have been tested against *Cryptococcus neoformans*. The combined effect of amphotericin B, flucytosine and fluconazole has been assessed in a mouse model of cryptococcal meningitis and in clinical settings,^{81,84,85} however, the triple combination did not outperform amphotericin B and flucytosine.

Candida

The echinocandins demonstrate fungicidal activity against most *Candida* species and are therefore generally recommended as primary therapy.^{25,86} Treatment with fluconazole is suggested as an alternative initial therapy. Combination treatment is not recommended for invasive candidiasis (IC) in the latest guideline published by the Infectious Diseases Society of America, since combinations of antimycotic agents are generally not more effective against *Candida* species compared to single agents.²⁵ The largest clinical trial involved 219 non-neutropenic patients with mostly *C. albicans* infections and found similar clinical outcomes for fluconazole alone compared with fluconazole and amphotericin B treatment.⁸⁷ There is limited evidence of effective combinations against IC. For example, amphotericin B in combination with flucytosine has been shown to be more effective than fluconazole at treating peritonitis.⁸⁸ Further, the combination of amphotericin B and flucytosine has also been shown to be more effective than amphotericin B alone for treatment of *Candida* meningitis, an uncommon manifestation of systemic candida infections.⁸⁹

Aspergillus

Voriconazole is currently the recommended first line therapy for invasive aspergillosis (IA),⁹⁰ improving clinical outcomes for patients with transplantation and leukemia (response rate 53%) while reducing severe side

effects from amphotericin B.⁹¹ Isavuconazole, a newer triazole, has been approved for treatment of IA and is better tolerable than voriconazole.⁹² Echinocandins have been approved for salvage therapy, but given their fungistatic activity against *Aspergillus*, efficacy is only 33% in severely immunocompromised patients.⁹³ The search for effective therapeutics is therefore ongoing. A systematic review of 7 observational studies and one randomized controlled trial (RCT) up until 2011 included data for 1071 patients and found evidence for combination anti-fungal therapy for primary IA to be conflicting and stresses the need for well-designed RCTs.⁹⁴ Retrospective reporting is susceptible to biases because outcomes in patients with different underlying diseases are compared, combination treatments are used in cases of severe disease, and selective reporting of positive outcomes.⁷³ The weak support for combination treatment is most likely due to the variety of antifungal combinations tested, the nature of IA investigated (6 studies looked at primary IA and 2 at salvage therapy for IA) and the heterogeneity of patient populations, treatment regimens and endpoints. Meta-analysis of 16 studies (including 1833 patients) comparing combinations of echinocandin with triazoles or amphotericin B against non-echinocandin monotherapy revealed that combination treatment improves clinical outcomes in salvage settings.⁹⁵ Due to the lack of definite proof that combination therapy is beneficial,^{96,97} initial combination therapy is not routinely recommended for IA, but is considered in high-risk patients.⁹⁰

Combinations of antifungal and non-antifungal drugs

Given the complex genetic landscape that exists in eukaryotic genomes coupled with the therapeutic superiority of the amphotericin B with flucytosine drug combination highlighted above, there is broad interest to identify other molecules that enhance the efficacy of our current antifungals. While none are yet in clinical use, a subset of the most promising drug combinations that show efficacy against pathogenic fungi are elaborated on below.

HSP90 inhibitors

One of the best-characterized examples of compounds capable of potentiating our current antifungals includes inhibitors of the essential molecular chaperone Hsp90. In pathogenic fungi, targeting Hsp90 function as an anti-fungal strategy holds considerable promise given that it governs crucial cellular responses to drug-induced stress.⁹⁸⁻¹⁰⁰ In *S. cerevisiae*, *C. albicans*, *C. neoformans* and *C. gattii* pharmacological inhibition of Hsp90 impairs the evolution of azole resistance and potentiates

azole activity *in vitro*.¹⁰¹⁻¹⁰³ Moreover, in *C. albicans*, *C. glabrata* and *A. fumigatus*, Hsp90 enables basal tolerance and resistance to the echinocandins.¹⁰⁴⁻¹⁰⁷ In *Caenorhabditis elegans* and murine models of fungal pathogenesis, combination therapy with Hsp90 inhibitors with caspofungin or fluconazole improves survival upon infection with *A. fumigatus* or *C. albicans* and *C. neoformans*, respectively.^{103,108} Furthermore, in mouse models of *C. albicans* disseminated disease, genetic compromise of fungal HSP90 expression reduces virulence¹⁰⁹ and enhances the therapeutic efficacy of fluconazole and caspofungin.^{104,108} Similarly, genetic compromise of *A. fumigatus* Hsp90 abrogates virulence in a murine model of invasive aspergillosis.¹⁰⁷ Finally, in an *in vivo* rat catheter model of *C. albicans* biofilm infection, clinically relevant Hsp90 inhibitors such as 17-AAG transform fluconazole from completely ineffective to highly efficacious without showing any signs of host toxicity.¹¹⁰ Thus, efforts to create fungal specific Hsp90 inhibitors holds considerable promise as a combination agent with multiple antifungals for the treatment of diverse fungal pathogens.

Calcineurin inhibitors

Hsp90 is one of the most highly connected hubs in cellular networks, interacting with an estimated 10% of the yeast proteome.^{111,112} Chemical genomic studies in *C. albicans* identified a multitude of Hsp90 genetic interactors important for cellular responses to azoles and echinocandins,¹¹³ suggesting that Hsp90 may have pleiotropic effects on circuitry governing drug resistance. One well-characterized Hsp90 client protein important for governing responses to antifungal-induced stress is the protein phosphatase calcineurin.¹⁰⁴ In pathogenic fungi, calcineurin regulates a myriad of physiological processes including cell cycle progression, cation homeostasis, morphogenesis, virulence, and antifungal drug responses.^{49,114} Pharmacological inhibition of calcineurin in *C. neoformans* abrogates growth at 37°C,¹¹⁵ potentiates the activity of both azoles and echinocandins in *C. albicans*,^{101,102,104,116,117} and renders the fungistatic azoles fungicidal against multiple *Candida* species.³⁶ Further, calcineurin inhibitors cyclosporine A and FK506 act synergistically with azoles against *C. albicans* biofilms both *in vitro* and *in vivo*, implicating calcineurin as a key modulator of azole resistance during biofilm growth.¹¹⁸ In *A. fumigatus*, calcineurin inhibitors enhance echinocandin activity, and transform the fungistatic activity of caspofungin to fungicidal.^{119,120} Calcineurin inhibitors also show potent activity against azole- and echinocandin-resistant strains of *A. fumigatus*.¹²¹ Recently, it was shown that calcineurin orchestrates dimorphic

transitions, antifungal drug responses, and virulence of the fungal pathogen *Mucor circinelloides*.^{122,123} The challenge that remains for antifungal drug discovery is to develop fungal specific calcineurin inhibitors or to identify fungal-specific calcineurin effectors that can be targeted pharmacologically.

Lysine deacetylases and lysine acetyltransferases

Additional cellular targets that have garnered considerable attention as an antifungal combination target are lysine deacetylases (KDACs) and lysine acetyltransferases (KATs). These enzymes catalyze the removal or addition of acetyl groups from lysine residues present on histones and other cellular proteins. The broad-spectrum KDAC inhibitor trichostatin A (TSA) potentiates azole activity in *C. albicans*.^{124,125} Further, a Hos2 KDAC inhibitor, MGCD290 (Mirati Therapeutics, San Diego, CA, USA), displays synergistic activity with azoles and echinocandins against diverse *C. albicans* drug-resistant clinical isolates *in vitro*.^{126,127} Additional *in vivo* studies coupled with promising preliminary clinical trials have supported the use of MGCD290 in combination with fluconazole to treat *C. albicans* infections.¹²⁸ KATs also modulate azole resistance. Genetic impairment of *C. albicans* ADA2, which encodes a component of the Spt-Ada-Gcn5-acetyltransferase (SAGA) coactivator complex, confers hypersensitivity to fluconazole due to impaired upregulation of efflux pumps Cdr1 and Mdr1.¹²⁹ Further, deletion of the *C. albicans* KAT gene *RTT109* confers hypersusceptibility to macrophages, altered metabolic gene expression, and induction of a weaker inflammatory response, culminating in attenuated virulence in a murine infection model.^{130,131} Deletion of *RTT109* also confers increased sensitivity to echinocandins, and other genotoxic stresses such as hydroxyurea and methyl methane sulfonate.¹³⁰ Thus, these KATs represent an excellent pharmacological target for future drug development.

Other targets

Other pharmacological inhibitors that target a variety of cellular processes have been reported to synergize with current antifungals. Many of these inhibitors are identified in the Antifungal Synergistic Drug Combination Database (ASDCD), a curated collection of 210 antifungal synergistic drug combinations involving 105 drugs from the literature (last updated 2013).¹³² Such compounds include, but are not limited to, pharmacological inhibitors of ADP-ribosylation factors,¹³³ protein kinase C¹³⁴, fungal sphingolipids,¹³⁵ and protein translation¹³⁶ in order to potentiate fluconazole. Further, tamoxifen and other triphenylethylene-based

estrogen receptor antagonists have antifungal activity against *C. neoformans*,^{137,138} and potentiate the activity of azoles, polyenes, and echinocandins in diverse fungal species.¹³⁹ Finally, the antidepressant sertraline combined with fluconazole provides improved activity relative to either drug alone in both an invertebrate model of cryptococcosis¹³⁸ and in a mouse model of disseminated cryptococcosis,¹³⁶ likely due to its membrane perturbing effects in fungi.¹³⁸

High-throughput screens

In addition to the several promising examples highlighted above, scientists in both academia and industry have leveraged the power of high-throughput screening methodologies in order to uncover novel antifungal combinations. Compounds that enhance the activity of known antimicrobials in *S. cerevisiae* have been identified both by focused small molecule library screens and by computational methods.^{140-142,51} Further, several high-throughput screening endeavors have searched for molecules that potentiate the azoles and overcome azole resistance in fungal pathogens themselves.^{138,139,143,144} Such compounds that increase azole efficacy are often found to target membrane function or sphingolipid biosynthesis.¹³⁸ One such example is the antimycobacterium compound clofazimine, which induces a cell membrane stress in fungi and enhances the efficacy of fluconazole in an *in vivo* *Galleria mellonella* model of fungal pathogenesis.¹³⁹ Further, screens performed as part of the NIH-Molecular Libraries and Probes Screening network identified a class of indole derivatives that restore fluconazole susceptibility to resistant *C. albicans* isolates.¹⁴³ Finally, high-throughput screens using microbial extracts identified the natural product cyclic hexadepsipeptide beauvericin as a potentiator of ketoconazole against *C. parapsilosis*.¹⁴⁵

In addition to employing high-throughput screens to look for agents that enhance the activity of the azoles, studies have identified molecules that potentiate the other antifungal drug classes. Recently, we screened sublethal concentrations of 6 known antifungals in combination with ~3,600 bioactive compounds against diverse fungal species to uncover synergistic drug combinations.¹³⁹ This dataset, termed the Antifungal Combination Matrix (ACM), identified ~1550 chemical combinations that abrogate fungal growth often in an antifungal- and species-specific manner.¹³⁹ The number of compounds that potentiated amphotericin B and caspofungin far exceeded the number of compounds that potentiated the azoles. Another study has employed high-throughput screens to assay >300,000 molecules in search for chemical entities that specifically perturb cell

wall integrity, the mode-of-action of the echinocandins.¹⁴⁶ Three benzothioureas were discovered that exhibited anticytotoxic activity and acted in an additive manner with caspofungin.¹⁴⁶ This was profound given that the echinocandins do not show activity against *Cryptococcus* species. The molecular mechanism by which these benzothioureas elicited activity was through the inhibition of the cell wall integrity MAP kinase cascade.¹⁴⁶

Targeting resistance determinants

Small molecules that target resistance determinants are often effective pairings with currently available antimicrobials to combat drug resistant isolates. Examples of such drug therapies include drug combinations of β -lactam drugs with inhibitors of β -lactamase resistance enzymes.³¹ A recent study with *C. glabrata* identified a small molecule (iKIX1) that inhibits azole efflux pump expression by disrupting the interaction between the Mediator complex and the transcriptional activator Pdr1.¹⁴⁷ Follow-up *in vivo* studies demonstrated that iKIX1 improved the therapeutic efficacy of fluconazole against fluconazole-resistant *C. glabrata* in both *G. mellonella* and mouse models of infection.¹⁴⁷ It is difficult to predict which, if any, of these antifungal combinations will represent a breakthrough in the way we treat fungal infections, but it is promising to see the diverse molecular targets that are being identified.

Conclusions

Antifungal combination therapy is well established for the treatment of cryptococcosis and it is recommended for some hard to treat invasive *Candida* and *Aspergillus* infections. New classes of antifungal agents will improve the chances that combination therapies are more effective than currently available monotherapies. There are several antifungal drugs in the pipeline that target the biosynthesis of chitin, GPI anchor and heme.¹⁴⁸ This comes at a time where legislation including the US Food and Drug Administration's (FDA) Generating Antibiotics Now (GAIN) and the Orphan Drug Acts has been passed by government agencies to push forward the development of antifungal agents that offer increased market exclusivity and several fungal pathogens qualify for both schemes. Many studies have targeted the fungal stress response in combination with antifungal agents and the results are promising. High-throughput screens for antifungal synergistic combinations are revealing further biological pathways and chemical scaffolds that can be targeted to yield effective antifungal drug combinations. The biggest challenge for targeting these pathways

is the identification of fungal-specific compounds or effectors. In addition to new drugs, better diagnostic tools need to be developed. This will improve therapeutic outcomes because of prompt diagnosis and early treatment.^{149,150} It will also allow for development of narrow-spectrum antifungals with improved antifungal activity.

Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed.

Funding

NR was supported by the Michael G. DeGroot Postdoctoral Fellowship Award and a CIHR Fellowship. Research was supported by a CIHR Grant (FRN119572) to GDW that focuses on chemical genomic strategies for new antifungal drug leads and by a Canada Research Chair in Antibiotic Biochemistry to GDW.

References

- [1] Roemer T, Boone C. Systems-level antimicrobial drug and drug synergy discovery. *Nat Chem Biol* 2013; 9:222-31; PMID:23508188; <http://dx.doi.org/10.1038/nchembio.1205>
- [2] Shapiro RS, Robbins N, Cowen LE. Regulatory circuitry governing fungal development, drug resistance, and disease. *Microbiol Mol Biol Rev* 2011; 75:213-67; PMID:21646428; <http://dx.doi.org/10.1128/MMBR.00045-10>
- [3] Brown ED, Wright GD. Antibacterial drug discovery in the resistance era. *Nature* 2016; 529:336-43; PMID:26791724; <http://dx.doi.org/10.1038/nature17042>
- [4] D'Costa VM, King CE, Kalan L, Morar M, Sung WW, Schwarz C, Froese D, Zazula G, Calmels F, Debruyne R, et al. Antibiotic resistance is ancient. *Nature* 2011; 477:457-61; PMID:21881561; <http://dx.doi.org/10.1038/nature10388>
- [5] Forsberg KJ, Reyes A, Wang B, Selleck EM, Sommer MO, Dantas G. The shared antibiotic resistome of soil bacteria and human pathogens. *Science* 2012; 337:1107-11; PMID:22936781; <http://dx.doi.org/10.1126/science.1220761>
- [6] Forsberg KJ, Patel S, Gibson MK, Lauber CL, Knight R, Fierer N, Dantas G. Bacterial phylogeny structures soil resistomes across habitats. *Nature* 2014; 509:612-6; PMID:24847883; <http://dx.doi.org/10.1038/nature13377>
- [7] Nordmann P, Poirel L, Walsh TR, Livermore DM. The emerging NDM carbapenemases. *Trends Microbiol* 2011; 19:588-95; PMID:22078325; <http://dx.doi.org/10.1016/j.tim.2011.09.005>
- [8] Du H, Chen L, Tang YW, Kreiswirth BN. Emergence of the mcr-1 colistin resistance gene in carbapenem-resistant Enterobacteriaceae. *Lancet Infect Dis* 2016; 3:287-8; [http://dx.doi.org/10.1016/S1473-3099\(16\)00056-6](http://dx.doi.org/10.1016/S1473-3099(16)00056-6)
- [9] Malhotra-Kumar S, Xavier BB, Das AJ, Lammens C, Butaye P, Goossens H. Colistin resistance gene mcr-1 harboured on a multidrug resistant plasmid. *Lancet Infect Dis* 2016; 3:283-4; [http://dx.doi.org/10.1016/S1473-3099\(16\)00012-8](http://dx.doi.org/10.1016/S1473-3099(16)00012-8)
- [10] Brown GD, Denning DW, Gow NA, Levitz SM, Netea MG, White TC. Hidden killers: human fungal infections. *Sci Transl Med* 2012; 4:165rv13; PMID:23253612; <http://dx.doi.org/10.1126/scitranslmed.3004404>
- [11] Day JN, Chau TT, Wolbers M, Mai PP, Dung NT, Mai NH, Phu NH, Nghia HD, Phong ND, Thai CQ, et al. Combination antifungal therapy for cryptococcal meningitis. *N Engl J Med* 2013; 368:1291-302; PMID:23550668; <http://dx.doi.org/10.1056/NEJMoa1110404>
- [12] CDC. Antibiotic Resistance Threats in the United States 2013. Atlanta, GA.
- [13] Zimmermann GR, Lehar J, Keith CT. Multi-target therapeutics: when the whole is greater than the sum of the parts. *Drug Discovery Today* 2007; 12:34-42; PMID:17198971; <http://dx.doi.org/10.1016/j.drudis.2006.11.008>
- [14] Hill JA, Cowen LE. Using combination therapy to thwart drug resistance. *Future Microbiol* 2015; 10:1719-26; PMID:26597425; <http://dx.doi.org/10.2217/fmb.15.68>
- [15] Baym M, Stone LK, Kishony R. Multidrug evolutionary strategies to reverse antibiotic resistance. *Science* 2016; 351:aad3292; PMID:26722002; <http://dx.doi.org/10.1126/science.aad3292>
- [16] Bock C, Lengauer T. Managing drug resistance in cancer: lessons from HIV therapy. *Nat Rev Cancer* 2012; 12:494-501; PMID:22673150; <http://dx.doi.org/10.1038/nrc3297>
- [17] Zumla A, Hafner R, Lienhardt C, Hoelscher M, Nunn A. Advancing the development of tuberculosis therapy. *Nat Rev Drug Discov* 2012; 11:171-2; PMID:22378254; <http://dx.doi.org/10.1038/nrd3694>
- [18] Eastman RT, Fidock DA. Artemisinin-based combination therapies: a vital tool in efforts to eliminate malaria. *Nat Rev* 2009; 7:864-74; PMID:19881520
- [19] Roemer T, Krysan DJ. Antifungal drug development: challenges, unmet clinical needs, and new approaches. *Cold Spring Harb Perspect Med* 2014; 4; pii: a019703; PMID:24789878; <http://dx.doi.org/10.1101/cshperspect.a019703>
- [20] Ostrosky-Zeichner L, Casadevall A, Galgiani JN, Odds FC, Rex JH. An insight into the antifungal pipeline: selected new molecules and beyond. *Nat Rev Drug Discov* 2010; 9:719-27; PMID:20725094; <http://dx.doi.org/10.1038/nrd3074>
- [21] Anderson TM, Clay MC, Cioffi AG, Diaz KA, Hisao GS, Tuttle MD, Nieuwkoop AJ, Comellas G, Maryum N, Wang S, et al. Amphotericin forms an extramembranous and fungicidal sterol sponge. *Nat Chem Biol* 2014; 10:400-6; PMID:24681535; <http://dx.doi.org/10.1038/nchembio.1496>
- [22] Perfect JR, Dismukes WE, Dromer F, Goldman DL, Graybill JR, Hamill RJ, Harrison TS, Larsen RA, Lortholary O, Nguyen MH, et al. Clinical practice guidelines for the management of cryptococcal disease: 2010 update by the infectious diseases society of america. *Clin Infect Dis* 2010; 50:291-322; PMID:20047480; <http://dx.doi.org/10.1086/649858>
- [23] Pappas PG, Kauffman CA, Andes D, Benjamin DK, Jr., Calandra TF, Edwards JE, Jr., Filler SG, Fisher JF, Kullberg BJ, Ostrosky-Zeichner L, et al. Clinical practice

- guidelines for the management of candidiasis: 2009 update by the Infectious Diseases Society of America. *Clin Infect Dis* 2009; 48:503-35; PMID:19191635; <http://dx.doi.org/10.1086/596757>
- [24] Vincent BM, Lancaster AK, Scherz-Shouval R, Whitesell L, Lindquist S. Fitness trade-offs restrict the evolution of resistance to amphotericin B. *PLoS Biol* 2013; 11: e1001692; PMID:24204207; <http://dx.doi.org/10.1371/journal.pbio.1001692>
- [25] Pappas PG, Kauffman CA, Andes DR, Clancy CJ, Marr KA, Ostrosky-Zeichner L, Reboli AC, Schuster MG, Vazquez JA, Walsh TJ, et al. Clinical practice guideline for the management of candidiasis: 2016 update by the infectious diseases society of America. *Clin Infect Dis* 2016; 62: e1-e50
- [26] Walsh TJ, Anaissie EJ, Denning DW, Herbrecht R, Kontoyannis DP, Marr KA, Morrison VA, Segal BH, Steinbach WJ, Stevens DA, et al. Treatment of aspergillosis: clinical practice guidelines of the Infectious Diseases Society of America. *Clin Infect Dis* 2008; 46:327-60; PMID:18177225; <http://dx.doi.org/10.1086/525258>
- [27] Denning DW. Echinocandin antifungal drugs. *Lancet* 2003; 362:1142-51; PMID:14550704; [http://dx.doi.org/10.1016/S0140-6736\(03\)14472-8](http://dx.doi.org/10.1016/S0140-6736(03)14472-8)
- [28] Arendrup MC, Perlin DS. Echinocandin resistance: an emerging clinical problem?. *Curr Opin Infect Dis* 2014; 27:484-92; PMID:25304391; <http://dx.doi.org/10.1097/QCO.0000000000000111>
- [29] Walker LA, Gow NA, Munro CA. Fungal echinocandin resistance. *Fungal Genet Biol* 2010; 47:117-26; PMID:19770064; <http://dx.doi.org/10.1016/j.fgb.2009.09.003>
- [30] Loyse A, Dromer F, Day J, Lortholary O, Harrison TS. Flucytosine and cryptococcosis: time to urgently address the worldwide accessibility of a 50-year-old antifungal. *J Antimicrob Chemother* 2013; 68:2435-44; PMID:23788479; <http://dx.doi.org/10.1093/jac/dkt221>
- [31] King AM, Reid-Yu SA, Wang W, King DT, De Pascale G, Strynadka NC, Walsh TR, Coombes BK, Wright GD. Aspergillomarasmine A overcomes metallo-beta-lactamase antibiotic resistance. *Nature* 2014; 510:503-6; PMID:24965651; <http://dx.doi.org/10.1038/nature13445>
- [32] Chait R, Craney A, Kishony R. Antibiotic interactions that select against resistance. *Nature* 2007; 446:668-71; PMID:17410176; <http://dx.doi.org/10.1038/nature05685>
- [33] Chevereau G, Bollenbach T. Systematic discovery of drug interaction mechanisms. *Mol Syst Biol* 2015; 11(4):807-
- [34] Wood KB, Wood KC, Nishida S, Cluzel P. Uncovering Scaling Laws to Infer Multidrug Response of Resistant Microbes and Cancer Cells. *Cell Rep* 2014; 6(6):1073-84; <http://dx.doi.org/10.1016/j.celrep.2014.02.007>
- [35] Imamovic L, Sommer MOA. Use of collateral sensitivity networks to design drug cycling protocols that avoid resistance development. *Sci Transl Med* 2013; 5(204):204ra132-204ra132
- [36] Onyewu C, Blankenship JR, Del Poeta M, Heitman J. Ergosterol biosynthesis inhibitors become fungicidal when combined with calcineurin inhibitors against *Candida albicans*, *Candida glabrata*, and *Candida krusei*. *Antimicrob Agents Chemother* 2003; 47:956-64; PMID:12604527; <http://dx.doi.org/10.1128/AAC.47.3.956-964.2003>
- [37] Winzeler EA, Shoemaker DD, Astromoff A, Liang H, Bussey H, Chu AM, Foury F, Friend SH, Giaever G, Lockhart DJ, et al. Functional characterization of the *S. cerevisiae* genome by gene deletion and parallel analysis. *Science* 1999; 285(5429):901-6; PMID:10436161; <http://dx.doi.org/10.1126/science.285.5429.901>
- [38] Costanzo M, Baryshnikova A, Bellay J, Kim Y, Spear ED, Sevier CS, Ding H, Koh JLY, Toufighi K, Mostafavi S, et al. The genetic landscape of a cell. *Science* 2010; 327(5964):425-31; <http://dx.doi.org/10.1126/science.1180823>
- [39] Mani R, St Onge RP, Hartman JL, Giaever G, Roth FP. Defining genetic interaction. *Proc Natl Acad Sci USA* 2008; 105(9):3461-6; PMID:18305163; <http://dx.doi.org/10.1073/pnas.0712255105>
- [40] Dobzhansky T. Genetics of Natural Populations. Xiii. Recombination and Variability in Populations of *Drosophila Pseudoobscura*. *Genetics* 1946; 31(3):269-90
- [41] Hillenmeyer ME, Fung E, Wildenhain J, Hoon S, Lee W, Proctor M, St Onge RP, Tyers M, Koller D, Altman RB, et al. The chemical genomic portrait of yeast: uncovering a phenotype for all genes. *Science* 2008; 320(5874):362-5; <http://dx.doi.org/10.1126/science.1150021>
- [42] Sharom JR, Bellows DS, Tyers M. From large networks to small molecules. *Curr Opin Chem Biol* 2004; 8(1):81-90; <http://dx.doi.org/10.1016/j.cbpa.2003.12.007>
- [43] Csermely P. Strong links are important, but weak links stabilize them. *Trends Biochem Sci* 2004; 29(7):331-4; PMID:15236738; <http://dx.doi.org/10.1016/j.tibs.2004.05.004>
- [44] Fitzgerald JB, Schoeberl B, Nielsen UB, Sorger PK. Systems biology and combination therapy in the quest for clinical efficacy. *Nat Chem Biol* 2006; 2(9):458-66; <http://dx.doi.org/10.1038/nchembio817>
- [45] Kitano H. A robustness-based approach to systems-oriented drug design. *Nat Rev Drug Discov* 2007; 6(3):202-10; PMID:17318209; <http://dx.doi.org/10.1038/nrd2195>
- [46] Lehár J, Krueger AS, Avery W, Heilbut AM, Johansen LM, Price ER, Rickles RJ, Short GF, Staunton JE, Jin X, et al. Synergistic drug combinations tend to improve therapeutically relevant selectivity. *Nat Biotechnol* 2009; 27(7):659-66; <http://dx.doi.org/10.1038/nbt.1549>
- [47] Sandovsky-Losica H, Shwartzman R, Lahat Y, Segal E. Antifungal activity against *Candida albicans* of nikkomycin Z in combination with caspofungin, voriconazole or amphotericin B. *J Antimicrob Chemother* 2008; 62:635-7; PMID:18490373; <http://dx.doi.org/10.1093/jac/dkn216>
- [48] Verwer PE, van Duijn ML, Tavakol M, Bakker-Woudenberg IA, van de Sande WW. Reshuffling of *Aspergillus fumigatus* cell wall components chitin and beta-glucan under the influence of caspofungin or nikkomycin Z alone or in combination. *Antimicrob Agents Chemother* 2012; 56:1595-8; PMID:22203603; <http://dx.doi.org/10.1128/AAC.05323-11>
- [49] Steinbach WJ, Reedy JL, Cramer RA, Jr., Perfect JR, Heitman J. Harnessing calcineurin as a novel anti-infective agent against invasive fungal infections. *Nature Reviews* 2007; 5:418-30; PMID:17505522
- [50] Cokol M, Chua HN, Tasan M, Mutlu B, Weinstein ZB, Suzuki Y, Nergiz ME, Costanzo M, Baryshnikova A, Giaever G, et al. Systematic exploration of synergistic drug pairs. *Mol Syst Biol* 2011; 8: 1-9

- [51] Wildenhain J, Spitzer M, Dolma S, Jarvick N, White R, Roy M, Griffiths E, Bellows DS, Wright GD, Tyers M. Prediction of Synergism from Chemical-Genetic Interactions by Machine Learning. *Cell Systems* 2015; 1:383-95; PMID:27136353; <http://dx.doi.org/10.1016/j.cels.2015.12.003>
- [52] Bansal M, Yang J, Karan C, Menden MP, Costello JC, Tang H, Xiao G, Li Y, Allen J, Zhong R, et al. A community computational challenge to predict the activity of pairs of compounds. *Nat Biotechnol* 2014; 32: 1-12; PMID:24406907
- [53] Johnson MD, MacDougall C, Ostrosky-Zeichner L, Perfect JR, Rex JH. Combination antifungal therapy. *Antimicrob Agents Chemother* 2004; 48:693-715; PMID:14982754; <http://dx.doi.org/10.1128/AAC.48.3.693-715.2004>
- [54] Toews ML, Bylund DB. Pharmacologic principles for combination therapy. *Proc Am Thorac Soc* 2005; 2: 282-9; discussion 90-1; <http://dx.doi.org/10.1513/pats.200504-037SR>
- [55] Jia J, Zhu F, Ma X, Cao Z, Cao ZW, Li Y, Li YX, Chen YZ. Mechanisms of drug combinations: interaction and network perspectives. *Nat Rev Drug Discov* 2009; 8: 111-28; <http://dx.doi.org/10.1038/nrd2683>
- [56] Yamamoto Y, Maesaki S, Takeya H, Yanagihara K, Ohno H, Ogawa K, Hirakata Y, Tomono K, Tashiro T, Kohno S. Combination therapy with fluconazole and flucytosine for pulmonary cryptococcosis. *Chemotherapy* 1997; 43: 436-41; <http://dx.doi.org/10.1159/000239603>
- [57] Polak A. Synergism of polyene antibiotics with 5-fluorocytosine. *Chemotherapy* 1978; 24(1):2-16; PMID:338264; <http://dx.doi.org/10.1159/000237753>
- [58] Barchiesi F, Di Francesco LF, Compagnucci P, Arzeni D, Giacometti A, Scalise G. *In-vitro* interaction of terbinafine with amphotericin B, fluconazole and itraconazole against clinical isolates of *Candida albicans*. *J Antimicrob Chemother* 1998; 41(1):59-65; PMID:9511038; <http://dx.doi.org/10.1093/jac/41.1.59>
- [59] Perea S, Gonzalez G, Fothergill AW, Sutton DA, Rinaldi MG. *In vitro* activities of terbinafine in combination with fluconazole, itraconazole, voriconazole, and posaconazole against clinical isolates of *Candida glabrata* with decreased susceptibility to azoles. *J Clin Microbiol* 2002; 40(5):1831-3; PMID:11980970; <http://dx.doi.org/10.1128/JCM.40.5.1831-1833.2002>
- [60] Yeh P, Tschumi AI, Kishony R. Functional classification of drugs by properties of their pairwise interactions. *Nat Genet* 2006; 38(4):489-94; <http://dx.doi.org/10.1038/ng1755>
- [61] Shalit I, Shadkchan Y, Samra Z, Osherov N. In Vitro Synergy of Caspofungin and Itraconazole against *Aspergillus* spp.: MIC versus Minimal Effective Concentration End Points. *Antimicrob Agents Chemother* 2003; 47(4):1416-8; PMID:12654681; <http://dx.doi.org/10.1128/AAC.47.4.1416-1418.2003>
- [62] Arikan S, Lozano-Chiu M, Paetznick V, Rex JH. In Vitro Susceptibility Testing Methods for Caspofungin against *Aspergillus* and *Fusarium* Isolates. *Antimicrob Agents Chemother* 2001; 45(1):327-30; PMID:11120990; <http://dx.doi.org/10.1128/AAC.45.1.327-330.2001>
- [63] Greco W, Bravo G, Parsons J. The search for synergy: a critical review from a response surface perspective. *Pharmacol Rev* 1995; 47(2):331
- [64] Loewe S. The problem of synergism and antagonism of combined drugs. *Arzneimittelforschung* 1953; 3(6):285-90; PMID:13081480
- [65] Bliss CI. The toxicity of poisons applied jointly. *Ann Appl Biol* 1939; 26: 585-615
- [66] Eliopoulos GM, Moellering RC. Antimicrobial combinations. *Antibiotics in Laboratory Med* 1996; 4:330-396
- [67] Rex JH, Pfaller MA, Walsh TJ, Chaturvedi V, Espinel-Ingroff A, Ghannoum MA, Gosey LL, Odds FC, Rinaldi MG, Sheehan DJ, et al. Antifungal Susceptibility Testing: Practical Aspects and Current Challenges. *Clin Microbiol Rev* 2001; 14: 643-58; <http://dx.doi.org/10.1128/CMR.14.4.643-658.2001>
- [68] Cantón E, Espinel-Ingroff A, Pemán J. Trends in antifungal susceptibility testing using CLSI reference and commercial methods. *Expert Rev Anti-Infective Ther* 2014; 7(1):107-19
- [69] Odds FC. Synergy, antagonism, and what the checkerboard puts between them. *J Antimicrob Chemother* 2003; 52(1):1; PMID:12805255; <http://dx.doi.org/10.1093/jac/dkg301>
- [70] Mukherjee PK, Sheehan DJ, Hitchcock CA, Ghannoum MA. Combination Treatment of Invasive Fungal Infections. *Clin Microbiol Rev* 2005; 18(1):163-94; <http://dx.doi.org/10.1128/CMR.18.1.163-194.2005>
- [71] Vazquez JA. Clinical practice: combination antifungal therapy for mold infections: much ado about nothing?. *Clin Infect Dis* 2008; 46(12):1889-901; <http://dx.doi.org/10.1086/588475>
- [72] Hatipoglu N, Hatipoglu H. Combination antifungal therapy for invasive fungal infections in children and adults. *Expert Rev Anti-Infective Ther* 2013; 11: 523-35; <http://dx.doi.org/10.1586/eri.13.29>
- [73] Steinbach WJ. Combination antifungal therapy for invasive aspergillosis – Is it indicated?. *Med Mycol* 2006; 44: 373-82; <http://dx.doi.org/10.1080/13693780600860953>
- [74] Maziarz EK, Perfect JR. Cryptococcosis. *Infect Dis Clin NA* 2016; 30(1):179-206
- [75] Loyse A, Wilson D, Meintjes G, Jarvis JN, Bicanic T, Bishop L, Rebe K, Williams A, Jaffar S, Bekker L-G, et al. Comparison of the early fungicidal activity of high-dose fluconazole, voriconazole, and flucytosine as second-line drugs given in combination with amphotericin B for the treatment of HIV-associated cryptococcal meningitis. *Clin Infect Dis* 2012; 54: 121-8; <http://dx.doi.org/10.1093/cid/cir745>
- [76] Perfect JR, Bicanic T. Cryptococcosis diagnosis and treatment: What do we know now. *Fungal Genet Biol* 2015; 78:49-54; <http://dx.doi.org/10.1016/j.fgb.2014.10.003>
- [77] Bennett JE, Dismukes WE, Duma RJ, Medoff G, Sande MA, Gallis H, Leonard J, Fields BT, Bradshaw M, Haywood H, et al. A comparison of amphotericin B alone and combined with flucytosine in the treatment of cryptococcal meningitis. *N Engl J Med* 1979; 301:126-31; PMID:449951; <http://dx.doi.org/10.1056/NEJM197907193010303>
- [78] Dismukes WE, Cloud G, Gallis HA, Kerkering TM, Medoff G, Craven PC, Kaplowitz LG, Fisher JF, Gregg CR, Bowles CA. Treatment of cryptococcal meningitis with combination amphotericin B and flucytosine for four as compared with six weeks. *N Engl J Med* 1987; 317(6):334-41; <http://dx.doi.org/10.1056/NEJM198708063170602>
- [79] Larsen RA, Leal MA, Chan LS. Fluconazole compared with amphotericin B plus flucytosine for cryptococcal

- meningitis in AIDS. A randomized trial. *Ann Intern Med* 1990; 113(3):183-7; <http://dx.doi.org/10.7326/0003-4819-113-3-183>
- [80] van der Horst CM, Saag MS, Cloud GA, Hamill RJ, Graybill JR, Sobel JD, Johnson PC, Tuazon CU, Kerkering T, Moskovitz BL, et al. Treatment of cryptococcal meningitis associated with the acquired immunodeficiency syndrome. National Institute of Allergy and Infectious Diseases Mycoses Study Group and AIDS Clinical Trials Group. *N Engl J Med* 1997; 337: 15-21; <http://dx.doi.org/10.1056/NEJM199707033370103>
- [81] Brouwer AE, Rajanu Wong A, Chierakul W, Griffin GE, Larsen RA, White NJ, Harrison TS. Combination antifungal therapies for HIV-associated cryptococcal meningitis: a randomised trial. *Lancet* 2004; 363: 1764-7; [http://dx.doi.org/10.1016/S0140-6736\(04\)16301-0](http://dx.doi.org/10.1016/S0140-6736(04)16301-0)
- [82] Utz JP, Garriques IL, Sande MA, Warner JF, Mandell GL, McGehee RF, Duma RJ, Shadomy S. Therapy of cryptococcosis with a combination of flucytosine and amphotericin B. *J Infect Dis* 1975; 132: 368-73; <http://dx.doi.org/10.1093/infdis/132.4.368>
- [83] O'Connor L, Livermore J, Sharp AD, Goodwin J, Gregson L, Howard SJ, Felton TW, Schwartz JA, Neely MN, Harrison TS, et al. Pharmacodynamics of liposomal amphotericin B and flucytosine for cryptococcal meningoencephalitis: safe and effective regimens for immunocompromised patients. *J Infect Dis* 2013; 208:351-61; PMID:23599314; <http://dx.doi.org/10.1093/infdis/jit164>
- [84] Diamond DM, Bauer M, Daniel BE, Leal MA, Johnson D, Williams BK, Thomas AM, Ding JC, Najvar L, Graybill JR, et al. Amphotericin B colloidal dispersion combined with flucytosine with or without fluconazole for treatment of murine cryptococcal meningitis. *Antimicrob Agents Chemother* 1998; 42: 528-33
- [85] Jackson AT, Nussbaum JC, Phulusa J, Namarika D, Chikasema M, Kanyemba C, Jarvis JN, Jaffar S, Hosseini-pour MC, van der Horst C, et al. A phase II randomized controlled trial adding oral flucytosine to high-dose fluconazole, with short-course amphotericin B, for cryptococcal meningitis. *AIDS* 2012; 26: 1363-70; <http://dx.doi.org/10.1097/QAD.0b013e328354b419>
- [86] McCarty TP, Pappas PG. Invasive Candidiasis. *Infect Dis Clin NA* 2016; 30: 103-24; <http://dx.doi.org/10.1016/j.idc.2015.10.013>
- [87] Rex JH, Pappas PG, Karchmer AW, Sobel J, Edwards JE, Hadley S, Brass C, Vazquez JA, Chapman SW, Horowitz HW, et al. A randomized and blinded multicenter trial of high-dose fluconazole plus placebo versus fluconazole plus amphotericin B as therapy for candidemia and its consequences in nonneutropenic subjects. *Clin Infect Dis* 2003; 36(10):1221-8; <http://dx.doi.org/10.1086/374850>
- [88] Abele-Horn M, Kopp A, Sternberg U, Ohly A, Dauber A, Russwurm W, Büchinger W, Nagengast O, Emmerling P. A randomized study comparing fluconazole with amphotericin B/5-flucytosine for the treatment of systemic *Candida* infections in intensive care patients. *Infection* 1996; 24(6):426-32; PMID:9007589; <http://dx.doi.org/10.1007/BF01713042>
- [89] Smego RA, Perfect JR, Durack DT. Combined therapy with amphotericin B and 5-fluorocytosine for *Candida* meningitis. *Rev Infect Dis* 1984; 6(6):791-801; PMID:6522917; <http://dx.doi.org/10.1093/clinids/6.6.791>
- [90] Cadena J, Thompson I, George R, Patterson TF. Invasive Aspergillosis. *Infect Dis Clin NA* 2016; 347: 125-42; <http://dx.doi.org/10.1016/j.idc.2015.10.015>
- [91] Herbrecht R, Denning DW, Patterson TF, Bennett JE, Greene RE, Oestmann JW, Kern WV, Marr KA, Ribaud P, Lortholary O, et al. Voriconazole versus amphotericin B for primary therapy of invasive aspergillosis. *N Engl J Med* 2002; 347:408-15; PMID:12167683; <http://dx.doi.org/10.1056/NEJMoa020191>
- [92] Ananda-Rajah MR, Kontoyiannis D. Isavuconazole: a new extended spectrum triazole for invasive mold diseases. *Future Microbiol* 2015; 10(5):693-708; <http://dx.doi.org/10.2217/fmb.15.34>
- [93] Viscoli C, Herbrecht R, Akan H, Baila L, Sonet A, Gallamini A, Giagounidis A, Marchetti O, Martino R, Meert L, et al. An EORTC Phase II study of caspofungin as first-line therapy of invasive aspergillosis in haematological patients. *J Antimicrob Chemother* 2009; 64(6):1274-81; <http://dx.doi.org/10.1093/jac/dkp355>
- [94] Garbati MA, Alasmari FA, Al-Tannir MA, Tleyjeh IM. The role of combination antifungal therapy in the treatment of invasive aspergillosis: a systematic review. *Int J Infect Dis* 2012; 16(2):e76-e81; <http://dx.doi.org/10.1016/j.ijid.2011.10.004>
- [95] Panackal AA, Parisini E, Proschan M. Salvage combination antifungal therapy for acute invasive aspergillosis may improve outcomes: a systematic review and meta-analysis. *Int J Infect Dis* 2014; 28:80-94; <http://dx.doi.org/10.1016/j.ijid.2014.07.007>
- [96] Raad II, El Zakhem A, El Helou G, Jiang Y, Kontoyiannis DP, Hachem R. Clinical experience of the use of voriconazole, caspofungin or the combination in primary and salvage therapy of invasive aspergillosis in haematological malignancies. *Int J Antimicrob Agents: Elsevier B.V.* 2015; 45: 283-8; <http://dx.doi.org/10.1016/j.ijantimicag.2014.08.012>
- [97] Marr KA, Schlamm HT, Herbrecht R, Rottinghaus ST, Bow EJ, Cornely OA, Heinz WJ, Jagannatha S, Koh LP, Kontoyiannis DP, et al. Combination Antifungal Therapy for Invasive Aspergillosis. *Ann Intern Med* 2015; 162: 81; <http://dx.doi.org/10.7326/M13-2508>
- [98] Cowen LE. The fungal Achilles' heel: targeting Hsp90 to cripple fungal pathogens. *Curr Opin Microbiol* 2013; 16:377-84; PMID:23588026; <http://dx.doi.org/10.1016/j.mib.2013.03.005>
- [99] Veri A, Cowen LE. Progress and prospects for targeting Hsp90 to treat fungal infections. *Parasitology* 2014; 141:1127-37; PMID:24560124; <http://dx.doi.org/10.1017/S0031182013002072>
- [100] Lamoth F, Juvvadi PR, Steinbach WJ. Heat shock protein 90 (Hsp90): A novel antifungal target against *Aspergillus fumigatus*. *Critical Rev Microbiol* 2014; 42: 1-12; <http://dx.doi.org/10.3109/1040841X.2014.947239>
- [101] Cowen LE, Carpenter AE, Matangkasombut O, Fink GR, Lindquist S. Genetic architecture of Hsp90-dependent drug resistance. *Eukaryot Cell* 2006; 5:2184-8; PMID:17056742; <http://dx.doi.org/10.1128/EC.00274-06>
- [102] Cowen LE, Lindquist S. Hsp90 potentiates the rapid evolution of new traits: drug resistance in diverse fungi.

- Science 2005; 309:2185-9; PMID:16195452; <http://dx.doi.org/10.1126/science.1118370>
- [103] Cordeiro RA, Evangelista AJ, Serpa R, Marques FJ, Melo CV, Oliveira JS, Franco JD, Alencar LP, Bandeira TJ, Brilhante RS, et al. Inhibition of heat shock protein 90 enhances the susceptibility to antifungals and reduces the virulence of *Cryptococcus neoformans/C. gattii* species complex. *Microbiology* 2016; 2:309-17; <http://dx.doi.org/10.1099/mic.0.000222>
- [104] Singh SD, Robbins N, Zaas AK, Schell WA, Perfect JR, Cowen LE. Hsp90 governs echinocandin resistance in the pathogenic yeast *Candida albicans* via calcineurin. *PLoS Pathog* 2009; 5:e1000532; PMID:19649312; <http://dx.doi.org/10.1371/journal.ppat.1000532>
- [105] Lamoth F, Juvvadi PR, Fortwendel JR, Steinbach WJ. Heat shock protein 90 is required for conidiation and cell wall integrity in *Aspergillus fumigatus*. *Eukaryot Cell* 2012; 11:1324-32; PMID:22822234; <http://dx.doi.org/10.1128/EC.00032-12>
- [106] Singh-Babak SD, Babak T, Diezmann S, Hill JA, Xie JL, Chen YL, Poutanen SM, Rennie RP, Heitman J, Cowen LE. Global analysis of the evolution and mechanism of echinocandin resistance in *Candida glabrata*. *PLoS Pathog* 2012; 8:e1002718; PMID:22615574; <http://dx.doi.org/10.1371/journal.ppat.1002718>
- [107] Lamoth F, Juvvadi PR, Gehrke C, Asfaw YG, Steinbach WJ. Transcriptional activation of heat shock protein 90 mediated via a proximal promoter region as trigger of caspofungin resistance in *Aspergillus fumigatus*. *J Infect Dis* 2014; 209:473-81; PMID:24096332; <http://dx.doi.org/10.1093/infdis/jit530>
- [108] Cowen LE, Singh SD, Köhler JR, Collins C, Zaas AK, Schell WA, Aziz H, Mylonakis E, Perfect JR, Whitesell L, et al. Harnessing Hsp90 function as a powerful, broadly effective therapeutic strategy for fungal infectious disease. *Proc Natl Acad Sci USA* 2009; 106:2818-23; PMID:19196973; <http://dx.doi.org/10.1073/pnas.0813394106>
- [109] Shapiro RS, Uppuluri P, Zaas AK, Collins C, Senn H, Perfect JR, Heitman J, Cowen LE. Hsp90 orchestrates temperature-dependent *Candida albicans* morphogenesis via Ras1-PKA signaling. *Curr Biol* 2009; 19:621-9; PMID:19327993; <http://dx.doi.org/10.1016/j.cub.2009.03.017>
- [110] Robbins N, Uppuluri P, Nett J, Rajendran R, Ramage G, Lopez-Ribot JL, Andes D, Cowen LE. Hsp90 governs dispersion and drug resistance of fungal biofilms. *PLoS Pathog* 2011; 7:e1002257; PMID:21931556; <http://dx.doi.org/10.1371/journal.ppat.1002257>
- [111] McClellan AJ, Xia Y, Deutschbauer AM, Davis RW, Gerstein M, Frydman J. Diverse cellular functions of the Hsp90 molecular chaperone uncovered using systems approaches. *Cell* 2007; 131:121-35; PMID:17923092; <http://dx.doi.org/10.1016/j.cell.2007.07.036>
- [112] Zhao R, Davey M, Hsu YC, Kaplanek P, Tong A, Parsons AB, Krogan N, Cagney G, Mai D, Greenblatt J, et al. Navigating the Chaperone Network: An Integrative Map of Physical and Genetic Interactions Mediated by the Hsp90 Chaperone. *Cell* 2005; 120:715-27; PMID:15766533; <http://dx.doi.org/10.1016/j.cell.2004.12.024>
- [113] Diezmann S, Michaut M, Shapiro RS, Bader GD, Cowen LE. Mapping the Hsp90 genetic interaction network in *Candida albicans* reveals environmental contingency and rewired circuitry. *PLoS Genetics* 2012; 8:e1002562; PMID:22438817; <http://dx.doi.org/10.1371/journal.pgen.1002562>
- [114] Fox DS, Heitman J. Good fungi gone bad: the corruption of calcineurin. *Bioessays* 2002; 24:894-903; PMID:12325122; <http://dx.doi.org/10.1002/bies.10157>
- [115] Odom A, Muir S, Lim E, Toffaletti DL, Perfect J, Heitman J. Calcineurin is required for virulence of *Cryptococcus neoformans*. *EMBO J* 1997; 16:2576-89; PMID:9184205; <http://dx.doi.org/10.1093/emboj/16.10.2576>
- [116] Cruz MC, Goldstein AL, Blankenship JR, Del Poeta M, Davis D, Cardenas ME, Perfect JR, McCusker JH, Heitman J. Calcineurin is essential for survival during membrane stress in *Candida albicans*. *EMBO J* 2002; 21:546-59; PMID:11847103; <http://dx.doi.org/10.1093/emboj/21.4.546>
- [117] Wiederhold NP, Kontoyiannis DP, Prince RA, Lewis RE. Attenuation of the activity of caspofungin at high concentrations against *Candida albicans*: possible role of cell wall integrity and calcineurin pathways. *Antimicrob Agents Chemother* 2005; 49:5146-8; PMID:16304189; <http://dx.doi.org/10.1128/AAC.49.12.5146-5148.2005>
- [118] Uppuluri P, Nett J, Heitman J, Andes D. Synergistic effect of calcineurin inhibitors and fluconazole against *Candida albicans* biofilms. *Antimicrob Agents Chemother* 2008; 52:1127-32; PMID:18180354; <http://dx.doi.org/10.1128/AAC.01397-07>
- [119] Kontoyiannis DP, Lewis RE, Oshero N, Albert ND, May GS. Combination of caspofungin with inhibitors of the calcineurin pathway attenuates growth *in vitro* in *Aspergillus* species. *J Antimicrob Chemother* 2003; 51:313-6; PMID:12562696; <http://dx.doi.org/10.1093/jac/dkg090>
- [120] Steinbach WJ, Schell WA, Blankenship JR, Onyewu C, Heitman J, Perfect JR. *In vitro* interactions between antifungals and immunosuppressants against *Aspergillus fumigatus*. *Antimicrob Agents Chemother* 2004; 48:1664-9; PMID:15105118; <http://dx.doi.org/10.1128/AAC.48.5.1664-1669.2004>
- [121] Lamoth F, Juvvadi PR, Gehrke C, Steinbach WJ. *In vitro* activity of calcineurin and heat shock protein 90 inhibitors against *Aspergillus fumigatus* azole- and echinocandin-resistant strains. *Antimicrob Agents Chemother* 2013; 57:1035-9; PMID:23165466; <http://dx.doi.org/10.1128/AAC.01857-12>
- [122] Lee SC, Li A, Calo S, Inoue M, Tonthat NK, Bain JM, Louw J, Shinohara ML, Erwig LP, Schumacher MA, et al. Calcineurin orchestrates dimorphic transitions, antifungal drug responses and host-pathogen interactions of the pathogenic mucoralean fungus *Mucor circinelloides*. *Mol Microbiol* 2015; 97:844-65; PMID:26010100; <http://dx.doi.org/10.1111/mmi.13071>
- [123] Lee SC, Li A, Calo S, Heitman J. Calcineurin plays key roles in the dimorphic transition and virulence of the human pathogenic zygomycete *Mucor circinelloides*. *PLoS Pathog* 2013; 9:e1003625; PMID:24039585; <http://dx.doi.org/10.1371/journal.ppat.1003625>
- [124] Smith WL, Edlind TD. Histone deacetylase inhibitors enhance *Candida albicans* sensitivity to azoles and related antifungals: correlation with reduction in *CDR* and *ERG* upregulation. *Antimicrob Agents Chemother*

- 2002; 46:3532-9; PMID:12384361; <http://dx.doi.org/10.1128/AAC.46.11.3532-3539.2002>
- [125] Robbins N, Leach MD, Cowen LE. Lysine Deacetylases Hda1 and Rpd3 Regulate Hsp90 Function thereby Governing Fungal Drug Resistance. *Cell Rep* 2012; 2:878-88; PMID:23041319; <http://dx.doi.org/10.1016/j.celrep.2012.08.035>
- [126] Pfaller MA, Messer SA, Georgopapadakou N, Martell LA, Besterman JM, Diekema DJ. Activity of MGCD290, a Hos2 histone deacetylase inhibitor, in combination with azole antifungals against opportunistic fungal pathogens. *J Clin Microbiol* 2009; 47:3797-804; PMID:19794038; <http://dx.doi.org/10.1128/JCM.00618-09>
- [127] Pfaller MA, Rhomberg PR, Messer SA, Castanheira M. In vitro activity of a Hos2 deacetylase inhibitor, MGCD290, in combination with echinocandins against echinocandin-resistant *Candida* species. *Diagnostic Microbiol Infect Dis* 2015; 81:259-63; PMID:25600842; <http://dx.doi.org/10.1016/j.diagmicrobio.2014.11.008>
- [128] Besterman J, Nguyen DT, Ste-Croix H. MGCD290, an Oral Fungal Hos2 Inhibitor, Enhances the Antifungal Properties of Fluconazole Following Multiple—or Single—Dose Oral Administration in Pre—and Post—Infection Settings. Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC). San Francisco, CA: MethylGene, Inc. 2012
- [129] Sellam A, Askew C, Epp E, Lavoie H, Whiteway M, Nantel A. Genome-wide mapping of the coactivator Ada2p yields insight into the functional roles of SAGA/ADA complex in *Candida albicans*. *Mol Biol Cell* 2009; 20:2389-400; PMID:19279142; <http://dx.doi.org/10.1091/mbc.E08-11-1093>
- [130] Wurtele H, Tsao S, Lepine G, Mullick A, Tremblay J, Drogaris P, Lee EH, Thibault P, Verreault A, Raymond M. Modulation of histone H3 lysine 56 acetylation as an antifungal therapeutic strategy. *Nat Med* 2010; 16:774-80; PMID:20601951; <http://dx.doi.org/10.1038/nm.2175>
- [131] Lopes da Rosa J, Boyartchuk VL, Zhu LJ, Kaufman PD. Histone acetyltransferase Rtt109 is required for *Candida albicans* pathogenesis. *Proc Natl Acad Sci USA* 2010; 107:1594-9; PMID:20080646; <http://dx.doi.org/10.1073/pnas.0912427107>
- [132] Chen X, Ren B, Chen M, Liu MX, Ren W, Wang QX, Zhang LX, Yan GY. ASDCD: antifungal synergistic drug combination database. *PloS One* 2014; 9:e86499; PMID:24475134; <http://dx.doi.org/10.1371/journal.pone.0086499>
- [133] Epp E, Vanier G, Harcus D, Lee AY, Jansen G, Hallett M, Sheppard DC, Thomas DY, Munro CA, Mullick A, et al. Reverse genetics in *Candida albicans* predicts ARF cycling is essential for drug resistance and virulence. *PLoS Pathog* 2010; 6:e1000753; PMID:20140196; <http://dx.doi.org/10.1371/journal.ppat.1000753>
- [134] LaFayette SL, Collins C, Zaas AK, Schell WA, Betancourt-Quiroz M, Gunatilaka AA, Perfect JR, Cowen LE. PKC signaling regulates drug resistance of the fungal pathogen *Candida albicans* via circuitry comprised of Mkc1, calcineurin, and Hsp90. *PLoS Pathog* 2010; 6:e1001069; PMID:20865172; <http://dx.doi.org/10.1371/journal.ppat.1001069>
- [135] Mor V, Rella A, Farnoud AM, Singh A, Munshi M, Bryan A, Naseem S, Konopka JB, Ojima I, Bullesbach E, et al. Identification of a New Class of Antifungals Targeting the Synthesis of Fungal Sphingolipids. *mBio* 2015; 6:e00647; PMID:26106079; <http://dx.doi.org/10.1128/mBio.00647-15>
- [136] Zhai B, Wu C, Wang L, Sachs MS, Lin X. The antidepressant sertraline provides a promising therapeutic option for neurotropic cryptococcal infections. *Antimicrob Agents Chemother* 2012; 56:3758-66; PMID:22508310; <http://dx.doi.org/10.1128/AAC.00212-12>
- [137] Butts A, DiDone L, Koselny K, Baxter BK, Chabrier-Rosello Y, Wellington M, Krysan DJ. A repurposing approach identifies off-patent drugs with fungicidal cryptococcal activity, a common structural chemotype, and pharmacological properties relevant to the treatment of cryptococcosis. *Eukaryot Cell* 2013; 12:278-87; PMID:23243064; <http://dx.doi.org/10.1128/EC.00314-12>
- [138] Spitzer M, Griffiths E, Blakely KM, Wildenhain J, Ejim L, Rossi L, De Pascale G, Curak J, Brown E, Tyers M, et al. Cross-species discovery of syncretic drug combinations that potentiate the antifungal fluconazole. *Mol Syst Biol* 2011; 7:499; PMID:21694716; <http://dx.doi.org/10.1038/msb.2011.31>
- [139] Robbins N, Spitzer M, Yu T, Cerone RP, Averette AK, Bahn YS, Heitman J, Sheppard DC, Tyers M, Wright GD. An antifungal combination matrix identifies a rich pool of adjuvant molecules that enhance drug activity against diverse fungal pathogens. *Cell Rep* 2015; 13:1481-92; PMID:26549450; <http://dx.doi.org/10.1016/j.celrep.2015.10.018>
- [140] Borisy AA, Elliott PJ, Hurst NW, Lee MS, Lehar J, Price ER, Serbedzija G, Zimmermann GR, Foley MA, Stockwell BR, et al. Systematic discovery of multicomponent therapeutics. *Proc Natl Acad Sci USA* 2003; 100:7977-82; PMID:12799470; <http://dx.doi.org/10.1073/pnas.1337088100>
- [141] Lehar J, Zimmermann GR, Krueger AS, Molnar RA, Ledell JT, Heilbut AM, Short GF, 3rd, Giusti LC, Nolan GP, Magid OA, et al. Chemical combination effects predict connectivity in biological systems. *Mol Syst Biol* 2007; 3:80; PMID:17332758; <http://dx.doi.org/10.1038/msb4100116>
- [142] Jansen G, Lee AY, Epp E, Fredette A, Surprenant J, Harcus D, Scott M, Tan E, Nishimura T, Whiteway M, et al. Chemogenomic profiling predicts antifungal synergies. *Mol Syst Biol* 2009; 5:338; PMID:20029371; <http://dx.doi.org/10.1038/msb.2009.95>
- [143] Youngsaye W, Dockendorff C, Vincent B, Hartland CL, Bittker JA, Dandapani S, Palmer M, Whitesell L, Lindquist S, Schreiber SL, et al. Overcoming fluconazole resistance in *Candida albicans* clinical isolates with tetracyclic indoles. *Bioorg Med Chem Lett* 2012; 22:3362-5; PMID:22497765; <http://dx.doi.org/10.1016/j.bmcl.2012.02.035>
- [144] Kaneko Y, Fukazawa H, Ohno H, Miyazaki Y. Combinatory effect of fluconazole and FDA-approved drugs against *Candida albicans*. *J Infect Chemother* 2013; 19:1141-5; PMID:23807392; <http://dx.doi.org/10.1007/s10156-013-0639-0>
- [145] Zhang L, Yan K, Zhang Y, Huang R, Bian J, Zheng C, Sun H, Chen Z, Sun N, An R, et al. High-throughput synergy screening identifies microbial metabolites as combination

- agents for the treatment of fungal infections. *Proc Natl Acad Sci USA* 2007; 104:4606-11; PMID:17360571; <http://dx.doi.org/10.1073/pnas.0609370104>
- [146] Hartland K, Pu J, Palmer M, Dandapani S, Moquist PN, Munoz B, DiDone L, Schreiber SL, Krysan DJ. High-throughput screen identifies a novel molecular scaffold that inhibits cell wall integrity pathway signaling. *ACS Infect Dis* 2016; 2:93-102; PMID:26807437; <http://dx.doi.org/10.1021/acsinfecdis.5b00111>
- [147] Nishikawa JL, Boeszoermyenyi A, Vale-Silva LA, Torelli R, Posteraro B, Sohn YJ, Ji F, Gelev V, Sanglard D, Sanguinetti M, et al. Inhibiting fungal multidrug resistance by disrupting an activator-Mediator interaction. *Nature* 2016; 530:485-9; PMID:26886795; <http://dx.doi.org/10.1038/nature16963>
- [148] Denning DW, Bromley MJ. Infectious disease. how to bolster the antifungal pipeline. *Science* 2015; 347:1414-6; PMID:25814567; <http://dx.doi.org/10.1126/science.aaa6097>
- [149] Jarvis JN, Bicanic T, Loyse A, Namarika D, Jackson A, Nussbaum JC, Longley N, Muzoora C, Phulusa J, Taseera K, et al. Determinants of mortality in a combined cohort of 501 patients with HIV-associated cryptococcal meningitis: implications for improving outcomes. *Clin Infect Dis* 2014; 58: 736-45; <http://dx.doi.org/10.1093/cid/cit794>
- [150] León C, Ostrosky-Zeichner L, Schuster M. What's new in the clinical and diagnostic management of invasive candidiasis in critically ill patients. *Intensive Care Med* 2014; 40(6):808-19; <http://dx.doi.org/10.1007/s00134-014-3281-0>