

RESEARCH ARTICLE

# Toxicity of spinosad to temephos-resistant *Aedes aegypti* populations in Brazil

Luciana dos Santos Dias<sup>1,2</sup>, Maria de Lourdes da Graça Macoris<sup>3</sup>, Maria Teresa Macoris Andrighetti<sup>3</sup>, Vanessa Camargo Garbeloto Otrera<sup>3</sup>, Adriana dos Santos Dias<sup>1,2</sup>, Luiz Guilherme Soares da Rocha Bauzer<sup>1,2</sup>, Cynara de Melo Rodvalho<sup>1,2</sup>, Ademir Jesus Martins<sup>1,2</sup>, José Bento Pereira Lima<sup>1,2\*</sup>

**1** Laboratório de Fisiologia e Controle de Artrópodes Vetores, Instituto Oswaldo Cruz, Fiocruz, Rio de Janeiro, RJ—Brasil, **2** Laboratório de Entomologia, Instituto de Biologia do Exército, Rio de Janeiro, RJ—Brasil, **3** Superintendência de Controle de Endemias, Marília, SP—Brasil

\* [jbento@ioc.fiocruz.br](mailto:jbento@ioc.fiocruz.br)



## Abstract

The mosquito *Aedes aegypti* is the primary vector of different arboviruses and represents a major public health problem. Several Brazilian populations of *Ae. aegypti* have developed resistance to temephos, the most used organophosphate larvicide. New tools which are less harmful to the environment and safer for humans are becoming increasingly important to control this insect vector. Spinosad, an aerobic fermentation product of a soil actinobacteria, has a favorable environmental profile. It presents selective insecticide properties, a mechanism of action that differs from those of many synthetic chemical insecticides. The toxicity of spinosad and temephos to *Aedes aegypti* populations from Brazil, which were previously exposed to temephos, were investigated in this study. Larval susceptibility (LC50) to temephos varied from 3µg/L for Rockefeller up to 260 µg/L for Santana do Ipanema field derived population. Larval susceptibility (LC50) to spinosad varied from 23µg/L for Rockefeller up to 93µg/L for Marília field derived population. In addition, a semi-field trial was performed to evaluate spinosad (Natular<sup>TM</sup> DT) initial efficacy and persistence toward four field-derived lineages and the Rockefeller lineage, used as an internal control. Spinosad was tested at 0.5mg active ingredient/L in 200L capacity water tanks. Mortality was recorded each 24 hours after exposition and tanks were further recolonized once per week with mortality being recorded daily for eight weeks. Spinosad provided a level equal or superior to 80% mortality during a seven to eight week evaluation period. The assessed populations did not present cross-resistance between spinosad and temephos in laboratory conditions. It demonstrates that spinosad may be a promising larvicide for the control of *Ae. aegypti*, especially for populations in which resistance to temephos has been detected.

## OPEN ACCESS

**Citation:** dos Santos Dias L, Macoris MdLdG, Andrighetti MTM, Otrera VCG, Dias AdS, Bauzer LGSdR, et al. (2017) Toxicity of spinosad to temephos-resistant *Aedes aegypti* populations in Brazil. PLoS ONE 12(3): e0173689. <https://doi.org/10.1371/journal.pone.0173689>

**Editor:** Pedro L. Oliveira, Universidade Federal do Rio de Janeiro, BRAZIL

**Received:** August 7, 2016

**Accepted:** February 26, 2017

**Published:** March 16, 2017

**Copyright:** © 2017 dos Santos Dias et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** This work was supported by Fundação Oswaldo Cruz (<http://portal.fiocruz.br/pt-br>) and Fundação coordenação de aperfeiçoamento de pessoal de nível superior (<http://www.capes.gov.br/>).

**Competing interests:** The authors have declared that no competing interests exist.

## Introduction

The *Aedes aegypti* (Linnaeus, 1762) mosquito, widely distributed in the tropical and subtropical regions of the planet, is highly adapted to the urban environment and is often found within and around households [1–3]. It is the main vector of dengue viruses and can play a role as vector of yellow fever virus in the urban cycle [4], which are considered two of the most

important viral diseases transmitted by arthropods [5]. Additionally, this insect is a potential vector transmitter of Chikungunya virus and Zika virus [6–8]. Efficient vaccines against dengue, chikungunya and zika are not currently available. Because of that, public health measures focus mainly on the vector control [9,10].

Control programs aim to reduce the populations of insects that transmit ethiological agents by eliminating potential breeding sites or by using insecticides [11–13]. In Brazil, the control of dengue vector larvae has been carried out for decades with the use of temephos, an organophosphate larvicide. Resistance to this compound has been reported since 1998 [14–16]. Nowadays the majority of the evaluated Brazilian populations is resistant to temephos [14,17–20].

Insecticide resistance is the heritable ability of the insect to survive to insecticide concentrations that are lethal to individuals of the same species (susceptible ones) [21]. The major mechanisms recorded for *Ae. aegypti* resistance to temephos has been associated with changes at its target site in the acetylcholinesterase. It has also been associated with metabolic mechanisms related with enzymes involved in the detoxification of xenobiotic compounds, named metabolic resistance [22–24]. The insecticide metabolic resistance can involve three major enzyme groups: esterases (ESTs), glutathione-S-transferases (GSTs) and mixed function oxidases (MFOs) [25]. Detoxifying enzymes can confer resistance to many chemicals belonging to the same group or to different groups, characterizing the cross resistance [26].

In this sense, alternatives to organophosphates for the control of *Ae. aegypti* and other insects of medical importance are essential [27,28]. Also, there are reports of *Ae. aegypti* resistance to the main insecticides in use [18]. Alternatives to organophosphates, such as, the insect growth regulators (IGRs), *Bacillus thuringiensis* serovar *israelensis* (*Bti*) and spinosad, which are environmental safe and innocuous for humans, have attracted growing interest. Spinosad is composed of two natural metabolites, spinosyns A and D [29], which are products of the aerobic fermentation of *Sacharopolyspora spinosa*, a soil actinobacteria [30,31]. Spinosad acts on both the nicotinic acetylcholine receptors and the  $\gamma$ -aminobutyric acid (GABA) receptors, causing excitation of the insect's nervous system, paralysis and death [32–34]. Therefore, although also neurotoxic, spinosad has a distinct molecular target when compared to organophosphates and pyrethroids, which target the enzyme acetylcholinesterase and the voltage gated sodium channel, respectively [23].

Tests of chronical toxicity demonstrate that spinosad does not present carcinogenic, teratogenic or neurotoxic effects in mammals [35], and has a favorable profile with low environmental persistence. It displays low toxicity for non-targeted insects and it is innocuous for fish and other aquatic vertebrates, when used in the recommended dosages [36–38]. Marcombe et al. [39] noticed that spinosad is a promising larvicide against *Ae. aegypti* on water reservoirs, where its residual efficacy lasted for up to 16 weeks. Moreover spinosad also showed high toxicity against other mosquitoes species as *Anopheles stephensi*, *Anopheles albimanus* and *Culex quinquefasciatus* [40–43]. Recently the World Health Organization Pesticide Scheme (WHOPES) approved spinosad for usage in drinking water [43]. However, possible cross-resistance selected by other compounds needs to be investigated before its wide utilization.

There is a high level of resistance to organophosphates and pyrethroids in worldwide *Ae. aegypti* populations. Moreover, spinosad and temephos have action on different target sites, which supports the lack of cross-resistance, considering this mechanism. However, we can not discard the hypothesis that cross-resistance could be found due to metabolic detoxification mechanism selected by temephos. Because of that it is important to evaluate the efficacy of spinosad over populations with distinct profiles of resistance to temephos. In this study, we assessed and compared the toxicity of spinosad and temephos insecticide in Brazilian *Ae. aegypti* populations with putative different genetic backgrounds, under laboratory and field (simulation) conditions.

## Materials and methods

1) Mosquitoes—Seven *Ae. aegypti* populations were collected in six different Brazilian states, representing four geopolitical regions of the country: Santarém and Marabá (State of Pará, Northern Brazil), Goiânia (State of Goiás, Midwestern Brazil), Cachoeiro de Itapemirim and Marília (State of Espírito Santo and State of São Paulo, respectively; Southeastern Brazil), Salvador and Santana do Ipanema (State of Bahia and State of Alagoas, respectively; Northeastern Brazil) (Fig 1). The eggs were collected in the field with the help of ovitraps containing hay infusion [44], which was distributed in a way to represent the whole geographical extension of the localities mentioned above [15]. These materials were collected by the municipalities' health secretaries and sent to the Laboratório de Fisiologia e Controle de Artrópodes Vetores (Fiocruz, Rio de Janeiro) by the Brazilian Health Ministry. The females were reared as follows: eggs collected in the field were hatched in the laboratory in order to obtain 1<sup>st</sup> instar larvae that developed to the adults of the parental generation of these strains. Approximately 1000 larvae were reared in plastic bowls containing 1L of dechlorinated water plus one gram of cat food (Friskies, Purina). The food was provided every three days until the larvae reached the pupae phase. The pupae were separated in plastic cups of 50ml and put inside cardboard cages until adults emergence. Adults were kept in a controlled environment with temperature of  $26^{\circ}\text{C} \pm 1$  and 70% humidity, and fed *ad libitum* with a solution of 10% sucrose. Guinea pigs *Cavia porcellus* (Linnaeus, 1758; Rodentia, Caviidae) were used for blood feeding, with approval of the Ethics Committee for Animal Use from Fundação Oswaldo Cruz (CEUA—Fiocruz) n° L -011/09.

2) Insecticides—Temephos (PESTANAL<sup>®</sup> - 95,6% temephos a.i., Sigma–Aldrich) and liquid spinosad (Natular<sup>™</sup> 20EC– 20.6% Spinosad a.i., Clarke Mosquito Control Products, Inc.) were used for laboratory assays. The insecticides were diluted so that a final concentration of 3000mg/L was achieved for both. After normalizing the concentration, we prepared the solutions for the bioassays. For the field simulations, one effervescent tablet of spinosad (Natular<sup>™</sup> DT– 7.48% Spinosad a.i., Clarke Mosquito Control Products, Inc.) was used per 200 L tank,



**Fig 1. Brazilian political map showing the states where collections were performed.**

<https://doi.org/10.1371/journal.pone.0173689.g001>

which provided a final concentration of 0.5 mg active ingredient/L, as recommended by WHO [45].

3) Laboratory tests—Assays were performed in the Laboratório de Fisiologia e Controle de Artrópodes Vetores, Instituto Oswaldo Cruz—Fiocruz, located at the Instituto de Biologia do Exército (IBEx), Rio de Janeiro, Brazil. The F1 and F2 generations descending from the seven field populations were used. The Rockefeller lineage, which is a reference of susceptibility to insecticides [46], was used as control for each experiment. This lineage was reared in the same conditions as described for the field colonies. The larvae used in the laboratory bioassays were reared as follow: 1,000 larvae from each population were kept in plastic bowls containing 1L of water and one gram of cat food (Friskies, Purina) until they reached L3 instar for bioassay performances.

Bioassay with temephos was conducted based on the World Health Organization (WHO) protocol [47]. For each dosage, 80 L3 larvae (four replicas containing 20 larvae each in white plastic cups containing 100ml of solution) were exposed to a minimum of 10 to 12 different doses of insecticide (between 1.05 and 720 $\mu$ g/L). The assays were conducted four times, on different days, for each of the seven populations and the Rockefeller lineage. The temephos solution was prepared with ethanol and dechlorinated water. For the control group, only ethanol (0.6%) was used. Mortality reading was performed after 24h of exposure.

Before the spinosad dose-response bioassays, we first evaluated the time of exposure to the product (24h, 48h or 72h). Third instar Rockefeller larvae were exposed to 10 spinosad concentrations, ranging from 10 to 70 $\mu$ g/L. For each concentration, four replicas were tested using 10 L3 larvae each, in transparent plastic cups containing 150mL of solution. The spinosad solution was prepared with ethanol and dechlorinated water. For the control group, only ethanol (0.4%) was used. Mortality was recorded at 24, 48 and 72 hours of exposure. The same experimental set was analyzed at different periods. 14mg of triturated cat meal (Friskies; Purina) per plastic cup was provided in the first day of the bioassays. The assays were conducted three times, on different days. Dose-response bioassays with natural populations were performed with 10 to 12 spinosad concentrations (between 10 and 300 $\mu$ g/L), as described above, with 24 hours of exposure.

4) Semi-field simulation assays—The experiments were conducted at the external area of SUCEN—Superintendência de Controle de Endemias—Laboratório de Entomologia Aplicada in Marília city, São Paulo State. The city climate is rainy tropical with a dry winter. The temperature ranges from 17 to 29°C [48].

Once the laboratory experiments were concluded, four natural populations were still available for carrying out field tests simulations. Those samples were sent to SUCEN, in Marília city, by the Laboratório de Fisiologia e Controle de Artrópodes Vetores (Fiocruz, Rio de Janeiro). The populations were divided in two groups, the first comprised Marília (F1 generation) and Salvador (F1 generation) populations which were tested from July 2011 to February 2012. The second group comprised Goiânia (F2 generation) and Santarém (F1 generation) populations, which were tested from May to November 2012. In both cases, Rockefeller reference population was used as a control.

Tests were performed outside the laboratory in a covered area to protect the tanks from environmental factors such as rain. Eighteen polyethylene water tanks filled with 200L of tap water were used to conduct the tests. In order to simulate residential usage situation, a third of the water volume of the tank was replaced three times a week.

For each test, 12 tanks were treated (E) and six remained untreated (C) as control. In order to avoid bias in the results related to tank position, larvae from a single population were tested in all tanks, placed in alternated positions, passing for each tank at least once. The day of the treatment was considered the "day zero", and 30 3rd instar larvae were used to colonize each

tank. The larvae were previously reared in plastic bowls containing 1L of water and one gram of cat food (Friskies, Purina). The larvae were placed into each water tanks. Soon after, one tablet of the Natular DT was added at the center of the 'E' tanks, as recommended by the manufacturer. Every week, surviving larvae were removed and a new batch of 30 3rd instar larvae was added to each tank. Food (Friskies; Purina) was provided for larvae in the first day and each three days of the experiment, if necessary.

The mortality was recorded 24 h after the exposure and then daily, at the same time. The water temperature and water pH of the tanks were monitored daily with the assistance of a thermo-hygrometer (Fisher Scientific<sup>®</sup>) and a Ph Meter (Orion Research Inc., Cambridge, MA).

The larvicide effect was evaluated for an eight-weeks period, which equals to the residences visiting cycle intervals of the National Dengue Control Program. Each test was repeated three times on different periods, thus completing one evaluation. The two first tests of one evaluation were conducted with covered water tanks, while in the third test the tanks were covered with nylon mesh to prevent entry of insects and eventual detritus. By using different covers, we wanted to evaluate eventual environmental influences over the residual effect of the product, such as the influence of indirect solar radiation in the tanks.

5) Statistical analysis—To estimate the lethal concentrations (LCs) for each type of treatment and their respective confidence interval, probit analysis was conducted using the Polo-PC software [49]. The resistance ratios (RR) were obtained through the division of LCs 50 and 95 of the field populations by the respective LCs of the Rockefeller lineage. The comparison of lethal concentration (LC<sub>95</sub>) between the different times of exposure to spinosad was conducted through the overlap of its 95% confidence intervals. A linear regression analysis was performed to check if there was a correlation between the RR<sub>95</sub> of the analysed populations to temephos and spinosad. The above mentioned analysis were all performed with the help of the Graph-Pad Prism software version 5.0 for Windows (GraphPad Software, San Diego, California, USA). A test t followed by Mann-Whitney test was performed to compare the effectiveness of spinosad in water tanks covered with a lid or with nylon mesh, with the help of the Graph-Pad Prism software version 5.0 for Windows (GraphPad Software, San Diego, California, USA).

## Ethics statements

No specific permissions were required as the mosquito collections were performed by the municipalities' health secretaries as requested by the Brazilian Health Ministry. The species used in this study (*Aedes aegypti*) is not an endangered or protected species. Mosquito blood-feeding on anesthetized guinea pigs (*Cavia porcellus*) was authorized by Fiocruz Ethical Committee (CEUA 011/09).

## Results

### Larval susceptibility

The comparisons of the LC<sub>95</sub> averages were performed by superimposing the 95% confidence intervals of lethal concentrations at each evaluated exposure time (S1 Table). An overlap occurred between 24, 48 and 72h exposure indicating no significant difference. Therefore, it was determined that the mortality reading could be done after 24 hours exposure of *Ae. aegypti* to spinosad.

The comparative analysis of the spinosad and temephos LCs toward the reference Rockefeller colony has shown that the latter product required a smaller dose to kill 50% of the larvae (Table 1). The lethal concentration (LC<sub>50</sub>) of the larvae bioassays for spinosad ranged from 23µg/L for the susceptible Rockefeller reference lineage up to 82µg/L for the field populations.

**Table 1. Toxicity of temephos and spinosad against 3rd instar *Aedes aegypti* larvae after a 24h exposure.**

Temephos								
Population	Number of larvae	Slope	LC50 (er of	CI 95%	RR50	LC95 (5% of	CI 95%	RR95
Rockefeller	2400	6.2	3	2.80–2.98	1	4.7	4.54–4.78	1
Santarém / PA	2640	4.3	19	18.24–19.37	6.5	45	42.93–48.49	8.5
Marabá / PA	2640	3.8	29	25.62–31.39	9.9	78	66.28–91.31	14.6
Goiânia / GO	2640	4.5	28	25.83–30.90	9.8	65	55.5–75.73	12.2
Cachoeiro de Itapemirim / ES	2640	5.6	65	61.86–69.33	22.7	129	118.68–140.97	24.3
Marília / SP	2400	4.5	6.9	6.60–7.30	2.3	17	15.00–18.00	3.6
Salvador / BA	2640	4.5	33	25.00–58.00	11.0	78.5	71.00–92.00	16.7
Santana do Ipanema / AL	2880	4.2	260	251.25–267.74	89.8	635	602.09–673.82	119.1
Spinosad								
Population	Number of larvae	Slope	LC50 (er of	CI 95%	RR50	LC95 (5% of	CI 95%	RR95
Rockefeller	1200	4.3	23	21.63–23.80	1	55	50.96–60.01	1
Santarém / PA	1440	3.9	57	53.89–60.66	2.5	152	139.62–168.65	2.8
Marabá / PA	1320	3.5	66	61.71–69.92	2.9	193	174.06–219.96	3.5
Goiânia / GO	1440	5.3	89	83.61–94.46	3.9	299	267.67–340.33	5.4
Cachoeiro Itapemirim / ES	1440	5.2	80	76.72–83.20	3.5	198	182.94–218.66	3.6
Marília / SP	1200	5.1	93	89.02–96.82	4.1	214	196.98–235.63	3.9
Salvador / BA	1440	4.2	80	76.15–83.96	3.5	195	181.42–213.27	3.6
Santana do Ipanema / AL	1440	4.6	82	78.63–85.17	3.6	187	173.58–203.52	3.4


RR = Resistance ratio LC = Lethal concentration, CI 95% = confidence interval 95% Resistance Ratio (RR) RR<5 = low resistance, RR between 5 and 10 = intermediate resistance, RR>10 = high resistance.

<https://doi.org/10.1371/journal.pone.0173689.t001>

For temephos the lethal concentration of the bioassays with larvae ranged from 3µg/L for Rockefeller up to 260µg/L for field populations. Although the LC50 for temephos was lower for the Rockefeller lineage, there was great variation in the LC50 for the field populations exposed to the product. For spinosad the LC50 in the Rockefeller lineage was higher but the variation in LC50 among the field populations was narrower. The field populations used in this study were tested to verify the susceptibility/resistance status to the larvicide organophosphate temephos and eventual cross-resistance to the spinosad. The amplitude of the dose-response effect to both insecticides can be observed in Fig 2. For organophosphates a classification of populations resistance can be build using the criteria proposed by Mazzarri and Georgiou [50]. In this criteria a RR (resistance ratio) lower than 5 indicates low resistance, RR between 5 and 10 indicates intermediate resistance and over 10 indicates a high resistance. The levels of spinosad toxicity were similar for all evaluated populations. RR<sub>50</sub> ranged from 2.5 (Santarem) to 4.1 (Marilia) and RR<sub>95</sub> ranged from 2.8 (Santarém) to 5.4 (Goiânia). Different levels of resistance were observed to temephos, with RR<sub>50</sub> ranging from 2.3 (Marília) to 89.8 (Santana do Ipanema) and RR<sub>95</sub> ranging from 3.6 (Marília) to 119.1 (Santana do Ipanema), as seen in Table 1. No significant correlation between the RR<sub>95</sub> for temephos and spinosad was observed (R<sup>2</sup> = 0.0368; p = 0.6803).

### Evaluation of spinosad persistence under semi-field conditions

The data related to the residual effect of spinosad (Natular DT) in simulated field are shown in Fig 3. Spinosad treatment showed persistence during a seven to eight-week period providing at least 80% of larvae mortality for two populations, Marilia (Fig 3A) and Goiânia (Fig 3B). For the remaining populations there was no decrease up to the eighth week. There was a small loss




**Fig 2. Mortality of *Ae. aegypti* larvae exposed for 24h to temephos (A) and spinosad (B) insecticides.**

<https://doi.org/10.1371/journal.pone.0173689.g002>

of the control population in one of the triplicates (S1C Fig) probably due to the high temperatures that occur in Marilia city in February.

The tanks were covered either with a lid (1<sup>st</sup> and 2<sup>nd</sup> replicates) or with a nylon mesh (3<sup>rd</sup> replicate). Regardless how the water tanks were covered, the mortality rate was not significantly different for each population, as revealed by the Mann Whitney tests ( $P > 0.05$  for all comparisons). The average water temperature was  $23.5^{\circ}\text{C} \pm 3.1$  and pH of the test was  $7.4 \pm 0.6$  (Fig 3C and 3D). The details of the tests for each population are shown in S1 Fig and S2 Fig.



**Fig 3. Field evaluation of Spinosad (Natular™ DT) against *Aedes aegypti* 3rd instar larvae.** The average mortality percentage is represented for three independent bioassays conducted in a row, A- Rockefeller, Marília and Salvador (tested between July 2011 and February 2012). B- Rockefeller, Goiânia and Santarém (tested between September 2011 and November 2012). C- pH and temperature of the solution in the tanks during the three assays for the Rockefeller, Marília and Salvador group. D- pH and temperature of the solution in the tanks during the three assays for the Rockefeller, Goiânia and Santarém group. The dose tested was 0.5mg/L. Every week the tanks were recolonized by removing the dead and the remaining living larvae and introducing 30 new 3rd instar larvae.

<https://doi.org/10.1371/journal.pone.0173689.g003>

The mortality effect of spinosad tended to be constant over the weeks, with reduction in mortality starting in the seventh week for the majority of populations. When the experiments were analyzed individually, small variations were observed. For example, there was a decline in mortality percentage (<80%) in the population of Salvador after the sixth week, increasing to rates above 80% in the following week (S1A Fig). In the third test (S2C Fig), Goiânia mortality rate was 72% in the seventh week.

## Discussion

The *Ae. aegypti* resistance to chemical products has been reported around the globe [50–56]. In Brazil, the detection of populations resistant to the temephos larvicide started in 1998 [14–16]. Because of this, additional to elimination or correct treatment of larvae breeding sites, there was a need to search for alternative compounds for *Ae. aegypti* control. Spinosad, with a low toxicity for mammals and the environment, and being neurotoxic for *Ae. aegypti*, seems to be a good option. However, there is a possibility it can be detoxified by cross-resistance.



Despite of this, it may yet be an important tool to combat vectors already resistant to other products.

According to WHO, studies to estimate the lethal concentration for *Ae. aegypti* larvae in the laboratory must be conducted using reference lineages. In addition, the larvae must remain exposed for over 24 hours to the larvicide [57]. Many authors have conducted studies of spinosad effect on *Ae. aegypti* larvae from laboratory colonies, however, the protocol and the time of exposure varied in each work [29,41,58–60]. In this study, procedures for experiment standardization using spinosad dose response curve were conducted with the Rockefeller lineage (L3 larvae) and mortality was scored after 24, 48 and 72 hours of exposure to the insecticide. This was performed as a preliminary step to confirm the correct period for mortality recording. Comparing the LC95 for each period, we noticed that there was an overlap in their interval of confidence indicating that the differences found were not significant, therefore leading us to keep 24 hours of exposure as ideal for dose-dependent bioassays with spinosad.

Studies performed with spinosad showed great variation on the lethal concentrations obtained using laboratory populations of *Ae. aegypti*: Darriet et al [29], Romi et al [58] and Paul et al [59] found a LC50 of 35µg/L, 10µg/L and 160µg/L respectively. However, Bond et al [41] and Pérez et al [60] used the same protocol and obtained similar LC50, 25µg/L and 26µg/L, respectively. These results are similar to the results we obtained in dose-response curve of spinosad over the susceptible Rockefeller lineage (LC50 = 23µg/L). It indicates that spinosad displayed high toxicity to *Ae. aegypti* reference colony tested and its lethal concentration is very low. However, the lethal concentration for the reference colony was not as low as that found for temephos (LC50 = 3.0µg/L). Here it is worth remembering that the temephos and spinosad used to determine toxicity in the bioassays have differences in their active ingredient contents, which are 95.6 and 20.6%, respectively. The discrepancy found by other authors may be attributed to differences in protocols used in each laboratory, which highlights the need to maintain a standard protocol that facilitates the comparison of results from different groups.

The field populations assessed in this study were characterized as resistant to the temephos organophosphate based on the criteria proposed by Mazzarri and Georghiou [50]. However, the RRs for spinosad in all populations were extremely low when compared to the RRs obtained for temephos. Only the population of Goiânia presented  $RR > 5$  ( $RR$  at  $LC_{95} = 5.4$ ) for spinosad. Since none of these populations have been treated with spinosad before, all the RRs observed seem to be variations of the populations susceptibility rather than resistance to spinosad. Interestingly, Goiânia population was not the one with the highest  $RR_{95}$  to the organophosphate temephos. The population of Santana do Ipanema showed the highest  $RR_{95}$  for temephos and a very low  $RR_{95}$  for spinosad. As expected, no significant correlation was noted between the  $RR_{95}$  of the populations to temephos and spinosad ( $R^2 = 0.0368$ ;  $p = 0.6803$ ), suggesting that the resistance mechanisms selected to temephos are not influencing the toxicity levels of spinosad. Some studies suggested that once spinosad has different targets in the insect's nervous system, there is a lower possibility of cross-resistance with other products used in the control of disease vectors [31,33,34], corroborating our results.

During the assessment of spinosad residual effect on semi-field simulation, the average temperature recorded in the recipients was  $23.5 \pm 3.1^\circ\text{C}$ , and the average of the pH was  $7.4 \pm 0.6$ . Other semi-field simulation study has shown that *Bti* larvicides efficiency had an inverse relationship between temperature (which ranged from 18.7 to 34.6°C during the simulation) and the persistence of the *Bti* on *Ae. aegypti* larvae [61]. This may be a problem, since the dengue vector proliferates in tropical and subtropical areas [62]. Fontoura et al [16], verified that the climatic variation, even in the period when elevated temperatures were not recorded on a field simulation, influenced negatively on the persistence of IGR Novaluron, with an increase in emergency of adults a week earlier if compared to simulations conducted in the periods when

the temperature remained stable. In our results, it was not possible to verify this influence, probably due to the low temperature variation. In addition, we also did not observe the interference of changes in the water pH, maybe because the variation during the assessments was very small. These alterations were not relevant, even in the recipients of the third assessment, which instead of a solid cover had a nylon mesh attached with a rubber band to seal the water tank. As observed in the mortality graphs, our results suggest that a higher exposure of the tank's content to the environmental variations, and a faster evaporation than that observed on the sealed recipients did not influence, in a significant manner, the residual effect of the product. A possible influence of the indirect insolation could potentially be detected since spinosad is vulnerable to photolysis [63]. Pérez et al. [60] verified that the half life of aqueous solutions of spinosad on *Ae. aegypti* larvae in plastic containers containing 1L of solution (10mg/L) differed according to the exposition of direct sun light or not. When exposed to sunlight conditions the half-life was 2.1 days. When allocated in the shadow the half life was 24.5 days. However, in our studies, since no statistic difference was observed between the two forms of treatment, the indirect insolation, if it occurred, did not have any critical effect.

A similar experiment conducted by Thavara et al. [64], showed that Spinosad DT, at 0.5mg/L in clay jugs, with 200L of tap water and weekly colonized with 25 third instar *Ae. aegypti* larvae, achieved a persistence period of 27 days. In our assessments with the same spinosad concentration, there was a minimum persistence of eight weeks (56 days) in every population (mortality higher than 80%). In other semi-field evaluation using plastic bowls also containing 1L of spinosad aqueous solution (1mg/L) the persistence was kept for eight weeks on *Ae. aegypti* larvae [41]. Differences in the nature of the recipients used on the field simulations may justify divergences on results.

In Brazil, the visit of health agents on households to check and treat possible foci happens every eight weeks [65]. Spinosad appeared to be efficient against Brazilian *Ae. aegypti* populations, presenting larvae mortality over 80% for seven to eight weeks. Therefore, schedule of mosquito control measures carried out by the health agents can be maintained. New assessments on a field simulation with reservoirs of different natures, expositions to different light and temperature conditions will be important to measure the product's effect on various environments.

## Conclusion

The data presented here suggests that there is no cross-resistance of the spinosad with the temephos, regardless the resistance level to the organophosphate. Spinosad is a less aggressive product towards the environment and presents low toxicity for non-targeted insects and low risk for humans and fauna. Also, it has long residual effect on field populations with different levels of resistance. Therefore, spinosad is showing to be a promising product for public health usage, especially if used carefully. However, deeper studies that better evaluate the action mechanisms of spinosad in Brazilian *Ae. aegypti* populations are necessary to foresee possible events of tolerance and resistance to the product and to avoid troubles arising from the inadequate use of this tool.

## Supporting information

**S1 Table. Lethal concentration average and Confidence Interval (CI) according to exposure time of Rockefeller larvae to spinosad.**

(DOCX)

**S1 Fig. Field simulation for evaluation of the spinosad effect over the mortality of *Ae. aegypti* larvae.** The average of mortality percentage for Rockefeller, Marilia and Salvador is represented for bioassays conducted in three distinct moments:

A–July to September 2011, B–September to November 2011 and C– December 2011 to February 2012. The red dotted line indicates the mortality level at 80%. The physicochemical variations, pH and temperature of the solution in the deposits during the three assays are represented in D. The dose tested was 0.5mg/L. Every week the tanks were recolonized by removing the dead and the remaining living larvae and introducing 30 new 3rd instar larvae. The tanks were covered either with a lid (1<sup>st</sup> and 2<sup>nd</sup> replicates) or with a nylon mesh (3<sup>rd</sup> replicate).  
(TIF)

**S2 Fig. Field simulation for evaluation of the spinosad effect over the mortality of *Ae. aegypti* larvae.** The average mortality percentage for Rockefeller, Goiânia and Santarém is represented for bioassays conducted in three distinct moments: A- May to June 2012, B–July to September 2012 and C–September to November 2012. The dotted red line indicates the mortality level at 80%. The physicochemical variations, pH and temperature of the solution in the deposits during the three assays are represented in D. The dose tested was 0.5mg/L. Every week the tanks were recolonized by removing the dead and the remaining living larvae and introducing 30 new 3rd instar larvae. The tanks were covered either with a lid (1<sup>st</sup> and 2<sup>nd</sup> replicates) or with a nylon mesh (3<sup>rd</sup> replicate).  
(TIF)

## Acknowledgments

The authors are thankful to the Brazilian Health Ministry for the spinosad samples and to Gilberto Couto Alcantara for technical contribution on the mosquito colony maintenance.

## Author Contributions

**Conceptualization:** MLGM LSD JBPL.

**Data curation:** LSD.

**Formal analysis:** LSD JBPL MLGM MTMA VGO ASD AJM LGSRB CMR.

**Funding acquisition:** JBPL MLGM.

**Investigation:** LSD MTMA VGO ASD.

**Methodology:** LSD MLGM MTMA VGO ASD JBPL.

**Project administration:** JBPL MLGM.

**Resources:** JBPL MLGM.

**Supervision:** JBPL MLGM.

**Validation:** JBPL MLGM.

**Visualization:** LSD LGSRB.

**Writing – original draft:** LSD JBPL MLGM AJM CMR LGSRB.

**Writing – review & editing:** JBPL MLGM AJM CMR LGSRB.

## References

1. Consoli RAG., Lourenço-de-Oliveira. Principais Mosquitos de Importância Sanitária no Brasil. Cadernos de Saúde Pública. 1st ed. Rio de Janeiro: Fiocruz; 1994.

2. Gubler DJ. Epidemic dengue/dengue hemorrhagic fever as a public health, social and economic problem in the 21st century. *Trends in Microbiology*. 2002 Feb; 10(2):100–3. PMID: [11827812](#)
3. Walker KR, Joy TK, Eilers-Kirk C, Ramberg FB. Human and environmental factors affecting *Aedes aegypti* distribution in an arid urban environment. *Journal of the American Mosquito Control Association*. 2011 Jun; 27(2):135–41. <https://doi.org/10.2987/10-6078.1> PMID: [21805845](#)
4. Pan American Health Organization (1994) Dengue and dengue hemorrhagic fever in the Américas: guidelines for prevention and control. (PAHO Publication 548) ISBN: 92 75 11548 6. Washington DC: Pan American Health Organization.
5. Gould EA, Solomon T. Pathogenic flaviviruses. *Lancet*. 2008 Feb 9; 371(9611):500–9. PMID: [18262042](#)
6. Honório NA, Câmara DCP, Calvet GA, Brasil P. Chikungunya: an arbovirus infection in the process of establishment and expansion in Brazil. *Cadernos de Saúde Pública*. 2015 May; 31(5):906–8. <https://doi.org/10.1590/0102-311XPE020515> PMID: [26083166](#)
7. Li MI, Wong PSJ, Ng LC, Tan CH. Oral susceptibility of Singapore *Aedes* (*Stegomyia*) *aegypti* (Linnaeus) to Zika virus. *PLoS Neglected Tropical Diseases*. 2012 Jan; 6(8):e1792. <https://doi.org/10.1371/journal.pntd.0001792> PMID: [22953014](#)
8. Wong P-SJ, Li MI, Chong C-S, Ng L-C, Tan C-H. *Aedes* (*Stegomyia*) *albopictus* (Skuse): a potential vector of Zika virus in Singapore. *PLoS Neglected Tropical Diseases*. 2013 Jan; 7(8):e2348. <https://doi.org/10.1371/journal.pntd.0002348> PMID: [23936579](#)
9. Tauil PL. [Urbanization and dengue ecology]. *Cadernos de Saúde Pública*. 2001 Jan; 17 Suppl:99–102.
10. Medronho R de A. Dengue no Brasil: desafios para o seu controle. *Cadernos de Saúde Pública*. 2008 May; 24(5):948–9. PMID: [18461223](#)
11. Attaran A, Roberts DR, Curtis CF, Kilama WL. Balancing risks on the backs of the poor. *Nature Medicine*. 2000 Jul; 6(7):729–31. <https://doi.org/10.1038/77438> PMID: [10888909](#)
12. Walker K. Cost-comparison of DDT and alternative insecticides for malaria control. *Medical and Veterinary Entomology*. 2000 Dec; 14(4):345–54. PMID: [11129697](#)
13. Ferreira BJ, Souza M de FM, Soares Filho AM, Carvalho AA. The historical evolution of dengue prevention and control programs in Brazil. *Ciência & Saúde Coletiva* 2009 Jun; 14(3):961–72.
14. Macoris M de LG, Andrighetti MTM, Takaku L, Glasser CM, Garbeloto VC, Cirino VC. Alteração de resposta de suscetibilidade de *Aedes aegypti* a inseticidas organofosforados em municípios do Estado de São Paulo, Brasil. *Revista de Saúde Pública*. 1999 Oct; 33(5):521–2. PMID: [10576756](#)
15. Lima JBP, Da-Cunha MP, Da Silva RCJ, Galardo AKR, Soares S da S, Braga IA, et al. Resistance of *Aedes aegypti* to organophosphates in several municipalities in the State of Rio de Janeiro and Espírito Santo, Brazil. *The American Journal of Tropical Medicine and Hygiene*. 2003 Mar; 68(3):329–33. PMID: [12685640](#)
16. Fontoura NG, Bellinato DF, Valle D, Lima JBP. The efficacy of a chitin synthesis inhibitor against field populations of organophosphate-resistant *Aedes aegypti* in Brazil. *Memórias do Instituto Oswaldo Cruz*. 2012 May; 107(3):387–95. PMID: [22510835](#)
17. Braga IA, Lima JBP, Soares S da S, Valle D. *Aedes aegypti* resistance to temephos during 2001 in several municipalities in the states of Rio de Janeiro, Sergipe, and Alagoas, Brazil. *Memórias do Instituto Oswaldo Cruz*. 2004 Mar; 99(2):199–203. PMID: [15250476](#)
18. Montella IR, Martins AJ, Viana-Medeiros PF, Lima JBP, Braga IA, Valle D. Insecticide resistance mechanisms of Brazilian *Aedes aegypti* populations from 2001 to 2004. *The American Journal of Tropical Medicine and Hygiene*. 2007 Sep; 77(3):467–77. PMID: [17827362](#)
19. Lima E, Paiva M, de Araújo A, da Silva É, da Silva U, de Oliveira L, et al. Insecticide resistance in *Aedes aegypti* populations from Ceará, Brazil. *Parasites and Vectors*. 2011 Jan; 4(1):5.
20. Chediak M, G Pimenta F Jr, Coelho GE, Braga IA, Lima JBP, Cavalcante KRL, et al. Spatial and temporal country-wide survey of temephos resistance in Brazilian populations of *Aedes aegypti*. *Memórias do Instituto Oswaldo Cruz*. 2016 May; 111(5):311–21.
21. Centers for Disease Control and Prevention—CDC. Insecticide Resistance. 2016 [cited 2016 Dec 14]. Available: <https://www.cdc.gov/zika/vector/insecticide-resistance.html>
22. Brogdon WG, McAllister JC. Insecticide resistance and vector control. *Emerging infectious diseases*. 1998 Jan; 4(4):605–13. <https://doi.org/10.3201/eid0404.980410> PMID: [9866736](#)
23. Hemingway J, Hawkes NJ, McCarroll L, Ranson H. The molecular basis of insecticide resistance in mosquitoes. *Insect biochemistry and molecular biology*. 2004 Jul; 34(7):653–65. <https://doi.org/10.1016/j.ibmb.2004.03.018> PMID: [15242706](#)
24. Araújo A, Araujo Diniz D, Helvecio E, de Barros R, de Oliveira CM, Ayres CF, et al. The susceptibility of *Aedes aegypti* populations displaying temephos resistance to *Bacillus thuringiensis israelensis*: a basis

- for management. *Parasites and Vectors* [Internet]. 2013 [cited 2016 Jul 20]; 6(1):297. Available from: <http://parasitesandvectors.biomedcentral.com/articles/10.1186/1756-3305-6-297> <https://doi.org/10.1186/1756-3305-6-297> PMID: 24499507
25. Hemingway J, Ranson H. Insecticide resistance in insect vectors of human disease. *Annual Review of Entomology*. 2000 Jan; 45(1):371–91.
  26. Bisset JA. Uso correcto de insecticidas: control de la resistencia. *Revista Cubana de Medicina Tropical*. 2002 Sep; 54(3):202–19. PMID: 15846946
  27. Zaim M, Guillet P. Alternative insecticides: an urgent need. *Trends Parasitol*. 2002 Apr; 18(4):161–3. PMID: 11998703
  28. Dias CN, Moraes DFC. Essential oils and their compounds as *Aedes aegypti* L. (Diptera: Culicidae) larvicides: review. *Parasitol Res*. [Internet] 2014 [cited 2016 Jul 20]; Feb; 113(2):565–92. Available: <http://www.ncbi.nlm.nih.gov/pubmed/24265058> <https://doi.org/10.1007/s00436-013-3687-6> PMID: 24265058
  29. Darriet F, Duchon S, Hougard JM. Spinosad: a new larvicide against insecticide-resistant mosquito larvae. *Journal of the American Mosquito Control Association*. 2005 Dec; 21(4):495–6. [https://doi.org/10.2987/8756-971X\(2006\)21\[495:SANLAI\]2.0.CO;2](https://doi.org/10.2987/8756-971X(2006)21[495:SANLAI]2.0.CO;2) PMID: 16506584
  30. Kirst HA, Michel KH, Mynderase JS, Chio EH, Yao RC, Nakasukasa WM, et al. Synthesis and Chemistry of Agrochemicals III. in: Baker DR, Fenyes JG, Steffens JJ, editors. Washington, DC: American Chemical Society; 1992. pp. 214–225.
  31. Sparks T, Thompson G, Larson L, Kirst H, Jantz O, Worden T, et al. Biological characteristics of the spinosyns: a new and naturally derived insect control agent. In: Proceedings, Beltwide Cotton Conference. San Antonio, Texas: National Cotton Council; 1995. p. 903–7.
  32. Thompson G., Busacca J., Jantz O., Kirst H., Larson L., Sparks T, et al. Spinosyns: na overview of new natural insect management systems. In: Beltwide Cotton Conference. San Antonio, Texas: National Cotton Council; 1995. p. 1039–43.
  33. Salgado VL. Studies on the Mode of Action of Spinosad: Insect Symptoms and Physiological Correlates. *Pesticide Biochemistry and Physiology*. 1998 Jul; 60(2):91–102.
  34. Watson GB. Actions of Insecticidal Spinosyns on  $\gamma$ -Aminobutyric Acid Responses from Small-Diameter Cockroach Neurons. *Pesticide Biochemistry and Physiology*. 2001 Sep; 71(1):20–8.
  35. Environmental Protection Agency. Spinosad; notice of filing a pesticide petition to establish a tolerance for a certain pesticide chemical in or on food. *Fed Regist* [Internet]. 2005 [cited 2016 Jul 20] 70 (138):41730–5. Available: <http://www.epa.gov/EPA-PEST/2005/July/Day-20/p13977.htm>
  36. Miles M, Dutton R. Spinosad—a naturally derived insect control agent with potential for use in glasshouse integrated pest management systems. In: Proceedings, 52nd International Symposium on Crop Protection, Gent, Belgium, 9 May 2000, Part I [Internet]. Faculteit Landbouwkundige en Toegepaste Biologische Wetenschappen; 2000 [cited 2016 Jul 22]. p. 393–400. Available: <http://www.cabdirect.org/abstracts/20013030388.html;jsessionid=6BECDB895B3B848F1FDDB6D31C0CB7C>
  37. Williams T, Valle J, Viñuela E. Is the Naturally Derived Insecticide Spinosad Compatible with Insect Natural Enemies? *Biocontrol science and technology*. 2003 Aug; 13(5):459–75.
  38. Pereira BB, Caixeta ES, Freitas PC, Santos VSV, Limongi JE, de Campos Júnior EO, et al. Toxicological assessment of spinosad: Implications for integrated control of *Aedes aegypti* using larvicides and larvivorous fish. *Journal of toxicology and environmental health. Part A*. 2016; 79(12):477–81. <https://doi.org/10.1080/15287394.2016.1176974> PMID: 27294296
  39. Marcombe S, Darriet F, Agnew P, Etienne M, Yp-Tcha M-M, Yébakima A, et al. Field efficacy of new larvicide products for control of multi-resistant *Aedes aegypti* populations in Martinique (French West Indies). *The American Journal of Tropical Medicine and Hygiene*. 2011 Jan; 84(1):118–26. <https://doi.org/10.4269/ajtmh.2011.10-0335> PMID: 21212213
  40. Prabhu K, Murugan K, Nareshkumar A, Bragadeeswaran S. Larvicidal and pupicidal activity of spinosad against the malarial vector *Anopheles stephensi*. *Asian Pacific Journal of Tropical Medicine*. 2011 Aug; 4(8):610–3. [https://doi.org/10.1016/S1995-7645\(11\)60157-0](https://doi.org/10.1016/S1995-7645(11)60157-0) PMID: 21914537
  41. Bond JG, Marina CF, Williams T. The naturally derived insecticide spinosad is highly toxic to *Aedes* and *Anopheles* mosquito larvae. *Medical and Veterinary Entomology*. 2004 Mar; 18(1):50–6. PMID: 15009445
  42. Sadanandane C, Boopathi Doss PS, Jambulingam P, Zaim M. Efficacy of two formulations of the bioinsecticide spinosad against *Culex quinquefasciatus* in India. *Journal of the American Mosquito Control Association*. 2009 Mar; 25(1):66–73. <https://doi.org/10.2987/08-5807.1> PMID: 19432070
  43. WHO. Spinosad DT in Drinking-water: Use for Vector Control in Drinking-water Sources and Containers [Internet]. Geneva, Switzerland: World Health Organization; 2010. 7 p. Available: [http://www.who.int/water\\_sanitation\\_health/dwq/chemicals/spinosadbg.pdf](http://www.who.int/water_sanitation_health/dwq/chemicals/spinosadbg.pdf)

44. Braga IA, Gomes A de C, Nelson M, Mello R de C, Bergamaschi DP, de Souza JM. Comparative study between larval surveys and ovitraps to monitor populations of *Aedes aegypti*. *Revista da Sociedade Brasileira de Medicina Tropical*. 2000 Jan; 33(4):347–53. PMID: [10936947](#)
45. WHO. WHOPES-Recommended Compounds and Formulations for Control of Mosquito Larvae [Internet]. 2016 [cited 2016 Dec 14]. Available: [http://www.who.int/whopes/Mosquito\\_larvicides\\_March\\_2016.pdf](http://www.who.int/whopes/Mosquito_larvicides_March_2016.pdf)
46. Kuno G. Early history of laboratory breeding of *Aedes aegypti* (Diptera: Culicidae) focusing on the origins and use of selected strains. *Journal of Medical Entomology*. 2010 Nov; 47(6):957–71. PMID: [21175042](#)
47. WHO. Instructions for determining the susceptibility or resistance of adult mosquito to organochlorine organophosphate and carbonate insecticides—diagnostic. (WHO/VBC/1981), editor. Geneva, Switzerland: World Health Organization; 1981.
48. CEPAGRI. Clima dos municípios paulistas [Internet]. 2012 [cited 2016 Jan 12]. Available from: [http://www.cpa.unicamp.br/outras-informacoes/clima\\_muni\\_329.html](http://www.cpa.unicamp.br/outras-informacoes/clima_muni_329.html)
49. Raymond M. Presentation d'une programme d'analyse logprobit pour microordinateur cahiers Orstom. *Series Entomology and Medical Parasitology*. 1985; 23(2):117–21.
50. Mazzarri MB, Georghiou GP. Characterization of resistance to organophosphate, carbamate, and pyrethroid insecticides in field populations of *Aedes aegypti* from Venezuela. *Journal of the American Mosquito Control Association*. 1995 Sep; 11(3):315–22. PMID: [8551300](#)
51. Hemingway J, Field L, Vontas J. An overview of insecticide resistance. *Science*. 2002 Oct 4; 298(5591):96–7. <https://doi.org/10.1126/science.1078052> PMID: [12364782](#)
52. Bisset JA, Marín R, Rodríguez MM, Severson DW, Ricardo Y, French L, et al. Insecticide resistance in two *Aedes aegypti* (Diptera: Culicidae) strains from Costa Rica. *Journal of Medical Entomology*. 2013 Mar; 50(2):352–61. PMID: [23540124](#)
53. Cui F, Raymond M, Qiao C-L. Insecticide resistance in vector mosquitoes in China. *Pest Management Science*. 2006 Nov; 62(11):1013–22. <https://doi.org/10.1002/ps.1288> PMID: [16953491](#)
54. Brown AW. The insecticide-resistance problem: a review of developments in 1956 and 1957. *Bull World Health Organ* [Internet]. 1958 Jan [cited 2016 Mar 21]; 18(3):309–21. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2537653&tool=pmcentrez&rendertype=abstract> PMID: [13536795](#)
55. Deming R, Manrique-Saide P, Medina Barreiro A, Cardeña EUK, Che-Mendoza A, Jones B, et al. Spatial variation of insecticide resistance in the dengue vector *Aedes aegypti* presents unique vector control challenges. *Parasites and Vectors*. 2016 Feb 4; 9(1):67.
56. Diniz DFA, Melo-Santos MAV de, Santos EM de M, Beserra EB, Helvecio E, de Carvalho-Leandro D, et al. Fitness cost in field and laboratory *Aedes aegypti* populations associated with resistance to the insecticide temephos. *Parasites and Vectors*. 2015 Jan; 8(1):662.
57. WHO. Guidelines for Laboratory and Field Testing of Mosquito Larvicides. World Health Organization Eradication, Dept. of Communicable Disease, Prevention, Control and Scheme and WHO Pesticide Evaluation; 2005. 39 p.
58. Romi R, Proietti S, Di Luca M, Cristofaro M. Laboratory evaluation of the bioinsecticide Spinosad for mosquito control. *Journal of the American Mosquito Control Association*. 2006 Mar; 22(1):93–6. [https://doi.org/10.2987/8756-971X\(2006\)22\[93:LEOTBS\]2.0.CO;2](https://doi.org/10.2987/8756-971X(2006)22[93:LEOTBS]2.0.CO;2) PMID: [16646328](#)
59. Paul A, Harrington LC, Scott JG. Evaluation of novel insecticides for control of dengue vector *Aedes aegypti* (Diptera: Culicidae). *Journal of Medical Entomology*. 2006 Jan; 43(1):55–60. [https://doi.org/10.1603/0022-2585\(2006\)043\[0055:EONIFC\]2.0.CO;2](https://doi.org/10.1603/0022-2585(2006)043[0055:EONIFC]2.0.CO;2) PMID: [16506447](#)
60. Pérez CM, Marina CF, Bond JG, Rojas JC, Valle J, Williams T. Spinosad, a naturally derived insecticide, for control of *Aedes aegypti* (Diptera: Culicidae): efficacy, persistence, and elicited oviposition response. *Journal of Medical Entomology*. 2007 Jul; 44(4):631–8. PMID: [17695018](#)
61. Lima JBP, Melo NV de, Valle D. Residual effect of two *Bacillus thuringiensis* var. *israelensis* products assayed against *Aedes aegypti* (Diptera: Culicidae) in laboratory and outdoors at Rio de Janeiro, Brazil. *Revista do Instituto de Medicina Tropical de São Paulo*. 2005 Jun; 47(3):125–30. PMID: [16021284](#)
62. WHO. Scientific working group on dengue [Internet]. Switzerland, Geneva: World Health Organization; 2000. p. 10. Available: <http://www.who.int/tdr/publications/documents/dengue-swg.pdf>
63. Cleveland Cheryl B., Bormett Gary A., Saunders Donald G., Powers Fred L., McGibbon Alec S., Reeves Graham L., et al. Environmental Fate of Spinosad. 1. Dissipation and Degradation in Aqueous Systems. *Journal of Agricultural and Food Chemistry*. 2002 May; 50: 3244–3256. PMID: [12009994](#)
64. Thavara U, Tawatsin A, Asavadachanukorn P, Mulla MS. Field evaluation in Thailand of spinosad, a larvicide derived from *Saccharopolyspora spinosa* (Actinomycetales) against *Aedes aegypti* (L.) larvae.

The Southeast Asian Journal of Tropical Medicine and Public Health. 2009 Mar; 40(2):235–42. PMID: [19323007](#)

65. Brasil, Ministério da Saúde, Secretaria de Vigilância em Saúde, Departamento de Vigilância Epidemiológica. Diretrizes nacionais para prevenção e controle de epidemias de dengue. Série A. N. Brasília: Ministério da Saúde; 2009. 160 p.