

ORIGINAL ARTICLE

Determinants of Quality of Life and Return to Work Following Acute Respiratory Distress Syndrome

A Systematic Review

Frank Dodoo-Schittko, Susanne Brandstetter, Sebastian Blecha, Kathrin Thomann-Hackner, Magdalena Brandl, Helge Knüttel, Thomas Bein, Christian Apfelbacher

SUMMARY

Background: Acute respiratory distress syndrome (ARDS) in adults is a consequence of lung damage caused by either pulmonary or extrapulmonary disease. Survivors often suffer from an impaired health-related quality of life (HRQoL), mental and physical impairments, and persistent inability to work.

Methods: In this systematic review of the literature, we consider the determinants of HRQoL and return to work (RtW). 24 observational studies showing a statistical association between one or more determinants and HRQoL or RtW were included. Because of the heterogeneity of these studies, no statistical aggregation of the individual effect estimates was carried out; instead, the results are summarized descriptively.

Results: Psychopathological manifestations, in particular, are associated with impaired quality of life. In contrast, many care- and disease-related determinants had only small, non-significant effects on HRQoL and RtW. The one-second capacity was found in all studies to be positively associated with the HRQoL. ARDS induced by sepsis seems to be a risk factor for a lower HRQoL in comparison to ARDS of other causes. A synthesis of the evidence is impeded both by the high level of heterogeneity of studies and by the high risk of selection bias in all studies.

Conclusion: The identification of determinants of impaired quality of life after ARDS is essential for the assessment of clinically relevant interventions. In multiple studies, major significant effects were only observed when determinants the content of which was closely related to the scales of the HRQoL instruments were measured at the same time as the HRQoL.

► Cite this as:

Dodoo-Schittko F, Brandstetter S, Blecha S, Thomann-Hackner K, Brandl M, Knüttel H, Bein T, Apfelbacher C: Determinants of quality of life and return to work following acute respiratory distress syndrome—a systematic review. *Dtsch Arztebl Int* 2017; 114: 103–9. DOI: 10.3238/arztebl.2017.0103

Acute respiratory distress syndrome (ARDS) in adults results in a life-threatening damage of the lung. The most important risk factor for ARDS is pneumonia, followed by sepsis and aspiration (1). The extent of injury and the resulting hypoxemia usually make mechanical ventilation necessary. The ARDS criteria established in 1994 by the American-European Consensus Conference (AECC) were long used as the standard for ARDS diagnosis (2). In 2011, the AECC criteria were revised by the Berlin Definition (3), which includes the criterion “acute onset” (<7 days) in addition to the presence of bilateral infiltrates that cannot be attributed exclusively to left-sided cardiac insufficiency. Classification of severity (mild, moderate, or severe) is based on the oxygenation index according to Horowitz.


With a prevalence of approximately 10% of all patients treated in an intensive care unit (ICU), and an in-hospital mortality of 35–46% (depending on severity), ARDS represents a major challenge for intensive medical care (1). An overall decrease in ARDS mortality could be achieved continuously until the 1990s (4). However, survivors of ARDS also suffer from persistent physical and psychological morbidity. For instance, the point prevalence of depressive symptoms (33%) or of generalized anxiety disorder (40%) is strongly increased in survivors two years after ARDS (5). An increased risk of post-traumatic stress disorder (PTSD) is present in 29% of ARDS patients (6). Further, a prospective cohort study showed that five years after the disease, the health-related quality of life (HRQoL) of survivors of ARDS was reduced by about one standard deviation as compared to a matched control population (7). Finally, a systematic review from 2006 (8) found that HRQoL of survivors of ARDS was significantly reduced in nearly all domain-specific pooled scores of the 36-item Short-Form Health Survey (SF-36) (9) as compared to population norms.

In addition to HRQoL, return to work (RtW) is an endpoint for survivors of ARDS, which is a good operationalization—indeed, better than clinical

Department for Epidemiology and Preventive Medicine/Medical Sociology, University of Regensburg: Dr. phil. Dipl.-Psych. Dodoo-Schittko, Dr. biol. hum. Dipl.-Psych. Brandstetter, Brandl, MPH Prof. Dr. sc. hum. Apfelbacher, PhD

Department of Anesthesiology, University Hospital Regensburg: Dr. med. Blecha, Dr. med. Thomann-Hackner, Prof. Dr. med. Bein, M.A.

University Library of Regensburg, University of Regensburg: Dr. rer. nat. Knüttel



Flow diagram of the study selection processes (according to the PRISMA Statement) ARDS, acute respiratory distress syndrome; HRQoL, health-related quality of life; RtW, return to work

parameters—for overcoming morbidity. In a German cohort of ARDS survivors, only 64% had returned to their previous work five years after ARDS (10).

Recognizing the factors associated with quality of life or RtW would help to identify risk groups and, notably, would serve as a starting point for intervention measures that could increase the success of treatment. Here, we provide, to our knowledge, the first systematic review of determinants of HRQoL and/or RtW in survivors of ARDS.

Methods

Protocol and registration

The presentation of this systematic review is based on the recommendations of the Preferred Reporting Items for Systematic Reviews (PRISMA) (e1). The protocol was registered in the International Prospective Register of Systematic Reviews PROSPERO (CRD42014014335) (e2).

Inclusion criteria

To ensure the highest possible external validity of the results found in the studies, only observational studies were included. Interventional studies, which often have highly selected populations, were excluded.

The examined population had to consist of survivors of ARDS or acute lung injury (ALI) (as defined by Berlin or AECC; deviating diagnostic criteria were accepted if described).

Statistical associations between determinants, and HRQoL and/or RtW, in survivors of ARDS or ALI had to be reported.

Information sources and searches

Searches of Cochrane Systematic Reviews, MEDLINE, PSYINDEX, PsycINFO, Embase, Science Citation Index Expanded, and Social Science Citation Index Expanded were conducted without date limitations, from the beginning of each database until 13 August 2014. A comprehensive, highly sensitive research strategy was developed by SB, FDS, and HK. The search strategy and documentation were based on latest recommendations (e3, e4).

Study selection

For study selection, the authors formed working pairs, each of which consisted of a clinical practitioner (SB, KTH) and a psychologist (SuB, FDS). Each pair then made a pre-selection based on titles and abstracts in a first step. The original articles to be included were then selected in a second step based on the full text.

Data collection

The following characteristics were extracted from the selected original articles by the first author (FDS):

- General information about the study (authors, year of publication, country, period of data collection, study design)
- Description of the sample/cohort (inclusion and exclusion criteria, with particular emphasis on the underlying diagnostic criteria of ARDS/ALI, sex, and age)
- Operationalization of the outcomes of HRQoL and RtW

Additionally, the effect sizes and significance of the investigated determinants were recorded whenever this had been documented or could be calculated from the indicated statistical values.

Risk of bias of individual studies

Since this systematic review was meant to include the full range of observational study designs, no instrument was suitable for determining a risk of bias applicable to all types of studies. Thus, an instrument was developed by three authors (FDS, SuB, and CA), based on the Newcastle-Ottawa Scale, for cohort and case-control studies (e5) (eBox). Using this instrument, each study was evaluated independently by two persons with regard to risk of selection and information bias.

Synthesis of results

Because of the heterogeneity of the studies (in study design, timing of measurements, determinants, and outcomes), the synthesis of results is a descriptive summary rather than a meta-analysis.

Results

The electronic search revealed 4559 studies. After the screening process, 24 studies met the inclusion criteria

(Figure 1). The size of the investigated cohorts/samples ranged from N = 15 to N = 152. Cross-sectional study designs (19 studies) outweighed designs with prospective (13 studies) and retrospective (2 studies) data collection approaches. In the majority of studies, HRQoL / RtW and determinants were obtained both prospectively and cross-sectionally.

Studies with follow-ups varied both in number (from one to four) and timing of follow-ups (Table, Figure 2). Statistical associations between determinants and HRQoL were reported in 23 of the 24 studies. Determinants for RtW were analyzed in six studies. HRQoL was recorded via six generic and two disease-specific instruments (Table). The risk of information bias is low due to the use of validated measurement instruments in almost all studies. In contrast, all studies have an unclear or high risk of selection bias.

Determinants of HRQoL

Overviews of the determinants investigated are given in eTables 1 and 2. In the category of sociodemographic determinants, only age seems to be associated with a reduced quality of life after surviving ARDS (7, 11, 12). The physical health component summary score of SF-36, for which a higher value indicates a higher quality of life (13), showed a strong negative correlation with age (ρ [rho] = -0.52).

The highest number of determinants was found among the disease-related characteristics. Results of pulmonary function testing were especially frequently examined at the same time as HRQoL and RtW. Among the different spirometric values and across studies, the Forced Expiratory Volume in 1 second (FEV1) is significantly positively associated with HRQoL (ρ = 0.16 to ρ = 0.46) (14–16). However, a high forced vital capacity (FVC) shows no significant correlation with HRQoL or RtW (12, 14–17). The effect sizes for the various quality of life instruments and their different domains range between ρ = -0.004 and ρ = 0.58.

In three studies, various aspects of morbidity during ARDS (based on Acute Physiology and Chronic Health Evaluation II, APACHE II; Lung Injury Score, LIS; Charlson–Deyo comorbidity score; extent of extrapulmonary organ failure) were examined as predictors of later HRQoL (12, 18, 19). Only low scores in LIS proved to be a significant determinant for reduced HRQoL measured using a dimension of the Sickness Impact Profile (SIP) modified for lung function (12, 20). A few studies examined the etiology of ARDS in more depth (14, 18, 19, 21). In the SF-36 domains, sepsis-induced ARDS, as compared to trauma-induced ARDS, was associated with small to moderate negative effects in the SF-36 domains as compared to trauma-induced ARDS (21). Larger effect sizes (Cohen's d = 0.57–0.65) resulted when the St. George's Respiratory Questionnaire (SGRQ) was used to determine quality of life (21).

The duration of medical care as a care-related determinant was examined in more detail in seven studies

(10, 13, 15, 19, 21–23). The total duration of treatment in both the ICU and the hospital (ρ = -0.34 to ρ = -0.45) (21, 22), as well as the duration of mechanical ventilation (ρ = -0.44 to ρ = 0.13) (10, 13, 15, 19, 21, 22) and extracorporeal membrane oxygenation (ECMO) (23), correlate negatively with some SF-36 domains. Of the supportive measures for ARDS treatment during the acute phase, ECMO was shown to have negative effects of up to medium strength (Cohen's d = 0.62) on the physical health component summary score of SF-36 (10, 24).

For the psychosocial determinants, the presence of depressive symptoms showed strong negative correlations, especially with the mental health component summary score of SF-36 (ρ = -0.64 to ρ = -0.94) (22, 25, 26). Symptoms of PTSD (6, 10, 27) and anxiety disorder (22, 28) also show strong negative associations with HRQoL. Likewise, cognitive deficits in the domains of memory, attention, and executive functions appear to be associated with the reduction of some domains of SF-36 and SIP (22, 29, 30). Finally, effect sizes of Cohen's d = 0.45 are observed for the physical role functioning of the SF-36 (22).

Determinants of return to work (RtW)

Of the care-related determinants, neither the duration of ICU treatment and total hospital stay (10) nor treatment with ECMO (10, 17, 24, 31) showed significant effects on RtW, although patients treated with ECMO have a twice-as-high chance of long-term incapacity for work as those not treated with ECMO (17). Survivors of ARDS with moderate to severe depressive symptoms have a 0.2-fold lower chance of return to work after two years (25). Aside from this, no significant effects were observed for disease-related (12) or sociodemographic (10) characteristics on RtW. For these variables, no effect sizes could be extracted from the original articles.

Discussion

While previous review articles have compiled evidence for a decreased quality of life in survivors of ARDS (8, 32), the explicit aim of this review was to provide a summary of the research on the determinants of HRQoL and RtW in ARDS survivors. Importantly, being able to identify determinants provides the opportunity to intervene during the disease phase and to reduce the incidence of physical and psychological impairments.


Evidence taken from the studies analyzed here is limited, due to the strong heterogeneity of these studies and their often low methodological quality (because of selection bias and insufficient statistical analyses). Nonetheless, for generic quality of life instruments, larger effects only exist when there is a narrow contextual relation (and probably also a close temporal link) between the investigated determinant and the quality of life instrument. Thus, no study shows that the initial disease severity predicts later HRQoL. Similarly, there is a lack of significant correlations between HRQoL

TABLE

Characteristics of the 24 studies included in the review

First author (year) (ref.)	Type of data collection	Country/region	ARDS criteria	HRQoL instrument	N (survivors)	Age
Adhikari et al. (2011) (26)	Cross-sectional	Canada	AECC	SF-36	48	Md = 48 IQR = 39–61
Adhikari et al. (2009) (25)	Cross-sectional	Canada	AECC	–	71	Md = 42 IQR = 35–56
Angus et al. (2001) (18)	Prospective	USA	AECC	QWB	104	M = 44.7 SD = 15.0
Briegel et al. (2013) (10)	Prospective/ cross-sectional	Germany	AECC	SF-36	125	–
Hopkins et al. (2004) (22)	Prospective/ cross-sectional	USA	„ARDS survivors“	SF-36	73	M = 45.8 SD = 16.4
Davidson et al. (1999) (21)	Prospective matched	USA	AECC	SF-36 SGRQ	77	Md = 40.6
Deja et al. (2006) (6)	Cross-sectional	Germany	„Severe ARDS“	SF-36	65	M = 39 SD = 15
Herridge et al. (2011) (7)	Prospective/ cross-sectional	Canada	AECC	SF-36	64	Md = 44
Kapfhammer et al. (2004) (27)	Cross-sectional	Germany	AECC	SF-36	46	Md = 36.5
Kim et al. (2004) (15)	Prospective/ cross-sectional	USA	AECC	CRQ, SQLI	29	–
Masclans et al. (2011) (14)	Prospective/ cross-sectional	Spain	AECC	NHP	38	Md = 50 IQR = 34–55
Li et al. (2006) (16)	Cross-sectional	Hong Kong	AECC „ARDS caused by SARS“	SF-36	36	M = 42.0 SD = 12.1
Linden et al. (2009) (11)	Cross-sectional	Sweden	„Severe ARDS and ECMO“	SGRQ	15	M = 40
Luyt et al. (2012) (24)	Prospective	France	„H1N1- associated ARDS“	SF-36	37	ECMO Md = 35.5 IQR = 30–39 no ECMO Md = 42 IQR = 32.75–51.25
McHugh et al. (1994) (12)	Prospective/ cross-sectional	USA	AECC	SIP	37	M = 41
Mikkelsen et al. (2009) (30)	Cross-sectional	USA	ARDS as self-reported	SF-36, SIP	79	M = 43.3 SD = 12.7
Rothenhausler et al. (2001) (29)	Cross-sectional	Germany	AECC	SF-36	46	M = 41.5 SD = 14.7
Schelling et al. (2000) (19)	Prospective/ cross-sectional	Germany	AECC	SF-36	50	Md = 34.5
Schelling et al. (1998) (17)	Prospective/ retrospective	Germany	AECC	SF-36	66	Md = 36
Schmidt et al. (2013) (23)	Prospective	France	ARDS and ECMO	SF-36, SGRQ	67	–
Stevenson et al. (2013) (28)	Cross-sectional	USA	AECC ALI patients	EQ-5D, SF-36	152	Md = 49 IQR = 40–57
Stoll et al. (1998) (31)	Prospective matched	Germany	AECC	SF-36	28	ECMO Md = 34 no ECMO Md = 38
Weinert et al. (1997) (13)	Retrospective/ cross-sectional	USA	AECC ALI patients	SF-36	24	M = 40 SD = 12
Wilcox et al. (2013) (40)	Cross-sectional	Canada	AECC	SF-36	24	–

AECC, American-European Consensus Conference; ALI, acute lung injury; ARDS, acute respiratory distress syndrome; CRQ, Chronic Respiratory Questionnaire; ECMO, extracorporeal membrane oxygenation; EQ-5D, EuroQol Five Dimensions Questionnaire; HRQoL, health-related quality of life; IQR, interquartile range; M, mean; Md, median; NHP, Nottingham Health Profile; QWB, quality of well-being; SARS, severe acute respiratory syndrome; SD, standard deviation; SF-36, 36-Item Short-Form Health Survey; SGRQ, St George's Respiratory Questionnaire; SIP, Sickness Impact Profile; SQLI, Spitzer's Quality of Life Index



Distribution of the follow-up periods in the studies with fixed survey time points (11 studies). If a study reported several follow-up periods, all were included in this Figure. Studies with varying follow-up periods across the participants (13 studies) were not considered.

and/or RtW, and care-related determinants, for almost every investigated supportive measure (including nitric oxide [NO] inhalation, corticosteroid delivery, and renal replacement therapy). The only exception is ECMO therapy, which shows a significant negative correlation with some SF-36 domains. The reason for this seemingly harmful effect of ECMO is most likely due to the lack of statistical adjustment for disease severity. In this context, it should be noted that the inclusion of confounding variables in statistical models, which is necessary for observational studies, was missing from almost all studies.

In contrast, significant associations with large effect sizes are reported for operationalizations of mental and physical morbidity, which are closely related to some scales of the HRQoL instruments and which were measured in a cross-sectional design together with the outcome. In particular, many studies report that physical morbidity, psychopathological symptomatology, or cognitive deficits after ICU treatment have adverse effects on HRQoL or RtW. For instance, strong correlations have been shown between the results of the 6-minute walk test or some spirometric values and the physical function domains of the SF-36. The same is true for the presence of a psychopathological symptomatology and the mental health component summary score of the SF-36.

In order to judge the importance of these results for survivors of ARDS, the high prevalence of persistent physical and psychological morbidity following an ICU stay must be considered additionally (33). In this context, the term ‘post-intensive care syndrome’ (PICS) has been coined to refer to new or worsening impairments that can be physical (pulmonary, neuromuscular, physical function), cognitive (executive functions, memory, attention, visuospatial processing), and mental (anxiety disorders, PTSD, and depression)

(34). Therefore, there is a need for interventions that go beyond the established rehabilitation measures—with the aim of positively influencing HRQoL and RtW, both directly and indirectly.

ICU follow-up clinics represent a complex intervention, even though evidence is insufficient. These clinics are medical institutions specially designed for the diagnosis and management of common impairments following ICU treatment (35).

A further intervention that is currently being discussed in the literature is the use of ICU diaries, which has shown to be effective in reducing psychopathological symptoms in some studies (36, 37). ICU diaries are chronological, daily records of the patient’s ICU stay. They are usually filled in by the nursing staff, but relatives and friends can also comment on their visits to the patient. These diaries aim to help the patient fill in memory gaps about the ICU treatment and thus come to terms with their experiences.

Finally, several randomized controlled trials (RCTs) have highlighted the positive effects of early mobilization of patients during ICU stay (38). However, these findings are limited by the fact that the strongest significant effects were reported only for short-term outcomes at the time of hospital discharge (e.g., 6-minute walk test [39]). To date, no significant effects have been shown for long-term outcomes at 12 months post-ICU discharge (38).

Limitations

The studies examined here are highly heterogeneous due to their use of different tools—generic and disease-specific—for HRQoL assessment, the use of different measurement time points for the outcomes of interest, and the high level of variations in diagnostic criteria for acute lung failure. In addition, the high risk of a selection bias found in all studies leads to a loss of internal and external validity.

Summary

Survival of acute lung failure is often associated with pronounced psychological and physical sequelae. Determinants of the development of such damage, which impairs the quality of life, have not yet been sufficiently investigated. Future research should focus on identifying predictors by using appropriate statistical analyses, which would enable risk groups to be identified and would allow targeted interventions to improve quality of life and facilitate RtW. Since the effects of the HRQoL measurement of determinants that are temporally distant are often small, sample size calculation should be based on a formula that takes into account relevant effect sizes.

The negative association between physical and psychological morbidity and HRQoL in survivors of ARDS shows that new effective interventions are needed to improve health after ICU treatment. Currently promising approaches include introducing mobilization during intensive care, keeping ICU diaries, and establishing ICU follow-up clinics.

Conflict of interest statement

Prof. Bein is a member of the Medical Advisory Board of Novalung (XENIOS AG), Heilbronn, for which he received attendance and speaking fees. The other authors declare that no conflict of interest exists.

Manuscript received on 13 May 2016, revised version accepted on 25 November 2016.

Translated from the original German by Veronica A. Raker, PhD

REFERENCES

1. Bellani G, Laffey JG, Pham T, et al.: Epidemiology, patterns of care, and mortality for patients with acute respiratory distress syndrome in intensive care units in 50 countries. *JAMA* 2016; 315: 788–800.
2. Bernard GR, Artigas A, Brigham KL, et al.: The American-European Consensus Conference on ARDS. Definitions, mechanisms, relevant outcomes, and clinical trial coordination. *Am J Respir Crit Care Med* 1994; 149: 818–24.
3. Force ADT, Ranieri VM, Rubenfeld GD, et al.: Acute respiratory distress syndrome: the Berlin Definition. *JAMA* 2012; 307: 2526–33.
4. Phua J, Badia JR, Adhikari NK, et al.: Has mortality from acute respiratory distress syndrome decreased over time? A systematic review. *Am J Respir Crit Care Med* 2009; 179(3): 220–7.
5. Bienvenu OJ, Colantuoni E, Mendez-Tellez PA, et al.: Cooccurrence of and remission from general anxiety, depression, and posttraumatic stress disorder symptoms after acute lung injury: a 2-year longitudinal study. *Crit Care Med* 2015; 43: 642–53.
6. Deja M, Denke C, Weber-Carstens S, et al.: Social support during intensive care unit stay might improve mental impairment and consequently health-related quality of life in survivors of severe acute respiratory distress syndrome. *Crit Care* 2006; 10: R147.
7. Herridge MS, Tansey CM, Matte A, et al.: Functional disability 5 years after acute respiratory distress syndrome. *N Engl J Med* 2011; 364: 1293–304.
8. Dowdy DW, Eid MP, Dennison CR, et al.: Quality of life after acute respiratory distress syndrome: a meta-analysis. *Intensive Care Med* 2006; 32: 1115–24.
9. McHorney CA, Ware JE, Jr., Raczek AE: The MOS 36-Item Short-Form Health Survey (SF-36): II. Psychometric and clinical tests of validity in measuring physical and mental health constructs. *Med Care* 1993; 31: 247–63.

KEY MESSAGES

- Patients who survive ARDS often have a reduced health-related quality of life (HRQoL) and show more psychopathological symptoms (depression, post-traumatic stress disorder, anxiety disorders).
- Return to work (RtW) has been an endpoint in only a few studies.
- Significant associations with HRQoL were shown in various studies only for determinants that had a narrow content and temporal relation to the scales of the HRQoL instruments. This applies in particular to some values of the pulmonary function testing (spirometry) and to the presence of psychiatric symptoms.
- Future research should focus on methodologically adequate investigation of determinants that enable the identification of risk groups or concrete preventive measures in terms of intervention options during the acute treatment phase.

10. Briegel I, Dolch M, Irlbeck M, et al.: [Quality of results of therapy of acute respiratory failure: changes over a period of two decades]. *Anaesthesist* 2013; 62: 261–70.
11. Linden VB, Lidegran MK, Frisen G, et al.: ECMO in ARDS: a long-term follow-up study regarding pulmonary morphology and function and health-related quality of life. *Acta Anaesthesiol Scand* 2009; 53: 489–95.
12. McHugh LG, Milberg JA, Whitcomb ME, et al.: Recovery of function in survivors of the acute respiratory distress syndrome. *Am J Respir Crit Care Med* 1994; 150: 90–4.
13. Weinert CR, Gross CR, Kangas JR, et al.: Health-related quality of life after acute lung injury. *Am J Respir Crit Care Med* 1997; 156: 1120–8.
14. Masclans JR, Roca O, Munoz X, et al.: Quality of life, pulmonary function, and tomographic scan abnormalities after ARDS. *Chest* 2011; 139: 1340–6.
15. Kim SJ, Oh BJ, Lee JS, et al.: Recovery from lung injury in survivors of acute respiratory distress syndrome: difference between pulmonary and extrapulmonary subtypes. *Intensive Care Med* 2004; 30: 1960–3.
16. Li TS, Gomersall CD, Joynt GM, et al.: Long-term outcome of acute respiratory distress syndrome caused by severe acute respiratory syndrome (SARS): an observational study. *Crit Care Resusc* 2006; 8: 302–8.
17. Schelling G, Stoll C, Haller M, et al.: Health-related quality of life and posttraumatic stress disorder in survivors of the acute respiratory distress syndrome. *Crit Care Med* 1998; 26: 651–9.
18. Angus DC, Musthafa AA, Clermont G, et al.: Quality-adjusted survival in the first year after the acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2001; 163: 1389–94.
19. Schelling G, Stoll C, Vogelmeier C, et al.: Pulmonary function and health-related quality of life in a sample of long-term survivors of the acute respiratory distress syndrome. *Intensive Care Med* 2000; 26: 1304–11.
20. Gilson BS, Gilson JS, Bergner M, et al.: The sickness impact profile. Development of an outcome measure of health care. *Am J Public Health* 1975; 65: 1304–10.
21. Davidson TA, Caldwell ES, Curtis JR, et al.: Reduced quality of life in survivors of acute respiratory distress syndrome compared with critically ill control patients. *JAMA* 1999; 281: 354–60.
22. Hopkins RO, Weaver LK, Chan KJ, et al.: Quality of life, emotional, and cognitive function following acute respiratory distress syndrome. *J Int Neuropsychol Soc* 2004; 10: 1005–17.

23. Schmidt M, Zogheib E, Roze H, et al.: The PRESERVE mortality risk score and analysis of long-term outcomes after extracorporeal membrane oxygenation for severe acute respiratory distress syndrome. *Intensive Care Med* 2013; 39: 1704–13.
24. Luyt CE, Combes A, Becquemin MH, et al.: Long-term outcomes of pandemic 2009 influenza A(H1N1)-associated severe ARDS. *Chest* 2012; 142: 583–92.
25. Adhikari NK, McAndrews MP, Tansey CM, et al.: Self-reported symptoms of depression and memory dysfunction in survivors of ARDS. *Chest* 2009; 135: 678–87.
26. Adhikari NK, Tansey CM, McAndrews MP, et al.: Self-reported depressive symptoms and memory complaints in survivors five years after ARDS. *Chest* 2011; 140: 1484–93.
27. Kapfhammer HP, Rothenhausler HB, Krauseneck T, et al.: Posttraumatic stress disorder and health-related quality of life in long-term survivors of acute respiratory distress syndrome. *Am J Psychiatry* 2004; 161: 45–52.
28. Stevenson JE, Colantuoni E, Bienvenu OJ, et al.: General anxiety symptoms after acute lung injury: predictors and correlates. *J Psychosom Res* 2013; 75: 287–93.
29. Rothenhausler HB, Ehrentraut S, Stoll C, et al.: The relationship between cognitive performance and employment and health status in long-term survivors of the acute respiratory distress syndrome: results of an exploratory study. *Gen Hosp Psychiatry* 2001; 23: 90–6.
30. Mikkelsen ME, Shull WH, Biester RC, et al.: Cognitive, mood and quality of life impairments in a select population of ARDS survivors. *Respirology* 2009; 14: 76–82.
31. Stoll C, Haller M, Briegel J, et al.: Health-related quality of life. Long-term survival in patients with ARDS following extracorporeal membrane oxygenation (ECMO). *Anaesthesist* 1998; 47: 24–9.
32. Davydow DS, Desai SV, Needham DM, et al.: Psychiatric morbidity in survivors of the acute respiratory distress syndrome: a systematic review. *Psychosom Med* 2008; 70: 512–9.
33. Schandl AR, Brattstrom OR, Svensson-Raskh A, et al.: Screening and treatment of problems after intensive care: a descriptive study of multidisciplinary follow-up. *Intensive Crit Care Nurs* 2011; 27: 94–101.
34. Needham DM, Davidson J, Cohen H, et al.: Improving long-term outcomes after discharge from intensive care unit: report from a stakeholders' conference. *Crit Care Med* 2012; 40: 502–9.
35. Lasiter S, Oles SK, Mundell J, et al.: Critical care follow-up clinics: A scoping review of interventions and outcomes. *Clin Nurse Spec* 2016; 30: 227–37.
36. Jones C, Backman C, Capuzzo M, et al.: Intensive care diaries reduce new onset post traumatic stress disorder following critical illness: a randomised, controlled trial. *Crit Care* 2010; 14: R168.
37. Garrouste-Orgeas M, Coquet I, Perier A, et al.: Impact of an intensive care unit diary on psychological distress in patients and relatives. *Crit Care Med* 2012; 40: 2033–40.
38. Castro-Avila AC, Seron P, Fan E, et al.: Effect of early rehabilitation during intensive care unit stay on functional status: Systematic review and meta-analysis. *PLoS One* 2015; 10: e0130722.
39. Denehy L, Skinner EH, Edbrooke L, et al.: Exercise rehabilitation for patients with critical illness: a randomized controlled trial with 12 months of follow-up. *Crit Care* 2013; 17: R156.
40. Wilcox ME, Patsios D, Murphy G, et al.: Radiologic outcomes at 5 years after severe ARDS. *Chest* 2013; 143: 920–6.

Corresponding author

Dr. phil. Dipl.-Psych. Frank Dodoo-Schittko
 Department for Epidemiology and Preventive Medicine
 Medical Sociology
 Universität Regensburg
 Dr.-Gessler-Str. 17
 93051 Regensburg, Germany
 frank.dodoo-schittko@klinik.uni-r.de

Supplementary material:

For eReferences please refer to:
www.aerzteblatt-international.de/ref0717

eBox, eTables:

www.aerzteblatt-international.de/17m103

Erratum

The article entitled “Regional Differences in the Prevalence of Cardiovascular Disease—Results from the German Health Update (GEDA) from 2009–2012” by Dornquast et al in issue 42 of *Deutsches Ärzteblatt* (21 October 2016) contains two numerical errors. In the methods section, the authors reported the response rates of the included GEDA studies. The values of 23.9% and 34.5% for GEDA 2012 and 2009 are incorrect. The correct proportions are 22.1% (GEDA 2012) and 29.1% (GEDA 2009).

MWR

Supplementary material to:

Determinants of Quality of Life and Return to Work Following Acute Respiratory Distress Syndrome
A Systematic Review

by Frank Dodoo-Schittko, Susanne Brandstetter, Sebastian Blecha, Kathrin Thomann-Hackner,
Magdalena Brandl, Helge Knüttel, Thomas Bein, and Christian Apfelbacher

Dtsch Arztebl Int 2017; 114: 103–9. DOI: 10.3238/arztebl.2017.0103

eREFERENCES

- e1. Moher D, Liberati A, Tetzlaff J, et al.: Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg* 2010; 8: 336–41.
- e2. Brandstetter S, Dodoo-Schittko F, Blecha S, et al.: Determinants of health-related quality of life and return to work in survivors of acute respiratory distress syndrome (ARDS): a protocol for a systematic review. PROSPERO 2014:CRD420140143352014. Available from: www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42014014335.
- e3. McGowan J, Sampson M, Salzwedel DM, et al.: PRESS Peer Review of Electronic Search Strategies: 2015 Guideline Statement. *J Clin Epidemiol* 2016; 75: 40–6.
- e4. Atkinson KM, Koenka AC, Sanchez CE, et al.: Reporting standards for literature searches and report inclusion criteria: making research syntheses more transparent and easy to replicate. *Res Synth Methods* 2015; 6: 87–95.
- e5. Wells GSB, O'Connell D, Peterson J, et al.: The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. www.ohri.ca/programs/clinical_epidemiology/oxford.asp (last accessed on 9/2016).

Instrument for assessing the risk of selection and information bias of the studies included in a systematic review

Selection bias

Description of the study population:

Sufficient specifications about the inclusion and exclusion criteria. This includes:

1. Clear definition of acute respiratory distress syndrome (ARDS): Berlin Definition or American-European Consensus Conference (AECC) definition; alternatively, criteria used for progression, oxygenation, and imaging should be specified;
2. Further inclusion and exclusion criteria, or a statement that clarifies that no further inclusion/exclusion criteria were considered (for instance: "The study included all patients which fulfilled the diagnostic criteria for ARDS according to the Berlin definition."). „Inclusion and/or exclusion criteria were used for patient recruitment“ should be chosen only if the conditions in 1. and 2. were met.

Categories:

- Inclusion and/or exclusion criteria were used for patient recruitment
- No inclusion and/or exclusion criteria were used for patient recruitment
- Unclear

Sampling method

Did all patients in the population have the same chance of entering the cohort/sample, or were the examined clusters (e.g. ICUs) randomly selected and was there a complete enumeration? These issues always refer to the (first) time of inclusion of patients, which can be the time of ICU admission for a prospective study, or the inclusion of survivors for a longitudinal study. For each case, it must be determined whether sampling was random at that time.

For example, if inclusion started at ARDS diagnosis, yet the statements are only given for follow-up of survivors from this sample at 6 months, the category „unclear“ must be selected. The same applies to the absence of specifications about the total population for which the studied cohort/sample should be representative.

Categories:

- The entire cohort/sample was drawn randomly from the population of interest
- The entire cohort/sample was not drawn randomly from the population of interest
- Unclear

Loss to follow-up in the cohort

In the case of a longitudinal study with several (>2) follow-ups, assessment of adequacy should only consider and evaluate the last follow-up. Furthermore, deceased patients should not be considered lost to follow-up. Thus, the number of deaths must be deducted from the total loss to follow-up. If this number is not documented, the 80%-rule still applies.

Categories:

- No follow-up
- Complete follow-up (for all recorded people)
- Loss to follow-up unlikely to cause bias (follow-up rate \geq 80%)
- Follow-up rate <80%
- No statements about loss to follow-up

Information bias

Classification of exposure

Categories:

- Secure records (e.g. records related to medical treatment, imaging, laboratory findings). This includes psychiatric diagnoses, which are performed by a doctor
- Validated psychometric instruments: all instruments with satisfactory objectivity, reliability, and validity, or instruments which were satisfactorily evaluated on the basis of probabilistic test theory
- Structured interview: systematic interview (questions and their sequence are fixed, with a mostly closed response format)
- Written self-reports: any unsystematic written documentation

Classification of outcomes

Categories:

- Secure records (e.g. records related to medical treatment, imaging, laboratory findings). This includes psychiatric diagnoses performed by a doctor
- Validated psychometric instruments: all instruments with satisfactory objectivity, reliability, and validity, or instruments which were satisfactorily evaluated on the basis of probabilistic test theory
- Structured interview: systematic interview (questions and their sequence are fixed, with a mostly closed response format)
- Written self-reports: any unsystematic written documentation

eTABLE 1

Determinants of health-related quality of life (HRQoL) described in the literature. Studies marked with an * show a significant association of a determinant from the corresponding. Effect sizes are given when they were either reported or could be calculated from the reported data.

Determinants of HRQoL	Study (Reference)	Effect size
Disease-related		
General morbidity measures		
APACHE II	Angus et al. (2001) (18) Schelling et al. (2000) (19)	–
Charlson–Deyo Comorbidity Score	Angus et al. (2001) (18)	–
Karnofsky Performance Status Scale	Weinert et al. (1997) (13)	SF-36 PCS: $\rho = -0.75$ SF-36 MCS: $\rho = -0.61$
	Angus et al. (2001)* (18)	QWB: $\rho = -0.27$
6-min walk test	Masclan et al. (2011) (14)	–
	Herridge et al. (2011)* (7)	–
	Li et al. (2006)* (16)	SF-36: $\rho = 0.52$ to $\rho = 0.75$
ARDS etiology		
	Angus et al. (2001) (18)	–
	Kim et al. (2004) (15)	–
	Masclans et al. (2011) (14)	–
	Davidson et al. (1999)* (21)	SF-36: Cohen's $d = 0.27$ to $d = 0.65$ SGRQ: Cohen's $d = 0.65$ (patients with sepsis had worse mean scores in all domains of SF-36 and SGRQ)
	Schelling et al. (2000) (19)	–
Extent of lung damage		
	Wilcox et al. (2013) (40)	SF-36-PCS: $\rho = -0.10$ SF-36-MCS: $\rho = -0.31$
	McHugh et al. (1994)* (12)	–
	Masclans et al. (2011) (14)	–
	Schelling et al. (2000)* (19)	–
Results of lung function testing		
Diffusion capacity for CO		
	Kim et al. (2004) (15)	SQLI: $\rho = -0.02$
	McHugh et al. (1994) (12)	SIP: $\rho = -0.44$ to $\rho = -0.46$
	Li et al. (2006)* (16)	SF-36: $\rho = 0.18$ to $\rho = 0.53$
	Schelling et al. (2000) (19)	–
FEV1 (forced expiratory volume in 1 second)		
	Masclans et al. (2011)* (14)	NHP: $\rho = -0.36$
	Kim et al. (2004) (15)	SQLI: $\rho = 0.16$
	Li et al. (2006)* (16)	SF-36: $\rho = 0.19$ to $\rho = 0.46$
FVC (forced vital capacity)		
	Masclans et al. (2011)* (14)	NHP: $\rho = -0.36$
	Kim et al. (2004) (15)	SQLI: $\rho = -0.004$
	McHugh et al. (1994) (12)	–
	Li et al. (2006)* (16)	SF-36: $\rho = 0.32$ to $\rho = 0.59$
	Schelling et al. (2000) (19)	–
FEV1/FVC		
	Schelling et al. (2000) (19)	–
Total lung capacity (TLC)		
	Schelling et al. (2000) (19)	–
	Kim et al. (2004) (15)	SQLI: $\rho = 0.08$
	Li et al. (2006)* (16)	SF-36 PF: $\rho = 0.14$ to $\rho = 0.36$
Residual lung volume		
	Li et al. (2006) (16)	SF-36: $\rho = -0.21$ to $\rho = 0.04$
CO transfer coefficient		
	Li et al. (2006) (16)	SF-36: $\rho = -0.23$ to $\rho = 0.1$

Determinants of HRQoL	Study (Reference)	Effect size
Maximum oxygen uptake	Kim et al. (2004)* (15)	SQLI: $\rho = 0.44$
Care-related		
Duration of intubation/ventilation	Hopkins et al. (2004)* (22)	SF-36 PCS: $\rho = -0.36$ SF-36 MCS: $\rho = -0.41$
	Schelling et al. (2000)* (19)	SF-36 PCS: $\rho = -0.44$ SF-36 MCS: $\rho = 0.13$
	Davidson et al. (1999)* (21)	–
	Kim et al. (2004) (15)	SQLI: $\rho = 0.01$
	Weinert et al. (1997) (13)	SF-36 PCS: $\rho = -0.30$ SF-36 MCS: $\rho = -0.25$
Duration of hospitalization	Hopkins et al. (2004)* (22)	SF-36 PF: $\rho = -0.35$ SF-36 PRF: $\rho = -0.45$
	Davidson et al. (1999)* (21)	SF-36 PF: Cohen's $d = 0.71$
Duration of ICU treatment	Hopkins et al. (2004)* (22)	SF-36 PF: $\rho = -0.34$ SF-36 PRF: $\rho = -0.43$
	Kim et al. (2004) (15)	SQLI: $\rho = -0.03$
Duration of ECMO treatment	Schmidt et al. (2013)* (23)	–
ECMO	Briegel et al. (2013) (10)	–
	Stoll et al. (1998)* (31)	–
	Luyt et al. (2012) (24)	SF-36-PCS: Cohen's $d = 0.25$ SF-36-MCS: Cohen's $d = 0.06$ (both scores showed better means for patients treated with ECMO)
Prone positioning	Masclans et al. (2011) (14)	–
Nitric oxide (NO) inhalation	Masclans et al. (2011) (14)	–
Administration of steroids	Masclans et al. (2011) (14)	–
Use of renal replacement therapy	Masclans et al. (2011) (14)	–
Psychosocial		
Depression	Adhikari et al. (2011)* (26)	SF-36 MCS: $\rho = -0.68$ SF-36 MH: $\rho = -0.82$ SF-36 EF: $\rho = -0.64$
	Hopkins et al. (2004)* (22)	SF-36: $\rho = -0.29$ to $\rho = -0.76$
	Weinert et al. (1997) (13)	SF-36 PCS: $\rho = -0.17$ SF-36 MCS: $\rho = -0.94$
Post-traumatic stress disorder (PTSD)	Deja et al. (2006)* (6)	–
	Briegel et al. (2013)* (10)	SF-36 PCS: Cohen's $d = 0.64$ SF-36 MCS: Cohen's $d = 0.76$ (survivors with PTSD had worse mean scores)
	Kapfhammer et al. (2004)* (27)	–
	Schelling et al. (1998)* (17)	–
Anxiety disorder	Hopkins et al. (2004)* (22)	SF-36: $\rho = -0.30$ to -0.59
	Stevenson et al. (2013)* (28)	SF-36 PCS: $\rho = -0.18$ SF-36 MCS: $\rho = -0.73$ EQ-5D VAS: $\rho = -0.34$
Cognition	Rothenhausler et al. (2001)* (29)	–
	Mikkelsen et al. (2009)* (30)	–

Determinants of HRQoL	Study (Reference)	Effect size
	Hopkins et al. (2004)* (22)	SF-36: Cohen's $d < 0.01$ to $d = 0.45$ (survivors with cognitive deficits had worse mean scores in all SF-36 domains)
Sociodemographic		
Age	Angus et al. (2001) (18)	–
	Weinert et al. (1997) (13)	SF-36 PCS: $\rho = -0.52$ SF-36 MCS: $\rho = 0.07$
	Linden et al. (2009)* (11)	–
	Herridge et al. (2011)* (7)	–
	McHugh et al. (1994)* (12)	–
Sex	Angus et al. (2001) (18)	–
	McHugh et al. (1994) (12)	–
Other		
Time elapsed since disease onset / follow-up period	Weinert et al. (1997) (13)	SF-36 PCS: $\rho = 0.29$ SF-36 MCS: $\rho = 0.37$
	Briegel et al. (2013) (10)	SF-36: Cohen's $d = 0.04$ to $d = 0.38$ (with the exception of general health perceptions, all SF-36 domains showed better values for patients in the decade 1995–2005 as compared to the decade 1985–1994).
	Hopkins et al. (2004)* (22)	–
	Masclans et al. (2011) (14)	NHP: $\rho = 0.68$
	Adhikiri et al. (2009) (25)	–
	Schmidt et al. (2013) (23)	–

APACHE, Acute Physiology And Chronic Health Evaluation; ARDS, acute respiratory distress syndrome; ECMO, extracorporeal membrane oxygenation; EF, emotional role functioning; EQ-5D VAS, EuroQol Five Dimensions Questionnaire visual analog scale; NHP, Nottingham Health Profile; MSC, mental health component summary score; MH, mental health; PF, physical functioning; PRF, physical role functioning; PSC, physical health component summary score; QWB, Quality of well-being; SF-36, 36-Item Short-Form Health Survey; SGRQ, St George's Respiratory Questionnaire; SIP, Sickness Impact Profile; SQLI, Spitzer's Quality of Life Index

eTABLE 2

Determinants of return to work (RtW) described in the literature. Studies marked with an * show a significant association between a determinant from the corresponding category with the respective outcome. Effect sizes are given when they were either reported or could be calculated from the reported data.

Determinants of RtW	Study	Effect size
Disease-related		
Extent of lung damage	McHugh et al. (1994) (12)	–
Results of lung function testing		–
Diffusion capacity for CO	McHugh et al. (1994) (12)	–
FVC (forced vital capacity)	McHugh et al. (1994) (12)	–
Care-related		
Duration of intubation/ventilation	Briegel et al. (2013) (10)	–
Duration of ICU treatment	Briegel et al. (2013) (10)	–
ECMO	Briegel et al. (2013) (10)	–
	Luyt et al. (2012) (24)	–
	Stoll et al. (1998) (31)	–
	Schelling et al. (1998) (17)	OR = 2.0 Long-term incapacity for work Category: ECMO Reference category: non-ECMO
Psychosocial		
Depression	Adhikari et al. (2009)* (25)	OR = 0.20 Return to work Category: moderate to severe depressive symptoms; reference category: minimal to mild depressive symptoms
Sociodemographic		
Age	Briegel et al. (2013) (10)	–
	McHugh et al. (1994) (12)	–
Sex	Briegel et al. (2013) (10)	–
Other		
Time between disease / follow-up period	Briegel et al. (2013) (10)	–

ECMO, extracorporeal membrane oxygenation; OR, odds ratio