

Review Article

The Effects of Modified Simiao Decoction in the Treatment of Gouty Arthritis: A Systematic Review and Meta-Analysis

Ya-Fei Liu,¹ Ying Huang,² Cai-Yu-Zhu Wen,³ Jun-Jun Zhang,¹ Guo-Lan Xing,¹ Sheng-Hao Tu,² and Zhe Chen²

¹Department of Nephrology, The First Affiliated Hospital of Zhengzhou University, 1 Jianshe East Road, Zhengzhou, Henan 450052, China

²Institute of Integrated Traditional Chinese and Western Medicine, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, 1095 Jiefang Avenue, Wuhan, Hubei 430030, China

³Hubei University of Chinese Medicine, 1 Huangjiahu West Road, Wuhan, Hubei 430065, China

Correspondence should be addressed to Ya-Fei Liu; yafeiliutjh@gmail.com and Sheng-Hao Tu; shtu@tjh.tjmu.edu.cn

Received 23 October 2016; Revised 31 January 2017; Accepted 2 February 2017; Published 8 March 2017

Academic Editor: Caigan Du

Copyright © 2017 Ya-Fei Liu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The modified Simiao decoctions (MSD) have been widely applied in the treatment of gouty arthritis in China. However, the evidence needs to be evaluated by a systematic review and meta-analysis. After filtering, twenty-four randomised, controlled trials (RCTs) comparing the effects of MSD and anti-inflammation medications and/or urate-lowering therapies in patients with gouty arthritis were included. In comparison with anti-inflammation medications, urate-lowering therapies, or coadministration of anti-inflammation medications and urate-lowering therapies, MSD monotherapy significantly lowered serum uric acid ($p < 0.00001$, mean difference = -90.62 , and 95% CI [$-128.38, -52.86$]; $p < 0.00001$, mean difference = -91.43 , and 95% CI [$-122.38, -60.49$]; $p = 0.02$, mean difference = -40.30 , and 95% CI [$-74.24, -6.36$], resp.). Compared with anti-inflammation medications and/or urate-lowering therapies, MSD monotherapy significantly decreased ESR ($p < 0.00001$; mean difference = -8.11 ; 95% CI [$-12.53, -3.69$]) and CRP ($p = 0.03$; mean difference = -3.21 ; 95% CI [$-6.07, -0.36$]). Additionally, the adverse effects (AEs) of MSD were fewer ($p < 0.00001$; OR = 0.08 ; 95% CI [$0.05, 0.16$]). MSD are effective in the treatment of gouty arthritis through anti-inflammation and lowering urate. However, the efficacy of MSD should be estimated with more RCTs.

1. Introduction

Gouty arthritis, one of the most common forms of inflammatory arthritis, is characterized by hyperuricemia and deposition of monosodium urate. The prevalence of gout among US men in 2007-2008 was 5.9%, and the prevalence among women was 2.0% [1]. Global epidemiology of gout indicated that the distribution of gout was uneven across the world, with prevalence being highest in Pacific countries. Developed countries incline to have a higher load of gout than developing countries, and the prevalence and incidence seem to be increasing [2].

Conventional therapies for gouty arthritis include anti-inflammation medications (colchicine, nonsteroidal anti-inflammatory drugs, and glucocorticoids) and urate-lowering therapies (allopurinol, benzbromarone, and febuxostat).

However, allopurinol could lead to severe hypersensitivity and was limited in patients with renal insufficiency [3, 4]. The Food and Drug Administration implemented enforcement action against companies illegally marketing unapproved single-ingredient oral colchicine due to its toxicity [5]. Meanwhile, the relatively high medical care cost of febuxostat restricts its application in the developing countries. Therefore, it is imperative to explore new available approaches for gouty arthritis, especially complementary and alternative medicine.

Plant-based medicines are widely employed in the treatment of gouty arthritis in China for thousands of years. In Chinese medicine, gouty arthritis is correlated with dampness, heat, sputum, and stasis. Among numerous effective prescriptions, Simiao pill, derived from Ermiao powder,

and described in a famous traditional Chinese medicine monograph Chengfang Biandu in Qing Dynasty of China, was widely applied for treatment of gouty arthritis. It is composed of four individual herbs: *Rhizoma Atractylodis*, *Cortex Phellodendri*, *Radix Achyranthis Bidentatae*, and *Semen Coicis*. Moreover, Simiao pill and its derivative prescriptions showed their beneficial efficacy in treating gouty arthritis and hyperuricemia in vitro and in vivo [6–9]. To cope with the intricate pathologic states of gouty arthritis in different stages, different modified Simiao decoctions (MSD) have been developed based on different syndromes and traditional Chinese medicine theory. Our previous studies also demonstrated that MSD and their major component berberine were of use in attenuating the monosodium urate crystals-induced inflammation [9, 10].

While MSD have been most frequently used for a long time in treating gouty arthritis, there exist a series of issues. In this regard, most of the clinical researches have arisen from uncontrolled clinical studies or from retrospective reports, and few multicentre clinical trials have been conducted to validate the effects of MSD in the treatment of gouty arthritis. In addition, the scientific evidence that MSD are as effective as other conventional treatments in treating gouty arthritis remains to be confirmed further. In regard to safety concerns, the safety of long-term MSD intake for gouty arthritis is uncertain. Given these uncertainties, it is essential to evaluate the pertinent studies to systematically review the potential effects and safety of the long-term intake of MSD in the treatment of gouty arthritis.

2. Materials and Methods

To confirm the accuracy of our systemic review and meta-analysis, we designed and performed our results applying a checklist of items that were as consistent as possible with the Preferred Reporting Items for Systemic Review and Meta-Analyses (PRISMA) statement [11].

2.1. Search Strategy. We searched the following foreign databases to identify trials: PubMed, the Cochrane Library, and Clinical Trials.gov. Meanwhile, we retrieved Chinese databases, such as the CNKI Database, WanFang Database, and Chinese Clinical Trial Register. All of the databases were searched from their available dates of inception to the latest issue (September 2016).

Different search strategies were combined as follows. For the English databases, we used free text terms, such as “simiao” and “gouty arthritis” or “gout”. For the Chinese databases, free text terms were applied, such as “simiao” and “Tong Feng” (which means gout in Chinese). A filter for clinical studies was applied. To collect sufficient trials, the reference lists of relevant articles were also searched to identify additional studies.

2.2. Selection Criteria. The randomised controlled trials (RCTs) were included regardless of blinding, publication status, or language. Studies were included for analysis if they satisfied the following criteria: (1) For the types of

interventions, treatments with MSD alone in RCTs were considered. The control groups consisted of treatments with anti-inflammation medications and/or urate-lowering therapies. (2) The study was an RCT with a parallel or crossover design. And (3) Patients were enrolled who were diagnosed as having gouty arthritis, according to the classified criteria for gouty arthritis by American Society of Rheumatism in 1997 or Chinese diagnostic criteria for gouty arthritis.

In this review, studies using any MSD combined with western medicine or drugs for external use or acupuncture were excluded. We also excluded case reports, reviews, retrospective studies, or studies without scheduled outcomes. For obviously repeated studies, the authors of the studies were contacted to clarify any ambiguities. If the authors could not be connected, the first published study was considered to be the original. Studies were also excluded if the control groups were not conventional therapies for gouty arthritis. RCTs were also eliminated from our analysis that lacked sufficient data to allow for calculating the mean changes from the baseline to the endpoints. Two reviewers selected the articles independently. We generated a flow diagram of the study selection in accordance with the PRISMA requirements.

2.3. Data Extraction and Management. The data were extracted by two independent reviewers, and any divergences were resolved by consensus or were arbitrated by a third reviewer. The studies' quality was assessed according to Cochrane handbook 5.3. The risk of bias included the following items: (A) random sequence generation (selection bias); (B) allocation concealment (selection bias); (C) blinding of participants and personnel (performance bias); (D) blinding of outcome assessment (detection bias); (E) incomplete outcome data (attrition bias); (F) selective reporting (reporting bias); (G) other bias.

The primary outcome was serum uric acid (SUA). The secondary outcomes consisted of erythrocyte sedimentation rate (ESR), C-reactive protein (CRP), and white blood cell (WBC). AEs were also collected from the studies. For the trials that applied a three-armed group design, the outcomes of the groups were extracted if they met the inclusion criteria. In cases in which the outcomes were vague or absent in the articles, we endeavoured to contact the authors. If the authors were not connected, we extracted the data by consensus.

2.4. Data Synthesis and Analysis. The effects of MSD intake on patients with gouty arthritis were calculated as differences between the MSD groups and the control groups, employing Review Manager meta-analysis software, version 5.3. To ensure the credibility of the results, the net changes in all of the outcomes were calculated as the mean differences (MSD minus control) in changes (endpoint minus baseline) for parallel trials. We calculated weighted mean differences and 95% confidence intervals (CIs) for continuous data. Heterogeneity was evaluated via the chi-square test and Higgins I^2 test. A fixed-effect model was employed when the studies in the group were sufficiently alike ($p > 0.10$); otherwise, a random-effects model was applied. A Z score was calculated to determine the overall effect, with significance set at $p <$




FIGURE 1: Study selection flow chart.

0.05. Publication bias was detected by Egger's regression asymmetry test and Begg's test when the number of included trials ≥ 5 (Stata software, version 12.0).

To minimise the clinical heterogeneity, in terms of SUA, we performed three subgroup analyses: MSD compared with anti-inflammation medications; MSD compared with urate-lowering therapies; and cointervention of anti-inflammation medications and urate-lowering therapies compared with MSD alone.

3. Results

3.1. Study Selection. The process of study selection was indicated in Figure 1. According to the prespecified selection

criteria defined in the Materials and Methods, 24 RCTs were included in the meta-analysis. In the anti-inflammation medications subgroup, we searched ten studies [13–16, 18, 20, 22, 24, 26, 33]. One study compared oral MSD and/or external use with colchicine and allopurinol, and oral MSD group was selected in comparison with colchicine and allopurinol [31]. The trial of Shi et al. compared three different MSD with indomethacin and benzbromarone, and we just extracted the data of Group III and control group [6]. In the urate-lowering therapies group, we searched three studies [7, 17, 23]. In the combined therapy subgroup, we searched eleven studies [6, 12, 19, 21, 25, 27–32]. The characteristics of the studies were summarised in Table 1. Together, those studies included a total of 1895 participants.

TABLE 1: Clinical and demographic characteristics of the patients with gouty arthritis.

Study (ref.)	Number of participants		Age (years)		Experimental	Control	Intervention	Duration (days)	Outcomes
	Experimental	Control	Experimental	Control					
Chen 2011 [12]	44	44	48.5 ± 9.26	49.01 ± 8.95	MSD		Colchicine; allopurinol	14	SUA; ESR; AEs
Fan and Weng 2016 [13]	30	30	48.0 ± 3.8	46.0 ± 3.6	MSD		Meloxicam	28	SUA; ESR; CRP; AEs
Fang 2008 [14]	20	20	57.35 ± 14.54	53.05 ± 14.47	MSD		Loxoprofen sodium	5	SUA; ESR; AEs
Gao et al. 2014 [15]	35	35	44.2 ± 7.2	39.4 ± 8.1	MSD		Colchicine; Etocoxib	7	SUA; ESR; CRP; AEs
Jia 2010 [16]	28	26	43.0 ± 10.8	44.0 ± 11.3	MSD		Colchicine	14	SUA; AEs
Li and Zhang 2007 [17]	40	40	47.5 ± 3.17	46.8 ± 2.74	MSD		Allopurinol	14	SUA
Li and Song 2013 [18]	35	35	NR	NR	MSD		Diclofenac sodium	7	SUA; AEs
Liu 2011 [19]	30	30	NR	NR	MSD		Allopurinol; nimesulide	21	SUA; ESR; CRP; WBC
Luo and Tang 2010 [20]	30	30	NR	NR	MSD		Colchicine; diclofenac sodium	3	SUA
Mi 2016 [21]	44	42	46.12 ± 8.21	45.42 ± 7.85	MSD		Allopurinol; nimesulide	21	SUA; ESR; CRP
Niu 2008 [22]	52	47	NR	NR	MSD		Colchicine	7	SUA
Renbin et al. 2008 [7]	60	60	46.9 ± 3.37	46.8 ± 3.24	MSD		Allopurinol	30	SUA; CRP
Shi et al. 2008 [6]	28	25	54.0 ± 12.5	52.9 ± 13.2	MSD		Indomethacin; benzbromarone	14	SUA; WBC; AEs
Sun 2006 [23]	36	28	53.2 ± 7.8	54.0 ± 8.3	MSD		Benzbromarone	28	SUA; ESR
Tang et al. 2008 [24]	30	26	NR	NR	MSD		Colchicine	15	SUA; AEs
Wang et al. 2009 [25]	90	88	NR	NR	MSD		Colchicine; benzbromarone	14	SUA; ESR; WBC; AEs
Wang et al. 2016 [26]	30	30	36.1 ± 2.8	36.7 ± 2.5	MSD		Colchicine	14	SUA; ESR; CRP
Wang et al. 2016 [27]	32	32	NR	NR	MSD		Allopurinol; NSAIDs	10	SUA; ESR; CRP; WBC; AEs
Yu and Huang 2008 [28]	40	40	48.2 ± 8.6	46.0 ± 9.7	MSD		Nimesulide; allopurinol	14	SUA; AEs
Zeng and Song 2013 [29]	60	58	NR	NR	MSD		Colchicine; allopurinol	14	SUA
Zhang and Chen 2012 [30]	36	28	45.2 ± 7.8	43.4 ± 8.3	MSD		Colchicine; allopurinol; Voltaren or celecoxib	7	SUA; ESR; WBC; AEs
Zhao and Mai 2008 [31]	41	40	NR	NR	MSD		Colchicine; allopurinol	30	SUA; ESR; WBC; AEs
Zhao 2014 [32]	48	45	NR	NR	MSD		Colchicine; allopurinol; Voltaren	7	SUA; ESR
Zhu 2016 [33]	49	48	45.77 ± 7.08	45.45 ± 7.18	MSD		Colchicine	15	SUA

Note: MSD: modified Simiao decoction; SUA: serum uric acid; ESR: erythrocyte sedimentation rate; CRP: C-reactive protein; WBC: white blood cell; AEs: adverse effects; NR, no report. Values are mean ± standard deviation (SD).

3.2. Study Descriptions. The included RCTs were published as full text between 2006 and 2016. The duration of intervention in the included trials ranged from 3 days to 30 days. All of the trials were originated from China. Two studies were published in English [6, 7], while 22 studies were published in Chinese. All of the RCTs were conducted as single-centre trials. One trial was a master's degree thesis [14], and the others were journal articles. All of the trials were performed in mainland China. The important sources and compositions of MSD were indicated in Table 2.

3.3. Quality of the Included Studies. As indicated in Figure 3, most of the included trials were of low quality due to unclear randomisation, deficient allocation concealment, inadequate blinding, and undescribed withdrawals and dropouts, compared with two trials [6, 21] that were of moderate quality. Meanwhile, high risk of other bias could exist in three trials [22, 25, 27].

3.4. Publication Bias. Egger's publication bias plots and Begg's test displayed that there were significant publication biases for three outcomes in terms of SUA (when compared with anti-inflammation medications), ESR, and AEs. Meanwhile, there were no publication biases for three outcomes in terms of SUA (with cointervention of anti-inflammation medications and urate-lowering therapies), CRP, and WBC. As presented in Figure 2, the calculated p values exceeded 0.05 for the three outcomes among the studies (SUA, $p = 0.056$; CRP, $p = 0.771$; WBC, $p = 0.453$), and the 95% CI for the intercept included zero. However, these results cannot be regarded as convincing except SUA (with cointervention of anti-inflammation medications and urate-lowering therapies), because there were fewer than ten trials.

3.5. Effects of Interventions

3.5.1. Effects of MSD on SUA

(1) MSD Compared with Anti-Inflammation Medications. Ten trials (involving 666 patients) compared the therapeutic effects of MSD and anti-inflammation medications [13–16, 18, 20, 22, 24, 26, 33]. The number of trial participants ranged from 20 to 52, with the trial duration varying from 3 days to 28 days. As illustrated in Figure 3, there was statistical heterogeneity between the studies. The MSD groups were superior to the anti-inflammation medication groups in terms of lowering the SUA ($p < 0.00001$; mean difference = -90.62 ; 95% CI [$-128.38, -52.86$]).

(2) MSD Compared with Urate-Lowering Therapies. Three trials (involving 264 patients) compared the therapeutic effects of MSD and urate-lowering therapies [7, 17, 23]. The number of trial participants ranged from 28 to 60, with the trial duration varying from 14 days to 30 days. As illustrated in Figure 3, there was statistical heterogeneity between the studies. The MSD groups were superior to the urate-lowering therapies groups with regard to lowering the SUA ($p < 0.00001$; mean difference = -91.43 ; 95% CI [$-122.38, -60.49$]).

(3) MSD Compared with Combined Therapies. Eleven trials (involving 965 patients) compared the therapeutic effects of MSD and combined therapies [6, 12, 19, 21, 25, 27–32]. The number of trial participants ranged from 25 to 90, with the trial duration varying from 7 days to 30 days. As illustrated in Figure 3, there was statistical heterogeneity between the studies. The MSD groups were superior to the combined therapies groups regarding lowering the SUA ($p = 0.02$; mean difference = -40.30 ; 95% CI [$-74.24, -6.36$]).

3.5.2. Effects of MSD on ESR. ESR was reported in thirteen trials (involving 1008 patients) [12–15, 19, 21, 23, 25–27, 30–32]. The number of trial participants ranged from 20 to 90, with the trial duration varying from 5 days to 30 days. As illustrated in Figure 3, there was statistical heterogeneity between the studies. The MSD groups were superior to the control groups regarding decreasing the ESR ($p < 0.00001$; mean difference = -8.11 ; 95% CI [$-12.53, -3.69$]).

3.5.3. Effects of MSD on CRP. CRP was determined in seven trials (involving 520 patients) [7, 13, 15, 19, 21, 26, 27]. The number of trial participants ranged from 30 to 60, with the trial duration varying from 7 days to 28 days. As illustrated in Figure 3, there was statistical heterogeneity between the studies. The MSD groups were superior to the control groups regarding reducing the CRP ($p = 0.03$; mean difference = -3.21 ; 95% CI [$-6.07, -0.36$]).

3.5.4. Effects of MSD on WBC. WBC was detected in five trials (involving 322 patients) [6, 19, 27, 30, 31]. The number of trial participants ranged from 25 to 41, with the trial duration varying from 7 days to 30 days. As illustrated in Figure 3, there was statistical heterogeneity between the studies. There was no significant reduction in terms of WBC between MSD and control groups ($p = 0.28$; mean difference = -0.72 ; 95% CI [$-2.04, 0.59$]).

3.6. The AEs of MSD. AEs were reported in twelve trials (involving 111 patients) [6, 12–16, 18, 24, 25, 27, 28, 30]. The number of cases ranged from 0 to 18. As illustrated in Figure 3, there was no statistical heterogeneity between the studies. The MSD groups were superior to the control groups regarding the AEs ($p < 0.00001$; OR = 0.08; 95% CI [0.05, 0.16]).

4. Discussion

Although several systematic reviews and meta-analyses concerning the efficacy of MSD in the treatment of gouty arthritis have been performed, the primary outcome of these systematic reviews was clinical response rate which was judged by Chinese criteria [34–37]. Except for the clinical response rate, the meta-analysis published by Xie (8 trials with 633 participants) also narrated the chemical profiles, such as SUA, ESR, and CRP [37]. Meanwhile, Xie divided the studies into two subgroups (with or without urate-lowering therapies) when analysing the effects of MSD on SUA [37]. The two systematic reviews published by Du et al. only reported the clinical response rate and safety [35, 36]. Unlike the previous

TABLE 2: The important sources and compositions of MSD.

Studies	Components of MSD
Chen 2011 [12]	<i>Rhizoma Atractylodis Macrocephalae</i> , <i>Rhizoma Atractylodis</i> , <i>Radix Angelicae Sinensis</i> , <i>Radix Saposhnikoviae</i> , <i>Radix Glycyrrhizae</i> , <i>Radix Puerariae</i> , <i>Radix Achyranthis Bidentatae</i> , <i>Radix Sophorae Flavescens</i> , <i>Radix Scutellariae</i> , <i>Cortex Phellodendri</i> , <i>Radix Ginseng</i> , <i>Rhizoma Cini</i> , <i>Radix Cnicifugae</i> , <i>Semen Coicis</i> , <i>Rhizoma Et Radix Notopterygii</i> , <i>Virgate Wormwood Herb</i> , <i>Anemarrhena asphodeloides</i> Bunge, <i>Polyporus umbellatus</i> , <i>Alisma orientalis</i>
Fan and Weng 2016	<i>Caulis Lonicerae</i> , <i>Caulis Trachelospermi</i> , <i>Gymnema alternifolium</i> , <i>Caulis Sinomenii</i> , <i>Smilax bockii</i> Warb, <i>Spatholobus suberectus</i> , <i>Cortex Phellodendri</i> , <i>Radix Achyranthis Bidentatae</i> , <i>Semen Coicis</i> , <i>Rhizoma Atractylodis</i>
Fang 2008	<i>Cortex Phellodendri</i> , <i>Radix Achyranthis Bidentatae</i> , <i>Semen Coicis</i> , <i>Rhizoma Atractylodis</i>
Gao et al. 2014	<i>Cortex Phellodendri</i> , <i>Radix Achyranthis Bidentatae</i> , <i>Rhizoma Atractylodis</i> , <i>Pseudobulbus Cremastrae</i> Seu <i>Pleiones</i> , <i>Rhizoma Smilacis Glabrae</i> , <i>Rhizome Dioscoreae Septemlobae</i> , <i>Polistes mandarinus</i> Saussure, <i>Herba Lysimachiae</i> , <i>Sargassum</i> , <i>Radix Stephaniae Tetrandrae</i> , <i>Semen Coicis</i>
Jia 2010	<i>Cortex Phellodendri</i> , <i>Radix Achyranthis Bidentatae</i> , <i>Semen Coicis</i> , <i>Rhizoma Atractylodis</i> , <i>Radix Stephaniae Tetrandrae</i> , <i>Alisma orientalis</i> , <i>Radix Angelicae Pubescentis</i> , <i>Radix Saposhnikoviae</i> , <i>Pheretima</i>
Li and Zhang 2007	<i>Cortex Phellodendri</i> , <i>Radix Achyranthis Bidentatae</i> , <i>Semen Coicis</i> , <i>Rhizoma Atractylodis</i> , <i>Caulis Lonicerae</i> , <i>Rhizome Dioscoreae Septemlobae</i> , <i>Fructus Chaenomelis</i> , <i>Cortex Fraxini</i> , <i>Alisma orientalis</i> , <i>Radix Angelicae Sinensis</i>
Li and Song 2013	<i>Cortex Phellodendri</i> , <i>Radix Achyranthis Bidentatae</i> , <i>Semen Coicis</i> , <i>Rhizoma Atractylodis</i> , <i>Alisma orientalis</i> , <i>Pheretima</i> , <i>Rhizome Dioscoreae Septemlobae</i> , <i>Rhizoma Smilacis Glabrae</i> , <i>Semen Strychni</i>
Liu 2011	<i>Cortex Phellodendri</i> , <i>Radix Achyranthis Bidentatae</i> , <i>Semen Coicis</i> , <i>Rhizoma Atractylodis</i> , <i>Virgate Wormwood Herb</i> , <i>Rhizome Dioscoreae Septemlobae</i> , <i>Rhizoma Smilacis Glabrae</i> , <i>Wild Chrysanthemum</i> , <i>Radix Clematidis</i> , <i>Fructus Chaenomelis</i> , <i>Portia cocos</i> , <i>Radix Angelicae Pubescentis</i> , <i>Radix Glycyrrhizae</i>
Luo and Tang 2010	<i>Cortex Phellodendri</i> , <i>Radix Achyranthis Bidentatae</i> , <i>Semen Coicis</i> , <i>Rhizoma Atractylodis</i> , <i>Rhizome Dioscoreae Septemlobae</i> , <i>Plantain Seed</i> , <i>Herba Plantaginis</i> , <i>Caulis Lonicerae</i> , <i>Caulis Trachelospermi</i>
Mi 2016	<i>Rhizoma Smilacis Glabrae</i> , <i>Semen Coicis</i> , <i>Rhizome Dioscoreae Septemlobae</i> , <i>Spatholobus suberectus</i> , <i>Rhizoma Atractylodis</i> , <i>Cortex Phellodendri</i> , <i>Radix Achyranthis Bidentatae</i> , <i>Radix Polygoni Multiflori</i> , <i>Rhizoma Corydalis</i> , <i>Radix Clematidis</i> , <i>Alisma orientalis</i> , <i>Pheretima</i> , <i>Eupolyphaga</i> Seu <i>Steleophaga</i> , <i>Herba Leonuri</i> , <i>Virgate Wormwood Herb</i> , <i>Radix Glycyrrhizae</i>
Niu 2008	<i>Cortex Phellodendri</i> , <i>Radix Achyranthis Bidentatae</i> , <i>Semen Coicis</i> , <i>Rhizoma Atractylodis</i> , <i>Rhizoma Smilacis Glabrae</i> , <i>Rhizome Dioscoreae Septemlobae</i> , <i>Pheretima</i> , <i>Radix Stephaniae Tetrandrae</i> , <i>Fructus Chaenomelis</i> , <i>Flos Lonicerae</i>
Renbin et al. 2008	<i>Cortex Phellodendri</i> , <i>Radix Achyranthis Bidentatae</i> , <i>Semen Coicis</i> , <i>Rhizoma Atractylodis</i> , <i>Caulis Lonicerae</i> , <i>Rhizoma Smilacis Glabrae</i> , <i>Anemarrhena asphodeloides</i> Bunge, <i>Radix Paeoniae Rubra</i> , <i>Radix Clematidis</i> , <i>Rhizome Dioscoreae Septemlobae</i> , <i>Alisma orientalis</i> , <i>Zaocys Dhumades</i>
Shi et al. 2008	<i>Cortex Phellodendri</i> , <i>Radix Achyranthis Bidentatae</i> , <i>Semen Coicis</i> , <i>Rhizoma Atractylodis</i> , <i>Caulis Lonicerae</i> , <i>Rhizoma Smilacis Glabrae</i> , <i>Radix Paeoniae Rubra</i> , <i>Gypsum</i> , <i>Anemarrhena asphodeloides</i> Bunge, <i>Ranulus Cinnamomi</i>
Sun 2006	<i>Cortex Phellodendri</i> , <i>Radix Achyranthis Bidentatae</i> , <i>Rhizoma Atractylodis</i> , <i>Rhizoma Smilacis Glabrae</i> , <i>Herba Siegesbeckiae</i> , <i>Radix Gentianae Macrophyllae</i> , <i>Semen Coicis</i> , <i>Herba Lysimachiae</i> , <i>Stigma Maydis</i> , <i>Cortex Erythrinae</i>
Tang et al. 2008	<i>Cortex Phellodendri</i> , <i>Radix Achyranthis Bidentatae</i> , <i>Semen Coicis</i> , <i>Rhizoma Atractylodis</i> , <i>Gypsum</i> , <i>Anemarrhena asphodeloides</i> Bunge, <i>Cortex Moutan</i> , <i>Radix Paeoniae Rubra</i> , <i>Radix Et Rhizoma Rhei</i> , <i>Alisma orientalis</i> , <i>Fructus Chaenomelis</i> , <i>Radix Clematidis</i>

TABLE 2: Continued.

Studies	Components of MSD
Wang et al. 2009	Cortex <i>Phellodendri</i> , Radix <i>Achyranthis Bidentatae</i> , Semen <i>Coicis</i> , Rhizoma <i>Atractylodis</i> , Rhizoma <i>Smilacis Glabrae</i> , Herba <i>Plantaginis</i> , Herba <i>Lysimachiae</i> , Caulis <i>Lonicerae</i> , Radix <i>Et Rhizoma Rhei</i> , Rhizome <i>Dioscoreae Septemlobae</i> , Rhizoma <i>Polygoni Cuspidati</i> , Radix <i>Clematidis</i> , Radix <i>Salviae Miltiorrhizae</i> , Radix <i>Angelicae Sinensis</i>
Wang et al. 2016	Cortex <i>Phellodendri</i> , Radix <i>Achyranthis Bidentatae</i> , Semen <i>Coicis</i> , Rhizoma <i>Atractylodis</i> , Ramulus <i>Cinnamomi</i> , Rhizoma <i>Atractylodis Macrocephalae</i> , Poria <i>Cocos</i> , <i>Polyporus umbellatus</i> , <i>Alisma orientalis</i> , Reed Rhizome, Lalang Grass Rhizome, Semen <i>Persicae</i> , Radix <i>Glycyrrhizae</i>
T. Wang et al. 2016	Ramulus <i>Cinnamomi</i> , Radix <i>Cyathulae</i> , Radix <i>Paeoniae Rubra</i> , Fructus <i>Chaenomelis</i> , <i>Anemarrhena asphodeloides</i> Bunge, Rhizoma <i>Atractylodis</i> , Cortex <i>Phellodendri</i> , Caulis <i>Lonicerae</i> , Rhizoma <i>Smilacis Glabrae</i> , Radix <i>Clematidis</i> , Gypsum, Semen <i>Coicis</i> , Herba <i>Plantaginis</i> , Herba <i>Lysimachiae</i> , Radix <i>Glycyrrhizae</i>
Yu and Huang 2008	Cortex <i>Phellodendri</i> , Radix <i>Achyranthis Bidentatae</i> , Rhizoma <i>Atractylodis</i> , Radix <i>Gentianae Macrophyllae</i> , Radix <i>Paeoniae Rubra</i> , Semen <i>Coicis</i> , Caulis <i>Lonicerae</i> , Rhizoma <i>Smilacis Glabrae</i> , Herba <i>Taraxaci</i>
Zeng and Song 2013	Cortex <i>Phellodendri</i> , Radix <i>Achyranthis Bidentatae</i> , Semen <i>Coicis</i> , Rhizoma <i>Atractylodis</i> , Rhizoma <i>Smilacis Glabrae</i> , Rhizome <i>Dioscoreae Septemlobae</i> , Radix <i>Paeoniae Rubra</i> , Caulis <i>Lonicerae</i> , Bombyx <i>Batryticatus</i> , <i>Eupatorium adenophorum</i> , Radix <i>Glycyrrhizae</i>
Zhang and Chen 2012	Cortex <i>Phellodendri</i> , Radix <i>Achyranthis Bidentatae</i> , Semen <i>Coicis</i> , Rhizoma <i>Atractylodis</i> , Caulis <i>Lonicerae</i> , Rhizoma <i>Smilacis Glabrae</i> , Rhizome <i>Dioscoreae Septemlobae</i> , Plantain Seed, Herba <i>Lysimachiae</i> , <i>Pseudobulbus Cremastrae</i> Seu <i>Pleitones</i> , Radix <i>Gentianae Macrophyllae</i> , Fructus <i>Chaenomelis</i>
Zhao and Mai 2008	Cortex <i>Phellodendri</i> , Radix <i>Achyranthis Bidentatae</i> , Semen <i>Coicis</i> , Rhizoma <i>Atractylodis</i> , Rhizoma <i>Atractylodis Macrocephalae</i> , Radix <i>Glycyrrhizae</i>
Zhao 2014	Cortex <i>Phellodendri</i> , <i>Alisma orientalis</i> , Radix <i>Cyathulae</i> , Radix <i>Angelicae Sinensis</i> , Rhizoma <i>Atractylodis</i> , Rhizome <i>Dioscoreae Septemlobae</i> , Fructus <i>Chaenomelis</i> , Cortex <i>Fraxini</i> , Semen <i>Coicis</i> , Caulis <i>Lonicerae</i>
Zhu 2016	<i>Pheretima</i> , Radix <i>Saposhnikoviae</i> , Radix <i>Angelicae Pubescentis</i> , <i>Alisma orientalis</i> , Radix <i>Stephaniae Tetrandrae</i> , Semen <i>Coicis</i> , Cortex <i>Phellodendri</i> , Radix <i>Achyranthis Bidentatae</i> , Rhizoma <i>Atractylodis</i>

Note: MSD: modified Simiao decoction.





FIGURE 2: Publication bias in the included trials. Egger’s linear regression test for detecting publication bias. Note: MSD: modified Simiao decoction; SUA: serum uric acid; ESR: erythrocyte sedimentation rate; CRP: C-reactive protein; WBC: white blood cell; AEs: adverse effects. “O” is a size graph symbol for the weight of each included study. The distance between two diamonds on the second vertical bar on the left represents the 95% CI for the intercept.

reviews, we included 24 trials and set three subgroups when determining the effects of MSD on SUA. However, the clinical response rate was not detected because it was easily influenced by subjective factors. Furthermore, we added more new trials published after the previous reviews. Thus, our systemic review is different from the previous reviews.

In terms of SUA, our results were consistent with Xie and Zhou et al. [34, 37]. These showed that MSD were of use in lowering SUA and they could be applied in the treatment of hyperuricemia and reduce the incidence of gouty arthritis.

Many of our results were in line with the results of Xie [37] between the MSD-treated groups and the control groups in


SUA



Risk of bias

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

ESR



Risk of bias

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

FIGURE 3: Continued.

terms of ESR and CRP. These indicated that MSD played an important role in anti-inflammation and decreasing disease activity of gouty arthritis, which further validated the effects of MSD in the treatment of gouty arthritis.

Unlike the previous review by Xie [37], we conducted three subgroups when comparing the effects of MSD and control groups on SUA. In addition, the WBC was observed in our systemic review. The results demonstrated that MSD had no beneficial effects in decreasing WBC, which was due to the aseptic inflammation of gouty arthritis.

Our results illustrated that MSD surpassed control groups with regard to AEs, which were in harmony with Xie and Du et al. [35–37]. The most common AEs with MSD were diarrhea, nausea, and vomit, and they could be alleviated with or without dose reductions. These showed that MSD were safe in the treatment of gouty arthritis.

However, several limitations of this meta-analysis should be noted. First, all of the included trials were performed in Chinese populations, which implied high risk of selection bias. This fact could have influenced the applicability of MSD to populations of other countries. Second, most of the studies published in Chinese were of poor quality concerning their designs, reporting, and methodologies. Third, the heterogeneity between the trials included in each subgroup was also significant. We believe that differences in control groups, components of MSD, doses, and durations of treatment were responsible for the heterogeneity. Fourth, the control groups, including anti-inflammation medications and/or urate-lowering therapies, were different. Hence, it is unsuitable to compare the effects of MSD and the control groups. Taking these facts into account, we should carefully interpret all of the conclusions due to the substantial methodological and clinical variety of the trials.

5. Conclusions

In summary, MSD monotherapy is superior to anti-inflammation medications and/or urate-lowering therapies in the treatment of gouty arthritis. Meanwhile, the AEs of MSD were mild. Based on their bioactivity, MSD function as anti-inflammation as well as lowering uric acid. Considering the low methodological quality of the included trials, more large and well-designed RCTs are needed before we can recommend MSD to replace western medicine.

Competing Interests

The authors declare no conflict of interests.

Authors' Contributions

Ya-Fei Liu, Cai-Yu-Zhu Wen, Ying Huang, and Zhe Chen extracted the data and carried out the data analysis. Jun-Jun Zhang and Guo-Lan Xing resolved divergences and encouraged achieving consensus. Sheng-Hao Tu and Zhe Chen participated in the design of the study and helped to revise the paper. All authors read and approved the paper.

Acknowledgments

The authors acknowledge the authors of the original studies that they included in their meta-analysis.

References

- [1] Y. Zhu, B. J. Pandya, and H. K. Choi, "Prevalence of gout and hyperuricemia in the US general population: the National Health and Nutrition Examination Survey 2007-2008," *Arthritis and Rheumatism*, vol. 63, no. 10, pp. 3136–3141, 2011.
- [2] C.-F. Kuo, M. J. Grainge, W. Zhang, and M. Doherty, "Global epidemiology of gout: prevalence, incidence and risk factors," *Nature Reviews Rheumatology*, vol. 11, no. 11, pp. 649–662, 2015.
- [3] K. R. Hande, R. M. Noone, and W. J. Stone, "Severe allopurinol toxicity. Description and guidelines for prevention in patients with renal insufficiency," *The American Journal of Medicine*, vol. 76, no. 1, pp. 47–56, 1984.
- [4] S. N. Ramasamy, C. S. Korb-Wells, D. R. W. Kannangara et al., "Allopurinol hypersensitivity: a systematic review of all published cases, 1950–2012," *Drug Safety*, vol. 36, no. 10, pp. 953–980, 2013.
- [5] M. Giouroukakis and M. Dryer, "Unapproved drugs—the drug information pharmacists' perspective," *Journal of Pharmacy Practice*, vol. 26, no. 2, pp. 112–119, 2013.
- [6] X. D. Shi, G. C. Li, Z. X. Qian, Z. Q. Jin, and Y. Song, "Randomized and controlled clinical study of modified prescriptions of Simiao Pill in the treatment of acute gouty arthritis," *Chinese Journal of Integrative Medicine*, vol. 14, no. 1, pp. 17–22, 2008.
- [7] Q. Renbin, S. Ruizi, L. Dejiu, C. Yuanlin, and Y. Hongping, "Treatment of 60 cases of gouty arthritis with modified Simiao Tang," *Journal of Traditional Chinese Medicine*, vol. 28, no. 2, pp. 94–97, 2008.
- [8] J. Hua, P. Huang, C.-M. Zhu, X. Yuan, and C.-H. Yu, "Anti-hyperuricemic and nephroprotective effects of Modified Simiao Decoction in hyperuricemic mice," *Journal of Ethnopharmacology*, vol. 142, no. 1, pp. 248–252, 2012.
- [9] Y.-F. Liu, S.-H. Tu, Z. Chen, Y. Wang, Y.-H. Hu, and H. Dong, "Effects of modified Simiao decoction on IL-1 β and TNF α secretion in monocytic THP-1 cells with monosodium urate crystals-induced inflammation," *Evidence-Based Complementary and Alternative Medicine*, vol. 2014, Article ID 406816, 7 pages, 2014.
- [10] Y. Liu, C. Wen, Z. Chen, Y. Wang, Y. Huang, and S. Tu, "Effects of berberine on NLRP3 and IL-1 β expressions in monocytic THP-1 cells with monosodium urate crystals-induced inflammation," *BioMed Research International*, vol. 2016, Article ID 2503703, 7 pages, 2016.
- [11] D. Moher, A. Liberati, J. Tetzlaff, and D. G. Altman, "Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement," *Annals of Internal Medicine*, vol. 151, no. 4, pp. 264–269, 2009.
- [12] B. Chen, "The clinical observation of treating 44 cases of acute gouty arthritis in angelica Niantong decoction plus Simiao san," *Clinical Journal of Chinese Medicine*, vol. 3, no. 13, pp. 25–27, 2011.
- [13] L. F. Fan and G. M. Weng, "Clinical observation of Liuteng Simiao decoction on the treatment of acute gouty arthritis," *Hebei Journal of Traditional Chinese Medicine*, vol. 38, no. 2, pp. 185–187, 2016.

- [14] X. Y. Fang, *Clinical Observation of Modified Simiao Decoction in the Treatment of Acute Gouty Arthritis*, Chengdu University of Traditional Chinese Medicine, 2008.
- [15] H. Gao, Z. G. Li, and H. B. Du, "Clinical observation of modified Simiao decoction in the treatment of acute gouty arthritis," *Journal of Qiqihar University of Medicine*, vol. 35, no. 18, pp. 2664–2665, 2014.
- [16] J. J. Jia, "Compared modified Simiao decoction with colchicine in the treatment of gout," *Journal of Emergency in Traditional Chinese Medicine*, vol. 19, no. 1, pp. 47–48, 2010.
- [17] S. X. Li and Z. Z. Zhang, "Clinical study of modified Simiao decoction in the treatment of 40 cases of gouty arthritis," *Journal of Traditional Chinese Medicine*, vol. 48, no. 6, pp. 521–523, 2007.
- [18] J. F. Li and T. B. Song, "The effects of Simiao Maqian decoction in the treatment of gouty arthritis," *Guangming Journal of Chinese Medicine*, vol. 28, no. 6, pp. 1161–1162, 2013.
- [19] M. Y. Liu, "Clinical research of modified Simiao decoction in the treatment of acute gouty arthritis and hyperuricemia," *Liaoning Journal of Traditional Chinese Medicine*, vol. 38, no. 4, pp. 675–677, 2011.
- [20] J. P. Luo and C. Q. Tang, "The effects of modified Simiao pill in the treatment of gout," *Journal of Liaoning University of Traditional Chinese Medicine*, vol. 12, no. 11, pp. 156–157, 2010.
- [21] J. G. Mi, "Treatment of 44 cases of gouty arthritis with Simiao decoction plus coix seed decoction," *Guangming Journal of Chinese Medicine*, vol. 31, no. 15, pp. 2223–2225, 2016.
- [22] M. Niu, "Treatment of 99 cases of acute gouty arthritis with modified Simiao decoction," *Heilongjiang Journal of Traditional Chinese Medicine*, vol. 37, no. 6, pp. 19–20, 2008.
- [23] J. S. Sun, "Clinical observation of modified Simiao decoction in the treatment of 36 cases with gouty arthritis," *Journal of Anhui Traditional Chinese Medical College*, vol. 25, no. 3, pp. 14–15, 2006.
- [24] C. L. Tang, W. L. Qian, and L. X. Zhu, "Compared modified Simiao decoction with colchicine in the treatment of acute gouty arthritis," *Heilongjiang Journal of Traditional Chinese Medicine*, vol. 51, no. 6, pp. 12–13, 2008.
- [25] H. J. Wang, G. H. Zhang, J. Zou, and X. Z. Fan, "Clinical research of modified Simiao decoction in the treatment of acute gouty arthritis and hyperuricemia," *Journal of Emergency in Traditional Chinese Medicine*, vol. 18, no. 10, pp. 1609–1610, 2009.
- [26] F. Wang, H. Q. Ren, X. B. Shen, L. Zhang, and X. Y. Cui, "Clinical observation of Simiao Wuling decoction in the treatment of acute gouty arthritis," *Journal of Hubei College of Traditional Chinese Medicine*, vol. 18, no. 3, pp. 81–84, 2016.
- [27] T. T. Wang, H. Zhu, and M. H. Zhang, "Analyzed the effects of Baihu plus cassia twig decoction and Simiao pill in the treatment of acute gouty arthritis," *Journal of New Chinese Medicine*, vol. 48, no. 1, pp. 63–64, 2016.
- [28] S. S. Yu and Y. R. Huang, "Observed the effects of qingre lishi jiedu method in the treatment of acute gouty arthritis," *Hubei Journal of Traditional Chinese Medicine*, vol. 30, no. 9, p. 32, 2008.
- [29] X. Z. Zeng and Y. M. Song, "Treatment of 60 cases of acute gouty arthritis with modified Simiao decoction," *Hunan Journal of Traditional Chinese Medicine*, vol. 29, no. 2, pp. 68–69, 2013.
- [30] Q. X. Zhang and Z. Q. Chen, "Clinical research of modified Simiao pill in the treatment of acute gouty arthritis," *Practical Clinical Journal of Integrated Traditional Chinese and Western Medicine*, vol. 12, no. 4, pp. 36–37, 2012.
- [31] B. Q. Zhao and M. Q. Mai, "Clinical observation of oral and external use of modified Simiao decoction in the treatment of gouty arthritis," *Journal of Emergency in Traditional Chinese Medicine*, vol. 17, no. 6, pp. 777–778, 2008.
- [32] J. Q. Zhao, "Clinical observation of modified Simiao decoction in the treatment of acute gouty arthritis," *Medicine and Health Care*, vol. 22, no. 4, p. 91, 2014.
- [33] J. X. Zhu, "Compared modified Simiao decoction with colchicine in the treatment of gout," *Chinese Journal of Modern Drug Application*, vol. 10, no. 4, pp. 266–267, 2016.
- [34] Q. Y. Zhou, L. Lu, J. K. Cui, W. W. Yu, X. L. Chen, and X. L. Liu, "A systematic review and meta analysis of Simiao powder in the treatment of gouty arthritis and uric acid," *Liaoning Journal of Traditional Chinese Medicine*, vol. 43, no. 7, pp. 1356–1360, 2016.
- [35] M. R. Du, Z. Z. Guo, and F. H. Feng, "The efficiency and safety of modified Simiao decoction in the treatment of acute gouty arthritis: a meta analysis," *Traditional Chinese Medicinal Research*, vol. 28, no. 8, pp. 66–70, 2015.
- [36] M. R. Du, Z. Z. Guo, and F. H. Feng, "The efficiency and safety of modified Simiao decoction in the treatment of acute gouty arthritis: a systematic review," *Chinese Journal of Experimental Traditional Medical Formulae*, vol. 21, no. 13, pp. 212–216, 2015.
- [37] J. S. Xie, *Modified Simiao Powder for Gout: A Systematic Review*, Chengdu University of Traditional Chinese Medicine, 2015.