

Article

Role of Pseudouridine Formation by Deg1 for Functionality of Two Glutamine Isoacceptor tRNAs

Roland Klassen* and Raffael Schaffrath*

Institut für Biologie, Fachgebiet Mikrobiologie, Universität Kassel, Heinrich-Plett-Str. 40, D-34132 Kassel, Germany

* Correspondence: roland.klassen@uni-kassel.de (R.K.); schaffrath@uni-kassel.de (R.S.)

Academic Editor: Valérie de Crécy-Lagard

Received: 16 December 2016; Accepted: 20 January 2017; Published: 26 January 2017

Abstract: Loss of Deg1/Pus3 and concomitant elimination of pseudouridine in tRNA at positions 38 and 39 (ψ 38/39) was shown to specifically impair the function of tRNA^{Gln}_{UUG} under conditions of temperature-induced down-regulation of wobble uridine thiolation in budding yeast and is linked to intellectual disability in humans. To further characterize the differential importance of the frequent ψ 38/39 modification for tRNAs in yeast, we analyzed the in vivo function of non-sense suppressor tRNAs SUP4 and sup70-65 in the absence of the modifier. In the tRNA^{Tyr}_{G ψ A} variant SUP4, UAA read-through is enabled due to an anticodon mutation (U ψ A), whereas sup70-65 is a mutant form of tRNA^{Gln}_{CUG} (SUP70) that mediates UAG decoding due to a mutation of the anticodon-loop closing base pair (G31:C39 to A31:C39). While SUP4 function is unaltered in deg1/pus3 mutants, the ability of sup70-65 to mediate non-sense suppression and to complement a genomic deletion of the essential SUP70 gene is severely compromised. These results and the differential suppression of growth defects in deg1 mutants by multi-copy SUP70 or tQ(UUG) are consistent with the interpretation that ψ 38 is most important for tRNA^{Gln}_{UUG} function under heat stress but becomes crucial for tRNA^{Gln}_{CUG} as well when the anticodon loop is destabilized by the sup70-65 mutation. Thus, ψ 38/39 may protect the anticodon loop configuration from disturbances by loss of other modifications or base changes.

Keywords: tRNA modification; PUS3/DEG1; translation; 5-methoxycarbonylmethyl-2-thiouridine; non-sense suppression; sup70-65

1. Introduction

In addition to standard nucleosides, tRNA is known to contain a large variety of modified nucleosides, formed by the addition of chemical groups or isomerization. Pseudouridine (ψ) is an isomer of uridine (5-ribosyl-uracil) and represents the initially discovered and most abundant modified nucleoside present in all three domains of life [1,2]. The model organism *Saccharomyces cerevisiae* has ten characterized pseudouridine synthases (Pus1–9 and Cbf5) that modify cytoplasmic and mitochondrial tRNAs and/or other types of RNA, including rRNA, small nuclear RNA (snRNA) and mRNA [2]. Despite its ubiquitous presence in tRNA, ψ appears to affect tRNA function rather subtly, since it is not generally required for cell viability, as demonstrated by the non-essential nature of all of the PUS genes in the yeast system [3]. However, the absence of PUS3/DEG1 in yeast uniquely causes slow growth, in particular at elevated temperature [4]. The general effects of ψ on tRNA function are likely mediated by increasing the rigidity of the sugar phosphate backbone and base stacking [2,5]. In humans, homozygous mutation of the DEG1 orthologue PUS3 is correlated with intellectual disability, suggesting a significant contribution to tRNA functioning not only in yeast but also in humans [6].

In yeast, Deg1 dependent formation of ψ occurs in a majority of tRNAs either at position 38 or 39 [4,7]. Slow growth of yeast mutants lacking Deg1 is strongly aggravated if parts of the

wobble uridine modification 5-methoxy-carbonylmethyl-2-thiouridine (mcm^5s^2U) are simultaneously absent [8,9]. The mcm^5 component of the mcm^5s^2U modification is formed by a pathway requiring the Elongator complex and a variety of accessory and potentially regulatory proteins [10–19]. The separate Urm1 pathway mediates the transfer of sulfur to the carbon 2 of the uracil base. This requires the sulfur carrier and ubiquitin-like modifier protein Urm1 and a number of additional proteins operating upstream (Nfs1, Tum1, Uba4) and downstream (Ncs2, Ncs6) of Urm1 resulting in sulfur flow from cysteine to mcm^5U [11,20–25]. The absence of either part alone (mcm^5U or s^2U) induces shared pleiotropic phenotypes, including growth defects at elevated temperature and sensitivity to a number of exogenous stressors including the target of rapamycin complex 1 (TORC1)-inhibiting drug rapamycin [26,27]. On a functional level, partial loss of mcm^5s^2U reduces binding of tRNA to the ribosomal A-site and in turn causes reading frame slippage [28,29]. Combined loss of mcm^5 and s^2U increases the severity of these phenotypes and causes a defect in protein homeostasis resulting in the accumulation of protein aggregates [30–32]. Similarly, combination of Elongator or Urm1 pathway mutations with a *deg1* deletion aggravates growth defects and induces protein aggregation, which is further linked to severe problems in cytoskeleton organization, cell polarity and nuclear segregation [9].

Pleiotropic phenotypes of mutants lacking $\psi38/39$ together with mcm^5s^2U can be specifically rescued by overexpressing $tRNA^{Gln}_{UUG}$, which is reasonable since this tRNA alone carries both modifications [8,9]. In addition to $tRNA^{Gln}_{UUG}$, there is an isoacceptor tRNA with the anticodon sequence CUG that is thought to decode the G-ending codon for Gln. Genetic data from yeast suggests that the U34-containing isoacceptor mainly decodes the A-ending codon and only inefficiently reads the G-ending codon [33]. Because of this, the single *tQ(CUG)* gene has been found to be essential for yeast cell viability [33]. Except for the wobble nucleoside and a variable base (A/G) in position 42, $tRNA^{Gln}_{CUG}$ contains an anticodon stem loop (ASL) identical with that of $tRNA^{Gln}_{UUG}$ and is also modified by Deg1 at position U38 [34]. Interestingly, however, growth defects of *deg1* single mutants were found to be suppressible by overexpression of $tRNA^{Gln}_{UUG}$, but not $tRNA^{Gln}_{CUG}$. This raises the possibility of differential functional dependencies of the two Gln isoacceptor tRNAs on the presence of $\psi38$. In this study, we investigated potential reasons for such divergent effects of one and the same modification on two highly similar tRNA species.

2. Results and Discussion

2.1. Phenotypic Rescue of *deg1* Mutants by tRNA Overexpression

$tRNA^{Gln}_{UUG}$ functionality has been shown to depend on $\psi38$, in particular under conditions of heat stress [8]. Since the $tRNA^{Gln}_{CUG}$ isoacceptor harbours an ASL almost identical to that of $tRNA^{Gln}_{UUG}$, we first investigated whether there are indeed differential dependencies of the two Gln isoacceptors on the presence of $\psi38$. The thermosensitive growth phenotype of *deg1* mutants was shown to be specifically suppressible by overexpression of $tRNA^{Gln}_{UUG}$ but not $tRNA^{Gln}_{CUG}$ or any other tRNA carrying U38 or U39, which is modified by Deg1 to ψ [8]. We confirm this result and find a clear suppression of *deg1*-induced thermosensitivity by multicopy $tRNA^{Gln}_{UUG}$ but not $tRNA^{Gln}_{CUG}$ or other tester tRNAs (Figure 1A). However, rescue by multicopy $tRNA^{Gln}_{UUG}$ is clearly incomplete as growth at 39 °C is not fully restored to wild type levels (Figure 1A). Therefore, we checked whether co-overexpression of the isoacceptor $tRNA^{Gln}_{CUG}$ together with $tRNA^{Gln}_{UUG}$ improves growth at elevated temperature (39 °C) compared to overexpression of $tRNA^{Gln}_{UUG}$ alone. We find that under these conditions, growth of *deg1* mutants already overexpressing $tRNA^{Gln}_{UUG}$ is not further improved by additional overexpression of $tRNA^{Gln}_{CUG}$ (Figure 1B). These results suggest that under heat stress conditions, growth defect of yeast cells lacking $\psi38$ results almost exclusively from malfunction of $tRNA^{Gln}_{UUG}$ rather than $tRNA^{Gln}_{CUG}$, despite containing almost identical ASLs.




Figure 1. Rescue of *deg1* growth defect by tRNA overexpression. **(A)** Single tRNA overexpression. Indicated tRNA species were expressed from multicopy plasmids and in *deg1* mutants and serial dilutions spotted on yeast peptone dextrose (YPD) plates that were incubated at 30 °C or 39 °C. In wild type cells, some tRNA species selected for overexpression carry 5-methoxy-carbonylmethyl-2-thiouridine (mcm^5s^2U), 5-carbamoylmethyluridine (ncm^5U), 5-methoxy-carbonylmethyluridine (mcm^5U), pseudouridine (ψ) 38/39 or combinations thereof, as indicated by (+); **(B)** Combined overexpression of $tRNA^{Gln}_{UUG}$ and $tRNA^{Gln}_{CUG}$. Cells carrying either a multicopy construct for overexpression of $tRNA^{Gln}_{UUG}$ ($tQ(UUG)$), $tRNA^{Gln}_{CUG}$ ($tQ(CUG)$) or both were assayed as in **(A)** for growth at 30, 37 and 39 °C. WT: wild type.

It should also be taken into account that copy numbers of the two Gln isoacceptors in yeast differ significantly due to the number of the respective tRNA genes (nine copies of *tQ(UUG)* versus one copy of *tQ(CUG)* (*Saccharomyces cerevisiae* genome database)). This correlates with a biased codon usage and may impact on the efficiency of phenotypic suppression by tRNA overexpression. Based on this consideration, phenotypic effects of the *DEG1* deletion may be rescuable by overexpression of $tRNA^{Gln}_{UUG}$, since this is the major Gln isoacceptor and consequently more important for overall translation as compared to the minor $tRNA^{Gln}_{CUG}$. However, the single *tQ(CUG)* gene is in fact known to be essential for yeast viability [33], indicating that this C34-containing tRNA is indeed indispensable for translation, despite its lower frequency of utilization compared to the one containing mcm^5s^2U . Hence, the differential dependency of both isoacceptors on ψ 38 as indicated by differential phenotypic suppression of the *deg1* phenotype is presumably, at least in part, due to the difference at the wobble base. Interestingly, $tRNA^{Gln}_{UUG}$ is known to harbour a hypomodified wobble uridine under mild heat stress conditions, which results from a destabilization of the *URM1* pathway and in turn suppresses the tRNA thiolation reaction that converts mcm^5U to mcm^5s^2U [8,25,35–37]. Thus, an explanation why $tRNA^{Gln}_{UUG}$ depends more than $tRNA^{Gln}_{CUG}$ on ψ 38 under heat stress conditions might be the presence of an additional functional impairment of the former tRNA due to wobble uridine hypomodification. $tRNA^{Gln}_{CUG}$ carries a different unmodified wobble base (C34) and therefore remains unaffected by the temperature effect on wobble uridine thiolation.

2.2. Genetic Interactions of *DEG1* under TORC1 Inhibition

In support of a specific requirement for ψ 38 in tRNA^{Gln}_{UUG} in the event of wobble base hypomodification, strong synthetic growth defects were observed upon combining the *deg1* mutation with mutations in Elongator or Urm1 pathway genes, indicative of functional crosstalk between the modifications at position 34 and 38 in tRNA^{Gln}_{UUG} [8,9,38]. Elongator and Urm1 pathway mutants are further known to exhibit shared phenotypes, including translational inaccuracy and sensitivity to TORC1 inhibiting agents such as caffeine or rapamycin and these are synergistically increased in double mutants lacking both modification activities [9,30–32]. To check whether negative genetic interactions between *DEG1*, Elongator and *URM1* can be extended to conditions of TORC1 inhibition, we scored rapamycin sensitivity of *deg1*, *urm1* and *elp3* single and all possible double mutant combinations. As shown in Figure 2, there is a clear rapamycin sensitivity in all three single mutants and this sensitivity is strongly aggravated in all the double mutants, with stronger effects seen in *elp3 deg1* and *urm1 deg1*, where U34 and U38/39 are hypomodified. Thus, there is an overlapping phenotype in *deg1*, *elp3* and *urm1* mutants and all affected modifications (mcm⁵/s²U, ψ 38/39) appear to mediate rapamycin tolerance independently of each other, possibly by maintaining a related aspect of tRNA function. In particular, the observed rapamycin sensitivity of *deg1* single mutants in the absence of heat stress suggests that ψ 38 alone becomes important for tRNA function under at least one stress condition that is different from heat. Since there is strong negative genetic interaction on rapamycin between *DEG1* and *URM1*/Elongator, the latter of which affect tRNA^{Gln}_{UUG} but not tRNA^{Gln}_{CUG}, we assume that these effects are caused by specific malfunction of tRNA^{Gln}_{UUG}.




Figure 2. Interactions between *DEG1* and genes coding for Elongator (*ELP3*) or Urm1 (*URM1*) pathway components under TORC1 inhibition. Indicated single and double mutants were spotted on drug-free yeast peptone dextrose (YPD) plates which were incubated at 30 °C or 37 °C or YPD medium containing 5 nM of the TORC1 inhibiting drug rapamycin.

2.3. Replacement of tQ(CUG) by the *sup70-65* Allele in Haploid Yeast

The finding that tRNA^{Gln}_{UUG} lacking ψ 38 becomes vulnerable to disturbances of the natural anticodon loop configuration (loss of U34 modification) suggested that tRNA^{Gln}_{CUG} devoid of ψ 38 might also be sensitized to destabilization of the ASL. However, as C34 is unmodified and no modification except ψ 38 is known for the ASL of tRNA^{Gln}_{CUG} [7,34], we utilized a variant of tRNA^{Gln}_{CUG} with a disturbance of the ASL due to a sequence change (G31A). This mutation, known as the *tQ(CUG)* allele *sup70-65*, replaces the Watson–Crick base pair G31:C39 normally closing the anticodon loop of tRNA^{Gln}_{CUG} by a mismatch (A31:C39) and reduces stability and charging efficiency of the mutant tRNA [39–41]. Despite these strong effects on the tRNA product of the essential single-copy gene *tQ(CUG)*, diploid yeast cells carrying the homozygous *sup70-65* mutation are viable [39–41] and may provide an opportunity to analyze the effects of ψ 38 removal on tRNA^{Gln}_{CUG} in the presence of a destabilized ASL. We utilized a haploid strain (S288C) lacking the single genomic

copy of *tQ(CUG)* and carrying the wild type *tQ(CUG)* gene on a plasmid that is counter-selectable by 5-fluoro-orotate (5-FOA) [31,40]. This strain was transformed with a second centromeric plasmid carrying the *sup70-65* allele of the *tQ(CUG)* gene. Subsequently, the former plasmid was removed by growing cells on 5-FOA medium, resulting in a strain carrying *sup70-65* as the sole genetic source of *tQ(CUG)* (Figure 3A). The viability of this strain suggests that even in haploid S288C cells, the mutant form of *tQ(CUG)* is active to maintain sufficient levels of translation. However, we find that in comparison to wild type, a strong thermosensitivity is caused by the mutation, while no significant growth defect was observed at 30 °C (Figure 3B). This finding possibly indicates that the destabilized form of tRNA^{Gln}_{CUG} (*sup70-65*) is sufficiently active in translation at normal, but not at elevated temperatures. The absence of defects in growth or CAG decoding in diploid *sup70-65* cells was already documented [39]. However, more recent work demonstrated a reduced expression of reporter genes with tandem CAG codons and several natural CAG-codon-containing mRNAs [40,41]. Hence, a translational defect is already present in *sup70-65* cells at 30 °C and this may be aggravated at elevated temperatures, as evidenced by the observed growth defect (Figure 3B). It is noteworthy that tRNA^{Gln}_{CUG} can be functionally replaced by the inefficient CAG decoder tRNA^{Gln}_{UUG} if the latter is present at elevated levels and carries the full mcm⁵s²U modification [33]. In such strain, CAG codons are exclusively decoded by tRNA^{Gln}_{UUG}, involving U-G wobble and the mcm⁵s²U modified wobble base. In particular, the absence of s²U in tRNA^{Gln}_{UUG} entirely prevents the suppression of lethal effects of a *tQ(CUG)* deletion by multi-copy *tQ(UUG)* [33]. Thus, the strong growth defect of haploid *sup70-65* cells at elevated temperatures may also indicate that tRNA^{Gln}_{UUG} contributes, to some extent, to CAG (Gln) decoding in this mutant, since tRNA^{Gln}_{CUG} is functionally impaired. If so, elevated temperature would repress thiolation and in turn prevent the engagement of tRNA^{Gln}_{UUG} in decoding of the G-ending Gln codon (CAG).




Figure 3. Role of ψ 38 for the *sup70-65* allele of the *tQ(CUG)* gene. (A) Experimental design for the construction of a haploid *sup70-65* strain and plasmid shuffle. A strain carrying a genomic deletion of the essential single-copy *tQ(CUG)* is kept alive by plasmid pAK1 (*tQ(CUG)*-*URA3*). A second plasmid is introduced carrying the mutant *sup70-65* allele of *tQ(CUG)* and pAK01 is eliminated from the strain by growth on 5-fluoro-orotate (5-FOA) media; (B) Drop-dilution assay as in Figure 1 with WT, *deg1* and the haploid strain carrying *sup70-65* as the sole genetic source of *tQ(CUG)*; (C) Plasmid shuffle assay to check for the ability to lose pAK01 (*tQ(CUG)*-*URA3*) in the presence (right) or absence (left) of an additional *deg1* deletion.

To test whether *sup70-65* is still able to replace wild type *tQ(CUG)* when ψ 38 is removed, we deleted the *DEG1* gene in a strain carrying a genomic deletion of *tQ(CUG)* and containing the wild type *tQ(CUG)* gene on the counter-selectable plasmid. Then, the ability to lose the *tQ(CUG)* plasmid in

the presence of the *sup70-65* plasmid was assayed in comparison to the strain lacking the *deg1* mutation. We find that colony formation on 5-FOA medium, which counter-selects the *tQ(CUG)* wild type allele, is severely reduced in the strain lacking *DEG1* (Figure 3C). This result supports the interpretation that $\psi38$ becomes critical for function of $\text{tRNA}^{\text{Gln}}_{\text{CUG}}$ in the presence of a destabilized ASL.

2.4. Comparison of Non-Sense Suppression by *sup70-65* and *SUP4* in the Presence and Absence of $\psi38/39$

Since *sup70-65* is known as an amber suppressor [39,40], we analyzed how loss of $\psi38$ influences the efficiency of UAG stop codon read-through by *sup70-65*. If $\psi38$ is indeed crucial for $\text{tRNA}^{\text{Gln}}_{\text{CUG}}$ function in the presence of a destabilized ASL, then stop codon read-through by *sup70-65* could be impaired by the *deg1* mutation. As a comparison, we scored the efficiency of *SUP4*, a distinct non-sense suppressor tRNA, derived from $\text{tRNA}^{\text{Tyr}}_{\text{G}\psi\text{A}}$ [10]. *SUP4* can read-through ochre (UAA) stop codons due to a base change at the wobble position and was already shown to crucially depend on the presence of the Elongator-mediated mcm^5U modification ($\text{mcm}^5\text{U}\psi\text{A}$) [10]. Since $\text{tRNA}^{\text{Tyr}}_{\text{G}\psi\text{A}}$ harbours a *DEG1* dependent ψ in position 39 [7], *SUP4* could in principle also be affected by the *deg1* mutation. To test this, we utilized a W303-1B-derived strain carrying *SUP4* and *ade2-1*, an allele of *ADE2* causing adenine auxotrophy due to an ochre mutation [10]. Non-sense suppression of the *ade2-1* transcript is known to cause adenine prototrophy and change of colony color from red (no read-through) to white (read-through) [10]. In this background, we deleted the *DEG1* gene and scored adenine auxotrophy as well as color colony in direct comparison with an *elp3* mutant that is known to affect *SUP4*-mediated read-through. While the presence of *SUP4* causes the expected change in color and adenine prototrophy, we find that this indeed depends on a functional *ELP3* gene but not on *DEG1* (Figure 4A). Since this assay might be too insensitive to detect smaller reductions in *SUP4*-mediated read-through and to confirm the obtained result with a premature ochre codon in another reporter gene, we utilized a quantitative assay relying on a *lacZ* reporter with an engineered ochre codon [33,42]. Again, read-through in this reporter was strongly enhanced by the presence of *SUP4* and suppressed by the *elp3* mutation, yet not by *deg1* (Figure 4B). Further, even application of mild temperature stress (37 °C) to cells carrying the reporter constructs did not significantly reduce read-through levels of the *deg1* mutant as compared to the wild type *SUP4* strain under the same conditions (Figure 4B). Thus, *Deg1*-mediated $\psi39$ in itself does not detectably impact on the function of *SUP4* in non-sense suppression in two different read-through contexts. This result is consistent with the absence of suppression of temperature sensitive growth of the *deg1* mutant by multi-copy $\text{tRNA}^{\text{Tyr}}_{\text{G}\psi\text{A}}$ (Figure 1A) and supports the conclusion that ψ s formed by *Deg1* in distinct tRNA species may have different functional consequences.

For *sup70-65*-mediated non-sense suppression, we utilized strain W303-1B, which carries the *trp1-1* allele that contains a premature amber (UAG) codon and is efficiently suppressible by *sup70-65* when the mutant tRNA gene is present on a multi-copy plasmid [40]. We deleted the *DEG1* gene in W303-1B and introduced *sup70-65* on a multi-copy construct. As a control, we used the parental *DEG1* wild type strain and introduced *sup70-65* on both, single and multi-copy vectors [40]. While there is no major difference in growth on tryptophan-supplemented minimal media, the presence of the *sup70-65* gene in both single and multi-copy conferred an ability to grow on tryptophan-free media specifically to the *DEG1* wild type. As expected, multi-copy expression of *sup70-65* was required for induction of efficient growth in the absence of tryptophan, due to relatively inefficient amber suppression by *sup70-65* via first base wobble (U1:G36/A2:U35/G3:C34) [40]. In the absence of the suppressor tRNA, no growth on tryptophan-free medium was observed and, most importantly, growth was also absent in the *deg1* mutant with the multi-copy *sup70-65* construct causing efficient read-through in the *DEG1* wild type strain. Thus, indeed *sup70-65*-mediated non-sense suppression critically depends on the presence of $\psi38$ formation by *Deg1*.




Figure 4. Role of ψ 38/39 for non-sense suppression by *SUP4* or *sup70-65*. (A) *SUP4*-mediated read-through of *ade2-1*. WT: Strain W303-1B, lacking *SUP4*; *deg1*: *DEG1* deletion in W303-1B; *SUP4*: strain UMY2893, carrying *SUP4*; *SUP4 deg1*: deletion of *DEG1* in UMY2893; *SUP4 elp3*: deletion of *ELP3* in UMY2893. Serial dilutions were spotted on adenine-free (-Ade) or adenine-containing medium (+Ade); (B) Quantitative *SUP4*-mediated *lacZ* read-through assay. Strains as described in (A) were transformed with pUKC815 and pUKC817 and relative read-through levels measured as described in Materials and Methods. 30 °C and 37 °C indicate growth temperatures. Averages of at least three biological replicates are shown along with standard deviation; (C) *sup70-65*-mediated read-through of *trp1-1*. Strain W303-1B or W303-1B *deg1* was transformed with empty vector (vector), single-copy (s.c.) or high-copy (h.c.) constructs with the *sup70-65* allele of *tQ(CUG)*. The ability of strains to grow on tryptophan-free (-Trp) medium by read-through of the *trp1-1* marker was scored by drop dilution assays. +Trp: tryptophan-containing medium condition.

3. Materials and Methods

3.1. Strains, Plasmids and General Methods

Yeast strains used are listed in Table 1. Standard methods were used in this study for yeast growth [43]. Yeast was routinely grown in complex yeast peptone dextrose (YPD) medium. To select transformants or maintain plasmids, either synthetic complete (SC) medium lacking specific nutrients or YPD medium containing $200 \mu\text{g}\cdot\text{mL}^{-1}$ G418 was used. Deletion of *DEG1* involved PCR-mediated synthesis of deletion cassettes using previously described oligonucleotides [9] and template plasmids pUG6 and pUG27 [44]. Transformation was done using the PEG/lithium acetate method [45]. Correct genomic replacement of the *DEG1* gene was verified by PCR using oligonucleotides described [9]. Multicopy plasmids for overexpression of $\text{tRNA}^{\text{Glu}}_{\text{UUC}}$, $\text{tRNA}^{\text{Lys}}_{\text{UUU}}$,

tRNA^{Arg}_{UCU}, tRNA^{Leu}_{UAA}, tRNA^{Tyr}_{GUA} and tRNA^{Gly}_{GCC} were described earlier [13] as well as the construct for tRNA^{Gln}_{UUG} [46]. Wild type tRNA^{Gln}_{CUG} overexpression used 2 μ plasmid pSUP70 [40]. pAK01 is a single copy (CEN/ARS) plasmid carrying the wild type *tQ(CUG)* gene as well as *URA3* as a counter-selectable marker [40]. Counter-selection was done using synthetic complete medium containing uracil and 5-FOA at 1 mg·mL⁻¹. Plasmids pSUP70-65-2 μ and pSUP70-65-CEN carry the *sup70-65* allele of *tQ(CUG)* and either the 2 μ origin or CEN-ARS, respectively [40].

Table 1. Strains used in this study.

Strain	Genotype	Reference/Source
<i>Saccharomyces cerevisiae</i> BY4741	MATa, <i>his3</i> Δ , <i>leu2</i> Δ , <i>met15</i> Δ , <i>ura3</i> Δ	Euroscarf, Frankfurt
<i>S. cerevisiae</i> <i>elp3</i>	BY4741 <i>elp3</i> Δ KanMX4	Euroscarf, Frankfurt
<i>S. cerevisiae</i> <i>urm1</i>	BY4741 <i>urm1</i> Δ KanMX4	Euroscarf, Frankfurt
<i>S. cerevisiae</i> <i>deg1</i>	BY4741 <i>deg1</i> Δ KanMX4	Euroscarf, Frankfurt
<i>S. cerevisiae</i> <i>elp3 urm1</i>	BY4741 <i>elp3</i> Δ KanMX4 <i>urm1</i> Δ HIS3	[31]
<i>S. cerevisiae</i> <i>deg1 urm1</i>	BY4741 <i>urm1</i> Δ KanMX4 <i>deg1</i> Δ SpHIS5	[9]
<i>S. cerevisiae</i> <i>deg1 elp3</i>	BY4741 <i>elp3</i> Δ KanMX4 <i>deg1</i> Δ SpHIS5	[9]
<i>S. cerevisiae</i> <i>sup70</i>	BY4741 pAK01 <i>tQ(CUG)</i> Δ KILEU2	[31]
<i>S. cerevisiae</i> <i>sup70-65-pAK01</i>	BY4741 pAK01 <i>tQ(CUG)</i> Δ KILEU2 pSUP70-65-CEN	this work
<i>S. cerevisiae</i> <i>sup70-65</i>	BY4741 <i>tQ(CUG)</i> Δ KILEU2 pSUP70-65 CEN	this work
<i>S. cerevisiae</i> <i>sup70-65-pAK01</i>	BY4741 pAK01 <i>tQ(CUG)</i> Δ KILEU2 <i>deg1</i> Δ KanMX4 pSUP70-65-CEN	this work
<i>S. cerevisiae</i> W303-1B	MAT α <i>leu2-3,112 trp1-1 can1-100 ura3-1 ade2-1 his3-11,15</i>	[47]
<i>S. cerevisiae</i> RK207	W303-1B <i>deg1</i> Δ SpHIS5	this work
<i>S. cerevisiae</i> RK273	W303-1B <i>deg1</i> Δ loxP	this work
<i>S. cerevisiae</i> UMY2893	MAT α SUP4 <i>leu2-3,112 trp1-1 can1-100 ura3-1 ade2-1 his3-11,15</i>	[10]
<i>S. cerevisiae</i> UMY2916	UMY2893 <i>elp3</i> Δ KanMX4	[10]
<i>S. cerevisiae</i> RK208	UMY2893 <i>deg1</i> Δ SpHIS5	this work

3.2. Phenotypic Assays

Yeast strains were either grown on YPD or appropriate SC solid medium to select for the presence of plasmids for 24 to 36 h. Cells were recovered from the plates and resuspended in sterile water. Cell densities were measured using absorbance at 600 nm and dilutions prepared from this suspension with final OD_{600nm} values of 0.15, 0.015, 0.0015 and 0.00015. These were spotted on either drug-free solid YPD plates or YPD plates supplemented with sterile rapamycin at a final concentration of 5 nM. Subsequently, plates were incubated at different temperatures for 36–72 h and photographed. The longer incubation time (72 h) was used for phenotypic assay with *deg1 elp3* double mutants. For qualitative read-through of *ade2-1* and *trp1-1* alleles, serial dilution were spotted on SC lacking adenine or tryptophan respectively and incubated for 48 to 60 h at 30 °C.

3.3. Quantitative Read-through Assay

Cells carrying pUKC815 (wild type *lacZ*) or pUKC817 (UAA ochre insertion) [42] were grown at 30 °C or 37 °C in liquid SC medium lacking uracil to OD₆₀₀ nm of 2–3 and harvested by centrifugation. Cells were washed and resuspended in Z-buffer (60 mM Na₂HPO₄, 40 mM NaH₂PO₄, 10 mM KCl, 50 mM 2-mercaptoethanol, pH 7). Cell density was measured by absorbance at 600 nm. To 500 μ L aliquots, two drops of 0.01% sodium dodecyl sulfate (SDS) solution and chloroform were added and samples mixed on a vortex for 30 s each. Following incubation at 37 °C for 5 min, 100 μ L of 4 mg·mL⁻¹ ortho-nitrophenyl- β -galactoside dissolved in Z-buffer was added. Reactions were stopped by the addition of 250 μ L 1 M Na₂CO₃ and absorbance at 420 nm measured. Activity units were calculated by employing Miller's formula [48]. Relative read-through efficiency (%) was calculated by dividing the beta galactosidase activity measured with the pUKC817 construct by the one measured with the pUKC815 construct [33,42]. For each strain, at least three independent cultures were measured with both constructs.

4. Conclusions

The emerging picture from the presented work and previous studies places Deg1-mediated ψ_{38} in a role to maintain the function of both Gln isoacceptor tRNAs. However, the presence of ψ_{38} becomes critical for tRNA function only in the presence of an additional disturbance of the ASL structure. In one case (tRNA^{Gln}_{UUG}), a strong growth defect indicative of a significant loss of function is observed specifically when *deg1* mutants are shifted to conditions known to downregulate the formation of a distinct modification (mcm⁵s²U) (Figures 1 and 2). In another case (tRNA^{Gln}_{CUG}) where this modification is absent per se, the introduction of a destabilization of the ASL can substitute for a hypomodified wobble base and sensitize this tRNA as well for the negative effects of the *deg1* mutation (Figures 3 and 4). It appears possible that other tRNA species such as tRNA^{Tyr}_{G ψ A} do not lose function in the absence of $\psi_{38/39}$ due to the existence of additional modifications that may stabilize the ASL. For example, tRNA^{Tyr}_{G ψ A}-derived suppressor tRNAs are known to carry N⁶-isopentenyl-adenosine (i⁶A) in position 37 [7]. The presence of i⁶A has been shown to support efficient ochre read-through [49] and therefore may protect the ASL from destabilization. We assume that multiple modifications of the ASL provide multiple layers of structural protection, which routinely are not ruptured in single tRNA modification mutants and this functional redundancy may explain the paradox situation of missing phenotypes for many of the evolutionary conserved tRNA modification genes.

Acknowledgments: We thank Ian Stansfield for providing plasmids and gratefully acknowledge funding support by the Deutsche Forschungsgemeinschaft (DFG) to R.S. (SCHA750/15) and their Priority Programme SPP1784 ‘Chemical Biology of Native Nucleic Acid Modifications’ to R.S. (SCHA750/20) and R.K. (KL2937/1).

Author Contributions: R.K. and R.S. conceived and designed the experiments; R.K. performed the experiments; R.K. and R.S. analyzed the data; R.K. and R.S. wrote the paper.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Yu, C.T.; Allen, F.W. Studies on an isomer of uridine isolated from ribonucleic acids. *Biochim. Biophys. Acta.* **1959**, *32*, 393–406. [[CrossRef](#)]
2. Spenkuch, F.; Motorin, Y.; Helm, M. Pseudouridine: Still mysterious, but never a fake (uridine)! *RNA Biol.* **2014**, *11*, 1540–1554. [[CrossRef](#)] [[PubMed](#)]
3. Giaever, G.; Chu, A.M.; Ni, L.; Connelly, C.; Riles, L.; Véronneau, S.; Dow, S.; Lucau-Danila, A.; Anderson, K.; André, B.; et al. Functional profiling of the *Saccharomyces cerevisiae* genome. *Nature* **2002**, *418*, 387–391. [[CrossRef](#)] [[PubMed](#)]
4. Lecointe, F.; Simos, G.; Sauer, A.; Hurt, E.C.; Motorin, Y.; Grosjean, H. Characterization of Yeast Protein Deg1 as Pseudouridine Synthase (Pus3) Catalyzing the Formation of ψ_{38} and ψ_{39} in tRNA Anticodon Loop. *J. Biol. Chem.* **1998**, *273*, 1316–1323. [[CrossRef](#)] [[PubMed](#)]
5. Charette, M.; Grey, M. Pseudouridine in RNA: What, where, how and why. *IUBMB Life* **2000**, *49*, 341–351. [[PubMed](#)]
6. Shaheen, R.; Han, L.; Faqeih, E.; Ewida, N.; Alobeid, E.; Phizicky, E.M.; Alkuraya, F.S. A homozygous truncating mutation in *PUS3* expands the role of tRNA modification in normal cognition. *Hum. Genet.* **2016**, *135*, 707–713. [[CrossRef](#)] [[PubMed](#)]
7. Machnicka, M.A.; Milanowska, K.; Osman Oglou, O.; Purta, E.; Kurkowska, M.; Olchowik, A.; Januszewski, W.; Kalinowski, S.; Dunin-Horkawicz, S.; Rother, K.M.; et al. MODOMICS: A database of RNA modification pathways—2013 update. *Nucleic Acids Res.* **2013**, *41*, D262–D267. [[CrossRef](#)] [[PubMed](#)]
8. Han, L.; Kon, Y.; Phizicky, E.M. Functional importance of ψ_{38} and ψ_{39} in distinct tRNAs, amplified for tRNA^{Gln}_{(UUG) by unexpected temperature sensitivity of the s²U modification in yeast. *RNA* **2015**, *21*, 188–201. [[CrossRef](#)] [[PubMed](#)]}
9. Klassen, R.; Ciftci, A.; Funk, J.; Bruch, A.; Butter, F.; Schaffrath, R. tRNA anticodon loop modifications ensure protein homeostasis and cell morphogenesis in yeast. *Nucleic Acids Res.* **2016**, *44*, 10946–10959. [[CrossRef](#)] [[PubMed](#)]

10. Huang, B.; Johansson, M.J.; Byström, A.S. An early step in wobble uridine tRNA modification requires the Elongator complex. *RNA* **2005**, *11*, 424–436. [[CrossRef](#)] [[PubMed](#)]
11. Huang, B.; Lu, J.; Byström, A.S. A genome-wide screen identifies genes required for formation of the wobble nucleoside 5-methoxycarbonylmethyl-2-thiouridine in *Saccharomyces cerevisiae*. *RNA* **2008**, *14*, 2183–2194. [[CrossRef](#)] [[PubMed](#)]
12. Kalhor, H.R.; Clarke, S. Novel methyltransferase for modified uridine residues at the wobble position of tRNA. *Mol. Cell. Biol.* **2003**, *23*, 9283–9292. [[CrossRef](#)] [[PubMed](#)]
13. Jablonowski, D.; Zink, S.; Mehlgarten, C.; Daum, G.; Schaffrath, R. tRNA^{Glu} wobble uridine methylation by Trm9 identifies Elongator's key role for zymocin-induced cell death in yeast. *Mol. Microbiol.* **2006**, *59*, 677–688. [[CrossRef](#)] [[PubMed](#)]
14. Fichtner, L.; Frohloff, F.; Bürkner, K.; Larsen, M.; Breunig, K.D.; Schaffrath, R. Molecular analysis of *KTI12/TOT4*, a *Saccharomyces cerevisiae* gene required for *Kluyveromyces lactis* zymocin action. *Mol. Microbiol.* **2002**, *43*, 783–791. [[CrossRef](#)] [[PubMed](#)]
15. Bär, C.; Zabel, R.; Liu, S.; Stark, M.J.; Schaffrath, R. A versatile partner of eukaryotic protein complexes that is involved in multiple biological processes: Kti11/Dph3. *Mol. Microbiol.* **2008**, *69*, 1221–1233. [[PubMed](#)]
16. Zabel, R.; Bär, C.; Mehlgarten, C.; Schaffrath, R. Yeast α -tubulin suppressor Ats1/Kti13 relates to the Elongator complex and interacts with Elongator partner protein Kti11. *Mol. Microbiol.* **2008**, *69*, 175–187. [[CrossRef](#)] [[PubMed](#)]
17. Mehlgarten, C.; Jablonowski, D.; Breunig, K.D.; Stark, M.J.; Schaffrath, R. Elongator function depends on antagonistic regulation by casein kinase Hrr25 and protein phosphatase Sit4. *Mol. Microbiol.* **2009**, *73*, 869–881. [[CrossRef](#)] [[PubMed](#)]
18. Abdel-Fattah, W.; Jablonowski, D.; di Santo, R.; Scheidt, V.; Hammermeister, A.; ten Have, S.M.; Thüning, K.L.; Helm, M.; Schaffrath, R.; Stark, M.J.R. Phosphorylation of Elp1 by Hrr25 is required for Elongator-dependent tRNA modification in yeast. *PLoS Genet.* **2015**, *11*, e1004931. [[CrossRef](#)] [[PubMed](#)]
19. Schaffrath, R.; Leidel, S.A. Wobble uridine modifications—A reason to live, a reason to die?! *RNA Biol.* **2016**. accepted.
20. Nakai, Y.; Nakai, M.; Hayashi, H. Thio-modification of yeast cytosolic tRNA requires a ubiquitin-related system that resembles bacterial sulfur transfer systems. *J. Biol. Chem.* **2008**, *283*, 27469–27476. [[CrossRef](#)] [[PubMed](#)]
21. Noma, A.; Sakaguchi, Y.; Suzuki, T. Mechanistic characterization of the sulfur-relay system for eukaryotic 2-thiouridine biogenesis at tRNA wobble positions. *Nucleic Acids Res.* **2009**, *37*, 1335–1352. [[CrossRef](#)] [[PubMed](#)]
22. Leidel, S.; Pedrioli, P.G.; Bucher, T.; Brost, R.; Costanzo, M.; Schmidt, A.; Aebersold, R.; Boone, C.; Hofmann, K.; Peter, M. Ubiquitin-related modifier Urm1 acts as a sulphur carrier in thiolation of eukaryotic transfer RNA. *Nature* **2009**, *458*, 228–232. [[CrossRef](#)] [[PubMed](#)]
23. Goehring, A.S.; Rivers, D.M.; Sprague, G.F., Jr. Urm1: A ubiquitin-like pathway that functions during invasive growth and budding in yeast. *Mol. Biol. Cell* **2003**, *14*, 4329–4341. [[CrossRef](#)] [[PubMed](#)]
24. Jüdes, A.; Ebert, F.; Bär, C.; Thüning, K.L.; Harrer, A.; Klassen, R.; Helm, M.; Stark, M.J.R.; Schaffrath, R. Urm1 and tRNA thiolation functions of ubiquitin-like Uba4•Urm1 systems are conserved from yeast to man. *FEBS Lett.* **2015**, *589*, 904–909. [[CrossRef](#)] [[PubMed](#)]
25. Jüdes, A.; Bruch, A.; Klassen, R.; Helm, M.; Schaffrath, R. Sulfur transfer and activation by ubiquitin-like modifier system Uba4•Urm1 link protein urmylation and tRNA thiolation in yeast. *Microb. Cell* **2016**, *3*, 423–433. [[CrossRef](#)]
26. Frohloff, F.; Fichtner, L.; Jablonowski, D.; Breunig, K.D.; Schaffrath, R. *Saccharomyces cerevisiae* Elongator mutations confer resistance to the *Kluyveromyces lactis* zymocin. *EMBO J.* **2001**, *20*, 1993–2003. [[CrossRef](#)] [[PubMed](#)]
27. Esberg, A.; Huang, B.; Johansson, M.J.O.; Byström, A.S. Elevated levels of two tRNA species bypass the requirement for elongator complex in transcription and exocytosis. *Mol. Cell* **2006**, *24*, 139–148. [[CrossRef](#)] [[PubMed](#)]
28. Rezgui, V.A.; Tyagi, K.; Ranjan, N.; Konevega, A.L.; Mittelstaet, J.; Rodnina, M.V.; Peter, M.; Pedrioli, P.G. tRNA tKUUU, tQUUG, and tEUUC wobble position modifications fine-tune protein translation by promoting ribosome A-site binding. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 12289–12294. [[CrossRef](#)] [[PubMed](#)]

29. Tükenmez, H.; Xu, H.; Esberg, A.; Byström, A.S. The role of wobble uridine modifications in +1 translational frameshifting in eukaryotes. *Nucleic Acids Res.* **2015**, *43*, 9489–9499. [[CrossRef](#)] [[PubMed](#)]
30. Nedialkova, D.D.; Leidel, S.A. Optimization of Codon Translation Rates via tRNA Modifications Maintains Proteome Integrity. *Cell* **2015**, *161*, 1606–1618. [[CrossRef](#)] [[PubMed](#)]
31. Klassen, R.; Grunewald, P.; Thüring, K.L.; Eichler, C.; Helm, M.; Schaffrath, R. Loss of Anticodon Wobble Uridine Modifications Affects tRNA^{Lys} Function and Protein Levels in *Saccharomyces cerevisiae*. *PLoS ONE* **2015**, *10*, e0119261. [[CrossRef](#)] [[PubMed](#)]
32. Klassen, R.; Bruch, A.; Schaffrath, R. Independent suppression of ribosomal +1 frameshifts by different tRNA anticodon loop modifications. *RNA Biol.* **2016**. [[CrossRef](#)] [[PubMed](#)]
33. Johansson, M.J.; Esberg, A.; Huang, B.; Björk, G.R.; Byström, A.S. Eukaryotic wobble uridine modifications promote a functionally redundant decoding system. *Mol. Cell. Biol.* **2008**, *28*, 3301–3312. [[CrossRef](#)] [[PubMed](#)]
34. Lecointe, F.; Namy, O.; Hatin, I.; Simos, G.; Rousset, J.P.; Grosjean, H. Lack of pseudouridine 38/39 in the anticodon arm of yeast cytoplasmic tRNA decreases in vivo recoding efficiency. *J. Biol. Chem.* **2002**, *277*, 30445–30453. [[CrossRef](#)] [[PubMed](#)]
35. Damon, J.R.; Pincus, D.; Ploegh, H.L. tRNA thiolation links translation to stress responses in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* **2015**, *26*, 270–282. [[CrossRef](#)] [[PubMed](#)]
36. Alings, F.; Sarin, L.P.; Fufezan, C.; Drexler, H.C.; Leidel, S.A. An evolutionary approach uncovers a diverse response of tRNA 2-thiolation to elevated temperatures in yeast. *RNA* **2015**, *21*, 202–212. [[CrossRef](#)] [[PubMed](#)]
37. Tyagi, K.; Pedrioli, P.G. Protein degradation and dynamic tRNA thiolation fine-tune translation at elevated temperatures. *Nucleic Acids Res.* **2015**, *43*, 4701–4712. [[CrossRef](#)] [[PubMed](#)]
38. Scheidt, V.; Jüdes, A.; Bär, C.; Klassen, R.; Schaffrath, R. Loss of wobble uridine modification in tRNA anticodons interferes with TOR pathway signaling. *Microb. Cell* **2014**, *1*, 416–424. [[CrossRef](#)]
39. Murray, L.E.; Rowley, N.; Dawes, I.W.; Johnston, G.C.; Singer, R.A. A yeast glutamine tRNA signals nitrogen status for regulation of dimorphic growth and sporulation. *Proc. Natl. Acad. Sci. USA* **1998**, *95*, 8619–8624. [[CrossRef](#)] [[PubMed](#)]
40. Kemp, A.J.; Betney, R.; Ciandrini, L.; Schwenger, A.C.; Romano, M.C.; Stansfield, I. A yeast tRNA mutant that causes pseudohyphal growth exhibits reduced rates of CAG codon translation. *Mol. Microbiol.* **2013**, *87*, 284–300. [[CrossRef](#)] [[PubMed](#)]
41. Gorgoni, B.; Ciandrini, L.; McFarland, M.R.; Romano, M.C.; Stansfield, I. Identification of the mRNA targets of tRNA-specific regulation using genome-wide simulation of translation. *Nucleic Acids Res.* **2016**, *44*, 9231–9244. [[CrossRef](#)] [[PubMed](#)]
42. Stansfield, I.; Akhmaloka; Tuite, M.F. A mutant allele of the *SUP45* (*SAL4*) gene of *Saccharomyces cerevisiae* shows temperature-dependent allosuppressor and omnipotent suppressor phenotypes. *Curr. Genet.* **2002**, *27*, 417–426. [[CrossRef](#)]
43. Sherman, F. Getting started with yeast. *Methods Enzymol.* **2002**, *350*, 3–41. [[PubMed](#)]
44. Gueldener, U.; Heinisch, J.; Koehler, G.J.; Voss, D.; Hegemann, J.H. A second set of *loxP* marker cassettes for Cre-mediated multiple gene knockouts in budding yeast. *Nucleic Acids Res.* **2002**, *30*, e23. [[CrossRef](#)] [[PubMed](#)]
45. Gietz, D.; Schiestl, R.H. High-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method. *Nat. Protocols* **2007**, *2*, 31–34. [[CrossRef](#)] [[PubMed](#)]
46. Klassen, R.; Paluszynski, J.P.; Wemhoff, S.; Pfeiffer, A.; Fricke, J.; Meinhardt, F. The primary target of the killer toxin from *Pichia acaciae* is tRNA^{Gln}. *Mol. Microbiol.* **2008**, *69*, 681–697. [[CrossRef](#)] [[PubMed](#)]
47. Fiorentini, P.; Huang, K.N.; Tishkoff, D.X.; Kolodner, R.D.; Symington, L.S. Exonuclease I of *Saccharomyces cerevisiae* Functions in Mitotic Recombination In Vivo and In Vitro. *Mol. Cell. Biol.* **1997**, *17*, 2764–2773. [[CrossRef](#)] [[PubMed](#)]
48. Miller, J.H. *Experiments in Molecular Genetics*; Cold Spring Harbor Laboratory Press: New York, NY, USA, 1972.
49. Laten, H.M. Antisuppression of class I suppressors in an isopentenylated-transfer RNA deficient mutant of *Saccharomyces cerevisiae*. *Curr. Genet.* **1984**, *8*, 29–32. [[CrossRef](#)] [[PubMed](#)]

