



Published in final edited form as:

Cancer Res. 2016 July 15; 76(14): 4032–4050. doi:10.1158/0008-5472.CAN-16-0887.

Efficacy and Mechanisms of Aerobic Exercise on Cancer Initiation, Progression, and Metastasis: A Critical Systematic Review of *In Vivo* Preclinical Data

Kathleen A. Ashcraft, PhD¹, Ralph M. Peace, DVM¹, Allison S. Betof, MD, PhD², Mark W. Dewhirst, DVM, PhD^{1,*}, and Lee W. Jones, PhD^{3,*}

¹Duke University Medical Center, Durham, NC, USA

²Massachusetts General Hospital, Boston, MA, USA

³Memorial Sloan Kettering Cancer Center, New York, NY, USA

Abstract

A major objective of the emerging field of exercise-oncology research is to determine the efficacy of, and biological mechanisms by which, aerobic exercise affects cancer incidence, progression and/or metastasis. There is a strong inverse association between self-reported exercise and the primary incidence of several forms of cancer; similarly, emerging data suggest that exercise exposure following a cancer diagnosis may improve outcomes for early-stage breast, colorectal, or prostate cancer. Arguably, critical next steps in the development of exercise as a candidate treatment in cancer control require preclinical studies to validate the biological efficacy of exercise, identify the optimal “dose”, and pinpoint mechanisms of action. To evaluate the current evidence base, we conducted a critical systematic review of *in vivo* studies investigating the effects of exercise in cancer prevention and progression. Studies were evaluated on the basis of tumor outcomes (e.g., incidence, growth, latency, metastasis), dose-response, and mechanisms of action, when available. A total of 53 studies were identified and evaluated on tumor incidence (n=24), tumor growth (n=33) or metastasis (n=10). We report that the current evidence base is plagued by considerable methodological heterogeneity in all aspects of study design, end points, and efficacy. Such heterogeneity precludes meaningful comparisons and conclusions at present. To this end, we provide a framework of methodological and data reporting standards to strengthen the field to guide the conduct of high-quality studies required to inform translational, mechanism-driven clinical trials.

Keywords

Exercise; Physical Activity; Mouse Models; Rat Models; Cancer; Carcinogenesis; Metastasis; Mechanisms

Correspondence: Lee W. Jones, PhD, Department of Medicine, Memorial Sloan Kettering Cancer Center, NY, NY, 10065, onesl3@mskcc.org, Mark W. Dewhirst, DVM, PhD, Gustavo S. Montana Professor of Radiation Oncology, Duke University School of Medicine, Durham, NC 27710, mark.dewhirst@duke.edu.

*denotes equal contribution

Conflict of Interest: The authors report no conflicts of interest

INTRODUCTION

Structured exercise training (hereto referred to as exercise) is considered an integral component of “standard of care” therapy in primary and secondary prevention of numerous common chronic conditions (1–4). In comparison, the role of exercise has received surprisingly little attention in individuals at high-risk or following a diagnosis of cancer. Over the past two decades however, an increasing number of groups are investigating the effects of general physical activity as well as exercise in the oncology setting, a field now commonly referred to as “Exercise-Oncology” (5).

A strong body of observational evidence indicates that higher levels of self-reported exercise, physical activity, as well as cardiorespiratory fitness (i.e., objective assessment of exercise exposure) are inversely associated with the primary incidence of several forms of cancer. This evidence base is summarized by several excellent systematic reviews and meta-analyses (6–9). For example, exercise participation consistent with the national guidelines (i.e., 150 minutes of moderate-intensity or 75 minutes of vigorous-intensity exercise per week) is associated, on average, with 25%, and 30% to 40% risk reductions in breast and colon cancers, respectively compared with inactivity. There is also evidence of a linear dose-response relationship in prevention of breast and colon cancer. On this basis, the evidence for the exercise-prevention relationship is categorized as “convincing” for breast and colon cancer by several national agencies, and regarding as “encouraging” or “promising” for the prevention of prostate, lung, and endometrial cancers (10, 11). On the basis of this data, several phase II randomized controlled trials (RCTs) were initiated, predominantly in breast cancer prevention, to investigate the effects of highly-structured exercise on modulation of host-related factors (e.g., adiposity, circulating factors including sex-steroid and metabolic sex hormones, and the immune-inflammatory axis) that may underpin the exercise-cancer prevention relationship (6, 12–14). In general, exercise was associated with modest changes in markers of adiposity and select circulating factors (6, 12, 13). Whether the observed alterations are of biological or clinical importance remains to be determined, as only one trial to date has investigated whether exercise-induced changes in circulating factors occur in conjunction with changes in factors/pathways in the organ/tissue of primary interest (i.e., colon crypts) (15). Whether exercise reduces cancer incidence or modulates established surrogate markers of cancer incidence has not been investigated.

In patients with cancer, a steadily growing and ever diversifying series of studies indicate that, in general, exercise is a tolerable adjunct therapy associated with significant benefit across a wide range of symptom control variables both during and after primary adjuvant therapy (5, 16, 17). This data combined with the powerful inverse relationships with primary cancer incidence has led researchers to investigate whether exercise influences disease outcomes after diagnosis (18, 19). Although not nearly as mature as the evidence in the prevention setting, emerging observational data suggest that regular exercise exposure is associated with between a 10%–50% reduction in the risk of recurrence and cancer-specific death in patients with colorectal, breast, and prostate cancer (reviewed in (20)), compared to inactivity.

In the development of potential drug candidates, data from observational studies is insufficient to support the initiation of human trials; appropriate preclinical evidence is first required prior to human testing (21). While exercise is not a drug and exhibits a markedly different safety profile than most anticancer agents, the use of preclinical models are of critical importance to confirm biological plausibility, establish the therapeutic window of efficacy, a biologically effective dose, and identify predictors of response (22). This evidence, combined with data from epidemiological and molecular epidemiological studies facilitates the design of early “signal-seeking” clinical studies and ultimately definitive RCTs. Accordingly, we conducted a critical systematic review of *in vivo* pre-clinical studies across the cancer continuum (i.e., prevention through metastasis). A secondary purpose was to provide recommendations to facilitate the standardization of the conduct of *in vivo* exercise-oncology studies as well as directions for future research.

METHODS

Search Strategy and Inclusion Criteria

A systematic literature search was conducted using OVID MEDLINE (1950 to May 2015), PUBMED (1962 to May 2015) and WEB OF SCIENCE (1950 to May 2015) with MeSH terms and keywords related to exercise, cancer and animals. Text words were searched and appropriate MeSH terms were found (Table S1). When possible, Boolean logic was used with MeSH terms to build searches. All results and reference lists were searched manually. Peer-reviewed research articles involving animals with cancer and exposed to chronic exercise (repeated bouts of more than 3 sessions) adopting either forced (i.e., treadmill running or swimming) endurance (aerobic) training or physical activity (i.e., voluntary wheel running) paradigms were considered eligible. All types of animal models of solid tumors were considered eligible including genetically predisposed models (transgenic, genetically engineered mouse models [GEMMs]), orthotopic, subcutaneous and intravenous injections, tumor transplant, and spontaneous or carcinogen-induced solid tumor models. Studies that included multiple treatments, such as study arms with dietary restrictions, were eligible, as long as it was possible to compare sedentary (control) and exercise alone groups. Studies that evaluated pre-neoplastic lesions (e.g., aberrant crypt foci, *Apc*^{Min}), or hematopoietic and ascites tumors were excluded. Only mouse and rat studies were included. Papers unavailable in English were excluded.

Study Selection and Classification

Four authors (KAA, RMP, ASB and LWJ) assessed study eligibility by reviewing the titles and abstracts of all potential citations according to the inclusion criteria. KAA, RMP, MWD and ASB extracted and interpreted data from published papers. When necessary, effort was made to contact authors and acquire publications for evaluation. Studies are summarized by type of exercise model and method of tumor initiation. Exercise characteristics are described using common prescription elements: modality, frequency, duration, intensity and total length of intervention.

Tumor initiation vs. growth

Papers were classified by the endpoints reported in the individual studies: (1) “Incidence” (tumor presence or multiplicity) (2) “Growth” (data on tumor growth, latency or survival), and (3) “Metastasis” (evaluation of tumors distinct from the primary tumor or developing following the common metastasis model of intravenous tumor cell injection). Study categorization was not always mutually exclusive; thus studies that included multiple endpoints applicable to more than one categorization were included in both.

Analysis of mechanistic findings

For evaluation of mechanistic findings, an alternative classification was applied based on the initiation of the exercise intervention relative to the time of tumor cell transplant or application of carcinogens. “Prevention” was defined as exercise initiated prior to tumor transplant/induction. “Progression” was defined as exercise initiated 3 days post-transplant/induction. Studies involving GEMMs were categorized as prevention studies. Distinguishing between exercise initiation and tumor cell inoculation is important because exercise-induced adaptations prior to tumor injection could ‘prime’ the host and/or tissue microenvironment, making it physiological or biologically distinct from tumor inoculation into a sedentary host. Mechanistic findings were classified as either intratumoral or systemic.

Interpretation and analysis

Studies were assessed for changes in tumor growth (as well as the related parameters of time to tumor-related endpoint, and tumor size/mass at the end of the study), incidence (tumor presence) and multiplicity (number of tumors/metastases). All study results that were reported to be “statistically significant” achieved $p < 0.05$ according to the authors of the original manuscripts. Preliminary analysis of included studies indicated considerable methodological heterogeneity in all aspects of study design, end points, and efficacy. As such, it was not possible to compare the efficacy of exercise across cancer setting (i.e., incidence, progression, metastasis), cancer histology, exercise paradigm (i.e., forced vs. voluntary), or exercise dose.

RESULTS

A total of 426 potential citations were initially identified using search terms. After secondary screening, 53 papers were deemed eligible and underwent full review (Figure S1).

Classification of papers was as follows: (1) incidence ($n=24$; 45.3%) (23–46), growth ($n=33$; 62.3%) (24, 26, 34, 37, 39, 40, 42, 44–69), and metastasis ($n=10$; 18.9%) (53, 55, 63, 64, 70–75).

Tumor Models and Exercise Prescription Characteristics

Included studies tested the effects of exercise on 15 different tumor types/cell lines, using six different tumor initiation methods (Table S2). Details of the exercise prescriptions are summarized in Tables 1, 2 and 3. The most common modalities included voluntary running ($n=22$; 41.5%) (24–28, 31, 33, 35, 42, 43, 51, 52, 55, 57, 64–66, 70, 71, 73–75) forced running ($n=25$; 47.2%) (26, 29, 36–41, 44–46, 48, 49, 54, 56, 58–60, 63, 67–70, 72, 74) and swimming ($n=10$; 18.9%) (23, 30, 32, 34, 47, 50, 53, 54, 61, 62).

Effect of exercise on tumor outcomes

In incidence studies, 58% reported that exercise inhibited tumor initiation or multiplicity (23–25, 27, 28, 31, 33, 35, 36, 41, 43–46), 8% reported that exercise accelerated tumor incidence (26, 42), and 3% found null effects (29, 30, 32, 34, 37–40). In the growth category, exercise was associated with tumor inhibition in 64% of studies (24, 39, 47–54, 58–63, 65–69), while 21% (37, 40, 44, 55–57, 64) and 9% (26, 34, 42) of studies reported null and accelerated tumor growth, respectively. Finally, in the tumor metastasis category, three studies utilized models in which metastases arose from a primary tumor; of these, two reported non-significant inhibition of tumor growth (55, 64), while the other found accelerated tumor growth with exercise (53). Eight studies utilized intravenously injected tumor cell model of metastases. MacNeil and Hoffman-Goetz found that exercise did not alter retention of lung tumor cells, but increased the median number of lung metastases (74). Conversely, Jadeski and Hoffman-Goetz reported that exercise decreased retention of lung tumor cells, but had no effect on the number of lung metastases (72). One additional paper reported no difference in tumor cell retention in the lungs (71), whereas three papers reported no effect of exercise on lung metastasis (64, 70, 73). Two papers reported that exercise inhibited lung metastasis multiplicity (63) or total volume (75).

Effects on the intratumoral microenvironment

Mechanistic findings are summarized in Table 4 and Figure 1. Studies were classified as prevention (n=20; 37.7%), progression (n=26; 49.1%) or metastasis (n=7; 13.2%). Eight prevention studies (40%) reported significant effects of exercise in modulation of local immune response, tumor metabolism and tumor physiology/angiogenesis. Ten progression studies (38.5%) examined mechanisms underlying the effects of exercise on tumor progression (32, 41, 42, 44, 52, 55, 57, 61, 68, 69). The mechanisms examined included multiple different factors/pathways such as apoptosis, perfusion and immune cell infiltration. Only one metastasis study examined changes in tumor biology: Higgins et al. found that exercise favored a pro-apoptotic environment (75), reflecting changes in immune cell populations/function, tumor physiology and signaling cascades.

Effects on systemic (host) pathways

Correlative systemic pathways examined are summarized in Table 4. Ten prevention (24, 28, 29, 40, 46–49, 59, 62), seven progression (23, 31, 37, 42, 51, 55, 64, 76) and five metastasis studies (70–72, 74, 75) reported significant effects of exercise on changes in systemic effectors, predominantly changes in factors involved in immune surveillance or metabolism.

DISCUSSION

The findings of this critical review indicate that the current evidence base is plagued by heterogeneity in all aspects of study methodology and data reporting. Unfortunately, such heterogeneity precludes rigorous evaluation of essential questions such as the biologically effective exercise dose to modulate specific tumor pathways or inhibit tumor growth, the effects of manipulating exercise intensity or duration or the differential impact of exercise across tumor subtypes (22). Nevertheless, our review did permit identification of the most salient methodological issues that prudent investigators may consider when designing *in-*

vivo pre-clinical exercise-oncology studies. These aspects are described in the proceeding sections and summarized in Table 5.

Heterogeneity in Exercise Prescription

The components of the exercise prescription being investigated in preclinical studies should mirror, as closely as possible, exercise prescription parameters tested in human studies (22). Key parameters include: (1) modality/paradigm (i.e., forced vs. voluntary), (2) dose (i.e., intensity, session duration, frequency of training sessions/week, and length of treatment), and (3) schedule (i.e., time of initiation).

Exercise Modality (Forced vs. Voluntary Paradigms)—A foremost decision is the use of forced versus voluntary exercise paradigms. An often overlooked key point is that voluntary wheel running is a model of physical activity whereas forced paradigms are a model of exercise training (i.e., structured and purposeful physical activity). Observational (epidemiological) studies typically measure both physical activity and exercise, whereas efficacy-based clinical trials largely examine highly structured exercise training. The decision of which exercise modality is selected should depend on whether the investigators envisage the study findings directly informing clinical translation or plausibility/mechanisms of a clinical observation.

An important caveat to consider in all exercise paradigms is the degree of associated negative stress, as evidenced by exercise-induced increases in serum corticosterone and fecal corticosterone metabolites (77, 78), as observed in both voluntary and forced exercise paradigms. Furthermore, voluntary running may result in “addictive behavior” with negative effects on the hypothalamic-pituitary-adrenal (HPA) axis and animal health (79). Human studies have taught us that exercise induces spikes in cortisol at the beginning of exercise and immediately after bout of exercise ceases (80). Stress-induced activation of the HPA axis may accelerate tumor growth (81), and have diverse effects on the immune response (reviewed in (82)), thereby confounding the efficacy of exercise on tumor growth characteristics. HPA-mediated changes on the immune response could contribute to discordance between studies with respect to immune function. For example Zhu et al. reported a decrease in TNF- α (43) whereas Abdalla et al. reported an increase (23).

Though changes in glucocorticoids have been seen using all exercise modes and intensities, such considerations may be especially important when investigating the effects of high-intensity or high-volume exercise treatment doses. This is relevant because, as reviewed here, studies have tested the effects of exhaustive exercise doses (40) (e.g., forced swimming for four hours followed by 12 hours of physical activity (54)), or forced swimming of animals loaded with external weights (30, 32, 47). The latter exercise paradigms may have additional safety and ethical concerns. Indeed, one study investigating the effect of forced swimming reported four animal deaths due to drowning (30). Researchers should also be cognizant of the impact of the exercise on the animals’ natural circadian rhythms. Fuss et al. demonstrated that voluntary wheel running occurs predominantly during the animals’ dark cycle (78), therefore it would be prudent to conduct exercise training during the dark cycle. Regardless of exercise prescription, corticosterone analysis would be helpful to include in all

study designs; such levels could be correlated with tumor growth end points. In this review, three studies analyzed systemic corticosterone levels (42, 48, 51); of these, two studies reported increases in corticosterone levels whereas the other study reported a decrease. Interestingly, tumor growth was inhibited in all studies.

Exercise Dose (Intensity, Session Duration, Frequency of Training, and Length of Treatment)—Exercise at different intensities confers remarkably different physiological and gene expression adaptations in mammals (83), however a key question is whether these differences translate into differential effects on tumor outcomes. In clinical trials, the efficacy of different exercise intensities varying from 50% to 100% of a patient-derived physiological parameter (e.g., age-predicted maximum heart rate, measured exercise capacity) have been investigated. Such an approach permits personalized exercise prescriptions, which is important because exercise abilities (and therefore appropriate exercise intensity) vary considerably in patients with cancer (83). Animal-specific exercise prescriptions are challenging to implement in pre-clinical studies, but could be important, as findings reviewed here suggest that exercise intensity may differentially modulate tumor end points. For example, using a transgenic model of p53+/- MMTV-Wnt-1, Colbert et al. found that exposure to forced treadmill training of different intensities accelerated tumor growth rate compared to sedentary controls, with the highest tumor multiplicity in the lowest intensity group (26). Conversely, Almeida et al. reported diminishing returns in the context of increasing exercise intensity using two different swimming intensities (50% or 80% of maximal workload; CT₅₀ and CT₈₀, respectively, defined using a maximum load test conducted one week prior to exercise initiation) (47). CT₅₀ caused significant tumor growth inhibition whereas CT₈₀ was ineffective. Using a model of forced treadmill running, Woods et al. compared the effects of “moderate” vs “exhaustive” exercise prescriptions, with no differences in mammary tumor growth or overall incidence (40). In contrast, Malicka et al. reported a trend towards a negative correlation between exercise intensity (varied by progressively adjusting the speed of the treadmill [low: 0.48–1.34 km/h, moderate: 0.6–1.68 km/h, high: 0.72–2.0 km/h]) and tumor incidence and multiplicity, as well as an increase in tumor cell apoptosis across all exercise prescriptions, compared to controls (45). Kato et al. examined duration of prescription of forced treadmill running using a nitrilotriacetate-induced renal carcinoma model. Rats were assigned to exercise exposure for either 12 weeks or 40 weeks (29). Twelve weeks of exercise increased microcarcinoma incidence and multiplicity compared to sedentary controls with no differences (compared to control) in 40 weeks of exercise.

Schedule of Exercise Exposure—A critical question is whether exercise exposure prior to diagnosis improves disease outcomes compared to inactivity (prior to diagnosis). Similarly, do previously sedentary patients initiating exercise after diagnosis have superior outcomes compared to those remaining sedentary or decreasing exercise levels after diagnosis? Exploratory findings from epidemiological studies suggest that timing or initiation of exercise exposure could be important (84). Two studies have empirically investigated this question: Betof et al. and Shalamzari et al. found that exercise initiated after tumor transplantation (equivalent to post-diagnosis setting) inhibited tumor growth, independent of exposure to exercise prior to transplantation (65, 67). Similarly, Radak et al.

found slower tumor growth with exercise pre- as well as post-tumor implantation in comparison with no exercise after transplant or sedentary control (61). Such findings stress the importance of maintaining exercise post-diagnosis. Finally, whether the effects of exercise are different across the steps in the metastatic cascade (altered adhesion [invasion]; intravasation; survival in the circulation; extravasation, and seeding at a distant site [metastatic colonization] (85)) has not been investigated. Addressing this question has significant clinical implications, and further research in this area is warranted.

Common Methodological/Experimental Weaknesses

Several salient methodological issues across studies were identified (see Table 5). Of these, two common issues that require particular attention regarded selection of appropriate *in vivo* tumor models, and lack of mechanistic studies.

In Vivo Tumor Models—Preclinical oncology studies have generally focused on subcutaneous tumor implantation models. These models permit monitoring of tumor growth kinetics but do not accurately recapitulate the tissue microenvironment of the ectopic (orthotopic) or distant organ or mimic the natural evolution of cancer. In addition, blood flow to subcutaneous tumors may not reflect that of orthotopic tumors. As the benefits/risks of using subcutaneous models are beyond the scope of this review, here we will simply caution investigators against using subcutaneous tumors. This model is not a perfect surrogate for spontaneously occurring tumors.

The majority of studies reviewed herein investigated the effects of exercise on tumor growth characteristics in the primary organ site. In contrast, studies of metastasis, the primary cause of cancer-related death, has received limited attention to date. Additionally, 73% of the metastasis studies we reviewed used an intravenous (tail-vein) injection of tumor cells as a model of metastasis which evaluates the ability of tumor cells to survive in circulation and colonize an organ but does not enable investigation of the early steps in the metastatic cascade.(86) Indeed, three studies reported on tumor cell retention in the lungs following intravenous injection (71, 72, 74), with two studies reporting a decrease in tumor cell retention in exercising animals, but no change in the final number of metastases. Interestingly, MacNeil et al. reported that exercise did not affect tumor cell retention in the lung. These three studies differed with respect to type of exercise and number of tumor cells injected; thus no meaningful comparisons can be made between them. The impact of the methodological differences can only be evaluated by completing the studies-side-by-side, comparing all combinations of exercise/tumor inoculation models.

An alternative, and arguably more appropriate model of metastasis, is surgical excision of the primary tumor which stimulates spontaneous metastasis, predominantly to the lungs; Only one study to date (Yan et al.) has adopted this model to examine exercise effects (64); they reported a trend towards an inverse relationship between distance run and metastatic burden. Conversely, Jones et al. examined the metastasis response in a prostate cancer model by quantifying the mass of “non-contiguous external masses that were grossly visible independent from the primary prostate tissue.” In this model, exercise was not associated with a significant reduction in the number or weight of metastases (55).

Emerging technology has provided researchers with a considerable number of *in vivo* models beyond cell line xenografts, such as patient-derived xenografts (PDXs), syngeneic allografts, and genetically-engineered mouse models (GEMMs) (87), as well as non-rodent models (e.g., zebrafish, *Drosophila*). The advantages and disadvantages of each model have been reviewed elsewhere (88–91). Ultimately, no one *in vivo* model will be appropriate to address all exercise-oncology questions, with model selection being contingent on the scientific question at hand as well as translational importance.

The Importance of Appropriate Control Groups—Consideration of the nature of control (non-exercising) groups should not be overlooked. To the extent possible, control animals should be exposed to the same variables as exercise counterparts including aspects related to housing, transportation to different facilities, or procedures to reinforce exercise behavior (i.e., prodding, shock). Taken even further, animals could be housed in cages with locked wheels, placed on stationary treadmills, or made to stand in very shallow pools of water, depending upon the exercise regimens used. We stress that exposing control and experimental mice to different housing or environmental conditions is a study weakness. For example, one study housed control animals in unusually small cages (5 inches in diameter and 6 inches high) to restrict activity (54), whereas exercise treatment animals were not confined. The differences in housing size and daily movement could have either induced a stress response or altered the animals' resting metabolism, thereby affecting tumor growth. We advise researchers reviewing extant publications for planning of their own exercise oncology studies to consider whether controls were properly handled before modeling their own work off of previous studies.

Mechanistic Studies/Analyses—The majority of studies reviewed here examined the effects of exercise on tumor growth characteristics as evaluated by tumor volume or growth rates. While such end points are clearly important, subtle but important modulations of intratumoral physiological or biological alterations can be masked. For instance, our group observed differences in perfusion and expression of key factors regulating metabolism and hypoxia, despite comparable primary tumor growth rates between exercised and sedentary animals (57). Elegant studies by McCollough et al. demonstrated that exercise improved blood flow and oxygen delivery to orthotopic prostate tumors in rats, but not to the normal prostate tissue in either tumor-bearing or control rats (92). These changes were reflected by decreased hypoxia within the tumor during exercise. While in-depth explications of the systemic or local molecular mechanisms underpinning the exercise-tumor prevention/progression relationship remains in its infancy, studies such as this one may set the precedent for future mechanistic studies in this field. The current working hypothesis is that exercise modulates tumor progression via modulation of the host-tumor interaction (19). Tumor progression is regulated by complex, multifaceted interactions between the systemic milieu (host), tumor microenvironment, and cancer cells (93). The microenvironments of primary and metastatic tumors are subject to modulation by systemic and local growth/angiogenic factors, cytokines, hormones, and resident cells (94, 95). Factors such as hepatocyte growth factor (HGF), tumor necrosis factor (TNF), interleukin (IL)-6, insulin, and leptin (96–98) have already been associated with higher risk of recurrence and cancer-

specific mortality in a number of solid malignancies.(99) Clearly, manipulation of such factors by physical activity could alter aspects of the cancer continuum (100).

As reviewed here, multiple systemic factors are perturbed by exercise, including metabolism, inflammatory-immune, and reactive oxygen species-mediated pathways (101). The breadth of these factors likely contributes to the pleiotropic benefits of exercise in health and disease (102, 103) and likely the potential antitumor effects of exercise. (19) Notably, most cancer studies investigate single pathways in isolation, without consideration of overlap/synergism between pathways. Because the host/tumor interaction is modulated by numerous host-related factors and multiple pathways (100), an ideal study approach would be to investigate exercise effects on multiple pathways simultaneously. This would fill in the missing gaps of the “multi-targeted” effects of exercise. A recent analysis of pre-clinical exercise oncology studies by Pedersen et al. reported that only 30.7% evaluated systemic changes in animals (104). In addition, intratumoral signaling, and changes in tumor vascularity were examined by only 29.5% and 6.8%, respectively (104).

Future Recommendations

With a view towards future studies, we encourage the exercise-oncology field to consider certain guidelines for preclinical exercise-oncology research (see Table 5). However, we realize that it is impractical for all exercise-oncology experiments to standardize all study procedures. Nevertheless, it may behoove the field to develop a means of quantifying the exercise prescription applied, which will then permit comparisons between studies. Such an approach is readily used in the fields of radiation oncology (i.e., the biologically equivalent dose [BED], which allows comparison of different dose/fractionation schemes (105)) and hyperthermia (i.e., the cumulative equivalent minutes at 43°C [CEM43], which standardizes the thermal killing effect of hyperthermia, regardless of variations in heating efficiencies between tumors (106)). Establishing a biological equivalent exercise dose (BEED) would facilitate comparisons across studies as well as provide an opportunity to test elements of prescriptions such as different schedules, timing, and duration.

Exercise investigations should also strive to adopt the experimental procedures and models being utilized in the general tumor biology literature. Indeed, it appears that more recent studies reviewed here have started to utilize clinically-relevant tumor models, favoring transgenic mice and orthotopic tumors over carcinogen-induction. The general cancer field is also starting to favor the use of patient-derived xenograft (PDX) models. PDX have certain weaknesses, including increased heterogeneity, and a requirement for immunocompromised mice, but could represent an important step towards personalized medicine. Furthermore, PDXs do not accrue additional mutations through *in vitro* culture and tend to be slower growing than murine-derived tumor lines. Delayed growth curves may highlight slight changes in tumor growth following manipulations of exercise dose. GEM models should also be considered, especially in mechanistic studies. A second trend in cancer biology is use of biomarkers. The discovery of blood, imaging, and/or genomic biomarker(s) to predict or monitor exercise response is of obvious importance.

Finally, researchers must be cognizant of clinical care, and design studies that reflect the clinical scenario. For example, the vast majority of the current literature investigates the

effects of exercise as monotherapy. The majority of clinical scenarios would be applying exercise as an adjuvant therapy to surgery, radiation, chemotherapy, or immunotherapies. As such, it is important for the next generation of preclinical studies aiming to study tumor progression to evaluate the interaction between exercise and the pharmacodynamic or pharmacokinetic activity of conventional and novel therapies to guide the design and interpretation of clinical studies. We advise researchers to proceed with caution and carefully include all possible controls groups, because the additional therapies could introduce new confounding factors that complicate data interpretations. Conversely, studies on tumor incidence may be strengthened by eliminating additional factors such as concomitant manipulation of dietary composition.

Conclusion

A sound foundation of basic and translational studies will optimize the therapeutic potential of exercise on symptom control and clinical outcomes across the cancer continuum. Despite its importance, we found that the current evidence base is plagued by considerable methodological heterogeneity in all aspects of study design, end points, and efficacy thereby precluding meaningful comparisons and conclusions. To this end, we have provided an overview of methodological and data reporting standards that we hope will set the platform for the next generation of preclinical studies required for the continued development and progression of exercise-oncology research.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

LWJ is supported in part by grants from the National Cancer Institute, AKTIV Against Cancer, and the Memorial Sloan Kettering Cancer Center Support Grant/Core Grant (P30 CA008748). MWD is supported by NIH CA40355.

REFERENCES

1. Booth FW, Gordon SE, Carlson CJ, Hamilton MT. Waging war on modern chronic diseases: primary prevention through exercise biology. *Journal of applied physiology*. 2000; 88(2):774–787. [PubMed: 10658050]
2. Giannuzzi P, Mezzani A, Saner H, Bjornstad H, Fioretti P, Mendes M, et al. Physical activity for primary and secondary prevention. Position paper of the Working Group on Cardiac Rehabilitation and Exercise Physiology of the European Society of Cardiology. *European journal of cardiovascular prevention and rehabilitation : official journal of the European Society of Cardiology, Working Groups on Epidemiology & Prevention and Cardiac Rehabilitation and Exercise Physiology*. 2003; 10(5):319–327.
3. Hawley JA, Hargreaves M, Joyner MJ, Zierath JR. Integrative biology of exercise. *Cell*. 2014; 159(4):738–749. [PubMed: 25417152]
4. Warburton DE, Nicol CW, Bredin SS. Health benefits of physical activity: the evidence. *CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne*. 2006; 174(6):801–809.
5. Jones LW, Alfano CM. Exercise-oncology research: past, present, and future. *Acta oncologica*. 2013; 52(2):195–215. [PubMed: 23244677]
6. Friedenreich CM, Woolcott CG, McTiernan A, Ballard-Barbash R, Brant RF, Stanczyk FZ, et al. Alberta physical activity and breast cancer prevention trial: sex hormone changes in a year-long

- exercise intervention among postmenopausal women. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2010; 28(9):1458–1466. [PubMed: 20159820]
7. Lynch BM, Friedenreich CM, Winkler EA, Healy GN, Vallance JK, Eakin EG, et al. Associations of objectively assessed physical activity and sedentary time with biomarkers of breast cancer risk in postmenopausal women: findings from NHANES (2003–2006). *Breast cancer research and treatment*. 2011; 130(1):183–194. [PubMed: 21553294]
 8. Gammon MD, Schoenberg JB, Britton JA, Kelsey JL, Coates RJ, Brogan D, et al. Recreational physical activity and breast cancer risk among women under age 45 years. *American journal of epidemiology*. 1998; 147(3):273–280. [PubMed: 9482502]
 9. Bernstein L, Henderson BE, Hanisch R, Sullivan-Halley J, Ross RK. Physical exercise and reduced risk of breast cancer in young women. *Journal of the National Cancer Institute*. 1994; 86(18):1403–1408. [PubMed: 8072034]
 10. Byers T, Nestle M, McTiernan A, Doyle C, Currie-Williams A, Gansler T, et al. American Cancer Society guidelines on nutrition and physical activity for cancer prevention: Reducing the risk of cancer with healthy food choices and physical activity. *CA: a cancer journal for clinicians*. 2002; 52(2):92–119. [PubMed: 11929008]
 11. Kushi LH, Doyle C, McCullough M, Rock CL, Demark-Wahnefried W, Bandera EV, et al. American Cancer Society Guidelines on nutrition and physical activity for cancer prevention: reducing the risk of cancer with healthy food choices and physical activity. *CA: a cancer journal for clinicians*. 2012; 62(1):30–67. [PubMed: 22237782]
 12. Friedenreich CM, Neilson HK, O'Reilly R, Duha A, Yasui Y, Morielli AR, et al. Effects of a High vs Moderate Volume of Aerobic Exercise on Adiposity Outcomes in Postmenopausal Women: A Randomized Clinical Trial. *JAMA oncology*. 2015
 13. Irwin ML, Yasui Y, Ulrich CM, Bowen D, Rudolph RE, Schwartz RS, et al. Effect of exercise on total and intra-abdominal body fat in postmenopausal women: a randomized controlled trial. *JAMA : the journal of the American Medical Association*. 2003; 289(3):323–330. [PubMed: 12525233]
 14. McTiernan A, Ulrich CM, Yancey D, Slate S, Nakamura H, Oestreicher N, et al. The Physical Activity for Total Health (PATH) Study: rationale and design. *Medicine and science in sports and exercise*. 1999; 31(9):1307–1312. [PubMed: 10487373]
 15. Campbell KL, McTiernan A, Li SS, Sorensen BE, Yasui Y, Lampe JW, et al. Effect of a 12-month exercise intervention on the apoptotic regulating proteins Bax and Bcl-2 in colon crypts: a randomized controlled trial. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2007; 16(9):1767–1774.
 16. Mishra SI, Scherer RW, Geigle PM, Berlanstein DR, Topaloglu O, Gotay CC, et al. Exercise interventions on health-related quality of life for cancer survivors. *The Cochrane database of systematic reviews*. 2012; 8:CD007566.
 17. Schmitz KH, Holtzman J, Courneya KS, Masse LC, Duval S, Kane R. Controlled physical activity trials in cancer survivors: a systematic review and meta-analysis. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2005; 14(7):1588–1595.
 18. Jones LW, Dewhirst MW. Therapeutic properties of aerobic training after a cancer diagnosis: more than a one-trick pony? *Journal of the National Cancer Institute*. 2014; 106(4) dju042.
 19. Betof AS, Dewhirst MW, Jones LW. Effects and potential mechanisms of exercise training on cancer progression: a translational perspective. *Brain, behavior, and immunity*. 2013; 30(Suppl):S75–S87.
 20. Ballard-Barbash R, Friedenreich CM, Courneya KS, Siddiqi SM, McTiernan A, Alfano CM. Physical activity, biomarkers, and disease outcomes in cancer survivors: a systematic review. *Journal of the National Cancer Institute*. 2012; 104(11):815–840. [PubMed: 22570317]
 21. Gorski DH. Integrative oncology: really the best of both worlds? *Nature reviews Cancer*. 2014; 14(10)
 22. Jones LW. Precision Oncology Framework for Investigation of Exercise as Treatment for Cancer. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2015

23. Abdalla DR, Murta EF, Michelin MA. The influence of physical activity on the profile of immune response cells and cytokine synthesis in mice with experimental breast tumors induced by 7,12-dimethylbenzanthracene. *European journal of cancer prevention : the official journal of the European Cancer Prevention Organisation*. 2013; 22(3):251–258.
24. Alessio HM, Schweitzer NB, Snedden AM, Callahan P, Hagerman AE. Revisiting influences on tumor development focusing on laboratory housing. *Journal of the American Association for Laboratory Animal Science : JAALAS*. 2009; 48(3):258–262. [PubMed: 19476713]
25. Andrianopoulos G, Nelson RL, Bombeck CT, Souza G. The influence of physical activity in 1,2-dimethylhydrazine induced colon carcinogenesis in the rat. *Anticancer research*. 1987; 7(4B):849–852. [PubMed: 3674772]
26. Colbert LH, Westerlind KC, Perkins SN, Haines DC, Berrigan D, Donehower LA, et al. Exercise effects on tumorigenesis in a p53-deficient mouse model of breast cancer. *Medicine and science in sports and exercise*. 2009; 41(8):1597–1605. [PubMed: 19568200]
27. Esser KA, Harpole CE, Prins GS, Diamond AM. Physical activity reduces prostate carcinogenesis in a transgenic model. *The Prostate*. 2009; 69(13):1372–1377. [PubMed: 19489028]
28. Ikuyama T, Watanabe T, Minegishi Y, Osanai H. Effect of voluntary exercise on 3'-methyl-4-dimethylaminoazobenzene-induced hepatomas in male Jc1:Wistar rats. *Proceedings of the Society for Experimental Biology and Medicine Society for Experimental Biology and Medicine (New York, NY)*. 1993; 204(2):211–215.
29. Kato T, Kawaguchi H, Miyoshi N, Aoyama K, Komatsu M, Horiuchi M, et al. Effect of habitual exercise on renal carcinogenesis by ferric nitrilotriacetate. *Environmental health and preventive medicine*. 2011; 16(4):232–238. [PubMed: 21431792]
30. Lunz W, Peluzio MC, Dias CM, Moreira AP, Natali AJ. Long-term aerobic swimming training by rats reduces the number of aberrant crypt foci in 1,2-dimethylhydrazine-induced colon cancer. *Brazilian journal of medical and biological research = Revista brasileira de pesquisas medicas e biologicas / Sociedade Brasileira de Biofisica [et al]*. 2008; 41(11):1000–1004.
31. Mann PB, Jiang W, Zhu Z, Wolfe P, McTiernan A, Thompson HJ. Wheel running, skeletal muscle aerobic capacity and 1-methyl-1-nitrosourea induced mammary carcinogenesis in the rat. *Carcinogenesis*. 2010; 31(7):1279–1283. [PubMed: 20299525]
32. Paceli RB, Cal RN, dos Santos CH, Cordeiro JA, Neiva CM, Nagamine KK, et al. The influence of physical activity in the progression of experimental lung cancer in mice. *Pathology, research and practice*. 2012; 208(7):377–381.
33. Reddy BS, Sugie S, Lowenfels A. Effect of voluntary exercise on azoxymethane-induced colon carcinogenesis in male F344 rats. *Cancer research*. 1988; 48(24 Pt 1):7079–7081. [PubMed: 3191484]
34. Saez Mdel C, Barriga C, Garcia JJ, Rodriguez AB, Ortega E. Exercise-induced stress enhances mammary tumor growth in rats: beneficial effect of the hormone melatonin. *Molecular and cellular biochemistry*. 2007; 294(1–2):19–24. [PubMed: 17136443]
35. Sugie S, Reddy BS, Lowenfels A, Tanaka T, Mori H. Effect of voluntary exercise on azoxymethane-induced hepatocarcinogenesis in male F344 rats. *Cancer letters*. 1992; 63(1):67–72. [PubMed: 1555208]
36. Thorling EB, Jacobsen NO, Overvad K. Effect of exercise on intestinal tumour development in the male Fischer rat after exposure to azoxymethane. *European journal of cancer prevention : the official journal of the European Cancer Prevention Organisation*. 1993; 2(1):77–82.
37. Westerlind KC, McCarty HL, Schultheiss PC, Story R, Reed AH, Baier ML, et al. Moderate exercise training slows mammary tumour growth in adolescent rats. *European journal of cancer prevention : the official journal of the European Cancer Prevention Organisation*. 2003; 12(4):281–287.
38. Whittal KS, Parkhouse WS. Exercise during adolescence and its effects on mammary gland development, proliferation, and nitrosomethylurea (NMU) induced tumorigenesis in rats. *Breast cancer research and treatment*. 1996; 37(1):21–27. [PubMed: 8750524]
39. Whittal-Strange KS, Chadan S, Parkhouse WS. Exercise during puberty and NMU induced mammary tumorigenesis in rats. *Breast cancer research and treatment*. 1998; 47(1):1–8. [PubMed: 9493970]

40. Woods JA, Davis JM, Kohut ML, Ghaffar A, Mayer EP, Pate RR. Effects of exercise on the immune response to cancer. *Medicine and science in sports and exercise*. 1994; 26(9):1109–1115. [PubMed: 7808244]
41. Zhu Z, Jiang W, McGinley JN, Thompson HJ. Energetics and mammary carcinogenesis: effects of moderate-intensity running and energy intake on cellular processes and molecular mechanisms in rats. *Journal of applied physiology*. 2009; 106(3):911–918. [PubMed: 19095749]
42. Zhu Z, Jiang W, Sells JL, Neil ES, McGinley JN, Thompson HJ. Effect of nonmotorized wheel running on mammary carcinogenesis: circulating biomarkers, cellular processes, and molecular mechanisms in rats. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2008; 17(8):1920–1929.
43. Zhu Z, Jiang W, Zacher JH, Neil ES, McGinley JN, Thompson HJ. Effects of energy restriction and wheel running on mammary carcinogenesis and host systemic factors in a rat model. *Cancer prevention research*. 2012; 5(3):414–422. [PubMed: 22246620]
44. Zielinski MR, Muenchow M, Wallig MA, Horn PL, Woods JA. Exercise delays allogeneic tumor growth and reduces intratumoral inflammation and vascularization. *Journal of applied physiology*. 2004; 96(6):2249–2256. [PubMed: 15020578]
45. Malicka I, Siewierska K, Pula B, Kobierzycki C, Haus D, Paslawska U, et al. The effect of physical training on the N-methyl-N-nitrosourea-induced mammary carcinogenesis of Sprague-Dawley rats. *Experimental biology and medicine* (Maywood, NJ). 2015
46. Piguat AC, Saran U, Simillion C, Keller I, Terracciano L, Reeves HL, et al. Regular exercise decreases liver tumors development in hepatocyte-specific PTEN-deficient mice independently of steatosis. *Journal of hepatology*. 2015; 62(6):1296–1303. [PubMed: 25623824]
47. Almeida PW, Gomes-Filho A, Ferreira AJ, Rodrigues CE, Dias-Peixoto MF, Russo RC, et al. Swim training suppresses tumor growth in mice. *Journal of applied physiology*. 2009; 107(1):261–265. [PubMed: 19478194]
48. Bacurau AV, Belmonte MA, Navarro F, Moraes MR, Pontes FL Jr, Pesquero JL, et al. Effect of a high-intensity exercise training on the metabolism and function of macrophages and lymphocytes of walker 256 tumor bearing rats. *Experimental biology and medicine* (Maywood, NJ). 2007; 232(10):1289–1299.
49. Bacurau RF, Belmonte MA, Seelaender MC, Costa Rosa LF. Effect of a moderate intensity exercise training protocol on the metabolism of macrophages and lymphocytes of tumour-bearing rats. *Cell biochemistry and function*. 2000; 18(4):249–258. [PubMed: 11180287]
50. Baracos VE. Exercise inhibits progressive growth of the Morris hepatoma 7777 in male and female rats. *Canadian journal of physiology and pharmacology*. 1989; 67(8):864–870. [PubMed: 2598121]
51. Daneryd P, Hafstrom L, Svanberg E, Karlberg I. Insulin sensitivity, hormonal levels and skeletal muscle protein metabolism in tumour-bearing exercising rats. *European journal of cancer*. 1995; 31A(1):97–103. [PubMed: 7535077]
52. Daneryd P, Westin T, Edstrom S, Soussi B. Tumour purine nucleotides and cell proliferation in response to exercise in rats. *European journal of cancer*. 1995; 31a(13–14):2309–2312. [PubMed: 8652261]
53. Gershbein LL, Benuck I, Shurrager PS. Influence of stress of lesion growth and on survival of animals bearing parenteral and intracerebral leukemia L1210 and Walker tumors. *Oncology*. 1974; 30(5):429–435. [PubMed: 4462061]
54. Hoffman SA, Paschkis KE, Debias DA, Cantarow A, Williams TL. The influence of exercise on the growth of transplanted rat tumors. *Cancer research*. 1962; 22:597–599. [PubMed: 14042313]
55. Jones LW, Antonelli J, Masko EM, Broadwater G, Lascola CD, Fels D, et al. Exercise modulation of the host-tumor interaction in an orthotopic model of murine prostate cancer. *Journal of applied physiology*. 2012; 113(2):263–272. [PubMed: 22604887]
56. Jones LW, Eves ND, Courneya KS, Chiu BK, Baracos VE, Hanson J, et al. Effects of exercise training on antitumor efficacy of doxorubicin in MDA-MB-231 breast cancer xenografts. *Clinical cancer research : an official journal of the American Association for Cancer Research*. 2005; 11(18):6695–6698. [PubMed: 16166449]

57. Jones LW, Viglianti BL, Tashjian JA, Kothadia SM, Keir ST, Freedland SJ, et al. Effect of aerobic exercise on tumor physiology in an animal model of human breast cancer. *Journal of applied physiology*. 2010; 108(2):343–348. [PubMed: 19959769]
58. Lira FS, Tavares FL, Yamashita AS, Koyama CH, Alves MJ, Caperuto EC, et al. Effect of endurance training upon lipid metabolism in the liver of cachectic tumour-bearing rats. *Cell biochemistry and function*. 2008; 26(6):701–708. [PubMed: 18636434]
59. Murphy EA, Davis JM, Barrilleaux TL, McClellan JL, Steiner JL, Carmichael MD, et al. Benefits of exercise training on breast cancer progression and inflammation in C3(1)SV40Tag mice. *Cytokine*. 2011; 55(2):274–279. [PubMed: 21600785]
60. Newton G. TUMOR SUSCEPTIBILITY IN RATS: ROLE OF INFANTILE MANIPULATION AND LATER EXERCISE. *Psychological reports*. 1965; 16:127–132. [PubMed: 14283948]
61. Radak Z, Gaal D, Taylor AW, Kaneko T, Tahara S, Nakamoto H, et al. Attenuation of the development of murine solid leukemia tumor by physical exercise. *Antioxidants & redox signaling*. 2002; 4(1):213–219. [PubMed: 11970855]
62. Sasvari M, Taylor AW, Gaal D, Radak Z. The effect of regular exercise on development of sarcoma tumor and oxidative damage in mice liver. *Journal of sports science & medicine*. 2011; 10(1):93–96. [PubMed: 24149300]
63. Uhlenbruck G, Order U. Can endurance sports stimulate immune mechanisms against cancer and metastasis? *International journal of sports medicine*. 1991; 12(Suppl 1):S63–S68. [PubMed: 1894399]
64. Yan L, Demars LC. Effects of non-motorized voluntary running on experimental and spontaneous metastasis in mice. *Anticancer research*. 2011; 31(10):3337–3344. [PubMed: 21965744]
65. Betof AS, Lascola CD, Weitzel D, Landon C, Scarbrough PM, Devi GR, et al. Modulation of murine breast tumor vascularity, hypoxia and chemotherapeutic response by exercise. *Journal of the National Cancer Institute*. 2015; 107(5)
66. Goh J, Endicott E, Ladiges WC. Pre-tumor exercise decreases breast cancer in old mice in a distance-dependent manner. *American journal of cancer research*. 2014; 4(4):378–384. [PubMed: 25057440]
67. Shalamzari SA, Agha-Alinejad H, Alizadeh S, Shahbazi S, Khatib ZK, Kazemi A, et al. The effect of exercise training on the level of tissue IL-6 and vascular endothelial growth factor in breast cancer bearing mice. *Iranian journal of basic medical sciences*. 2014; 17(4):231–258. [PubMed: 24904714]
68. Aveseh M, Nikooie R, Aminaie M. Exercise-induced changes in tumour LDH-B and MCT1 expression are modulated by oestrogen-related receptor alpha in breast cancer-bearing BALB/c mice. *The Journal of physiology*. 2015
69. Guerit J, Lefeuvre-Orfila L, Vincent S, Cretual A, Ravanat JL, Gratas-Delamarche A, et al. Exercise training combined with antioxidant supplementation prevents the antiproliferative activity of their single treatment in prostate cancer through inhibition of redox adaptation. *Free radical biology & medicine*. 2014; 77:95–105. [PubMed: 25236740]
70. Hoffman-Goetz L, May KM, Arumugam Y. Exercise training and mouse mammary tumour metastasis. *Anticancer research*. 1994; 14(6B):2627–2631. [PubMed: 7872692]
71. Hoffmann-Goetz L, MacNeil B, Arumugam Y. Tissue distribution of radiolabelled tumor cells in wheel exercised and sedentary mice. *International journal of sports medicine*. 1994; 15(5):249–253. [PubMed: 7960319]
72. Jadeski L, Hoffman-Goetz L. Exercise and in vivo natural cytotoxicity against tumour cells of varying metastatic capacity. *Clinical & experimental metastasis*. 1996; 14(2):138–144. [PubMed: 8605727]
73. MacNeil B, Hoffman-Goetz L. Exercise training and tumour metastasis in mice: influence of time of exercise onset. *Anticancer research*. 1993; 13(6A):2085–2088. [PubMed: 8297117]
74. MacNeil B, Hoffman-Goetz L. Effect of exercise on natural cytotoxicity and pulmonary tumor metastases in mice. *Medicine and science in sports and exercise*. 1993; 25(8):922–928. [PubMed: 8371653]

75. Higgins KA, Park D, Lee GY, Curran WJ, Deng X. Exercise-induced lung cancer regression: mechanistic findings from a mouse model. *Cancer*. 2014; 120(21):3302–3310. [PubMed: 24989479]
76. Zhao G, Zhou S, Davie A, Su Q. Effects of moderate and high intensity exercise on T1/T2 balance. *Exercise immunology review*. 2012; 18:98–114. [PubMed: 22876723]
77. Roberts CJ, Stuhr KL, Hillard CJ. Swim stress differentially affects limbic contents of 2-arachidonoylglycerol and 2-oleoylglycerol. *Neuroscience*. 2012; 204:74–82. [PubMed: 22192839]
78. Fuss J, Ben Abdallah NM, Vogt MA, Touma C, Pacifici PG, Palme R, et al. Voluntary exercise induces anxiety-like behavior in adult C57BL/6J mice correlating with hippocampal neurogenesis. *Hippocampus*. 2010; 20(3):364–376. [PubMed: 19452518]
79. Richter SH, Gass P, Fuss J. Resting Is Rusting: A Critical View on Rodent Wheel-Running Behavior. *The Neuroscientist : a review journal bringing neurobiology, neurology and psychiatry*. 2014; 20(4):313–325.
80. Hackney AC, Viru A. Twenty-four-hour cortisol response to multiple daily exercise sessions of moderate and high intensity. *Clinical physiology*. 1999; 19(2):178–182. [PubMed: 10200900]
81. Thaker PH, Han LY, Kamat AA, Arevalo JM, Takahashi R, Lu C, et al. Chronic stress promotes tumor growth and angiogenesis in a mouse model of ovarian carcinoma. *Nature medicine*. 2006; 12(8):939–944.
82. Zen M, Canova M, Campana C, Bettio S, Nalotto L, Rampudda M, et al. The kaleidoscope of glucorticoid effects on immune system. *Autoimmunity reviews*. 2011; 10(6):305–310. [PubMed: 21224015]
83. Sasso JP, Eves ND, Christensen JF, Koelwyn GJ, Scott J, Jones LW. A framework for prescription in exercise-oncology research. *Journal of cachexia, sarcopenia and muscle*. 2015; 6(2):115–124.
84. Irwin ML, Smith AW, McTiernan A, Ballard-Barbash R, Cronin K, Gilliland FD, et al. Influence of pre- and postdiagnosis physical activity on mortality in breast cancer survivors: the health, eating, activity, and lifestyle study. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2008; 26(24):3958–3964. [PubMed: 18711185]
85. Mina LA, Sledge GW Jr. Rethinking the metastatic cascade as a therapeutic target. *Nature reviews Clinical oncology*. 2011; 8(6):325–332.
86. Chambers AF, Groom AC, MacDonald IC. Dissemination and growth of cancer cells in metastatic sites. *Nature reviews Cancer*. 2002; 2(8):563–572. [PubMed: 12154349]
87. Day CP, Merlino G, Van Dyke T. Preclinical mouse cancer models: a maze of opportunities and challenges. *Cell*. 2015; 163(1):39–53. [PubMed: 26406370]
88. Gonzalez C. *Drosophila melanogaster*: a model and a tool to investigate malignancy and identify new therapeutics. *Nature reviews Cancer*. 2013; 13(3):172–183. [PubMed: 23388617]
89. Gould SE, Junttila MR, de Sauvage FJ. Translational value of mouse models in oncology drug development. *Nature medicine*. 2015; 21(5):431–439.
90. Sharpless NE, Depinho RA. The mighty mouse: genetically engineered mouse models in cancer drug development. *Nature reviews Drug discovery*. 2006; 5(9):741–754. [PubMed: 16915232]
91. White R, Rose K, Zon L. Zebrafish cancer: the state of the art and the path forward. *Nature reviews Cancer*. 2013; 13(9):624–636. [PubMed: 23969693]
92. McCullough DJ, Stabley JN, Siemann DW, Behnke BJ. Modulation of blood flow, hypoxia, and vascular function in orthotopic prostate tumors during exercise. *Journal of the National Cancer Institute*. 2014; 106(4) dju036.
93. Straussman R, Morikawa T, Shee K, Barzily-Rokni M, Qian ZR, Du J, et al. Tumour micro-environment elicits innate resistance to RAF inhibitors through HGF secretion. *Nature*. 2012; 487(7408):500–504. [PubMed: 22763439]
94. McAllister SS, Gifford AM, Greiner AL, Kelleher SP, Saelzler MP, Ince TA, et al. Systemic endocrine instigation of indolent tumor growth requires osteopontin. *Cell*. 2008; 133(6):994–1005. [PubMed: 18555776]
95. McAllister SS, Weinberg RA. Tumor-host interactions: a far-reaching relationship. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2010; 28(26):4022–4028. [PubMed: 20644094]

96. Goodwin PJ, Ennis M, Pritchard KI, Trudeau ME, Koo J, Taylor SK, et al. Insulin- and obesity-related variables in early-stage breast cancer: correlations and time course of prognostic associations. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2012; 30(2):164–171. [PubMed: 22162568]
97. Sheen-Chen SM, Chen WJ, Eng HL, Chou FF. Serum concentration of tumor necrosis factor in patients with breast cancer. *Breast cancer research and treatment*. 1997; 43(3):211–215. [PubMed: 9150900]
98. Sheen-Chen SM, Liu YW, Eng HL, Chou FF. Serum levels of hepatocyte growth factor in patients with breast cancer. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2005; 14(3):715–717.
99. Hursting SD, Berger NA. Energy balance, host-related factors, and cancer progression. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2010; 28(26):4058–4065. [PubMed: 20697088]
100. Glass OK, Inman BA, Broadwater G, Courneya KS, Mackey JR, Goruk S, et al. Effect of aerobic training on the host systemic milieu in patients with solid tumours: an exploratory correlative study. *British journal of cancer*. 2015; 112(5):825–831. [PubMed: 25584487]
101. Joyner MJ, Green DJ. Exercise protects the cardiovascular system: effects beyond traditional risk factors. *The Journal of physiology*. 2009; 587(Pt 23):5551–5558. [PubMed: 19736305]
102. Kodama S, Saito K, Tanaka S, Maki M, Yachi Y, Asumi M, et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA : the journal of the American Medical Association*. 2009; 301(19):2024–2035. [PubMed: 19454641]
103. Lee IM, Shiroma EJ, Lobelo F, Puska P, Blair SN, Katzmarzyk PT, et al. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet*. 2012; 380(9838):219–229. [PubMed: 22818936]
104. Pedersen L, Christensen JF, Hojman P. Effects of exercise on tumor physiology and metabolism. *Cancer journal*. 2015; 21(2):111–116.
105. Fowler JF. 21 years of biologically effective dose. *The British journal of radiology*. 2010; 83(991):554–568. [PubMed: 20603408]
106. Dewhirst MW, Viglianti BL, Lora-Michiels M, Hanson M, Hoopes PJ. Basic principles of thermal dosimetry and thermal thresholds for tissue damage from hyperthermia. *International journal of hyperthermia : the official journal of European Society for Hyperthermic Oncology, North American Hyperthermia Group*. 2003; 19(3):267–294.




Figure 1. Postulated intratumoral and systemic mechanisms underlying exercise oncology across the cancer continuum. Note that few reports combined measurements in tumor with systemic changes. The link between intratumoral changes and systemic changes is largely unknown.




Figure 2. Hypothesized pathways by which endurance exercise may impact tumor progression and metastasis. Key host tissues such as skeletal muscle, adipose tissue, bone marrow, and the liver mediate the effect of exercise on a variety of systemic pathways. Exercise-induced alterations in systemic and circulating factors, in turn, influences ligand availability in the tumor microenvironment which alters cellular signaling modulating the hallmarks of cancer.

Table 1

Tumor Incidence Studies

Study	Reference	Rodent Model	Tumor Type / Induction model	METHODS			RESULTS
				Exercise Modality	Exercise Protocol		
					Exercise Prescription Freq/week Dur Intens Length	Exercise Initiation	
Andrianopoulos, 1987	25	5w old male Sprague-Dawley rats	Intestinal / DMH, i.p. q1w for 6w	Voluntary wheel running	Wheel running	1w prior to first injection	Tumor Incidence Results -Tumors present in 18/20 SED rats and 6/11 EX rats
Reddy, 1988	33	Male F344 rats	Intestinal / Subcutaneous AOM 15mg/kg BW, q1wx2w at 7 wk of age.	Voluntary wheel running	Wheel running for 38 weeks	4d post-AOM	-EX ↓ incidence and multiplicity adenocarcinomas, and liver foci. of colon and
Sugie, 1992	35	5w old F-344 rats	Hepatocellular / 15 mg/kg AOM s.c. q1w for 2w	Voluntary wheel running	38w	4d post-injection.	-Liver tumors noted in 7% of AOM treated SED only.
Ikuyama, 1993	28	Jcl:Wistar rats	Hepatoma / 0.0177 g/day/kg BW dietary 3'-Me-DAB for 35 weeks.	Voluntary wheel running	Wheel running for 62 weeks, using food as a reward for achieving specified distances	17w prior to dietary 3'-Me-DAB	-65% reduction in tumor incidence.
Zhu, 2008	42	21d old female Sprague-Dawley rats	Mammary / 25 or 50 mg/kg MNU, i.p.	Voluntary wheel running	Voluntary wheel running for 8w	1w post-injection	-84.5% incidence vs 98.1%, (SED vs. EX)
Esser, 2009	27	C3(1)Tag mice	Prostate / Transgenic mouse model	Voluntary wheel running	Wheel running for 10 weeks Data analyzed based on mice that ran >5K or <5K a day	10w of age	-18% (>5K) and 55% (<5K) of animals with high grade neoplasia at 20 weeks -57% reduction in incidence of high grade neoplasia in >5K vs <5K mice
Alessio, 2009	24	3w old female Sprague-Dawley rats	Spontaneous tumors	Voluntary wheel running or activity box	Wheel: every other day, 24h access throughout animals' life Activity box (PA): 1h in large activity box twice a week for life.	Spontaneous tumors, animals followed for life.	-During weeks 60-120, 38% of EX rats were tumor-bearing animals (vs. 42% PA rats and 54% SED rats). -At week 88, tumor multiplicity was 0.69 for EX animals, 0.75 for PA, and 0.96 for SED.
Colbert, 2009	26	Female heterozygous (p53+/-); MMTV-Wnt-1 transgenic mice	Mammary / Transgenic	Voluntary wheel running and forced treadmill running	Treadmill running: TREX 1: 5x(45min) 20m/min at 5% grade until completion TREX 2: 5x(45min) 24m/min at 5% grade until completion	11w of age	-Tumor incidence ↑ in wheel running mice by 32%

Study	Reference	Rodent Model	Tumor Type / Induction model	METHODS			RESULTS
				Exercise Modality	Exercise Protocol		
					Exercise Prescription Freq/week Dur Intensity Length	Exercise Initiation	
Mann, 2010	31	21d old female Sprague-Dawley rats	Mammary / 50 mg/kg MNU i.p.	Voluntary non-motorized and motorized wheel running	Voluntary wheel running with 24 hour access.	1w post-injection	-96%, 74% and 70% incidence in controls, non-motorized and motorized mice, respectively
Zhu, 2012	43	21d old female Sprague-Dawley rats	Mammary / 50 mg/kg MNU i.p.	Voluntary wheel running at a fixed daily distance	Three levels: WR-High: maximum 3500m/d WR-Low: maximum 1750m/d. SED control.	1w post-injection	-97% tumor incidence in controls, 80% tumor incidence in WR-High and -Cannot compare WR-High and WR-Low, because dietary energy restriction was applied to WR-Low only
Thorling, 1993	36	5w old male Fischer rats	Intestinal / 15mg/kg AOM s.c. on Days 1, 4 and 8.	Forced treadmill running	5x(2km/day) 7m/min 38w. First week was acclimatization.	3d after last injection	-EX ↓ colon neoplasia incidence, 53% vs 78% in
Woods, 1994	40	6w old male C3H/HeN mice	Mammary / 2.5 × 10 ⁵ mammary SCA-1 cells s.c. in the back.	Forced treadmill running	Treadmill running: Moderate: 7x(30min) 18m/min at 5% grade 1w. Exhaustive: 7x(varied 18m/min for 30min, then 3m/min↑ every 30min until exhausted) 1w	3d prior to injection.	-Increase in tumor incidence at Day 7 in both EX groups, but no differences in any subsequent time points.
Whittal, 1996	38	21d old female Sprague-Dawley Rats	Mammary / 50 mg/kg MNU i.p. at 50d of age	Forced treadmill running	Progressive training to 5x(60min) 18 m/min at 15% grade 4w	21d of age (29d prior to injection)	-Incidence/multiplicity at 24w post-NMU: 58 tumors in SED rats, 33 tumors in EX rats -No significant difference in latency or incidence
Whittal-Strange, 1998	39	21d old female Sprague-Dawley rats	Mammary / 37.5mg/kg MNU i.p. at 50d of age, (1 day after last bout of EX)	Forced treadmill running	Progressive training to 5x(60min) 18 m/min at 15% grade 4w	21d of age (29d prior to injection)	-Incidences of carcinomas, high grade and low grade tumors were 29.2%, 10.4%, and 25% in SED, and 38%, and 14.3% and 21.4% in EX
Westerlind, 2003	37	21d old female Sprague-Dawley rats	Mammary / 25 or 50 mg/kg MNU, i.p.	Forced treadmill running	Week 1: 5x 10-15min 30m/min 1w. Week 2-9: 5x(30min 23-25m/min 8w	1w post-injection	-↑ Latency in EX (35.8d vs. 33.1d). -No difference in median tumor-free survival time was observed in the EX versus sham-EX (SHAM), nor

Study	Reference	Rodent Model	Tumor Type / Induction model	METHODS			RESULTS
				Exercise Modality	Exercise Protocol		
					Exercise Prescription Freq/week Dur Intens Length	Exercise Initiation	
Zielinski, 2004	44	6–8w old female BALB/c mice	Neoplastic Lymphoid cells / 2×10^7 EL-4 cells s.c. in the back behind the neck	Forced treadmill running	7× 3h or until volitional fatigue/gradually increasing speed, 20–40m/min, 5% grade 5–14d	First session immediately before injection.	were there any differences in multiplicity at either a high or moderate dose of MNU -EX ↓ tumor appearance
Zhu, 2009	41	21d old female Sprague-Dawley rats	Mammary / 50 mg/kg MNU, i.p.	Forced wheel running	Motorized running wheel; details not provided	1w post-injection	-66.7% and 92.6% incidence in EX and SED
Kato, 2011	29	5w old male Fischer 344 rats	Renal / 5mg/kg BW Fe-NTA i.p. once a day for 7 days, then 10mg/kg Fe-NTA 3×/wk for 11w.	Forced treadmill running	Short-term: 15m 8m/min, 0% 12w. Long-term: 15m 8m/min, 0% 12w; then 5× 30m 8m/min, 0% 12w for a total of 24w training Exercise continued until 40 weeks, but at lower intensity to account for the decline in rat health.	Training done 15m before each injection	-No differences in number of rats with nodules, nodules/rat or mean area of nodules. -Short-term EX ↑ rats with microcarcinomas, - Long-term EX ↓ rats cinomas, karyomegalic cells and degenerative tubules compared to with microcar short-term.
Malicka, 2015	45	4w old female Sprague Dawley rats	Mammary / 180 mg/kg MNU i.p.	Forced treadmill running	Low intensity (LIT): 5× 10–35min 0.48–1.34 km/h 12w Moderate intensity (MIT): 5× 10–35min 0.6–1.68 km/h 12w High intensity (HIT): 5× 10–35min 0.72–2.0 km/h 12w	Immediately after MNU injection	-Incidence 64%, 67%, 40% and 43% in SED, LIT, MIT and HIT groups, respectively (<i>Not significant</i>) -Multiplicity: Rats with tumors had an average of 2.4, 1.6, 1 and 1 tumors in SED, LIT, MIT and HIT groups, respectively (<i>Statistics not performed</i>)
Pignat, 2015	46	7–9w old male AlbCrePten flox/flox mice	Hepatocellular carcinoma / Transgenic	Forced treadmill running	5w acclimation period followed by: 5× 60m 12.5m/min 27w	7–9w of age	-Tumor incidence 100% in SED vs. 71% in EX
Lunz, 2008	30	11w old male Wistar Rats	Intestinal / 4 s.c. injections DMH 3 days	Forced swimming with 0%,	Week 1–2: 5× 5–20min load 2w	24h post first injection	-No difference in tumor incidence

Study	Reference	Rodent Model	Tumor Type / Induction model	METHODS			RESULTS
				Exercise Modality	Exercise Prescription Freq/week Dur Intens Length	Exercise Initiation	
			apart.	2% or 4% BW load	Week 3-5: 5x[5-20min] + load 3w Week 6-35: 5x[20min] +load 30w		Tumor Incidence Results -Aerobic swimming training with 2% body weight of load protected against the DMH-induced preneoplastic colon lesions, but not against tumor development in the rat
Paceli, 2012	32	Adult male Balb/c mice	Lung / 1.5mg/kg BW urethane i.p. twice, 2 days apart	Forced swimming	Aerobic: 4x[20m]-[19w in 5 days Anaerobic: 3x[20m (2m swimming/2m resting)]/progressive loading of 5-20%BW 20w	Within a week after injection	-No significant effects of aerobic training on lung cancer incidence. -Aerobic training resulted in 8 lung nodules per animal vs 52 in the control. <i>Not significant</i> . -Median control nodules was 2.0, median aerobic control nodules was 0.0. <i>Not significant</i> .
Abdalla, 2013	23	8w old female Balb/c mice	Mammary / 1mg/ml DMBA p.o. once weekly for 6 weeks	Forced swimming	5x[45m]-[8 weeks Water temperature 30+/-4°C	Same as tumor initiation	-EX ↓ tumor incidence
Sáez Mdel, 2007	34	50d old female Sprague-Dawley rats	Mammary / 5mg/w DMBA gastric intubation for 4w	Forced swimming	30m/d, 5d/w for 38-65d.	1d after appearance of first tumor	-No difference in survival time or tumor multiplicity

EX = exercise or activity groups; SED = Sedentary controls; NMU = nitrosomethylurea; MNU = 1-methyl-1-nitrosourea; AOM= azoxymethane; DMBA= 7,12-dimethylbenz(a)anthracene; DMH= 1,2-dimethylhydrazine; 3'-Me-DAB= 3'-methyl-4-dimethylaminoazobenzene; Fe-NTA= ferric nitrilotriacetate; s.c.= sub-cutaneous; i.v.= intravenous

Table 2

Tumor Growth Studies

Study	Reference	Rodent Model	Tumor Type / Induction model	METHODS			RESULTS
				Exercise Modality	Exercise Protocol		
					Exercise Prescription Freq/week Dur Intens Length	Exercise Initiation	
Daneryd, 1995	51	Female Wistar Furth rats	Leydig cell / 1.5mm ³ injection of Leydig cell sarcoma (LTW)	Voluntary wheel running	13d	Immediately after injection	-EX ↓ tumor volume by 3.4%.
Daneryd, 1995	52	Female Wistar Furth rats	Leydig cell / 1.5mm ³ injection of Leydig cell sarcoma (LTW) or Nitrosoguanine-induced adenocarcinoma s.c. into each flank	Voluntary wheel running	32d	Immediately after injection	- 13.4g vs 16.4g EX vs SED final LTW weights
Zhu, 2008	42	21d old female Sprague-Dawley rats	Mammary / 25 or 50 mg/kg MNU, i.p.	Voluntary wheel running	Voluntary wheel running for 8w	1w post-injection	-Tumor weight 0.62g vs. 1.16g (SED vs. EX)
Jones, 2010	57	3-4w old female athymic mice	Mammary / 1×10 ⁶ MDA-MB-231 cells, injected orthotopically	Voluntary wheel running	Voluntary wheel running for 41-48d	2d post-implant	No change in primary tumor growth (EX ↑21%)
Yan, 2011	64	3w old male C57BL/6 mice	Lung / 2.5×10 ⁵ /50 µl/mouse Lewis lung carcinoma cells s.c. lower dorsal region	Voluntary wheel running	Primary tumors excised at 1cm diameter, access to wheels continued for additional 2w.	9w before tumor implantation	-No difference in tumor cross-sectional area and tumor volume.
Jones, 2012	55	6-8w old male C57BL/6 mice	Prostate / 5×10 ⁵ mouse prostate C-1 cells, orthotopically	Voluntary wheel running	Wheel running for 8 weeks	14d after tumor transplant	-Primary tumor growth was comparable between groups
Goh, 2014	66	18m old Balb/cBy mice	Mammary / 1×10 ⁴ cells in 4 th mammary fat pad	Voluntary wheel running	Voluntary wheel running for 90 days	60 days prior to tumor transplant, followed by 30 days post-transplant	-Inverse relationship between distance run and final tumor mass
Betof, 2015	65	Female Balb/c or female C57Bl/6 mice	Mammary / 5×10 ⁵ 4T1-luc or 2.5×10 ⁵ E0771 cells in dorsal mammary fat pad	Voluntary wheel running	Voluntary wheel running beginning either 9 weeks prior to tumor transplant, or at the time of tumor transplant	SS: SED before and after transplant RS: EX for 9 weeks prior to transplant; SED after transplant	- Growth rates of SS and RS were similar, and growth rates of SR and RR were similar

Study	Reference	Rodent Model	Tumor Type / Induction model	METHODS			RESULTS
				Exercise Modality	Exercise Protocol		
					Exercise Prescription Freq/week Dur Intens Length	Exercise Initiation	
Alessio, 2009	24	3w old female Sprague-Dawley rats	Spontaneous tumors	Voluntary wheel running or activity box	Wheel: every other day, 24h access throughout animals' life. Activity box (PA): 1h in large activity box twice a week for life.	SR: SED before transplant; EX after transplant RR: EX 9 weeks prior to transplant, continuing after transplant	- EX ↓ tumor growth compared to SED (both tumor models)
Colbert, 2009	26	Female heterozygous (p53+/-); MMTV-Wnt-1 transgenic mice	Mammary / Transgenic	Voluntary wheel running and forced treadmill running	Treadmill running: TRES 1: 5×(45min) 20m/min at 5% grade until completion TRES 2: 5×(45min) 24m/min at 5% grade until completion Voluntary wheel running with 24 hour access.	11w of age	- Time to tumor size of 1.5cm: 24.8d, 13.8d, and 19.5d in control, TRES1 and TRES2 animals. - Treadmill running led to faster tumor growth, no difference due to voluntary wheel running - Treadmill running ↓ survival
Newton, 1965	60	45d old Sprague-Dawley rats	Carcinoma / Equal volumes of Walker-256 cells s.c. into the right flank.	Forced treadmill running	Pre-Tumor: 50h over 5d at 950ft/hr. Post-Tumor: 138h over 10d at 950 ft/hr.	5d before tumor implantation +/- 4d after tumor implantation	-EX ↓ final tumor weight vs SED control. -Early life manipulation + EX ↓ final tumor weight vs EX alone and SED.
Uhlenbruck, 1991	63	BALB/c mice	Sarcoma / 2.4×10 ⁴ L-1 cells s.c.	Forced treadmill running	7×distances of 200m/400m/800m(0.3m/s 2w	4w before and 2w after injection	-200m group ↓ tumor weight
Woods, 1994	40	6w old male C3H/HeN mice	Mammary / 2.5 × 10 ⁵ mammary SCA-1 cells s.c. in the back.	Forced treadmill running	Treadmill running: Moderate: 7×(30min) 18m/min at 5% grade 1w. Exhaustive: 7×(varied) 18m/min for 30min, then 3m/min [†] every 30min until exhausted 1w	3d prior to injection.	No difference in growth rate of tumor size at time of euthanasia (two weeks)
Whittall-Strange, 1998	39	21d old female Sprague-Dawley rats	Mammary / 37.5mg/kg NMU i.p. at 50d of age, (1 day after last bout of EX)	Forced treadmill running	Progressive training to 5×(60min) 18 m/min at 15% grade 4w	21d of age (29d prior to injection)	-Tumor growth rate at 22w post-NMU: 0.043g/day in SED vs. 0.107g/day in EX -Final tumor weight: (3.2g in SED vs. 1.2g in EX)

Study	Reference	Rodent Model	Tumor Type / Induction model	METHODS			RESULTS	
				Exercise Modality	Exercise Prescription Freq/week Dur Intens Length	Exercise Initiation	Tumor Progression Results	
Bacurau, 2000	49	12w old male Wistar rats	Carcinoma / 2×10^7 Walker-256 cells s.c. in the flank	Forced treadmill running	$5 \times [60\text{min}]60\%$ of VO ₂ peak 10w	8w prior to injection	-Day 15 tumor weight 1.82% vs 19% of BW (EX vs SED) -EX prolonged survival by 1.9 fold	
Westerlind, 2003	37	21d old female Sprague-Dawley rats	Mammary / 25 or 50 mg/kg MNU, i.p.	Forced treadmill running	Week 1: $5 \times [10-15\text{min}]30\text{m}/\text{min} 1\text{w}$ Week 2-9: $5 \times [30\text{min}]23-25\text{m}/\text{min} 8\text{w}$	1w post-injection	-↑ Latency in EX (35.8d vs. 33.1d). -No difference in median tumor-free survival time was observed in the EX versus sham-EX (SHAM), nor were there any differences in multiplicity at either a high or moderate dose of MNU	
Zielinski, 2004	44	6-8w old female BALB/c mice	Neoplastic lymphoid cells / 2×10^7 EL-4 cells s.c. in the back behind the neck	Forced treadmill running	$7 \times [3\text{h or until volitional fatigue}]$ gradually increasing speed, 20-40m/min, 5% grade 5-14d	First session immediately before injection.	-No difference in tumor density	
Jones, 2005	56	3-4w old female athymic mice	Mammary / Flank injection of 5×10^6 MDA-MB-231 cells	Forced treadmill running	$5\text{d}/\text{w} 10\text{m}/\text{min}$ for 10min up to $18\text{m}/\text{min}$ for 45 min 0% grade 8w	14d post-injection	-No change in tumor growth.	
Bacurau, 2007	48	8w old male Wistar rats	Carcinoma / 2×10^7 Walker-256 cells s.c. in the flank	Forced treadmill running	$5 \times [30\text{min}]85\%$ of VO ₂ max 10w	8w prior to injection	-Survival: 16d for SED, 45d for EX -EX tumors were 6.9% of final body mass vs. 17.33% for SED control.	
Lira, 2008	58	Male Wistar rats	Carcinoma / 2×10^7 Walker-256 cells s.c. in the flank	Forced treadmill running	$5 \times [60\text{min}]60-65\%$ of VO ₂ peak 10w 2w pre-training period: rats ran progressively from 15 to 60min at 10m/min. Running was increased to 20m/min for two weeks after injection.	8w prior to injection	-Tumor weight: 17.2g in SED; 1.9g in EX	
Murphy, 2011	59	4w old C3(1)SV40Tag mice	Mammary / Transgenic (tumors began developing at 12w of age).	Forced treadmill running	$6 \times [60\text{min}]20\text{m}/\text{min}$ at 5% 20w	4w of age	-Tumor volume: ↓ in EX at 21 and 22w.	
Gueritat, 2014	69	10-12w old Copenhagen rats	Prostate / surgical s.c. implantation of R3527 Dunning AT1	Forced treadmill running	$5 \times [15-60\text{min}]20-25\text{m}/\text{min} 5\text{w}$	15d after tumor implantation	- EX rats had smaller tumors at 14 and 21 days compared to SED controls	

Study	Reference	Rodent Model	Tumor Type / Induction model	METHODS			RESULTS
				Exercise Modality	Exercise Protocol		
					Exercise Prescription Freq/week Dur Intens Length	Exercise Initiation	
Shalamzari, 2014	67	4–6w old Balb/c mice	Mammary / 1×10^6 MC4-L2 s.c. in the flank	Forced treadmill running	- 20–40min 6–20 m/min 15w	RTR: Sedentary before and after transplant RTE: EX after tumor transplant only ETR: EX before tumor transplant only ETE: EX before and after transplant 9w before tumor transplant, and/or 6w after transplant	- Tumor doubling time was significantly slower in EX vs. SED (6.19d vs. 8.81d) - ETE had significantly slower growth compared to RTR - No difference in final tumor volume of RTE and ETR groups
Avesch, 2015	68	5w old female Balb/c mice	Mammary / 1.2×10^6 MC4-L2 cells in right dorsal mammary fat pad	Forced treadmill running	7x 20–55min 10–20 m/min 7w	10d after tumor transplant	-EX decreased tumor volume
Malicka, 2015	45	4w old female Sprague Dawley rats	Mammary / 180 mg/kg MNU i.p.	Forced treadmill running	Low intensity (LIT): $5 \times 10-35\text{min} 0.48-1.34 \text{ km/h} 12\text{w}$ Moderate intensity (MIT): $5 \times 10-35\text{min} 0.6-1.68 \text{ km/h} 12\text{w}$ High intensity (HIT): $5 \times 10-35\text{min} 0.72-2.0 \text{ km/h} 12\text{w}$	Immediately after MNU injection	-No difference in final volume of total tumor volume
Pignet, 2015	46	7–9w old male AlbCrePten flox/flox mice	Hepatocellular carcinoma / Transgenic	Forced treadmill running	5w acclimation period followed by: $5 \times 60\text{min} 2.5\text{m/min} 27\text{w}$	7–9w of age	-EX decreased total volume of liver tumors
Hoffman, 1962	54	Wistar rats	Carcinoma / 2mL Walker 256 cell suspension s.c. into the right thigh.	Continuous running on a 20ft runway + swimming + revolving drum	21d of EX, all EX did all 3 modalities each day: Runway: continuous running on 20ft runway, duration and intensity not clear Swimming: increasing 20min/day to 4h/day Revolving drum: 5.4 mi in 12h	Immediately after injection	-97% inhibition of tumor growth. -Tumor weight ↓ in EX group
Gershbein, 1974	53	Holtzman rats	Carcinoma / Carcinoma	Forced swimming	$10 \times 15\text{min} - 10\text{d}$	Immediately after injection.	-EX ↓ tumor size.

Study	Reference	Rodent Model	Tumor Type / Induction model	METHODS			RESULTS
				Exercise Modality	Exercise Prescription Freq/week Dur Intens Length	Exercise Initiation	
Baracos, 1989	50	Sprague-Dawley rats	Walker-256 tumor i.m. into both hindlimbs. Hepatoma / 20uL Morris hepatoma 777 s.c.	Forced swimming	Low: 5x 5min/d, increased by 5min/d for 3w. Medium: 5x 10min/d, increased by 10min/d for 3w. High: 5x 15min/d, increased by 15min/d for 3w.	2 Groups - 3w of swimming, tumor transplant, then 3w additional swimming. - 3w of swimming beginning 3d post-transplant	-No change in survival rates. -EX ↓ final tumor weight, both groups.
Radak, 2002	61	Adult female hybrid BDF1 mice	Solid leukemia / 5x10 ⁶ P-388 lymphoid leukemia cells s.c.	Forced swimming	5x 60min - 10w ET: training terminated at tumor implantation. EC: training continued for 18d after tumor implantation.	10w before injection.	-EC animals had slower tumor growth than ET and control, (endpoints were ~1.24cm ³ vs. 2cm ³ and 2.4cm ³)
SáezMdel, 2007	34	50d old female Sprague-Dawley rats	Mammary / 5mg/w DMBA gastric intubation for 4w	Forced swimming	30m/d, 5d/w for 38-65d.	1d after appearance of first tumor	-EX ↑ tumor growth by 200%.
Almeida, 2009	47	7w old male Swiss mice	Carcinoma / 2x10 ⁶ Ehrlich tumor cells s.c. in the dorsum.	Forced swimming	5x 60min 50/80% max test 6w. Progressive load test began after 1w: load increased by 2% BW every 3min until exhaustion.	4w before injection	-50% workload ↓ tumor weight and tumor volume (0.18mg/g and 0.11mm ³) vs control (0.55mg/g and 0.48mm ³) -Tumor volume and weight were 270% and 280% ↑ in SED mice.
Sasvari, 2011	62	Adult female BDF1 mice	Sarcoma / 5x10 ⁶ S-180 cells s.c. injected.	Forced swimming	5x 60min - 10w ET: cells injected after EX. ETC: EX (10w), cells injected, EX (18 additional days)	10w before tumor implantation.	-ETC ↓ final tumor weight vs. SED control.

EX = exercise or activity groups; SED = Sedentary controls; NMU = nitrosomethylurea; MNU = 1-methyl-1-nitrosourea; AOM = azoxymethane; DMBA = 7,12-dimethylbenz(a)anthracene; DMH = 1,2-dimethylhydrazine; 3-Me-DAB = 3-methyl-4-dimethylaminoazobenzene; Fe-NTA = ferric nitrilotriacetate; s.c. = sub-cutaneous; i.v. = intravenous

Table 3

Tumor Metastasis Studies

	Study	Reference	Rodent Model	Tumor Type / Induction model	METHODS			RESULTS
					Exercise Modality	Exercise Protocol		
						Exercise Prescription Freq/week Dur Intens Length	Exercise Initiation	
Naturally occurring	Gershbein, 1974	53	Holtzman rats	Carcinoma / Walker-256 tumor i.m. into both hindlimbs.	Forced swimming	10x 15min ~ 10d	Immediately after injection.	Tumor Metastasis Results -Older EX rats ↑ lower abdominal and inguinal secondary tumor nodules.
	Yan, 2011	64	3w old male C57BL/6 mice	Lung / 2.5x10 ⁵ /50 µl/mouse Lewis lung carcinoma cells s.c. lower dorsal region	Voluntary wheel running	Tumors excised at 1cm diameter, access to wheels continued for additional 2w.	9w before tumor implantation	-Trend to inverse relationship between running distance and metastatic tumor yield
	Jones, 2012	55	6–8w old male C57BL/6 mice	Prostate / 5x10 ⁵ mouse prostate C-1 cells, orthotopically	Voluntary wheel running	Wheel running for 8 weeks	14d after tumor transplant	-EX ↓ tumor nodal involvement by 36%, metastases weight by 88% and number of metastases by 34% (<i>None were significant</i>)
	Uhlenbruck, 1991	63	BALB/c mice	Sarcoma / 2.4x10 ⁴ L-1 cells i.v.	Forced treadmill running	7x predetermined distances of 200–850m 0.3–0.5m/s 4–6w	4w before injection, followed by either rest or an additional 2w running	-EX decreased lung tumor multiplicity -Differences in running prescriptions make it difficult to determine if exercise cessation after tumor cell transplant was different from continuous exercise
	MacNeil, 1993a	74	4–5w old male C3H/He mice	Transformed fibroblasts / 1.5x10 ⁶ CIRAS 1 tumor cells i.v. via tail vein.	Forced treadmill and voluntary wheel running	Treadmill: 5x 30min 15m/min, 0% 9w. Voluntary Wheel: 24h access to wheel for 9w. All animals remained SED after injection for 3w.	9w before injection	-No significant effect of activity on lung tumor retention. -Significant but weak correlation of distance run and multiplicity of lung metastases. -Median number of lung metastases per animal was greater in EX mice (21 vs 10 SED control).
	MacNeil, 1993b	73	C3H/He mice	Transformed fibroblasts / 3x10 ⁵ CIRAS1 cells i.v.	Voluntary wheel running	3–12w.	9w prior to and/or 3w after injection	-EX had no effect on lung tumor density -EX prior to tumor injection ↑ incidence in the lowest tertile of tumor distribution vs SED controls.
Artificial (intravenous injection of tumor cells)	Hoffmann-Goetz, 1994a	71	C3H/He-bg2J/+ and C3H/HeJ mice	Transformed fibroblasts / 5x10 ⁵ CIRAS 1 or CIRAS 3 radiolabeled transformed fibroblast cells i.v. via tail vein.	Voluntary wheel running	24h access to running wheel for 9w.	9w before injection	-EX ↓ retention of radioactivity in lungs 5min and 30min post-injection.
	Hoffmann-Goetz, 1994b	70	3w old female BALB/c mice	Mammary / Lateral tail vein injection of 1x10 ⁴ MMT 66 cells	Forced treadmill running or voluntary wheel running	Treadmill running: 5x 30min 18m/min at 0% 8w or 3w. Groups: WS: wheel mice, 24h access for 8w, then SED for 3w. TS: treadmill mice, then SED for 3w.	8w prior or 36h after injection for 3w	-No EX effect on number of lung tumors. -TT tended to have ↑ tumor multiplicity. -ST and SW tended to have ↓ tumor multiplicity.

Study	Reference	Rodent Model	Tumor Type / Induction model	METHODS			RESULTS
				Exercise Modality	Exercise Protocol		
					Exercise Prescription Freq/week Dur Intens Length	Exercise Initiation	
Jadeski, 1996	72	4–9w old C3H/He-bg2J/+ and C3H/HeJ mice	Transformed fibroblasts / 5×10^5 CIRAS 1 or CIRAS 3 transformed fibroblast cells i.v. via tail vein.	Forced treadmill running	TT/WT: Continuous EX for 11w total. ST or SW: SED for 8w then treadmill (ST) or wheel (SW) for 3w. SS: SED for 11w total.	9w before injection	-EX ↓ tumor cell lung retention (44.2 vs 52.8% control). -EX ↓ lung retention of the less aggressive CIRAS 1, no difference in CIRAS 3 cells. -EX did not alter final tumor lung metastases numbers, (subgroup evaluated 3w after EX and injection).
Yan, 2011	64	3w old male C57BL/6 mice	Melanoma / $0.75 \times 10^5/200$ μ l/mouse B16BL/6 cells in lateral tail vein	Voluntary wheel running	Melanoma: 24h access to wheels, continued 2w post-implantation.	9w before tumor implantation	-No difference in number of lung metastasis
Higgins, 2014	75	6w old male scid-beige mice	Lung / 5×10^5 A549-luc-C8 cells i.v. via tail vein	Voluntary wheel running	28d Mice averaged 600–1200 m/day	After lung tumors were detectable	- EX decreased tumor volume, as measured by bioluminescent imaging

EX= exercise or activity groups; SED= Sedentary controls; i. v.= intravenous

Table 4

Mechanistic results

Study	Reference	Tumor type	Exercise Modality	Mechanistic findings		
				Intratumoral	Systemic	Potential Implications
Ikuyama, 1993	28	Carcinogen induced hepatoma	-None reported.	-EX ↓ gamma-glutamyl transpeptidase and ↑ ALP.	Reduced liver damage	
Alessio, 2009	24	Spontaneous tumors	Voluntary wheel running	-None reported.	-Mean serum prolactin levels were ↓ in exercising rats and ↑ in SED rats at 24 and 52w of age.	Prolactin has been associated with tumor growth
Goh, 2014	66	Orthotopic mammary tumor	Voluntary wheel running	-Inverse correlation between distance run and mitotic index within tumors	-EX increased VO ₂ and respiratory exchange rate	Decreased tumor cell proliferation
Betof, 2015	65	Orthotopic mammary tumor	Voluntary wheel running	-EX increased colocalization of desmin and CD31 and decreased tumor hypoxia and necrosis -EX increased expression of Fas, caspase 8 and cleaved caspase 3	-None reported	Increased microvessel density and maturity, which leads to improved tumor perfusion; increased tumor cell apoptosis
Woods, 1994	40	s.c. mammary tumor transplant	Forced treadmill running	-Moderate EX: ↑ phagocytic cells -Exhaustive EX: ↓ total and highly phagocytic cells	-None reported	Increased tumoricidal immune response
Bacurau, 2000	48	s.c. carcinoma transplant	Forced treadmill running	-None reported	-EX ↓ glucose consumption and lactate production, and ↑ glutamine consumption of macrophages -EX ↓ H ₂ O ₂ production by	Increased tumoricidal immune response

Study	Reference	Tumor type	Exercise Modality	Intratumoral	Mechanistic findings	
					Systemic	Potential Implications
Bacurau, 2007	49	s.c. carcinoma transplant	Forced treadmill running	-EX impairs tumor cell glucose and glutamine metabolism.	-EX, non-tumor ↑ plasma corticosterone vs. control. -EX prevents tumor-induced reduction in body weight and food intake, activation of glutamine metabolism in macrophages and lymphocytes.	Modulation of physiology
Lira, 2008	58	s.c. carcinoma transplant	Forced treadmill running	-None reported.	-EX ↓ liver triacylglycerol (TAG) content compared to SED. -SED tumor animals had ↑ serum and liver TAG, and ↓ rate of VLDL secretion, apoB expression and microsomal TAG transfer protein compared to control.	Associated with reduced cachexia
Murphy, 2011	59	Transgenic mammary tumors	Forced treadmill running	-None reported.	→ spleen weight compared to wild-type mice. -No difference in spleen weight due to EX. -EX ↓ plasma IL-6 and MP-1.	Increased tumoricidal immune response
Kato, 2011	29	Carcinogen-induced renal tumors	Forced treadmill running	-None reported.	-Long-term EX and Fe-NTA ↓ renal brown droplets compared to SED, ↑ adrenal weight	Brown droplets could reflect renal damage caused by carcinogen applied. Increases in adrenal weight have been linked to psychological stress.

Study	Reference	Tumor type	Exercise Modality	Mechanistic findings		
				Intratumoral	Systemic	Potential Implications
Shalamzari, 2014	67	s.c. mammary tumor transplant	Forced treadmill running	- EX after tumor transplant decreased tumor IL-6 and VEGF - IL-6 and VEGF levels in ETR mice were not different from RTR (sedentary) mice.	-None reported	Reduced inflammation and subsequent angiogenesis
Piguet, 2015	46	Transgenic hepatocellular carcinoma	Forced treadmill running	- EX decreased proliferation (Ki67 staining) in tumor nodules >15mm ³ .	- EX altered gene expression of fatty acid metabolism pathways	Altered lipogenesis pathways are prognostic indicators following HCC surgical removal
Mallick, 2015	45	Carcinogen-induced mammary tumor	Forced treadmill running	-EX increased TUNEL positive cells	- None reported	
Almeida, 2009	47	s.c. carcinoma transplant	Swimming	-50% workload ↓ macrophage infiltration and neutrophil accumulation.	-80% workload induced cardiac hypertrophy vs. 50%.	Altered immune response; future analysis of macrophage subsets would be beneficial to the field
Sasvari, 2011	62	s.c. sarcoma transplant	Swimming	-None reported.	-EX ↓ oxidative modification of protein in the liver as measured by protein carbonyls.	Reduced oxidative stress or improved anti-oxidant activity
Daneryd, 1995	51	Carcinogen-induced or s.c. leydig cell transplant	Voluntary wheel running	-None reported	- EX normalized insulin sensitivity compared to SED - EX ↑ skeletal muscle metabolism - EX attenuated ↓ of reverse triiodothyronine secretion by the thyroid - EX ↑ insulin:glucagon ratio - EX ↓ corticosterone	Reflects metabolic adaptations to prevent hypo- or hyperglycemia, which may develop in cancer patients.
Progression						

Study	Reference	Tumor type	Exercise Modality	Mechanistic findings		
				Intratumoral	Systemic	Potential Implications
Daneryd, 1995	52	s.c. leydig cell transplant	Voluntary wheel running	-EX 1.31-fold ↑ in tumor cell energy charge and uric acid content	-None reported.	Uric acid has since been shown to stimulate dendritic cell activation, and is elevated in tumors with anti-tumor immune responses
Zhu, 2008	42	Carcinogen-induced mammary tumor	Voluntary wheel running	-EX activated AMPK -EX ↓ VEGF -EX ↓ Bcl-2, X-linked inhibitor of apoptosis pathway (XIAP) while ↑ Bax, apoptosis peptidase-activating factor-1	↓ Insulin, IGF-1, CRP leptin and estradiol ↑ Corticosterone	EX increases cell metabolism and skews the BCL-2 family member protein profile pro-apoptotic.
Jones, 2010	57	Orthotopic mammary tumor	Voluntary wheel running	-EX ↑ perfused vessels and HIF-1 protein levels	-None reported	Improved tumor perfusion, oxygenation
Yan, 2011	64	s.c. lung tumor transplant	Voluntary wheel running	-None reported.	-Tumor presence ↑ plasma VEGF, PDGF-AB, MCP-1. -Running: ↓ plasma insulin and leptin, ↑ adiponectin, ↑ plasma VEGF.	VEGF expression has been linked to worse outcome in patients, but could instead represent improved microvessel density and vascular normalization in tumors.
Jones, 2012	55	Orthotopic prostate tumor	Voluntary wheel running	-EX ↑vascularization, stabilization of HIF-1α → 40% ↑regulation of VEGF -Expression of prometastatic genes was significantly modulated in exercising animals with a shift toward reduced metastasis	↑ activity of protein kinases within MEK/MAPK and PI3/mTOR	Improved tumor perfusion/oxygenation
Mann, 2010	31	Carcinogen-induced mammary tumor	Voluntary non-motorized or motorized wheel running	-None reported	-Induced citrate synthase activity	Reflects training response
Zhu, 2012	43	Carcinogen-induced mammary tumor	Voluntary wheel running	-None reported	↓ Insulin-like growth factor-1	Increased tumoricidal immune response

Study	Reference	Tumor type	Exercise Modality	Mechanistic findings		
				Intratumoral	Systemic	
					Potential Implications	
Westerlind, 2003	37	Carcinogen-induced mammary tumor	running at a fixed daily distance	-None reported	(IGF-1), insulin, interleukin-6, serum amyloid protein, TNF- α , and leptin - \uparrow IGF-binding protein 3 (IGFBP-3) and adiponectin - \uparrow heart and soleus weights	Reflects training response
Zielinski, 2004	44	s.c. neoplastic lymphoid cell transplant	Forced treadmill running	-No difference in the fluid content of tumor -EX \downarrow in vessel density -EX \downarrow inflammatory cells (macrophages and neutrophils) but increased lymphocytes	-None reported	Altered immune response; future analysis of macrophage subsets would be beneficial to the field
Zhu, 2009	41	Carcinogen-induced mammary tumor	Forced wheel running	-Apoptosis induced via mitochondrial pathway in EX -Cell cycle progression suppressed at G(1)/S transition in EX -EX \downarrow blood vessel density	-None reported	EX promotes tumor cell apoptosis and prevents proliferations and angiogenesis
Gueritai 2014	69	s.c. prostate tumor transplant	Forced treadmill running	- EX decreased tumor cell proliferation (Ki67 staining), p-ERK/ERK ratio and 8-OHdG - No difference in tumor SOD, protein carbonylation or lipid oxidation	-None reported	ERK phosphorylation is increased by oxidative stress
Aveseh, 2015	68	Orthotopic mammary tumor	Forced treadmill running	- EX shifted tumor lactate dehydrogenase expression towards the LDHI isoenzyme	-None reported	A shift towards LDH-1 may signal reduced lactate concentrations in the tumor. Lack of MCT1 may result in the tumor cells utilizing glucose instead of

Study	Reference	Tumor type	Exercise Modality	Mechanistic findings		
				Intratumoral	Systemic	Potential Implications
				- EX decreased tumor monocarboxylate transporter 1 (MCT1) and CD147 expression.		lactose, and eventually starving the tumor cells. CD147 is required for MCT1 expression
Radak, 2002	61	s.c. solid leukemia transplant	Swimming	-EC tumors had ↑ Ras protein compared to control. -EC tumors had ↑ I-kappaB protein than ET and control.	-None reported	May be associated with lymphocyte proliferation and activity, but additional analyses are needed
Abdalla, 2013	23	Carcinogen-induced mammary tumor	Swimming	-None reported.	-Physical activity ↑ lymphocytes producing IFN- γ , IL-2, IL-12, and TNF- α . -Physical activity promoted immune system polarization toward an antitumor Th1 response pattern profile.	Increased tumoricidal immune response
MacNeil, 1993a	74	i.v. injection of transformed fibroblasts	Forced treadmill or voluntary wheel running	-None reported.	-Wheel group ↑ splenic NK response vs control.	Increased tumoricidal immune response
Hoffmann-Goetz, 1994a	71	i.v. injection of transformed fibroblasts	Voluntary wheel running	-None reported.	-Running ↓recovery of radioactivity from liver, spleen and kidney at 30min and 90min post-injection.	May reflect decreased retention of circulating tumor cells in organs, thereby lowering the risk of metastasis.
Hoffman-Goetz, 1994b	70	i.v. injection of mammary tumor cells	Forced treadmill running or voluntary wheel running	-None reported	-Pre-tumor EX ↑ LAK cell activity. -NK activity lower in animals that stopped EX at tumor injection.	Increased tumoricidal immune response
Jadeski, 1996	72	i.v. injection of transformed fibroblasts	Forced treadmill running	-None reported	-EX ↑ citrate synthase activity in the soleus.	Indicates a training effect developed

Metastasis

Study	Reference	Tumor type	Exercise Modality	Mechanistic findings		
				Intratumoral	Systemic	Potential Implications
Higgins, 2014	75	i.v. injection of lung tumor cells	Voluntary wheel running	-EX tumors had increased levels of p53, Bax and Bak. -EX tumors had increased staining for cleaved caspase 3	-EX increased serum BUN and decreased ALT, but both were still within normal ranges	Supports increased apoptosis of tumor cells

Table 5

Recommendations for Preclinical Exercise Oncology Research

Concern	Recommendation
Use of xenograft models in immune deficit animals	Orthotopic implantation of syngeneic tumor cell lines or induction of orthotopic tumors via transgenic or chemical methods in immune competent animals
Poor description of exercise intervention characteristics	Describe frequency, intensity, duration, and progression, as appropriate. Avoid vague terms such as “exercise to exhaustion”. Confirmation of ‘training’ effect via muscle fiber or mitochondrial function analysis
Handling of control (sedentary) animals	Handling, social interaction, and environment should be similar to animals randomized to exercise conditions. This includes differences in cage size and social housing. If possible, animals should be acclimated to exercise, or introduced to the activity gradually.
Tail vein models of metastasis	Consider using orthotopic implantation of syngeneic tumor cell lines or transgenic models that spontaneously develop metastasis. However, tail vein models of metastasis may still be useful for assessing the effects of exercise at later time points in the metastatic cascade.
Lack of assessment of systemic and molecular mechanisms	Investigate effects on systemic mechanisms (metabolic and sex hormones, inflammation, immunity, and products of oxidation) postulated to underlie effects of exercise on tumorigenesis as well as potential mediating molecular mechanisms (e.g., cell signaling pathways, angiogenesis, metabolism, migration). Findings should be validated by the use of knock-out/knock-in transgenic animals.
Lack of assessment of tumor biology beyond tumor incidence, weight, or volume	Report on other common markers of the neoplastic phenotype (e.g., apoptosis, proliferation, microvessel density, necrosis, angiogenesis)
Lack of concern regarding the psychological differences between voluntary physical activity vs. forced exercise	Comprehensive study on hypothalamic- pituitary- adrenal axis activation in response to different exercise prescriptions and the effects that associated hormones have on tumor progression/prevention in sedentary controls.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript