

Prioritizing US Cervical Cancer Prevention With Results From a Geospatial Model

Jonathan K. Kish
 Alicia I. Rolin
 Zhaohui Zou
 James E. Cucinelli
 Zaria Tatalovich
 Mona Saraiya
 Sean F. Altekruise

Jonathan K. Kish, Alicia I. Rolin, Zaria Tatalovich, and Sean F. Altekruise, National Cancer Institute, Rockville; Zhaohui Zou and James E. Cucinelli, Information Management Services, Silver Spring, MD; and Mona Saraiya, Centers for Disease Control and Prevention, Atlanta, GA.

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

Authors' disclosures of potential conflicts of interest and contributions are found at the end of this article.

Corresponding author: Sean F. Altekruise, DVM, MPH, PhD, 9609 Medical Center Dr, Room 4E536, Rockville, MD 20850; e-mail: altekruksesf@mail.nih.gov.

abstract

Purpose To determine if differences in screening and vaccination patterns across the population may accentuate ethnic and geographic variation in future burden of disease.

Methods Using Cancer in North America data provided by the North American Association of Central Cancer Registries, county cervical cancer incidence trends from 1995 to 2009 were modeled for the entire United States using ecologic covariates. Rates for health service areas were also modeled by ethnicity. State-level incidence was mapped together with Papanicolaou (Pap) screening, past 3 years (women ≥ 18 years old), and three-dose human papillomavirus (HPV) vaccine coverage (girls 13 to 17 years old) to identify potential priority areas for preventive services.

Results US cervical cancer incidence decreased more during the periods 1995 to 1999 and 2000 to 2004 than during the period 2005 to 2009. During these 15 years, the most affected areas became increasingly confined to Appalachia, the lower Mississippi Valley, the Deep South, Texas, and Florida. Hispanic and black women experienced a higher incidence of cervical cancer than both white and Asian and Pacific Islander women during each period. Women in 10 of 17 states/districts with a high incidence ($\geq 8.14/100,000$) reported low Pap testing ($< 78.5\%$), HPV vaccine coverage ($< 33.9\%$), or both prevention technologies.

Conclusion The decline in cervical cancer incidence has slowed in recent years. Access to HPV vaccination, targeted screening, and treatment in affected populations is needed to reduce cervical cancer disparities in the future.

J Glob Oncol 2. © 2016 by American Society of Clinical Oncology Licensed under the Creative Commons Attribution 4.0 License

INTRODUCTION

Cervical cancer incidence and mortality in the United States have declined steadily in the decades since the adoption of cytology-based Papanicolaou (Pap) testing.¹ Recently, cervical cancer death rates seem to have plateaued, and a slight decrease has been reported in the use of Pap testing.² Cervical cancer disparities (the uneven distribution of burden to areas of lower socioeconomic status and minority ethnic groups) are increasing.³⁻⁶ Cervical cancer incidence is elevated among older and black women^{7,8} and in impoverished areas such as Appalachia, the lower Mississippi Valley, and the United States–Mexico border area.⁹ Reduced access to screening services is a driving factor in these areas. Nationally, 3-year Pap screening completion is $> 80\%$ for women 21 to 65 years of age, with lower levels of screening among immigrants, minorities, the poor, and

older women.¹⁰ Approximately 47% of girls born in the United States in 2000 had received at least one dose of human papillomavirus (HPV) vaccine by age 13 years.¹¹ Safety concerns and views on sexual activity influence parental decisions regarding the vaccination of adolescent girls.¹¹ Although the safety and efficacy of the HPV vaccination is well established, in 2013, 38% of girls 13 to 17 years of age in the United States had completed the three-dose HPV vaccine series,¹¹ far less than optimal.¹² Because the reduction in cervical cancer incidence as a result of HPV vaccination will take decades to be realized,¹² surveillance is needed to identify communities with a higher burden of cervical cancer and to direct cervical cancer screening and vaccination services where they are needed most.¹

To define high-priority areas for preventive services, we estimated small-area cervical cancer

incidence, including by ethnicity, and mapped state-level incidence with percentage screening and HPV vaccination coverage.

METHODS

Incidence Data

The North American Association of Central Cancer Registries (NAACCR) provided a Cancer in North America data set with county-level incidence and population denominators for this analysis. A total of 36 of 50 state registries, all of which met NAACCR silver or gold standards for tumor registration data quality, gave active consent to include their incident cases in the analytic data set. For the years 1995 to 2009, complete or partial county-level cervical cancer incidence data were reported by NAACCR registries in areas covering 74% of the United States population. This included complete data from 1995 to 2009 for 22 states encompassing 56% of the population (Arizona, California, Colorado, Connecticut, Hawaii, Idaho, Illinois, Iowa, Kentucky, Louisiana, Maine, Michigan, Nebraska, New Jersey, New Mexico, New York, Pennsylvania, Rhode Island, Texas, Utah, Washington, and Wyoming) and multiple years of data for 14 states encompassing 18% of the population (Alaska, Arkansas, Georgia, Massachusetts, Mississippi, Montana, Nevada, North Carolina, North Dakota, South Carolina, South Dakota, Tennessee, Virginia, and West Virginia). In these 14 states, data were missing for a median of 2 reporting years, generally the earliest reporting years. Two lower Mississippi Valley states were missing data for 1995 to 2003. The remaining 14 states and the District of Columbia, which did not provide data, accounted for 26% of the US population.

Spatiotemporal Model

We developed a model similar in composition to one used commonly to predict current-year cancer incidence in the United States¹³ to estimate county-level cervical cancer incidence in 5-year intervals from 1995 to 2009.¹⁴ The multivariable logistic regression model included a set of covariates selected through a forward and backward process. A generalized linear model was used (PROC GLIMMIX, SAS 9.3, SAS Institute, Cary, NC) with three random terms to account for spatial autocorrelation (longitude and latitude of the county), temporal autocorrelation (year of diagnosis), and residual autocorrelation of covariates. Missing cervical cancer incidence at the county level was modeled on the basis of the reported incidence in counties with comparable attributes. The incidence for counties with observed data was

also updated according to model predictions to slightly adjust reported rates. The time periods of interest for modeled all-ethnicity county-level incidence were 1995 to 1999, 2000 to 2004, and 2005 to 2009. To present incidence among ethnic groups, data were aggregated at the health service area (HSA) level,¹⁵ reducing instability from small counts at the county level. HSAs are either a single county or a cluster of contiguous counties that are relatively self-contained with respect to hospital care. Data were suppressed when there were 16 or fewer modeled incident cases. The accuracy of predicted rates for areas with missing data depends on how well covariates in the model predict the actual incidence.

Model Inputs

Demographic inputs of cervical incident cases were non-Hispanic ethnicity (hereafter referred to as white, black, American Indian and Alaska Native, and Asian and Pacific Islander), Hispanic ethnicity (all ethnicities), and age. County of residence was geocoded on the basis of latitude and longitude. County-level population estimates were obtained from the US Census Bureau Summary File for each year from 1995 through 2009. Incidence data, stratified by age (< 44, 45 to 64, and ≥ 65 years) and year of diagnosis were retrieved using SEER*Stat 8.1.2 (Information Management Services, Calverton, MD).

County-level covariates included in the model were county-level rural-urban density data,¹⁶ an Area Health Resources covariate enumerating the number of hospital-based physicians at the county level,¹⁵ and data on the percentages of the county population who were black, Asian and Pacific Islander, and American Indian and Alaskan Native. Cervical cancer mortality data reported by the National Center for Health Statistics were also included for each county in the United States. Socioeconomic covariates incorporated into the model were the percentage of the county population with income below the poverty level and the percentage of the population ≥ 25 years of age with ≥ 4 years of college education. The model also contained a variable indicating whether the county was in a National Program of Cancer Registries–funded area. The model provided estimates of county-level cervical cancer incidence for the entire United States, including areas and years with missing data as well as those with reported data. To illustrate geographic distributions in rates, modeled county estimates were mapped using 2000 Census county designations (ArcGIS 10.1; ESRI, Redlands, CA). In the all

ethnicities combined model uncertainty related to small numbers was addressed with spatial smoothing. The population-weighted, nonparametric algorithm used universal Kriging after detrending.¹⁷ Data were approximately normally distributed and no transformations were applied, although first-order surface trend was removed. Smoothing was not applied to incidence maps of specific ethnic groups because of the potential for instability.

State-Level Incidence Trends

State-level modeled incidence trends on the basis of estimated age-standardized incidence were analyzed using joinpoint regression analysis (Joinpoint 3.5.0; Information Management Services). The technique fit a series of joined straight lines on a logarithmic scale for annual age-standardized rates.¹⁸ Using data from the model for each year, state trends were estimated for 5-year fixed intervals, or annual average percent change (AAPC), using weighted annual percent changes from joinpoint models.

Identification of Priority Areas

Three state-level variables were visualized: modeled cervical cancer incidence during the period 2005 to 2009, state-level proportions of women 21 to 65 years of age who had had a Pap test during the past 3 years¹⁹ estimated from pooled 2008 to 2010 responses to the National Health Interview

Survey and Behavioral Risk Factor Surveillance System, and HPV vaccination coverage by state (obtained from the National Immunization Survey of Teens performed during 2013)²⁰ defined as the percentage of girls 13 to 17 years old who had received three doses or more of either the bivalent or the quadrivalent vaccine.

State-level cervical cancer incidence was mapped in tertiles (low, midlevel, and high incidence). Low incidence was 5.17 to 6.75, midlevel incidence ranged from 6.76 to 8.13, and high incidence was 8.14 to 9.76 cases per 100,000 women. Low state-level Pap screening coverage was defined as < 78.5%, midlevel as 78.5% to 81.4%, and high as 81.5% to 88.5%. Low state-level HPV vaccination uptake was defined as < 33.9%, midlevel as 33.9% to 40.1%, and high as 40.2% to 56.5%. States in the bottom third distribution for three-dose HPV vaccine receipt (< 33.9%), Pap screening (< 78.5%), or both vaccine and Pap screening were depicted with horizontal, vertical, and cross-hatched lines, respectively. The status of state Medicaid expansion as of January 2016 was also assessed.²¹

RESULTS

Modeled cervical cancer incidence and observed death rates per 100,000 women from 1995 to 2009 are presented in Table 1. Incidence and death rates were highest among non-Hispanic black followed by Hispanic women in each time

Table 1 – Age-Adjusted Cervical Cancer Incidence and Mortality by Selected Characteristics, 1995 to 2009

Characteristic	1995-1999				2000-2004				2005-2009				Overall			
	Modeled Incidence		Observed Death		Modeled Incidence		Observed Death		Modeled Incidence		Observed Death		Modeled Incidence		Observed Death	
	N	Rate	N	Rate	N	Rate	N	Rate	N	Rate	N	Rate	N	Rate	N	Rate
Overall	75,085	10.8	22,086	3.1	66,406	8.9	20,013	2.6	63,174	7.9	19,838	2.4	204,665	9.1	61,937	2.7
Ethnicity																
White	51,100	9.7	13,951	2.6	42,942	7.9	12,323	2.2	38,879	6.9	11,897	2.1	132,922	8.1	38,171	2.3
Black	11,891	15.7	4,272	6.3	10,629	12.6	3,796	5.0	10,243	11.0	3,723	4.4	32,763	12.9	11,791	5.1
Hispanic	9,280	17.1	1,817	3.8	9,872	14.3	2,047	3.4	10,891	12.8	2,325	3.0	30,042	14.4	6,189	3.3
API	2,369	10.5	597	3.0	2,532	8.7	647	2.4	2,745	7.6	648	1.9	7,646	8.7	1,892	2.3
AI/AN	444	9.2	99	3.2	430	7.8	107	2.7	416	6.9	134	2.8	1,291	7.9	340	2.8
Age, y																
≤ 44	30,434	6.7	5,294	1.2	25,104	5.5	4,470	1.0	21,875	4.9	3,977	0.9	77,413	5.7	13,741	1.0
45-64	29,757	20.5	8,357	5.7	28,726	16.9	8,306	4.9	29,363	15.2	9,079	4.6	87,845	17.3	25,742	5.0
≥ 65	14,895	14.6	8,435	8.2	12,576	11.9	7,237	6.8	11,936	10.7	6,782	6.1	39,407	12.3	22,454	7.0

NOTE. Incidence and death rates are per 100,000 women. Ethnic mortality statistics exclude deaths in the District of Columbia, Maine, Minnesota, New Hampshire, North Dakota, Oklahoma, and South Carolina because of instability in Hispanic population estimates.²² Data sources: North American Association of Central Cancer Registries for cervical cancer incidence, National Center for Health Statistics for deaths, 1995 to 2009. Incident case counts estimated using spatiotemporal model; observed death counts, cervical cancer. Abbreviations: API, non-Hispanic Asian/Pacific Islander; AI/AN, non-Hispanic American Indian/Alaska native; y, years.


period. Incidence and death rates decreased from the period 1995 to 1999 to the period 2005 to 2009 among all ethnic groups. Age-specific incidence per 100,000 women was highest among 45- to 64-year-old women, averaging 17.3 and decreasing from 20.5 during the period 1995 to 1999 to 15.2 during the period 2005 to 2009. The highest death rate was also seen among women ≥ 65 years of age, averaging 7.0 and decreasing from 8.2 during the period 1995 to 1999 to 6.1 during the period 2005 to 2009.

Figure 1 displays the smoothed, county-level modeled estimates of cervical cancer incidence by time period. From 1995 to 1999, the highest incidence was found in counties extending south and west from Appalachia to southeastern Colorado, northeastern New Mexico, and Texas, with a low incidence found in southwestern New England, the Northern Plains, and Mountain West. During the period 2000 to 2004, elevated rates became more localized to central Appalachia, the Ohio River Valley, the Lower Mississippi River Valley, rural Texas including the Mexican border area, and the rural Southeast. During the period 2005 to 2009, areas with the highest incidence were even more contained within Appalachia, the lower Mississippi Valley, the Deep South, Texas, and Florida.

Overall, in the United States during 1995 to 1999, 2000 to 2004, and 2005 to 2009, the AAPC in modeled cervical cancer incidence declined significantly by -3.5 , -3.1 , and -1.1 percent per year over successive periods (Data Supplement). This pattern of slowing but still statistically significant decreasing incidence trends in recent years was seen in 27 states. In nine states and in the District of Columbia, trends did not change significantly over the three time periods. In three states (Georgia, Virginia, and Louisiana) there was a non-significant increase in cervical cancer incidence between 2005 and 2009. In nine other states (Alaska, Tennessee, Maryland, New Jersey, South Carolina, South Dakota, Illinois, Iowa, and Oregon), the decrease in cervical cancer incidence during the period 2005 to 2009 was not statistically significant. In Utah and Oklahoma the AAPC during the period 1995 to 1999 was not statistically significant; however, incidence decreased significantly during more recent time periods.

Figure 2 presents modeled HSA-level incidence by ethnicity during the period 2005 to 2009. Color ranges vary by map. Higher cervical cancer incidence was seen among Hispanic and black than among white and Asian and Pacific Islander women. Among whites, incidence was highest

Fig 1 –
Smoothed, county-level modeled US cervical cancer incidence per 100,000 women, 5-year intervals, 1995 to 2009.




in Appalachia, the Mississippi and Ohio River valleys, Indiana, and rural parts of Illinois, Texas, Louisiana, Arkansas, Oklahoma, Tennessee, Alabama, and Georgia. Isolated areas with high incidence were found in central California, northwestern Arizona, central Pennsylvania, southern New Jersey, and northern and central Florida. High modeled incidence areas for blacks were found in eastern Texas and adjacent areas of Oklahoma, Louisiana, and Arkansas. The incidence was also elevated along the Mississippi River Valley from Illinois to Tennessee, Arkansas, and Mississippi. Affected rural areas included adjacent areas of Mississippi and Alabama, and Georgia and north central Florida. A high incidence was also seen in northern and central Indiana, rural South Carolina, southern New Jersey, New York City, and southeastern Florida.

Of areas with large Hispanic populations, the regions with the highest modeled incidence included west Texas, central and southern Florida, and metropolitan Houston, Chicago, and New York.

Among Asian and Pacific Islanders, a high modeled incidence was seen in central California and Houston, Texas; central and south Florida; and metropolitan New York. The incidence for American Indians/Alaska Natives was not presented because of low counts.

Figure 3 presents state-level modeled cervical cancer incidence during the period 2005 to 2009. States are classified into three categories: low, midlevel, and high cervical cancer incidence states. States in the bottom tertile of Pap screening use and HPV vaccine uptake were classified as having low adoption of these preventive services. Sixteen states plus the District of Columbia, shown in red, had a high cervical cancer incidence. In two high-incidence states, both Pap screening and HPV vaccine uptake were low (Arkansas and Nevada). There was a low proportion of Pap screening in three high-incidence states (Tennessee, West Virginia, and Indiana) and of HPV vaccine uptake in five high-incidence jurisdictions (District of Columbia, Mississippi, Illinois,

Fig 2 –
Modeled health services area-level cervical cancer incidence by ethnicity, 2005 to 2009.






Fig 3 –

States with high, midlevel, and low cervical cancer incidence during the period 2005 to 2009, further depicting states with low percentages of Papanicolaou (PAP) screening or human papillomavirus (HPV) vaccination. Less than 78.5% of women > 18 years old reported PAP screening in the past 3 years during the period 2008 to 2010, and < 33.9% of girls 13 to 17 years of age had received three doses of HPV vaccine as of 2013.

Kentucky, and New Jersey). Many of the 16 states with a midlevel cervical cancer incidence (shown in orange) were adjacent to high-incidence states. One midlevel-incidence state (Missouri) had low uptake of both Pap screening and HPV vaccination. Among the midlevel-incidence states, five (New Mexico, Wyoming, Nebraska, Oklahoma, and Ohio) had a low percentage of Pap screening, and five (Alaska, Kansas, Maryland, North Carolina, and Georgia) had low vaccine uptake. States with the lowest cervical cancer incidence (shown in gold) were located in New England, the northern tier of states, the four corners region, and Virginia. In three low-incidence states (Montana, Idaho, and Utah), a low percentage of women received Pap screening, and a low percentage of girls received three doses of HPV vaccine. There was also a low percentage of Pap screening in Arizona, North Dakota, and South Dakota, and the percentage of girls in Virginia who received HPV vaccination was less than 33.9%.

As of January 2016, six of 17 jurisdictions with the highest cervical cancer incidence had not

expanded Medicaid coverage (Alabama, Florida, Mississippi, South Carolina, Tennessee, and Texas). Seven of 16 states with a midlevel cervical cancer incidence had not expanded Medicaid (Georgia, Kansas, Missouri, North Carolina, Nebraska, Oklahoma, and Wyoming). Among the 18 states with the lowest cervical cancer incidence, three had not expanded Medicaid (Idaho, Montana, and Utah).

DISCUSSION

This study suggests that the long-term decrease in US cervical cancer incidence is slowing down. This finding is consistent with the nonsignificant decreasing trend reported during the period 2005 to 2009 in areas in the United States with high-quality incidence data.¹² High-incidence geographic areas were confined increasingly to rural areas within Appalachia, Texas, the lower Mississippi Valley, and the southeastern United States. Hispanic and black women had a higher cervical cancer incidence than did white and Asian and

Pacific Islander women. In 10 of 17 states/districts with a high cervical cancer incidence, there was low use of Pap screening or HPV vaccination. Locally tailored cervical cancer vaccination, screening, and treatment efforts that target poor women living in medically underserved geographic areas are needed to maintain progress in reducing cervical cancer disparities.

As recently as the 1970s, cervical cancer was a leading cause of cancer among US women; however, the incidence has decreased in subsequent decades.⁸ Trends in this study differ from sustained decreasing cervical cancer incidence trends over the past half-century, but are consistent with recent findings describing a leveling off of cervical cancer mortality² and incidence¹² in the United States. A possible explanation for the slowing of the decreasing trend is that women with access to health care are benefiting from preventive services such as Pap screening and HPV testing,² to a greater extent than are women in medically underserved groups. These underserved women, who experience a higher burden of cervical cancer,²³ include ethnic minorities,⁵ women from low socioeconomic backgrounds,⁵ and women living in impoverished geographic areas.¹⁰ Future progress to reduce the burden of cervical cancer depends on access to vaccination, screening, and treatment of these hard-to-reach groups.²⁴ Of note, some states with the highest incidence of cervical cancer have low percentages of Pap screening and HPV vaccine uptake. Provisions of the Affordable Care Act, which require most health insurance plans to cover cervical cancer screening and HPV vaccination with no cost sharing, could improve cervical cancer prevention among low-income women.²⁵

In this study spanning the years 1995 through 2009, there were progressively smaller areas with an elevated cervical cancer incidence over time. Regions with the highest burden of disease during the period 2005 to 2009 were largely contained to economically deprived counties within Appalachia, Texas, the lower Mississippi Valley, and the southeastern United States. The limited progress in reducing the incidence of this cancer in areas with slow economic development or an influx of immigrant populations is consistent with findings reported in Mexico,²⁶ Brazil,²⁷ and England.²⁸

Lack of awareness, lack of access to health care, and cultural beliefs are barriers to cervical cancer prevention within population subgroups.²⁹⁻³¹ For instance, in Connecticut,³² heterogeneity in the occurrence of cervical cancer precursors is reported.

Culturally competent locally targeted outreach needs to be part of cervical cancer control programs. One study of African-American women living in the high cervical cancer mortality area of Sunflower County, Mississippi, indicated that door-to-door visits to offer home self-collection HPV test kits increased participation in cervical cancer screening almost four-fold compared with clinic-based Pap testing alone.³³ In a national study, predictors of not being screened for cervical cancer included not having made a physician office visit within the past 12 months because of cost, minority ethnicity, lack of a high school diploma among residents of metropolitan areas, and self-reported fair or poor general health among non-metropolitan area residents.³⁴ The heterogeneity of underserved women suggests a need for screening and HPV vaccination outreach across broad areas.³⁵

Other researchers have reported ethnic and geographic disparities in cervical cancer screening³⁴ and incidence.³⁶ In the United States–Mexico border area, Hispanic women were less likely than other women to have had a recent Pap test,³⁷ and white women in Appalachia had higher rates of HPV infection compared with the US population.³⁸ Although provider recommendation improves acceptance of HPV vaccination, minority and low-income women are least likely to receive such recommendations.³⁹ Cervical cancer prevention can be advanced through community-based interventions,⁴⁰ particularly in communities with limited access to a formal health care system. These community-based interventions may be more effective than one-size-fits-all approaches.⁴¹⁻⁴³

This study identified populations that would benefit from cervical cancer outreach by ethnicity, geography, and access to screening and HPV vaccination. Ongoing spatial analysis is recommended to monitor cervical cancer trends in the HPV vaccine era.⁴⁴ Study limitations include missing data for some states, which was partially addressed with geospatial modeling. Furthermore, HPV vaccination data were available only at the state level. County-level data would improve prioritization of outreach to areas with a high cervical cancer burden. Despite limitations, the analysis identifies priority areas for interventions to improve screening and vaccination rates. Although progress has been made in reducing the incidence of cervical cancer, outreach is needed in low-socioeconomic areas of the United States. Provisions of the Affordable Care Act that eliminate cost sharing for cervical cancer screening and HPV vaccination in most health plans should reduce cost as a barrier to receiving these prevention services.

DOI: [10.1200/JGO.2015.001677](https://doi.org/10.1200/JGO.2015.001677)

Published online on jgo.ascopubs.org on April 11, 2016.

AUTHOR CONTRIBUTIONS

Conception and design: Jonathan K. Kish, Mona Saraiya, Sean F. Altekruise

Administrative support: Mona Saraiya, Sean F. Altekruise

Collection and assembly of data: Jonathan K. Kish, Alicia I. Rolin, James E. Cucinelli, Zaria Tatalovich

Data analysis and interpretation: Jonathan K. Kish, Alicia I. Rolin, Zhaohui Zou, Mona Saraiya

Manuscript writing: All authors

Final approval of manuscript: All authors

Jonathan K. Kish

Employment: Cardinal Health

Consulting or Advisory Role: Cardinal Health (Inst)

Travel, Accommodations, Expenses: Cardinal Health

Alicia I. Rolin

No relationship to disclose

Zhaohui Zou

No relationship to disclose

James E. Cucinelli

No relationship to disclose

Zaria Tatalovich

No relationship to disclose

Mona Saraiya

No relationship to disclose

Sean F. Altekruise

No relationship to disclose

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or jco.ascopubs.org/site/ifc.

REFERENCES

1. Smith RA, Brooks D, Cokkinides V, et al: Cancer screening in the United States, 2013: A review of current American Cancer Society guidelines, current issues in cancer screening, and new guidance on cervical cancer screening and lung cancer screening. *CA Cancer J Clin* 63:88-105, 2013
2. Benard VB, Thomas CC, King J, et al: CDC vital signs, Cervical cancer incidence, mortality, and screening—United States, 2007–2012. *MMWR. Morb Mortal Wkly Rep* 63:1004-1009, 2014
3. Akers AY, Newmann SJ, Smith JS: Factors underlying disparities in cervical cancer incidence, screening, and treatment in the United States. *Curr Probl Cancer* 31:157-181, 2007
4. Saraiya M, Ahmed F, Krishnan S, et al: Cervical cancer incidence in a prevaccine era in the United States, 1998-2002. *Obstet Gynecol* 109:360-370, 2007
5. Simard EP, Fedewa S, Ma J, et al: Widening socioeconomic disparities in cervical cancer mortality among women in 26 states, 1993-2007. *Cancer* 118:5110-5116, 2012
6. Yabroff KR, Lawrence WF, King JC, et al: Geographic disparities in cervical cancer mortality: What are the roles of risk factor prevalence, screening, and use of recommended treatment? *J Rural Health* 21:149-157, 2005
7. DeSantis C, Naishadham D, Jemal A: Cancer statistics for African Americans, 2013. *CA Cancer J Clin* 63:151-166, 2013
8. Howlader N, Noone AM, Krapcho M, et al (eds): SEER cancer statistics review, 1975–2010. Bethesda, MD: National Cancer Institute. http://seer.cancer.gov/csr/1975_2010/
9. Horner MJ, Altekruise SF, Zou Z, et al: U.S. geographic distribution of prevaccine era cervical cancer screening, incidence, stage, and mortality. *Cancer Epidemiol Biomarkers Prev* 20:591-599, 2011
10. Centers for Disease Control and Prevention; National Center for Health Statistics: Health, United States, 2010: With Special Feature on Death and Dying. 2011. <http://www.cdc.gov/nchs/data/abus/abus10.pdf>
11. Stokley S, Jeyarajah J, Yankey D, et al: Human papillomavirus vaccination coverage among adolescents, 2007-2013, and postlicensure vaccine safety monitoring, 2006-2014—United States. *MMWR Morb Mortal Wkly Rep* 63:620-624, 2014
12. Jemal A, Simard EP, Dorell C, et al: Annual Report to the Nation on the Status of Cancer, 1975-2009, featuring the burden and trends in human papillomavirus(HPV)-associated cancers and HPV vaccination coverage levels. *J Natl Cancer Inst* 105:175-201, 2013
13. American Cancer Society; Cancer facts & figures 2015. <http://www.cancer.org/research/cancerfactsstatistics/cancerfactsfigures2015/index>
14. Pickle LW, Hao Y, Jemal A, et al: A new method of estimating United States and state-level cancer incidence counts for the current calendar year. *CA Cancer J Clin* 57:30-42, 2007
15. US Department of Health and Human Services; Health Resources and Services Administration (HRSA). Area Health Resources Files (AHRF). <http://ahrh.hrsa.gov/>
16. United States Department of Agriculture, Economic Research Service: Rural-urban continuum codes. www.ers.usda.gov/data-products/rural-urban-continuum-codes.aspx
17. Goovaerts P. Geostatistics for Natural Resources Evaluation. New York, NY, Oxford University Press, 1997

18. Kim HJ, Fay MP, Feuer EJ, et al: Permutation tests for joinpoint regression with applications to cancer rates. *Stat Med* 19:335-351, 2000
19. National Cancer Institute: Small area estimates for states, counties, & health service areas. <http://sae.cancer.gov/>
20. Centers for Disease Control and Prevention: 2013 NIS-Teen vaccination coverage table data. <http://www.cdc.gov/vaccines/imz-managers/coverage/nis/teen/data/tables-2013.html>
21. Kaiser Family Foundation: Status of state action on the Medicaid expansion decision. <http://kff.org/health-reform/state-indicator/state-activity-around-expanding-medicaid-under-the-affordable-care-act/>
22. National Cancer Institute Surveillance, Epidemiology, and End Results Program: Policy for calculating Hispanic mortality for 1990+ data. http://seer.cancer.gov/seerstat/variables/mort/origin_recode_1990/
23. Freeman HP, Wingrove BK: Excess Cervical Cancer Mortality: A Marker for Low Access to Health Care in Poor Communities. Rockville, MD: National Cancer Institute, Center to Reduce Cancer Health Disparities, May 2005. NIH Pub. No. 05-5282, 2007 <http://www.cancer.gov/about-nci/organization/crhd/about-health-disparities/resources/excess-cervical-cancer-mortality.pdf>
24. Scarinci IC, Garcia FA, Kobetz E, et al: Cervical cancer prevention: New tools and old barriers. *Cancer* 116:2531-2542, 2010
25. The Henry J. Kaiser Family Foundation: Women's Health Policy. Women's Health Insurance Coverage. <http://kff.org/womens-health-policy/fact-sheet/womens-health-insurance-coverage-fact-sheet/>
26. Sánchez-Barriga JJ: Mortality trends from cervical cancer in the seven socioeconomic regions and the thirty two federative entities of Mexico, 2000-2008 [in Spanish]. *Gac Med Mex* 148:42-51, 2012
27. Gonzaga CM, Freitas-Junior R, Barbaresco AA, et al: Cervical cancer mortality trends in Brazil: 1980-2009. *Cad Saude Publica* 29:599-608, 2013
28. Shack L, Jordan C, Thomson CS, et al: Variation in incidence of breast, lung and cervical cancer and malignant melanoma of skin by socioeconomic group in England. *BMC Cancer* 8:271, 2008
29. Lai JY, Tinker AV, Cheung WY: Factors influencing the willingness of US women to vaccinate their daughters against the human papillomavirus to prevent cervical cancer. *Med Oncol* 30:582, 2013
30. Williams WW, Lu PJ, Saraiya M, et al: Factors associated with human papillomavirus vaccination among young adult women in the United States. *Vaccine* 31:2937-2946, 2013
31. Downs LS Jr, Scarinci I, Einstein MH, et al: Overcoming the barriers to HPV vaccination in high-risk populations in the US. *Gynecol Oncol* 117:486-490, 2010
32. Niccolai LM, Julian PJ, Bilinski A, et al: Geographic poverty and racial/ethnic disparities in cervical cancer precursor rates in Connecticut, 2008-2009. *Am J Public Health* 103:156-163, 2013
33. Castle PE, Rausa A, Walls T, et al: Comparative community outreach to increase cervical cancer screening in the Mississippi Delta. *Prev Med* 52:452-455, 2011
34. Stanley SL, Thomas CC, King JB, et al: Predictors of never being screened for cervical cancer by metropolitan area. *J Community Health* 39:400-408, 2014
35. Bach PB: Gardasil: From bench, to bedside, to blunder. *Lancet* 375:963-964, 2010
36. Downs LS, Smith JS, Scarinci I, et al: The disparity of cervical cancer in diverse populations. *Gynecol Oncol* 109:S22-S30, 2008
37. Coughlin SS, Uhler RJ, Richards T, et al: Breast and cervical cancer screening practices among Hispanic and non-Hispanic women residing near the United States-Mexico border, 1999-2000. *Fam Community Health* 26:130-139, 2003
38. Reiter PL, Katz ML, Ruffin MT, et al: HPV prevalence among women from Appalachia: results from the CARE project. *PLoS One* 8:e74276, 2013 [Erratum: *PLoS One* 8, 2013]
39. Jeudin P, Liveright E, Del Carmen MG, et al: Race, ethnicity, and income factors impacting human papillomavirus vaccination rates. *Clin Ther* 36:24-37, 2014
40. Kobetz E, Menard J, Barton B, et al: Patnè en Aksyon: Addressing cancer disparities in Little Haiti through research and social action. *Am J Public Health* 99:1163-1165, 2009
41. Gwede CK, Ashley AA, McGinnis K, et al: Designing a community-based lay health advisor training curriculum to address cancer health disparities. *Health Promot Pract* 14:415-424, 2013
42. Wells KJ, Luque JS, Miladinovic B, et al: Do community health worker interventions improve rates of screening mammography in the United States? A systematic review. *Cancer Epidemiol Biomarkers Prev* 20:1580-1598, 2011
43. Zapka J, Taplin SH, Ganz P, Grunfeld E, Sterba K. Multilevel factors affecting quality: examples from the cancer care continuum. *J Natl Cancer Inst Monogr* 2012:11-19, 2012
44. Richardson DB, Volkow ND, Kwan MP, et al: Medicine. Spatial turn in health research. *Science* 339:1390-1392, 2013