

Deletion of the Three Distal S1 Motifs of *Saccharomyces cerevisiae* Rrp5p Abolishes Pre-rRNA Processing at Site A₂ without Reducing the Production of Functional 40S Subunits

Harmjan R. Vos, Alex W. Faber, Maaike D. de Gier, Jan C. Vos,* and Hendrik A. Raué

Section of Biochemistry and Molecular Biology, Department of Chemistry, Faculty of Exact Sciences and Institute of Molecular Biological Science, BioCenter Amsterdam, Vrije Universiteit, Amsterdam, The Netherlands

Received 3 May 2004/Accepted 17 August 2004

Yeast Rrp5p, one of the few *trans*-acting proteins required for the biogenesis of both ribosomal subunits, has a remarkable two-domain structure. Its C-terminal region consists of seven tetratricopeptide motifs, several of which are crucial for cleavages at sites A₀ to A₂ and thus for the formation of 18S rRNA. The N-terminal region, on the other hand, contains 12 S1 RNA-binding motifs, most of which are required for processing at site A₃ and thus for the production of the short form of 5.8S rRNA. Yeast cells expressing a mutant Rrp5p protein that lacks S1 motifs 10 to 12 (mutant *rrp5Δ6*) have a normal growth rate and wild-type steady-state levels of the mature rRNA species, suggesting that these motifs are irrelevant for ribosome biogenesis. Here we show that, nevertheless, in the *rrp5Δ6* mutant, pre-rRNA processing follows an alternative pathway that does not include the cleavage of 32S pre-rRNA at site A₂. Instead, the 32S precursor is processed directly at site A₃, producing exclusively 21S rather than 20S pre-rRNA. This is the first evidence that the 21S precursor, which was observed previously only in cells showing a substantial growth defect or as a minor species in addition to the normal 20S precursor, is an efficient substrate for 18S rRNA synthesis. Maturation of the 21S precursor occurs via the same endonucleolytic cleavage at site D as that used for 20S pre-rRNA maturation. The resulting D-A₃ fragment, however, is degraded by both 5'→3' and 3'→5' exonuclease digestions, the latter involving the exosome, in contrast to the exclusively 5'→3' exonucleolytic digestion of the D-A₂ fragment. We also show that *rrp5Δ6* cells are hypersensitive to both hygromycin B and cycloheximide, suggesting that, despite their wild-type growth rate, their preribosomes or ribosomes may be structurally abnormal.

Ribosome biogenesis in eukaryotes largely takes place in the nucleolus, a specialized compartment of the nucleus, and proceeds via a complex series of steps, including transcription, modification, and processing of the pre-rRNA, integrated with the ordered assembly of various precursor species with ribosomal proteins.

Pre-rRNA transcription involves multiple copies of two types of transcriptional units, one encoding 5S rRNA and the other containing the genes for 18S, 5.8S, and 25S/28S rRNAs. The latter species therefore are transcribed as a single large precursor, which contains external transcribed spacers at either end (5'-ETS and 3'-ETS) as well as two internal transcribed spacers (ITS1 and ITS2) separating the mature rRNA sequences (Fig. 1A). This primary transcript is modified extensively by ribose methylation as well as pseudouridylation (14, 15), followed by removal of the spacers in a series of endonucleolytic and exonucleolytic processing reactions.

Eukaryotic pre-rRNA maturation has been characterized in most detail in the yeast *Saccharomyces cerevisiae* (reviewed in references 20, 28, 38, and 52) (Fig. 1B), where the first detectable pre-rRNA molecule is the 35S species, generated through endonucleolytic cleavage by Rnt1p in the 3'-ETS shortly downstream from the 3' end of mature 25S rRNA (1, 29). Cleavages at sites A₀, A₁, and A₂ then lead to the formation of the 20S

and 27S A₂ (27SA₂) pre-rRNAs. Endonucleolytic cleavage at site D, which occurs in the cytoplasm in yeast cells but not in other eukaryotic cells, removes the remaining ITS1 sequences from 20S pre-rRNA to form the 3' end of mature 18S rRNA (36, 40, 49).

The 27SA₂ pre-rRNA is processed into two forms of 27SB pre-rRNA via different pathways. The major one involves endonucleolytic cleavage at site A₃ by RNase MRP, resulting in 27SA₃ pre-rRNA (13, 31, 32, 39); the latter is then rapidly converted into 27SB short (27SB_S) pre-rRNA by 5'→3' exonucleolytic digestion to site B1_S, the 5' end of short-form mature 5.8S rRNA (25). The minor pathway involves purported endonucleolytic cleavage at site B1_L, located 6 or 7 nucleotides (nt) upstream of B1_S, to give 27SB long (27SB_L) pre-rRNA having the 5' end of long-form mature 5.8S rRNA (38, 52). Both 27SB species are cleaved at site C₂ within ITS2 to give the 7S and 25.5S precursors, from which the remaining ITS2 sequences are removed exonucleolytically (19, 22, 33, 34, 48).

In addition to the ribosomal proteins, a large number of nonribosomal factors play a crucial role in eukaryotic ribosome biogenesis. These *trans*-acting factors include numerous small nucleolar ribonucleoproteins as well as a host of non-small nucleolar ribonucleoproteins. With very few exceptions, inactivation of a specific *trans*-acting factor affects the biogenesis of only one of the ribosomal subunits and not the other (20, 28, 38, 52). One such exception is nucleolar protein Rrp5p, which was shown to be involved in the formation of both subunits (18, 45, 51); another is the recently identified protein Rrp12p (37).

* Corresponding author. Mailing address: Section of Biochemistry and Molecular Biology, FEW, Vrije Universiteit, de Boelelaan 1083, 1081 HV Amsterdam, The Netherlands. Phone: 31 20 444 7569. Fax: 31 20 444 7553. E-mail: vos@few.vu.nl.




FIG. 1. Processing of pre-rRNA in *S. cerevisiae*. (A) Structure of rDNA transcription unit. Thick bars represent the mature 18S, 5.8S, and 25S rRNA sequences, and thin bars represent the spacer sequences. Processing sites and locations of oligonucleotides used in this study are depicted. ETS, external transcribed spacer. (B) pre-rRNA processing pathway.

Rrp5p is an essential 193-kDa protein having a remarkable two-domain structure (18, 45) (Fig. 2). The N-terminal two thirds of the protein contain 12 S1 RNA-binding motifs, while the C-terminal portion consists of seven tetratricopeptide (TPR) repeats, motifs thought to be involved in protein-protein interactions (9, 30). Mutational analysis indicated that the two domains are also functionally distinct; deletion of various combinations of the first eight S1 RNA-binding motifs of the protein blocked processing at site A₃ and shifted the production of 5.8S rRNA from the short form to the long form as the predominant species (17, 18). Moreover, in the deletion mutants, a new processing site in ITS1, named A₄, was found midway between sites A₂ and A₃ (17). Deletion mutations in several of the TPR motifs, on the other hand, inhibited the early cleavage steps at sites A₀ to A₂, resulting in the depletion of 20S pre-rRNA and thus 18S rRNA (18, 45).

Yeast cells dependent upon mutant Rrp5p that lacks S1 RNA-binding motifs 10 to 12 (amino acids 1132 to 1354) grow normally and produce wild-type levels of all mature rRNAs (18), suggesting that S1 motifs 10 to 12 are irrelevant for ribosome biogenesis. However, detailed analysis of pre-rRNA

processing was not carried out with this *rrp5Δ6* mutant to exclude the possibility of a deviation that would not affect its ultimate outcome. The data reported in this article demonstrate that such a deviation does indeed occur in mutant cells. The absence of S1 motifs 10 to 12 leads to an Rrp5p protein that no longer supports cleavage at site A₂. Consequently, in mutant cells, the 32S precursor is cleaved directly at site A₃, giving rise to a 3'-extended 21S intermediate rather than the normal 20S pre-rRNA. The 21S pre-rRNA is then converted into mature 18S rRNA by normal cleavage at site D. The use of this alternative pathway does not significantly affect the levels of mature rRNAs or the rates of growth of the cells, demonstrating that 21S pre-rRNA is an efficient substrate for 18S rRNA synthesis. However, mutant cells are more sensitive to several antibiotics, possibly reflecting a structural abnormality in their ribosomes.

MATERIALS AND METHODS

Plasmids and strains. *Escherichia coli* strain MH1 was used for the cloning and propagation of plasmids. Yeast strains used in this study are listed in Table 1. Plasmids pTRP1-*ProtA::rrp5* and pTRP1-*ProtA::rrp5Δ6* were described by Eppens et al. (18). Mutant ribosomal DNA (rDNA) plasmids pT and pTM3 were described by Van Beekvelt et al. (46). Yeast transformation was performed as described by Gietz et al. (24). Strains YJV163 and YJV207 were obtained by transformation of strain YJV148 (*rrp5::HIS3* pURA-*RRP5*) with pTRP1-*ProtA::rrp5* and pTRP1-*ProtA::rrp5Δ6*, respectively (18). Transformants were grown on selective plates containing 2% glucose and then transferred to plates containing 5-fluoroorotic acid to select against plasmid pURA-*RRP5* and thus effect the plasmid shuffle (10). Positive colonies were checked for uracil auxotrophy.

For the integration of the *ProtA::rrp5Δ6* allele, we first cloned the 5.8-kb XbaI-PstI fragment from plasmid pTRP1-*ProtA::rrp5Δ6* into pUC19. The *LEU2* marker from pYDp-L (8) was cloned into the SacII site of the upstream sequence of the *RRP5* promoter of the plasmid-encoded mutant allele. The resulting plasmid was used to create a linear XbaI-PstI integration fragment, which was then transformed into strains YJV140 and YCA20 (the latter a kind gift from D. Tollervey). *Leu*⁺ prototrophic colonies were selected and screened by PCR to check for correct replacement of the genomic wild-type *RRP5* gene by the *rrp5Δ6* allele, resulting in strains YHV124 and YCA20Δ6. Proper expression of the mutant protein in these strains was checked by Western analysis.

Antibiotic sensitivity assay. Freshly grown colonies of YJV163 and YJV207 were resuspended in sterile water and 10-fold serial dilutions were spotted onto yeast extract-peptone-dextrose (YPD) plates containing no antibiotics, 400 μg of paromomycin/ml, 50 μg of hygromycin B/ml, or 200 μg of cycloheximide/ml. Plates were incubated at 30°C for 3 days.

RNA analysis. Cells were grown in the appropriate selective medium lacking methionine but containing either 2% glucose or 1% galactose–1% raffinose–1% sucrose (GRS) as the carbon source. To inhibit 5'→3' exonucleases, we incubated cells with 0.2 M LiCl for 1 or 2 h before harvesting the cultures. Cells were harvested at approximately 20 units of optical density at 600 nm. Isolation of RNA, Northern hybridization, and primer extension analysis were carried out as described previously (53). Oligonucleotides used as probes in these analyses are listed in Table 2, and their locations are depicted in Fig. 1A.

RESULTS

The *rrp5Δ6* deletion mutant shows antibiotic hypersensitivity. To investigate whether there is a previously unreported role for S1 motifs 10 to 12 in ribosome biogenesis, we decided first to test *rrp5Δ6* mutant cells for possible antibiotic sensitivity. Several reports have shown that yeast strains defective in the synthesis of either the large or the small ribosomal subunit are hypersensitive to the aminoglycoside antibiotic paromomycin (26, 27, 54). Furthermore, hygromycin B sensitivity has been reported for several mutants that are impaired in normal 60S subunit production, while cycloheximide sensitivity has




FIG. 2. Structure and function of Rrp5p. (A) The positions of the S1 RNA-binding motifs and the TPR motifs are indicated by triangles and pentagrams, respectively. The region removed by the $\Delta 6$ mutation and the fragments expressed in the bipartite *rrp5N2-rrp5C1* mutant (43) (see Discussion) are shown below the structure. (B) Previously constructed deletion mutants and their processing and growth phenotypes (18).

also been used as an indicator for altered 60S subunit maturation (7, 26, 35).

Using the *rrp5*-null yeast strain YJV148 (Table 1), we replaced its *URA3* plasmid containing the wild-type *RRP5* gene with a *TRP1* plasmid encoding the *rrp5* $\Delta 6$ deletion mutant and spotted the resulting strain, together with a control expressing the wild-type gene, in 10-fold serial dilutions on YPD plates containing paromomycin (400 $\mu\text{g/ml}$), hygromycin B (50 $\mu\text{g/ml}$), or cycloheximide (200 $\mu\text{g/ml}$). At these concentrations, the wild-type cells showed no or at most only a very limited reduction in growth (Fig. 3, upper rows). Growth of the mutant cells, however, was significantly retarded by all three antibiotics (Fig. 3, lower rows), the strongest effects being seen for hygromycin B (Fig. 3C) and cycloheximide (Fig. 3D). These results suggest that the *rrp5* $\Delta 6$ mutation does affect ribosome biogenesis and prompted us to analyze the pre-rRNA processing phenotype of this mutant in more detail.

S1 RNA-binding motifs 10 to 12 are essential for the processing of 32S pre-rRNA at site A₂. To monitor pre-rRNA

processing in the *rrp5* $\Delta 6$ mutant, we performed Northern analysis of total RNA isolated from exponentially growing cells that depend on either Rrp5 $\Delta 6$ p or wild-type Rrp5p. As observed previously (18), the *rrp5* $\Delta 6$ strain accumulates mature 18S and 25S rRNAs to wild-type levels (Fig. 4A). However, analysis of the various precursor rRNA species demonstrated that pre-rRNA processing in the mutant strain did differ from that in its wild-type counterpart. We were unable to detect 27SA₂ pre-rRNA in the *rrp5* $\Delta 6$ cells by using probe 3, which is complementary to the pre-rRNA just upstream from site A₃ (Fig. 4B). Instead, this probe visualized a product that migrated slightly more slowly than the 20S pre-rRNA (Fig. 4B, lane 6). This result can be seen more clearly by comparing lanes 3 and 4 of Fig. 4B, where hybridization was carried out with probe 2, which is located shortly downstream from site D. This probe detected the 20S rRNA band in the wild-type cells (Fig. 4B, lane 3) and the slightly longer product in the mutant cells (lane 4). Since the latter band was not visualized by probe 1, located just upstream from site A₁ (Fig. 4B, lane 2), or probe 4, which

TABLE 1. Yeast strains used in this study

Strain	Genotype	Reference or source
YJV140	<i>MATa ade2 his3 leu2 trp1 ura3</i>	51
YJV148	<i>MATa ade2 his3 leu2 trp1 ura3 rrp5::HIS3 pURA3-RRP5</i>	18
YJV163	<i>MATa ade2 his3 leu2 trp1 ura3 rrp5::HIS3 pTRP1-ProtA::rrp5</i>	18
YJV207	<i>MATa ade2 his3 leu2 trp1 ura3 rrp5::HIS3 pTRP1-ProtA::rrp5$\Delta 6$</i>	This study
YHV124	<i>MATa ade2 his3 leu2 trp1 ura3 ProtA::rrp5$\Delta 6$ (LEU2)</i>	This study
YHV134	YHV124 + pT	This study
YHV136	YHV124 + pTM3	This study
YCA20	<i>MATa his3$\Delta 200$ leu2$\Delta 1$ trp1 ura3-52 gal2 gal$\Delta 108$ GAL10::RRP45 RRP5</i>	5
YCA20 $\Delta 6$	<i>MATa his3$\Delta 200$ leu2$\Delta 1$ trp1 ura3-52 gal2 gal$\Delta 108$ GAL10::RRP45 ProtA::rrp5$\Delta 6$ (LEU2)</i>	This study

TABLE 2. Oligonucleotides used as probes in this study

Probe	Nucleotide sequence
1	5'-TAAAAGAAGAAGCAACAAGCAG-3'
2	5'-GCTCTCATGCTCTTGCCAAAAC-3'
3	5'-TGTTACCTCTGGGCCC-3'
4	5'-CCAGTTACGAAAATCTTGTTTTGAC-3'
5	5'-GAATGTTTGAGAAGGAAATGACGCTC-3'

is complementary to the pre-rRNA sequence between sites A₃ and B₁ (lane 8), its 5' and 3' ends probably coincide with sites A₁ and A₃, respectively. Together, these results indicate that in the *rrp5Δ6* mutant, processing at site A₂ is strongly impaired and the 32S pre-rRNA is instead cleaved directly at site A₃. This alternative processing route appears to have somewhat slower kinetics than the normal one, as concluded from the increased level of 32S pre-rRNA in the mutant cells (Fig. 4B). No 27SA₃ precursor was detected by probe 4 (Fig. 4B, lane 8), because this precursor is rapidly converted into the 27SB₅ species and is barely detectable even in wild-type cells (25) (Fig. 5, lane 1).

In order to document the inhibition of 32S pre-rRNA cleavage at site A₂ more quantitatively, we performed a reverse transcription assay with probe 5, which is complementary to the sequence that spans the 3' end of 5.8S rRNA and thus hybridizes to the 27SA and 27SB as well as the 7S precursor species. A prominent stop corresponding to site A₂ was visible when RNA from the wild-type strain was used as the template (Fig. 5, lane 1), but there was no detectable signal at this position upon analysis of RNA from the mutant strain (lane 2). Signals corresponding to site A₃ were generated with RNA




FIG. 3. Comparison of antibiotic sensitivities of strains YJV163 (*rrp5::HIS3* pTrp-*ProtA::RRP5*; indicated as wt) and YJV207 (*rrp5::HIS3* pTrp-*ProtA::rrp5Δ6*; indicated as Δ6). Ten-fold serial dilutions of equal numbers of freshly grown cells of the two strains were spotted onto YPD plates containing no antibiotic, paromomycin (400 μg/ml), hygromycin B (50 μg/ml), or cycloheximide (200 μg/ml) and incubated at 30°C for 3 days.




FIG. 4. Effect of the *rrp5Δ6* mutation on pre-rRNA processing. Total RNA was isolated from exponentially growing YJV163 or YJV207 cells expressing ProtA-Rrp5p (lanes 1, 3, 5, 7, and 9) or ProtA-Rrp5Δ6p (lanes 2, 4, 6, 8, and 10), respectively. RNA was separated on a 1.2% agarose gel containing formaldehyde and subjected to Northern analysis with the indicated probes. (A) Ethidium bromide staining to visualize mature 18S and 25S rRNAs. (B) Northern analysis. Figure 1A shows the locations of the probes. The band labeled 27S corresponds to 27SA₂ pre-rRNA for probe 3, to 27SA₂ and 27SA₃ pre-rRNAs for probe 4, and to 27SA₂, 27SA₃, 27SB₅, and 27SB_L pre-rRNAs for probe 5.

from the wild-type and mutant cells, although in the latter the level of the 27SA₃ precursor seemed to be somewhat lower, indicating an even more rapid conversion into 27SB₅ pre-rRNA in the mutant cells than in the wild-type cells. In summary, therefore, the data demonstrate a crucial role for the region of Rrp5p that encompasses S1 motifs 10 to 12 in the processing of 32S pre-rRNA at site A₂.

The 21S pre-rRNA species was observed previously in mutant yeast strains in which cleavage at site A₂ was inhibited by inactivation of the genes encoding ribosomal proteins Rps0p and Rps21p (41), by depletion of various *trans*-acting factors, including Rrp7p (6) and Rrp8p (12), or by expression of two noncontiguous fragments of Rrp5p, consisting of S1 motifs 1 to 9 and TPR motifs 1 to 7 (43). Although the collective data did indicate that 21S pre-rRNA can mature, the severely impaired growth of all of these mutants suggested it to be a rather poor substrate for 18S rRNA synthesis. The complete lack of cleavage of 32S pre-rRNA at site A₂ in the *rrp5Δ6* mutant, however, had no significant effect on the rate of growth of the cells, demonstrating that 21S pre-rRNA can be processed into mature 18S rRNA with an efficiency similar to that of the normal 20S precursor. In fact, the recent detection of low levels of 21S pre-rRNA even in wild-type cells (21) suggests that direct




FIG. 5. Reverse transcription analysis of total RNA isolated from exponentially growing YJV163 (lane 1) or YJV207 (lane 2) cells expressing ProtA-Rrp5p or ProtA-Rrp5 Δ 6p, respectively, with probe 5 (Fig. 1A). The lower panel shows a longer exposure of the area of the same gel containing the stop at site A₃. The signal marked with an asterisk is an artificial stop always observed with these strains (17).




FIG. 6. Analysis of unstable low-molecular-weight RNA molecules. Total RNA was isolated from exponentially growing cells expressing ProtA-Rrp5p (lanes 1 and 2) or ProtA-Rrp5 Δ 6p (lanes 3 and 4) with or without the addition of LiCl to a final concentration of 0.2 M and further incubation for 2 h. RNA was separated on a denaturing 8% polyacrylamide gel and subjected to Northern analysis with probe 2 (Fig. 1A).

cleavage of 32S pre-rRNA at site A₃ may be used as a minor alternative pathway for pre-rRNA processing in yeast strains.

A comparison of the polysome profile of the *rrp5* Δ 6 mutant with that of its wild-type parent did not reveal any significant differences (data not shown), a finding which we take as further evidence that pre-rRNA processing via 21S and 27SA₃ pre-rRNAs is a bona fide alternative pathway.

Maturation of 21S pre-rRNA. Regarding the manner in which 21S pre-rRNA matures, it should be noted that our data do not exclude the possibility that the absence of S1 motifs 10 to 12 disturbs the correct relative timing of the cleavages at sites A₂ and A₃ rather than inhibiting cleavage at site A₂. Thus, 21S pre-rRNA may still be a substrate for the endonuclease cutting at site A₂, resulting in the formation of the normal 20S precursor species. To assess the validity of this scenario, we performed Northern analysis with probe 2 of total RNA from exponentially growing cells expressing Rrp5 Δ 6p or wild-type Rrp5p, both before and after treatment of the cells with 0.2 M LiCl. In the absence of methionine, the addition of LiCl indirectly inhibits the 5'→3' exonucleases Rat1p and Xrn1p (16), the latter of which is primarily responsible for the degradation of the D-A₂ fragment produced by the conversion of 20S pre-rRNA into mature 18S rRNA (40). The D-A₂ fragment was readily detectable in the RNA sample from wild-type cells treated with LiCl (Fig. 6, lane 2). In accordance with earlier results (46), we noted some heterogeneity in this fragment which was likely due to incomplete inhibition of the exonucleases. No D-A₂ fragment could be detected in either untreated or LiCl-treated *rrp5* Δ 6 cells (Fig. 6, lanes 3 and 4), suggesting that 21S pre-rRNA was not cleaved at site A₂. However, there also was no trace of a D-A₃ fragment, a finding which may indicate that the remaining portion of ITS1 is removed from 21S pre-rRNA by exonucleolytic digestion rather than endonucleolytic cleavage.

To determine whether cleavage at site D is required for the maturation of 21S pre-rRNA, we used a previously created




FIG. 7. Effect of mutation of site D on the maturation of 21S pre-rRNA. Strain YJV207 expressing Rrp5 Δ 6p was transformed with either plasmid pT, encoding a tagged, wild-type rDNA unit, or pTM3, encoding a tagged, mutant rDNA unit that carries a 4-nt substitution mutation across site D which inhibits cleavage at this site (46). Total RNA was isolated from both types of transformants (lanes 1 and 3 or lanes 2 and 4, respectively) and separated on an 8% polyacrylamide gel. The gel was stained with ethidium bromide (EtBr) to visualize total mature rRNA (lanes 1 and 2). Northern analysis was performed with a probe complementary to the tag in the 18S sequence (50) to detect plasmid-derived 21S pre-rRNA and 18S rRNA (lanes 3 and 4).

cis-acting substitution mutation of the 4-nt region spanning site D (mutation M3), which completely inhibits this cleavage (46). The rrp5 Δ 6 strain was transformed with an rDNA plasmid (pTM3) that contains mutation M3 as well as a neutral tag in the 18S rRNA sequence (50), allowing discrimination between plasmid-derived and endogenous pre-rRNAs or rRNAs. Total RNA was isolated from the resulting strain as well as a control strain transformed with a similarly tagged plasmid lacking the M3 mutation (pT) and was subjected to Northern analysis with a probe complementary to the tag. Whereas rrp5 Δ 6 cells transformed with either pTM3 or pT contained comparable levels of endogenous 18S and 25S rRNAs, as visualized by ethidium bromide staining (Fig. 7, lanes 1 and 2), tagged 18S rRNA was visible only in cells containing pT (lane 3). In cells transformed with pTM3, the plasmid-derived 21S pre-rRNA containing the mutant site D region completely failed to mature (Fig. 7, lane 4). These results demonstrate that cleavage at site D remains a crucial step in the conversion of the 21S precursor into mature 18S rRNA. However, the absence of the expected D-A₃ fragment in LiCl-treated rrp5 Δ 6 cells (Fig. 6) indicates that the fragment is degraded in a manner different from that established for its normal D-A₂ counterpart. An obvious candidate for this degradation is the exosome, the major 3'→5' exonuclease complex in eukaryotic cells, for which a role in the removal of other spacer fragments has already been established (3, 4, 33).

To investigate the possible involvement of the exosome in 21S pre-rRNA processing, we constructed an rrp5 Δ 6 strain in which this exonuclease complex can be conditionally inactivated. Yeast strain YCA20 (Table 1) possesses a gene encoding the essential exosome component Rrp45p under the con-

trol of the repressible *GAL10* promoter. This strain was transformed with a linear construct containing the *ProtA::rrp5 Δ 6* gene as well as a *LEU2* marker. Genomic DNA was isolated from Leu⁺ colonies, and the successful replacement of the wild-type *RRP5* gene with its *ProtA::rrp5 Δ 6* counterpart was confirmed by PCR. Positive strains were checked for the expression of ProtA-Rrp5 Δ 6p by Western blot analysis (data not shown). The resulting double-mutant strain YCA20- Δ 6 (*GAL::rrp45 ProtA::rrp5 Δ 6*) and the wild-type control strain YCA20 (*GAL::rrp45 RRP5*) were grown in GRS medium and then shifted to glucose-based medium for 8 h, resulting in the depletion of Rrp45p and thus in the inactivation of the exosome (3). At 1 h before harvest, LiCl was added to portions of both GRS- and glucose-grown cultures to also block Rat1p and Xrn1p activities. Total RNA then was isolated from both LiCl-treated and untreated cultures and was subjected to Northern hybridization (Fig. 8).

In accordance with the data shown in Fig. 6, the D-A₂ fragment was detected by probe 2 in LiCl-treated wild-type YCA20 cells both before and after the shift to glucose (Fig. 8A, lanes 2 and 4), whereas YCA20- Δ 6 cells completely lacked this fragment, irrespective of the growth conditions (lanes 5 to 8). However, when we depleted the exosome from the latter mutant and, at the same time, inhibited 5'→3' exonuclease activity by LiCl treatment, probe 2 detected a fragment that migrated more slowly than the D-A₂ fragment (Fig. 8A, lane 8). The same fragment also hybridized with probe 3, located between sites A₂ and A₃ (Fig. 8B), but not with probe 4, located downstream from A₃ (Fig. 8C). Together with the estimated size of ~280 nt, these data indicate that the product in question corresponds to the D-A₃ fragment. We conclude that the aberrant 21S pre-rRNA is efficiently converted into 18S rRNA in the same manner as the normal 20S precursor. In contrast to the D-A₂ fragment, however, which is primarily degraded by 5'→3' exonuclease digestion (16), the D-A₃ fragment is subject to efficient degradation from either end, since it can only be detected when the activity of the 5'→3' exonucleases as well as that of the exosome is inhibited.

In this experiment, we also detected small amounts of products corresponding to A₂-E and A₃-E fragments (Fig. 8), which arose from a reversed order of ITS1 and ITS2 processing, respectively. These products were observed previously in wild-type and various mutant yeast strains (17).

DISCUSSION

Rrp5p occupies an exceptional position among the many *trans*-acting factors that participate in eukaryotic ribosome biogenesis, because contrary to the large majority of these factors, it is involved in the formation of both subunits. Previous mutational analysis indicated close agreement of this functional dichotomy of Rrp5p with its division into two distinct structural domains (Fig. 2). The C-terminal TPR domain was found to contain several elements essential for the early cleavages at sites A₀ to A₂ and thus the formation of the small subunit. The eight proximal N-terminal S1 motifs, on the other hand, appeared to be important for cleavage at site A₃ and the consequent predominant production of the short form of the large-subunit 5.8S rRNA. For the four distal S1 copies (9 to 12; amino acids 977 to 1354), no functional role was assigned,




FIG. 8. Effect of inactivation of exonucleases on the accumulation of unstable, low-molecular-weight processing products. YCA20 cells (*GAL::rrp45 Prot::rrp5*) (lanes 1 to 4) and YCA20 Δ 6 cells (*GAL::rrp4 ProtA::rrp5 Δ 6*) (lanes 5 to 8) were harvested immediately before and 8 h after a shift from GRS- to glucose-based medium (odd-numbered lanes). Equal parts of the cultures were also treated with 0.2 M LiCl for 1 h before harvest (even-numbered lanes). RNA was separated on a denaturing 8% polyacrylamide gel, and Northern analysis was performed. Figure 1A shows the locations of the probes.

because their removal did not have any noticeable effect upon growth or the steady-state levels of the mature rRNA species (18). However, the results reported in this article necessitate a refinement of this view of structure-function relationships in Rrp5p. As is evident from both Northern and primer extension analyses (Fig. 5 and 6) of the *rrp*- Δ 6 mutant, the region containing S1 motifs 10 to 12 (amino acids 1132 to 1354) is essential for cleavage at site A₂, the first of the two processing steps that normally produce the mature 3' end of 18S rRNA. Consequently, this portion of Rrp5p should be grouped functionally with the adjacent TPR domain, rather than the S1 domain, although it should be stressed that both A₀ and A₁ cleavages are not affected by deletion of S1 motifs 10 to 12. Interestingly, the revised partitioning of Rrp5p into functional domains conforms well to our finding that two contiguous fragments of the protein that abut at amino acid 1131 supply full Rrp5p functionality when expressed in *trans* (18). Individual deletion of S1 motif 9 (mutant *rrp5 Δ 5*) (18) had no effect on the levels of 27SA₂ and 20S pre-rRNAs (H. R. Vos, unpublished data); therefore, this motif is the only one of the 12 that remains functionless.

The last S1 motif (12) of Rrp5p is separated from the first TPR motif by a region of about 100 amino acids (residues 1345 to 1456) that does not contain any identifiable structural motifs and that has not been individually subjected to mutational analysis (Fig. 2). However, Torchet and Hermann-Le Denmat have shown that yeast cells expressing a bipartite *RRP5* allele that encodes two noncontiguous fragments of the protein containing S1 motifs 1 to 9 (amino acids 1 to 1131; Rrp5N2) and TPR motifs 1 to 7 (amino acids 1457 to 1729; Rrp5C1) (Fig. 2) are viable, although they bypass processing at site A₂ and show a severe growth defect resulting from the underproduction of functional 40S subunits (43). Since, compared to the Rrp5 Δ 6 protein, the bipartite Rrp5N2-Rrp5C1 protein contains an extra deletion of amino acids 1345 to 1456, it seems likely that the impaired growth is due to the lack of the latter region. Interestingly, the growth defect can be partially suppressed by the

overexpression of snR10 snoRNA (44). Since this overexpression enhances polysome formation without restoring cleavage at site A₂ or increasing the total amount of 40S subunits, it was suggested that snR10 overexpression improves 40S subunit assembly in the bipartite mutant, resulting in a higher percentage of functionally competent subunits. Thus, amino acids 1345 to 1456 of Rrp5p may constitute a further functional domain that is not involved in pre-rRNA processing but that assists in the correct assembly of 40S subunits.

Partial or complete bypassing of site A₂ and the consequent formation of 21S pre-rRNA were observed before upon mutation or depletion of several other *trans*-acting factors (6, 12, 42), as well as some small-subunit ribosomal proteins (41). However, the severe growth defects or even lethality as well as the significant reduction in 40S subunit formation invariably resulting from these mutations suggested that the 21S pre-rRNA is a considerably poorer substrate for 18S rRNA synthesis than its normal 20S counterpart. A similar conclusion was drawn for the bipartite *rrp5N2-rrp5C1* mutant discussed above (43). The recent detection of a small amount of 21S pre-rRNA in cells showing no growth defect (21) cannot be taken as evidence for its efficient conversion into 18S rRNA, since the large majority of 32S pre-rRNA in these cells is processed via the initial cleavage at site A₂. The *rrp5 Δ 6* mutant is the first strain that shows a complete lack of cleavage at site A₂ without detectable reductions in the amount of 18S rRNA and growth rate, demonstrating that cleavage at site A₂ can be bypassed without any negative consequences for the production of functional ribosomes. We therefore conclude that the alternative processing route involving the direct cleavage at site A₃ of 32S pre-rRNA into 21S and 27SA₃ rather than 20S and 27SA₂ pre-rRNAs is as effective as the normal one.

Maturation of 21S pre-rRNA proceeds via the same mechanism as that used for the conversion of the 20S precursor into 18S rRNA; i.e., it depends upon endonucleolytic cleavage at site D, the mature 3' end of 18S rRNA. This conclusion follows from the complete inhibition of 18S rRNA production by the

M3 mutation (Fig. 7), which inhibits cleavage at site D (46), as well as the detection of a D-A₃ fragment in *rrp5Δ6* mutant cells (Fig. 8). Since cleavage at site D is a cytoplasmic event (36, 40, 49), the presence of the extra ITS1 sequence does not appear to hamper the nuclear export of 21S pre-rRNA, which is part of a 43S preribosome (43). However, whereas the D-A₂ fragment is degraded exclusively from the 5' end (40), the D-A₃ fragment can be efficiently eliminated by digestion from the 5' as well as the 3' terminus, since this fragment can be detected only when both the exosome and the major 5'→3' exonucleases are inhibited (Fig. 8). The reason for the different sensitivities of the two fragments to attack by the exosome remains to be established. However, 3'-shortened forms of 21S but not 20S pre-rRNAs were detected previously (2), indicating that the 3' end of the former molecule is accessible to the exosome within the 43S preribosomal particle, whereas the 3' end of 20S pre-rRNA is not. It is possible that the exosome, which is a processive enzyme, can be loaded onto the precursor before the cleavage at site D takes place.

In multicellular organisms, the existence of two alternative, efficient routes for pre-rRNA processing is a well-established fact, even to the extent that both can operate simultaneously in the same cell (23). Like ITS1 processing in the *rrp5Δ6* mutant, either pathway includes only a single internal cleavage within ITS1 (site 3), which is thought to be carried out by RNase MRP (47) but also to require U3, although this small nucleolar ribonucleoprotein plays only a supportive and not an essential role (11). On the other hand, generation of the mature 3' end of 18S rRNA (site 2) in higher eukaryotes, unlike processing at yeast site D, is completely dependent upon U3 (23), as is cleavage at yeast site A₂. Thus, traits of yeast site A₂ recur in both site 2 and site 3 of higher eukaryotic ITS1, and the latter also may be similar to yeast site A₃. Since the data presented in this article demonstrate that, in terms of the production of functional ribosomes, cleavage at site A₂ is fully dispensable, the question arises as to why this cleavage still is part of the normal pre-rRNA processing pathway in yeast cells. A possible answer to this question may lie in the higher sensitivity of *rrp5Δ6* cells to antibiotics which, in a natural habitat, would be a disadvantage. Even though the ribosomes produced in the mutant cells are fully functional in the absence of antibiotics, shifting a portion of ITS1 from 66S to 43S preribosomes may have subtle effects on subunit assembly that increase the sensitivity of either the preribosomal particles or the completed subunits to these antibiotics. The hypersensitivity of *rrp5Δ6* cells to hygromycin B and cycloheximide indicates that the assembly of the large subunit may be affected in particular (7, 26, 35).

REFERENCES

1. Abou-Elela, S., H. Igel, and M. Ares, Jr. 1996. RNAase III cleaves eukaryotic preribosomal RNA at a U3 snoRNP-dependent site. *Cell* **85**:115–124.
2. Allmang, C., Y. Henry, H. Wood, J. P. Morrissey, E. Petfalski, and D. Tollervey. 1996. Recognition of cleavage site A2 in the yeast pre-rRNA. *RNA* **2**:51–62.
3. Allmang, C., J. Kufel, G. Chanfreau, P. Mitchell, E. Petfalski, and D. Tollervey. 1999. Functions of the exosome in rRNA, snoRNA and snRNA synthesis. *EMBO J.* **18**:5399–5410.
4. Allmang, C., P. Mitchell, E. Petfalski, and D. Tollervey. 2000. Degradation of ribosomal RNA precursors by the exosome. *Nucleic Acids Res.* **28**:1684–1691.
5. Allmang, C., E. Petfalski, A. Podtelejnikov, M. Mann, D. Tollervey, and P. Mitchell. 1999. The yeast exosome and human PM-Scl are related complexes of 3'→5' exonucleases. *Genes Dev.* **13**:2148–2158.
6. Baudin-Baillieu, A., D. Tollervey, C. Cullin, and F. Lacroute. 1997. Functional analysis of Rrp7p, an essential yeast protein involved in pre-rRNA processing and ribosome assembly. *Mol. Cell. Biol.* **17**:5023–5032.
7. Benard, L., K. K. Carroll, R. C. P. Valle, and R. B. Wickner. 1998. Ski6p is a homolog of RNA-processing enzymes that affects translation of non-poly(A) mRNAs and 60S ribosomal subunit biogenesis. *Mol. Cell. Biol.* **18**:2688–2696.
8. Berben, G., J. Dumont, V. Gilliquet, P. A. Bolle, and F. Hilger. 1991. The YDp plasmids: a uniform set of vectors bearing versatile gene disruption cassettes for *Saccharomyces cerevisiae*. *Yeast* **7**:475–477.
9. Blatch, G. L., and M. Lassle. 1999. The tetratricopeptide repeat: a structural motif mediating protein-protein interactions. *Bioessays* **21**:932–939.
10. Boeke, J. D., F. La Croute, and G. R. Fink. 1984. A positive selection for mutants lacking orotidine-5'-phosphate decarboxylase activity in yeast: 5-fluoro-orotic acid resistance. *Mol. Gen. Genet.* **197**:345–346.
11. Borovjagin, A. V., and S. A. Gerbi. 1999. U3 small nucleolar RNA is essential for cleavage at sites 1, 2 and 3 in pre-rRNA and determines which rRNA processing pathway is taken in *Xenopus* oocytes. *J. Mol. Biol.* **286**:1347–1363.
12. Bousquet-Antonelli, C., E. Vanrobays, J.-P. Gélugne, M. Caizergues-Ferrer, and Y. Henry. 2000. Rrp8p is a yeast nuclear protein functionally linked to Gar1p and involved in pre-rRNA cleavage at site A2. *RNA* **6**:826–843.
13. Chu, S., R. H. Archer, J. M. Zengel, and L. Lindahl. 1994. The RNA of RNase MRP is required for normal processing of ribosomal RNA. *Proc. Natl. Acad. Sci. USA* **91**:659–663.
14. Decatur, W., and M. J. Fournier. 2002. rRNA modifications and ribosome function. *Trend Biochem. Sci.* **27**:344–351.
15. Decatur, W. A., and M. J. Fournier. 2003. RNA-guided nucleotide modification of ribosomal and other RNAs. *J. Biol. Chem.* **278**:695–698.
16. Dichtl, B., A. Stevens, and D. Tollervey. 1997. Lithium toxicity in yeast is due to the inhibition of RNA processing enzymes. *EMBO J.* **16**:7184–7195.
17. Eppens, E. A., A. W. Faber, M. Rondajir, R. S. Jahangir, S. van Hemert, J. C. Vos, J. Venema, and H. A. Raué. 2002. Deletions in the S1 domain of Rrp5p cause processing at a novel site in ITS1 of yeast pre-rRNA that depends on Rex4p. *Nucleic Acids Res.* **30**:4222–4231.
18. Eppens, E. A., S. Rensen, S. Granneman, H. A. Raué, and J. Venema. 1999. The roles of Rrp5p in the synthesis of yeast 18S and 5.8S rRNA can be functionally and physically separated. *RNA* **5**:779–793.
19. Faber, A. W., M. van Dijk, H. A. Raué, and J. C. Vos. 2002. Ngl2p is a Cer4p-like RNA nuclease essential for the final step in 3'-end processing of 5.8S rRNA in *Saccharomyces cerevisiae*. *RNA* **8**:1095–1101.
20. Fatica, A., and D. Tollervey. 2002. Making ribosomes. *Curr. Opin. Cell Biol.* **14**:313–318.
21. Gallagher, J. E. G., and S. J. Baserga. 2004. Two-hybrid Mpp10p interaction-defective Imp4 proteins are not interaction defective in vivo but do confer specific pre-rRNA processing defects in *Saccharomyces cerevisiae*. *Nucleic Acids Res.* **32**:1404–1413.
22. Geerlings, T., J. C. Vos, and H. A. Raué. 2000. The final step in the formation of 25S rRNA in *Saccharomyces cerevisiae* is performed by 5'→3' exonucleases. *RNA* **6**:1698–1703.
23. Gerbi, S. A., and A. V. Borovjagin. 2004. Pre-ribosomal RNA processing in multicellular organisms, p. 170–198. *In* M. O. Olson (ed.), *The nucleolus*. Kluwer Academic/Plenum Publishers, New York, N.Y.
24. Gietz, D., A. S. Jean, R. A. Woods, and R. H. Schiestl. 1992. Improved methods for high efficiency transformation of intact yeast cells. *Nucleic Acids Res.* **20**:1425.
25. Henry, Y., H. Wood, J. P. Morrissey, E. Petfalski, S. Kearsey, and D. Tollervey. 1994. The 5' end of yeast 5.8S rRNA is generated by exonucleases from an upstream cleavage site. *EMBO J.* **13**:2452–2463.
26. Ho, J. H.-N., and A. W. Johnson. 1999. *NMD3* encodes an essential cytoplasmic protein required for stable 60S ribosomal subunits in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **19**:2389–2399.
27. Kressler, D., J. De La Cruz, M. Rojo, and P. Linder. 1997. Fall1p is an essential dead-box protein involved in 40S ribosomal subunit biogenesis in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **17**:7283–7294.
28. Kressler, D., P. Linder, and J. De La Cruz. 1999. Protein *trans*-acting factors involved in ribosome biogenesis in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **19**:7897–7912.
29. Kufel, J., B. Dichtl, and D. Tollervey. 1999. Yeast Rnt1p is required for cleavage of the pre-ribosomal RNA in the 3' ETS but not the 5' ETS. *RNA* **5**:909–917.
30. Lamb, J. R., S. Tugendreich, and P. Hieter. 1995. Tetratricopeptide repeat interactions: to TPR or not to TPR? *Trends Biochem. Sci.* **20**:257–259.
31. Lygerou, Z., C. Allmang, D. Tollervey, and B. Séraphin. 1996. Accurate processing of a eukaryotic precursor ribosomal RNA by ribonuclease MRP *in vitro*. *Science* **272**:268–270.
32. Lygerou, Z., P. Mitchell, E. Petfalski, B. Séraphin, and D. Tollervey. 1994. The *POPI* gene encodes a protein component common to the RNase MRP and RNase P ribonucleoproteins. *Genes Dev.* **8**:1423–1433.
33. Mitchell, P., E. Petfalski, A. Shevchenko, M. Mann, and D. Tollervey. 1997. The exosome—a conserved eukaryotic RNA processing complex containing multiple 3'→5' exoribonucleases. *Cell* **91**:457–466.
34. Mitchell, P., E. Petfalski, and D. Tollervey. 1996. The 3' end of yeast 5.8S

- rRNA is generated by an exonuclease processing mechanism. *Genes Dev.* **10**:502–513.
35. Moy, T. I., D. Boettner, J. C. Rhodes, P. A. Silver, and D. A. Askew. 2002. Identification of a role for *Saccharomyces cerevisiae* Cgr1p in pre-rRNA processing and 60S subunit biosynthesis. *Microbiology* **148**:1081–1090.
 36. Moy, T. I., and P. A. Silver. 1999. Nuclear export of the small ribosomal subunit requires the Ran-GTPase cycle and certain nucleoporins. *Genes Dev.* **13**:2118–2133.
 37. Oeffinger, M., M. Dlakic, and D. Tollervey. 2004. A pre-ribosome-associated HEAT-repeat protein is required for export of both ribosomal subunits. *Genes Dev.* **18**:196–209.
 38. Raué, H. A. 2004. Pre-ribosomal RNA processing and assembly in *Saccharomyces cerevisiae*, p. 199–222. In M. O. Olson (ed.), *The nucleolus*. Kluwer Academic/Plenum Publishers, New York, N.Y.
 39. Schmitt, M. E., J. L. Bennett, D. J. Dairaghi, and D. A. Clayton. 1993. Secondary structure of RNase MRP RNA as predicted by phylogenetic comparison. *FASEB J.* **7**:208–213.
 40. Stevens, A., C. L. Hsu, K. R. Isham, and F. W. Larimer. 1991. Fragments of the internal transcribed spacer 1 of pre-rRNA accumulate in *Saccharomyces cerevisiae* lacking 5'→3' exoribonuclease 1. *J. Bacteriol.* **173**:7024–7028.
 41. Tabb-Massey, A., J. M. Caffrey, P. Logsdan, S. Taylor, J. O. Trent, and S. R. Ellis. 2003. Ribosomal proteins Rps0 and Rps21 of *Saccharomyces cerevisiae* have overlapping functions in the maturation of the 3'-end of 18S rRNA. *Nucleic Acids Res.* **31**:6798–6805.
 42. Tollervey, D. 1987. A yeast small nuclear RNA is required for normal processing of pre-ribosomal RNA. *EMBO J.* **6**:4169–4175.
 43. Torchet, C., and S. Hermann-Le Denmat. 2000. Bypassing the rRNA processing endonucleolytic cleavage at site A2 in *Saccharomyces cerevisiae*. *RNA* **6**:1498–1508.
 44. Torchet, C., and S. Hermann-Le Denmat. 2002. High dosage of the small nucleolar RNA snR10 specifically suppresses defects of a yeast *rrp5* mutant. *Mol. Genet. Genomics* **268**:70–80.
 45. Torchet, C., C. Jacq, and S. Hermann-Le Denmat. 1998. Two mutant forms of the S1/TPR-containing protein Rrp5p affect the 18S rRNA synthesis in *Saccharomyces cerevisiae*. *RNA* **4**:1636–1652.
 46. Van Beekvelt, C. A., R. E. Jeeninga, J. Van 't Riet, J. Venema, and H. A. Raué. 2001. Identification of *cis*-acting elements involved in 3'-end formation of *Saccharomyces cerevisiae* 18S rRNA. *RNA* **7**:896–903.
 47. Van Eenennaam, H., N. Jarrous, W. J. van Venrooij, and G. J. M. Pruijn. 2000. Architecture and function of the human endonucleases RNase P and RNase MRP. *IUBMB Life* **49**:265–272.
 48. Van Hoof, A., P. Lennertz, and R. Parker. 2000. Three conserved members of the RNase D family have unique and overlapping functions in the processing of 5S, 5.8S, U4, U5, RNase MRP and RNase P RNAs in yeast. *EMBO J.* **19**:1357–1365.
 49. Vanrobays, E., P.-E. Gleizes, C. Bousquet-Antonelli, J. Noaillac-Depeyre, M. Caizergues-Ferrer, and J.-P. Gélugne. 2001. Processing of 20S pre-rRNA to 18S ribosomal RNA in yeast requires Rrp10p, an essential non-ribosomal cytoplasmic protein. *EMBO J.* **20**:4204–4213.
 50. Venema, J., A. Dirks-Mulder, A. W. Faber, and H. A. Raué. 1995. Development and application of an *in vivo* system to study yeast ribosomal RNA biogenesis and function. *Yeast* **11**:145–156.
 51. Venema, J., and D. Tollervey. 1996. *RRP5* is required for formation of both 18S and 5.8S rRNA in yeast. *EMBO J.* **15**:5701–5714.
 52. Venema, J., and D. Tollervey. 1999. Ribosome synthesis in *Saccharomyces cerevisiae*. *Annu. Rev. Genet.* **33**:216–311.
 53. Venema, J., H. Vos, A. W. Faber, W. J. van Venrooij, and H. A. Raué. 2000. Yeast Rrp9p is an evolutionarily conserved U3 snoRNP protein essential for the early pre-rRNA processing cleavages and requires box C for its association. *RNA* **6**:1660–1671.
 54. Zabetakis, D. 2000. Inheritance of suppressors of the drug sensitivity of a NSR1 deleted yeast strain. *Yeast* **16**:147–159.