



The cytoplasmic tail of L-selectin interacts with the adaptor-protein complex AP-1 subunit μ 1A via a novel basic binding motif

Received for publication, November 18, 2016, and in revised form, January 30, 2017. Published, Papers in Press, February 24, 2017. DOI 10.1074/jbc.M116.768598

✉ Karim Dib^{‡S1}, Irina G. Tikhonova[¶], Aleksandar Ivetic^{||}, and Peter Schu^{**}

From the [‡]Max Planck Institute for Biochemistry, Department of Molecular Medicine, 82152 Martinsried, Germany, the ^SCenter for Experimental Medicine, Queen's University Belfast, Belfast BT9 7BL, Northern Ireland, United Kingdom, the [¶]School of Pharmacy, Queen's University Belfast, Belfast BT9 7BL, Northern Ireland, United Kingdom, the ^{||}BHF Center for Research Excellence, Cardiovascular Division, Faculty of Life Sciences and Medicine, King's College London, London SE5 9NU, United Kingdom, and ^{**}University Medical Center Göttingen, Department of Cellular Biochemistry, Georg-August University Göttingen, D-37073 Göttingen, Germany

Edited by Luke O'Neill

L-selectin regulates leukocyte adhesion and rolling along the endothelium. Proteins binding to the cytoplasmic tail of L-selectin regulate L-selectin functions. We used L-selectin cytoplasmic tail peptide pulldown assays combined with high sensitivity liquid chromatography/mass spectrometry to identify novel L-selectin tail-binding proteins. Incubation of the L-selectin tail with cell extracts from phorbol 12-myristate 13-acetate-stimulated Raw 264.7 macrophages resulted in the binding of μ 1A of the clathrin-coated vesicle AP-1 complex. Furthermore, full-length GST- μ 1A and the GST- μ 1A C-terminal domain, but not the GST- μ 1A N-terminal domain, bind to L-selectin tail peptide, and the intracellular pool of L-selectin colocalizes with AP-1 at the *trans*-Golgi network. We identified a novel basic protein motif consisting of a cluster of three dibasic residues (³⁵⁶RR³⁵⁷, ³⁵⁹KK³⁶⁰, and ³⁶²KK³⁶³) in the membrane-proximal domain of the L-selectin tail as well as a doublet of aspartic acid residues (³⁶⁹DD³⁷⁰) in the membrane-distal end of the L-selectin tail involved in μ 1A binding. Stimulation of Raw 264.7 macrophages with PMA augmented the amount of μ 1A associated with anti-L-selectin immunoprecipitates. However, full-length GST- μ 1A did not bind to the phospho-L-selectin tail or phospho-mimetic S364D L-selectin tail. Accordingly, we propose that phosphorylation of μ 1A is required for interaction with the L-selectin tail and that L-selectin tail phosphorylation may regulate this interaction *in vivo*. Molecular docking of the L-selectin tail to μ 1A was used to identify the μ 1A surface domain binding the L-selectin tail and to explain how phosphorylation of the L-selectin tail abrogates μ 1A interaction. Our findings indicate that L-selectin is transported constitutively by the AP-1 complex, leading to the formation of a *trans*-Golgi

network reserve pool and that phosphorylation of the L-selectin tail blocks AP-1-dependent retrograde transport of L-selectin.

Leukocyte trafficking is an essential mechanism of immune surveillance. The migration to and from peripheral tissues enables leukocytes to patrol the entire body, providing immune responses as needed. The exit of leukocytes from the circulation and migration into tissues is regulated at the level of cell interactions with the vascular endothelium. At least two families of adhesion receptors participate in this interaction: the integrins and the selectins (1). Selectins mediate the initial phase of leukocyte recognition of endothelium, and this takes the form of leukocyte rolling along the vessel wall (1). There are only three types of selectins: L, E, and P. L-selectin is expressed exclusively in leukocytes and acts as a major contributor of leukocyte tethering and rolling (2). Indeed, mice deficient in L-selectin display defects in lymphocytes homing to lymphoid organs and in neutrophil accumulation in inflamed tissues (3). Unlike E- and P-selectins, L-selectin is constitutively expressed on the membrane and possesses a membrane-proximal cleavage site that is hydrolyzed after leukocyte activation, leading to shedding of its extracellular domain (4). L-selectin shedding plays a key role in the regulation of leukocyte functions. It impairs the homing of activated lymphocytes to lymph nodes (5), augments neutrophil's rolling velocity (4), controls transendothelial migration of monocytes (6), and confers lytic activity to T lymphocytes (7).

L-selectin is composed of an N-terminal lectin domain, binding endothelial ligands, followed by an epidermal growth factor-like domain, a short consensus repeat domain, a transmembrane domain, and a short cytoplasmic tail of 17 amino acids (8). The cytoplasmic tail of L-selectin is essential for L-selectin function and localization on the tips of microvilli. Indeed, L-selectin mutants lacking the membrane-distal 11 amino acid residues cannot support leukocyte rolling along endothelial venules and adhesion (9). This defect in L-selectin function cannot be explained by changes in ligand binding and is instead due to a defect in the recruitment of binding partners to L-selectin cytoplasmic tail.

This work was supported by the 7th Framework Programme for Research of the European Commission through the award of an individual Intra-European Marie Curie Fellowship (to K. D.) (project 326812). This work was also supported by Deutsche Forschungsgemeinschaft (DFG) Grant Schu802/2-4 (to P. S.) and grants from the British Heart Foundation (BHF) (to A. I.). The authors declare that they have no conflicts of interest with the contents of this article.

¹ To whom correspondence should be addressed: Center for Experimental Medicine, Queen's University Belfast, Medical Biology Center (MBC), Fl. 3, 97 Lisburn Rd., Belfast BT9 7BL, Northern Ireland, United Kingdom. Tel.: 44-28-90972634; E-mail: k.dib@qub.ac.uk.

Interaction of L-selectin with μ 1A-adaptin

A

L-selectin tail peptide: Desthiobiotin-GAhxRRLKKGKKSQERMDDPY-OH

Scrambled control peptide: Desthiobiotin-GAhxEGRMKQPDKRDSKLRKY-OH

B




Figure 1. Identification of μ 1A as a novel protein interacting with L-selectin cytoplasmic tail. A, peptide sequences used to identify proteins interacting with the L-selectin cytoplasmic tail. B, Raw 264.7 macrophages were stimulated for 1 min with PMA (1 μ g/ml), after which cells were lysed. Lysates were incubated with synthetic desthiobiotinylated peptides (scrambled control or L-selectin tail peptides) immobilized on Dynabeads Myone streptavidin. Proteins bound to the peptides were eluted, acetone-precipitated, and digested with trypsin and Lys-C. Quantitative mass spectrometric analysis was performed on a LTQ Orbitrap mass spectrometer and analyzed using the MaxQuant Software. Peptides deriving from the bait (*Sell* for L-selectin) and L-selectin-binding proteins are indicated on the graph. The AP1m1 gene encodes for the μ 1A subunit of the AP-1 complex.

To date, only a limited number of L-selectin tail binding partners are known. These include calmodulin (10), the ERM (ezrin-radixin-moesin) family of proteins (11), α -actinin (12), Lck (13), and protein kinase C (PKC) family members (14). Regions of the L-selectin tail involved in the interaction with these proteins have been mapped. For instance, calmodulin binds the unusual highly positively charged membrane-proximal N-terminal motif of dibasic residues, ³⁵⁶RRLKKGKKS³⁶⁴. Moesin binds arginine 357 and lysine 362 of this domain (15). On the other hand, the more acidic C-terminal sequence of 11 amino acids of L-selectin cytoplasmic tail binds the cytoskeletal protein α -actinin (12) and the tyrosine kinase Lck (13), whereas the binding sites for PKC family members have not yet been identified. Binding of moesin to the RRLKK motif facilitates the anchor of L-selectin to microvilli, which is critical for leukocyte adhesion and rolling (15). Calmodulin binding is thought to induce a conformational change in the extracellular lectin domain of L-selectin that renders the cleavage site resistant to proteolysis (10). Interaction of α -actinin with the C-terminal region of the tail is important for leukocyte tethering and rolling but is not essential for ligand recognition (12). Finally, the recruitment of Lck to the C-terminal tyrosine residue (13) and binding of PKC family members to the L-selectin tail (14) are essential for L-selectin outside-in signaling.

Despite this gained knowledge, several aspects of L-selectin functions remain poorly understood. For example, the mechanism controlling the replenishment of L-selectin on the membrane surface and the clustering of L-selectin on the tips of microvilli are not well known. Identification of novel L-selectin cytoplasmic tail-interacting proteins may shed light on the mechanisms regulating L-selectin functions.

The murine L-selectin tail contains only one serine residue at position 364, which is conserved in human L-selectin. L-selectin

is phosphorylated in response to T lymphocyte receptor cross-linking or treatment with PMA (14) (an activator of PKC) or engagement of chemokine receptors (16). It was proposed, but not proven, that such phosphorylation may be essential for the dynamic association/dissociation of proteins to the L-selectin tail relevant to the regulation of L-selectin functions. Indeed, moesin (11) and PKC family members (14) bind to the L-selectin tail in lymphocytes stimulated with PMA, but not in resting cells, whereas calmodulin binds L-selectin tail only in resting cells (10).

In this study, we aimed at identifying novel proteins interacting with L-selectin cytoplasmic tail domain by using sophisticated mass-spectrometry-based proteomics. We report the binding of the μ 1A-adaptin subunit of the AP-1 complex (17–19) to the L-selectin tail and the regulation of this binding by phosphorylation of L-selectin cytoplasmic tail and μ 1A. We propose that the AP-1-dependent transport of L-selectin may not be constitutive but regulated by the dynamic interaction between L-selectin tail and μ 1A in specific cellular compartments.

Results

Identification of novel proteins interacting with the L-selectin cytoplasmic domain

To identify novel proteins binding to L-selectin cytoplasmic tail, we synthesized a desthiobiotinylated peptide comprising the entire 17-amino acid-long cytoplasmic tail of murine L-selectin. As a control, a desthiobiotinylated scrambled peptide of similar size was designed. A linker between the desthiobiotin group and the first amino acid of the two peptides was added (Fig. 1A). This was to ensure that the desthiobiotin group, when coupled to Dynabeads MyOne streptavidin C1 beads, does not

interfere with the binding of proteins to the N-terminal membrane-proximal residues of L-selectin tail peptides.

To perform peptide pulldown assays, Raw 264.7 macrophages were stimulated with PMA (1 min, 1 μ g/ml) and lysed in mammalian protein extraction reagent (MPER)² buffer. Cleared lysates were incubated with either the L-selectin tail peptide (L-sel peptide) or a scrambled peptide (sc peptide) pre-coupled to Dynabeads MyOne streptavidin C1 beads. Cell lysis and experimental conditions (incubation time, washing conditions) were optimized to allow identification of novel L-selectin tail-interacting proteins. Our liquid chromatography (LC)/mass spectrometry (MS) analysis was validated by the identification of L-selectin peptides (*Sell* in Fig. 1), which derived from the bait used. We identified four novel proteins interacting specifically with L-selectin tail peptide. These include μ 1A-adaptin (AP1m1 of the adaptor-protein complex 1 AP-1, ID: 11767), Cfh (complement component factor h, ID: 12628), Grn (granulin, ID: 14824), and Lrrc48 (leucine-rich repeat containing 48, ID: 74665) (Fig. 1B). The highest MS signal intensity associated with L-selectin tail peptide was obtained for μ 1A; therefore, we decided to analyze this interaction in more detail.

μ 1A is one of the four adaptin subunits of the adaptor-protein complex 1 (AP-1). AP-1 controls the sorting of transmembrane proteins into clathrin-coated transport vesicles, mediating transport between the *trans*-Golgi network (TGN) and endosomes (17, 18, 20). This way, this vesicular carrier regulates the transport of cargo proteins (including receptors) to the plasma membrane (21).

The C-terminal domain of μ 1A interacts with L-selectin

To confirm the μ 1A/L-selectin tail interaction, we investigated the binding of full-length μ 1A as well as of μ 1A subdomains, expressed in *E. coli* as GST-fusion proteins, to L-selectin tail peptide. μ 1A consists of a N-terminal domain (aa 1–153), positioned close to the membrane, and a C-terminal domain (aa 154–423), which binds the canonical YXX \emptyset sorting motif (where \emptyset is a bulky hydrophobic residue) (17, 18, 22, 23). Expression of cDNAs encoding for full-length μ 1A (GST-FL- μ 1A), the C- (GST-C- μ 1A) or the N-terminal domain (GST-N- μ 1A) of μ 1A *in E. coli* resulted in the synthesis of GST-fusion proteins with predicted molecular weights as shown on the polyacrylamide gel stained with Coomassie Blue (Fig. 2A).

We next incubated the L-sel peptide, or the sc peptide with purified GST-FL- μ 1A. Thereafter, the beads were collected, washed, and resuspended in Laemmli buffer. Proteins were resolved on 10% SDS-PAGE, transferred onto a PVDF membrane, and subjected to Western blotting analysis using an anti-GST Ab (Fig. 2, B and C) or an anti- μ 1A Ab (Fig. 2D). As shown in Fig. 2B, GST-FL- μ 1A binds to the L-sel peptide but not to the sc peptide. GST (added in excess) did not bind to the L-sel peptide, confirming binding specificity. In addition, we found

that GST-C- μ 1A, but not GST-N- μ 1A, binds the L-sel peptide (Fig. 2, C and D). Thus, the C-terminal domain of μ 1A binds L-selectin cytoplasmic tail, as it has been demonstrated for proteins with canonical YXX \emptyset sorting motifs, whereas, as expected, the N-terminal domain, which binds the β 1 subunit of the AP-1 complex (23), did not bind to the L-selectin tail.

μ 1A is associated with anti-L-selectin receptor immunoprecipitates

We next sought to characterize the interaction between μ 1A, as part of the AP-1 complex, and L-selectin tail in cells in culture. To this end, Raw 264.7 macrophages were stimulated or not for different time periods with PMA (1 μ g/ml), before being lysed. Cell lysates were incubated with either an anti-L-selectin Ab or a control isotype-matched IgG2a. μ 1A (AP-1) association with anti-L-selectin immunoprecipitates was investigated by Western blotting analysis using an anti- μ 1A Ab.

We found little association between L-selectin and μ 1A (AP-1) at steady state in resting cells. However, we observed a time-dependent increase in the amount of μ 1A (AP-1) associated with anti-L-selectin immunoprecipitates upon stimulation of the cells with PMA (1 μ g/ml) (Fig. 3A, top). As a control, we showed that μ 1A (AP-1) is not associated with IgG2a immunoprecipitates (Fig. 3B, top). We also verified that anti-L-selectin Abs precipitated L-selectin (Fig. 3, A and B, bottom panels), as evidenced by the detection of proteins with molecular masses of 75–120 kDa representing the different glycosylated forms of L-selectin (24). However, due to its low level of expression in Raw 264.7 macrophages, the presence of L-selectin in crude cell lysates could not be detected under these experimental conditions.

L-selectin and AP-1 *in vivo* colocalization

To test for *in vivo* L-selectin and AP-1 colocalization by confocal microscopy, we used an established THP-1 cell line expressing wild type, full-length L-selectin green fluorescent protein (GFP) (6). Endogenous AP-1 was labeled with an antibody directed against the γ 1 adaptin AP-1 subunit (20). This antibody recognizes the C-terminal flexible region of γ 1, which is even accessible when AP-1 is part of a membrane coat. Actin staining (*red*) served as control. Images show two pools of L-selectin (*green*), one at the plasma membrane and one in intracellular vesicles (Fig. 4, A and B). The intracellular pool is clustered around large membrane domains that have AP-1 bound (*blue*) and therefore represent TGN membranes.

Mander's colocalization coefficient was measured to prove L-selectin/ μ 1A (AP-1) colocalization in intracellular vesicles (overlap coefficient between *blue* and *green* signals). We compared the colocalization coefficient measured in an area including intracellular vesicles with the colocalization coefficient measured in an area containing the plasma membrane (Fig. 4B). Because there is no AP-1 at the plasma membrane, the colocalization coefficient measured in this area is not a real colocalization signal and is used as a background reference.

We found that the colocalization coefficients were 0.80 ± 0.09 and 0.15 ± 0.01 for intracellular vesicles and plasma membrane areas, respectively (Fig. 4C) ($p < 0.0001$). This result

²The abbreviations used are: MPER, mammalian protein extraction reagent; Ab, antibody; CCV, clathrin-coated vesicles; EE, early endosomes; PMA, phorbol 12-myristate 13-acetate; TBS, Tris-buffered saline; TGN, *trans*-Golgi network; MPR, mannose 6-phosphate receptor(s); Ahx, hexanoic acid; TRITC, tetramethylrhodamine isothiocyanate.

Interaction of L-selectin with μ 1A-adaptin




Figure 2. μ 1A binds L-selectin cytoplasmic tail peptide *in vitro*. A, BL21 bacteria were transformed with pGEX-5X3 plasmids encoding for the different GST forms of μ 1A proteins, including full-length μ 1A (GST-FL- μ 1A) or the N-terminal part (GST-N- μ 1A) or C-terminal part (GST-C- μ 1A) of μ 1A. After induction with isopropyl β -D-thiogalactopyranoside, bacteria were collected and lysed, and lysates were incubated with glutathione-Sepharose beads. After 1 h, the beads were collected and washed, and 2 \times Laemmli buffer containing 1 mM DTT was added. Samples were run on 10% SDS-PAGE, and proteins were stained with Coomassie Brilliant Blue. The positions of GST-FL- μ 1A, GST-N- μ 1A, GST-C- μ 1A, and GST are indicated on the left by arrows. B, purified GST-FL- μ 1A or GST were incubated with a scrambled peptide (sc peptide) or L-selectin cytoplasmic tail peptide (L-sel tail peptide) coupled to streptavidin-coated beads. After 1 h, the collected beads were washed and resuspended in Laemmli buffer. The proteins were separated on 10% SDS-PAGE and transferred onto a PVDF membrane, which was subjected to Western blotting analysis using an anti-GST Ab (top). The positions of GST-FL- μ 1A and GST are indicated on the left by arrows. Loading controls (which correspond to the amount of GST-FL- μ 1A or GST used for the peptide pulldown assay) are shown below. Note that an excess of GST versus GST-FL- μ 1A has been used. C, binding of GST-N- μ 1A or GST-C- μ 1A to a scrambled peptide (sc peptide) or L-selectin tail peptide (L-sel tail peptide) was carried out as described in B. D, binding of GST-FL- μ 1A or GST-C- μ 1A to a scrambled peptide (sc peptide) or L-selectin tail peptide (L-sel tail peptide) was carried out as described in B and C with the exception that Western blotting analysis was carried out with an anti- μ 1A Ab.

demonstrates that L-selectin and AP-1 (μ 1A) colocalize in intracellular vesicles.

The limited L-selectin/AP-1 colocalization is in line with the transient AP-1/cargo protein interactions during protein sorting and transport via clathrin-coated vesicles (CCV). Thus, TGN-associated L-selectin may serve as a reserve pool. These proteins will be transported to the plasma membrane when this pool needs to be replenished. One function of AP-1 is the sorting of mannose 6-phosphate receptors (MPR), which facilitate the transport of soluble lysosomal enzymes from the TGN to endosomes. AP-1-dependent MPR sorting ensures that 50% of the cellular MPR pool resides in the TGN (awaiting newly syn-

thesized lysosomal enzymes), whereas only 10% of the cellular pool is at the plasma membrane due to their highly efficient AP-2-mediated clathrin-dependent endocytosis (20, 21). Thus, AP-1-dependent L-selectin sorting might be responsible for the formation of a localized L-selectin reserve pool in the TGN.

Phosphorylation of Ser-364 of L-selectin tail peptide prevents binding of μ 1A

To investigate whether phosphorylation of serine 364 of the L-selectin tail affects μ 1A binding, we synthesized an L-selectin tail peptide, in which serine 364 was replaced by a phosphoserine (L-sel-p-peptide). We found that GST-FL- μ 1A, GST-C-




Figure 3. μ 1A is associated with anti-L-selectin immunoprecipitates. Raw 264.7 macrophages, stimulated or not with PMA (1 μ g/ml) for different time periods, were lysed, and lysates were subjected to immunoprecipitation (IP) with an anti-L-selectin Ab (A) or a control IgG2a (B). After 1 h, protein A-Sepharose beads were added for an additional 1 h. The beads were then collected by centrifugation, washed, and resuspended in 2 \times Laemmli buffer. The immunoprecipitates were subjected to 10% SDS-PAGE and transferred to a PVDF membrane, which was cut into two pieces. The lower part of the membrane was immunoblotted with an anti- μ 1A Ab (A and B, top), and the top part of the membrane was immunoblotted with an anti-L-selectin Ab (A and B, bottom), as described under "Experimental procedures." The arrows indicate the position of μ 1A, L-selectin, and IgG.

μ 1A, or murine μ 1A (AP-1) from cell lysates bound the L-sel peptide but not the L-sel-p-peptide (Fig. 5A). We also designed a L-selectin peptide in which serine 364 was replaced by an aspartic acid residue (L-sel S364D) to imitate a constitutively phosphorylated L-selectin tail. Again, we found that GST-FL- μ 1A, GST-C- μ 1A, or murine μ 1A (AP-1) from a cell lysate bound the L-sel peptide but not the L-sel S364D peptide (Fig. 5A). To ensure that this effect was due to the introduction of a negative charge at position 364 of the L-selectin tail and not to the replacement of serine 364 by another amino acid, we also conducted experiments with a control peptide in which serine 364 was replaced by an alanine residue (L-sel S364A). We found that GST-FL- μ 1A (Fig. 5B, left) or GST-C- μ 1A (Fig. 5B, right) bound equally well to L-sel and L-sel S364A peptides. Thus, we concluded that phosphorylation of the L-selectin tail on serine 364 abrogates its interaction with μ 1A (AP-1).

Clusters of positively charged amino acids are essential for μ 1A interaction with the L-selectin tail

The membrane-proximal domain of the L-selectin tail contains an unusual cluster of three dibasic (positively charged) amino acids, ³⁵⁶RR³⁵⁷, ³⁵⁹KK³⁶⁰, and ³⁶²KK³⁶³. To investigate whether this RRLKKGKK sequence of the L-selectin tail is involved in the interaction with μ 1A, we designed peptides in which one arginine or one lysine residue within any of the three dibasic motifs was exchanged by an alanine residue (Fig. 6A). These peptides were incubated with GST-FL- μ 1A or GST-C- μ 1A, and an interaction between the proteins and peptides was analyzed as described above.

We found that replacing a single arginine or a single lysine residue within any of the three dibasic clusters was sufficient to prevent GST-FL- μ 1A (Fig. 6B, left) or GST-C- μ 1A (Fig. 6B, right) from interacting with the L-selectin tail. We also investigated whether the C-terminal, negatively charged aspartic acid residues Asp-369 and Asp-370 of the L-selectin tail have a func-




Figure 4. Subcellular distribution of L-selectin and AP-1 in monocyte-like THP-1 cell line. THP-1 cells stably expressing wild type L-selectin, C-terminally tagged with green fluorescent protein (WT L-selectin-GFP), were prepared for confocal microscopy as described under "Experimental procedures." Cells were labeled for endogenous AP-1 (anti- γ 1 subunit, blue) and actin (TRITC-phalloidin, red). A, L-selectin (green) is present at the plasma membrane and within discrete punctae that colocalize with AP-1-positive signal within the TGN. B, the rectangles show the areas used to perform quantification of L-selectin/ μ 1A colocalization. A region of interest of similar dimension was used to measure intracellular regions and at the plasma membrane. C, the Volocity software was used to measure the overlap coefficient between blue and green signals corresponding to AP-1 and L-selectin, respectively. Each signal was thresholded to remove background signals before analysis. Threshold parameters were maintained in all of the analysis. Instrument parameter settings remained constant between image acquisitions. The data represent means of colocalization co-efficiency \pm S.E. (error bars) of 22 cells prepared over three independent experiments. ****, $p < 0.0001$ (unpaired Student's t test). Scale bar, 7.5 μ m. PM, plasma membrane; IC, intracellular pool.

tion in μ 1A binding. We replaced separately each aspartic acid residue by an asparagine residue (D369N or D370N), or both were substituted with asparagine residues (D369N/D370N) (Fig. 6C). We found that replacement of one or two aspartic acid residues by asparagine augmented the binding of GST-C- μ 1A to these L-selectin mutated tails (Fig. 6D). Thus, the highly positively charged membrane-proximal region of the L-selectin tail is essential for μ 1A binding, whereas its membrane-distal part, with its doublet of negatively charged amino acids, weakens the interaction with μ 1A.

Molecular docking of the L-selectin tail to μ 1A

To identify the μ 1A surface domain binding L-selectin cytoplasmic tail and to interpret the results of μ 1A binding to the L-selectin tail with substituted amino acids, we explored docking of the L-selectin tail to the crystal structure of μ 1A, as determined in the tetrameric AP-1 complex. To do this, we took into consideration the fact that the L-selectin tail binding surface domain of μ 1A has to be accessible and not sterically blocked by the interactions of μ 1A with the other subunits of the AP-1

Interaction of L-selectin with μ 1A-adaptin




Figure 5. Phosphorylation of serine 364 of L-selectin cytoplasmic tail prevents the binding of μ 1A. *A*, purified GST-FL- μ 1A or GST-C- μ 1A was incubated with a scrambled peptide (sc peptide), L-selectin tail (L-sel tail peptide), a phosphorylated L-selectin tail peptide (L-sel-p-peptide), or the L-selectin tail peptide in which serine 364 was replaced by an aspartic acid (L-sel S364D peptide). Thereafter, association of GST-FL- μ 1A (*A*, left), GST-C- μ 1A (*A*, middle) or μ 1A assembled into an AP-1 complex (*A*, right) with the different L-selectin tail peptides was determined as described in the legend to Fig. 2. *B*, purified GST-FL- μ 1A or GST-C- μ 1A was incubated with sc peptide, L-selectin tail peptide, or L-sel S364A peptide. Thereafter, association of GST-FL- μ 1A (*B*, left) or GST-C- μ 1A (*B*, right) with the different peptides was measured as described in *A*.

complex. Fig. 7A shows the favorable docking pose, which explains the binding specificities of μ 1A to L-selectin tail peptides. The L-selectin tail/ μ 1A interaction is predicted to have a strong electrostatic nature generated by two complementary charged surface areas at the binding interface. The positively charged cluster of arginine and lysine residues of the N-terminal region of the L-selectin tail are matched by several negatively charged residues, namely the aspartate and glutamate residues Glu-311, Glu-337, Glu-340, and Asp-417 of μ 1A. In addition, the positively charged residues of the L-selectin tail form hydrogen bonds with residues Asn-318, Asn-338, and Gln-419 of μ 1A. The interactions cited above are disrupted when the positively charged residues of the RRLKKGKK motif of the L-selectin tail are substituted to alanine, and this explains the loss of μ 1A binding. Ser-364 of the L-selectin tail does not form a specific interaction with μ 1A, in line with the unaltered binding of the S364A tail to μ 1A, but it is in close proximity to the negatively charged residues Glu-311 and Glu-340 of μ 1A. Therefore, introducing a negatively charged group at position 364 of the L-selectin tail by the addition of a phosphate group or by substituting serine 364 by an aspartic acid residue will cause electrostatic repulsion, thus preventing binding of μ 1A to the L-selectin tail.

The aspartic acid residues at positions 369 and 370 of the L-selectin tail form polar interactions with Lys-199 and Arg-410 of μ 1A, respectively, which would stabilize the binding of μ 1A. However, Lys-199 and Arg-410 have also polar inter-

actions with Glu-261 and Asp-174 of μ 1A. The aspartate residues at positions 369 and 370 of the L-selectin tail could have some repulsion with Glu-261 and Asp-174 as well as the nearby Glu-381, which would weaken the binding. Asparagine residues at these positions could form interactions not only with the positively charged Lys-199 or Arg-410, but also with the negatively charged Glu-261, Asp-174, or Glu-381 residues of μ 1A, thereby enhancing the binding of the two proteins. This explains the augmented binding of μ 1A to the L-selectin tail peptides in which the aspartic acid residues 369 and 370 were replaced by asparagine residues. Our predicted binding mode is in good agreement with our biochemical binding experiments.

Since μ 1A (AP-1) and μ 2 (AP-2) are highly homologous and because both bind YXX Φ -based sorting motifs and have overlapping cargo specificities, we also modeled the L-selectin tail onto the C-terminal μ 2 domain to examine whether a binding mode could permit an interaction, although μ 2 was not identified in our L-selectin tail pulldown experiments (Fig. 1). Docking predicts an interaction energy of 150–275 kcal/mol for L-selectin/ μ 2 binding poses, contrasting with the 400–500 kcal/mol for L-selectin/ μ 1A binding poses. Thus interactions of the L-selectin tail with μ 2 are much less favorable, confirming the pulldown experiment. It is likely that L-selectin is not endocytosed by the AP-1-homologous plasma membrane AP-2 complex.




Figure 6. Replacement of arginine or lysine residues within the clusters of dibasic residues prevents binding of μ 1A. *A*, amino acid sequence of L-selectin cytoplasmic tail and the different substitutions in the three clusters of dibasic residues is shown. *B*, purified GST-FL- μ 1A or GST-C- μ 1A were incubated with a scrambled peptide (sc peptide), L-selectin cytoplasmic tail (L-sel tail peptide), or peptides in which one amino acid within the clusters of dibasic residues was replaced by an alanine residue. These include L-sel R357A peptide, L-sel R360A peptide, and L-sel R363A peptide. Thereafter, association of GST-FL- μ 1A (*B*, left) or GST-C- μ 1A (*B*, right) with the different peptides was determined as described in the legend to Fig. 2*D*. *C*, amino acid sequence of L-selectin cytoplasmic tail with substitutions of aspartic acid residues by asparagine. *D*, purified GST-C- μ 1A was incubated with sc peptide, L-sel tail peptide, or peptides in which one or two of the aspartic acid residues was replaced by an asparagine residue. These include L-sel D3369N peptide, L-sel D370N peptide, and the double-substituted L-sel DDNN peptide. Thereafter, association of GST-C- μ 1A with the different peptides was determined as described above.

Discussion

In this study, we performed L-selectin cytoplasmic tail pull-down assays combined with a high sensitivity LC/MS platform to identify novel L-selectin tail-binding proteins in cell extracts from stimulated Raw 264.7 macrophages. By using LC/MS-based peptide sequencing, we identified the AP-1 adaptor-protein complex subunit μ 1A as a novel binding protein of the L-selectin tail. We confirmed this interaction by showing that 1) μ 1A expressed as a GST-fusion protein binds to the L-selectin tail, 2) μ 1A as part of a fully assembled AP-1 complex is associated with anti-L-selectin immunoprecipitates, 3) L-selectin

is present as an intracellular pool, which colocalizes with AP-1, and 4) the C-terminal μ 1A domain binds selectively the non-phosphorylated L-selectin tail.

μ 1A is one of four adaptin subunits of the AP-1 complex. AP-1 selects the cargo proteins for CCV, and it coordinates CCV formation as well as CCV uncoating, a prerequisite for vesicle-organelle membrane fusion. Besides μ 1A, the AP-1 complex contains the adaptin subunits γ , β 1, and σ 1A (18, 19). To incorporate cargo proteins into transport vesicles, μ 1A binds to specific sequences (called sorting motifs or cargo-binding domains) within the cytoplasmic domains of such proteins. μ 1A is composed of two domains. The N-terminal domain (amino acids 1–153) binds the β 1 subunit of the AP-1 complex and thus contributes to the assembly of the AP-1 complex. The C-terminal μ 1A domain (amino acids 153–422) binds sorting motifs of cargo proteins incorporated into CCV (23). In keeping with these structure/function studies of the AP-1 complex, we discovered that GST-C- μ 1A but not GST-N- μ 1A binds to the L-selectin tail.

Two canonical sorting motifs in cytosolic tails of transmembrane proteins have been described for AP-1: the tyrosine-based motif YXX Φ , which is bound by the μ 1A C-terminal domain, and the dileucine-based motifs (e.g. (D/E)XXXL(L/I)), which are bound by σ 1A and/or σ 1B adaptins (25, 26). L-selectin cytoplasmic tail does not contain either of these two sorting motifs. This means that other sequences must be involved in the interaction with μ 1A.

In addition to these two canonical motifs, non-canonical binding motifs have been identified that interact with AP-1 as well. For example, yeast AP-1 binds a 12-residue domain of the pheromone receptor Ste13p, which contains a tribasic sequence, KRK (27). In addition, AP-1 binds to the highly basic RARHRRNVDR sequence of the polymeric immunoglobulin receptor (28). Based on these examples, the three dibasic clusters of amino acids in L-selectin tail most likely represent a novel, basic, non-canonical binding motif sequence for μ 1A. To investigate this possibility, we designed L-selectin tail peptides in which one arginine or lysine residue within any of the three dibasic motifs was exchanged by an alanine residue (R357A, K360A, or K363A). We showed that GST-FL- μ 1A or GST-C- μ 1A did not bind R357A, K360A, or K363A L-selectin tail peptides. This experiment demonstrated that each dibasic residue within the N-terminal membrane-proximal region of the L-selectin tail is essential for binding μ 1A. To understand the role of residues Arg-357, Lys-360, and Lys-363 for μ 1A interaction, we used molecular docking to predict the predominant binding mode of the L-selectin tail with the 3D structure of μ 1A. We found that the positively charged cluster of arginine and lysine residues of the N-terminal region of the L-selectin tail ³⁵⁶RRLKKGKKS³⁶⁴ are matched by aspartate and glutamate residues Glu-311, Glu-337, Glu-340, and Asp-417 of μ 1A and form hydrogen bonds with residues Asn-318, Asn-338, and Gln-419 of μ 1A. Whereas this L-selectin tail region also binds moesin/ezrin, it has been shown that only Arg-357, and not Lys-360 or Lys-363, is required for this interaction (15). Thus, the motifs mediating the binding of moesin/ezrin and μ 1A overlap, but they involve different amino acid residues. It was reported that association of calmodulin with the L-selectin tail

Interaction of L-selectin with μ 1A-adaptin




Figure 7. Molecular model of L-selectin/ μ 1A interactions. *A, left*, L-selectin tail docked to the surface of μ 1A. The surface of μ 1A is colored based on the electrostatic potential; red and blue show negatively and positively charged areas, correspondingly. Residues of the L-selectin tail used in mutagenesis are labeled. *A, right*, binding interactions between L-selectin tail and μ 1A. Hydrogen bonds are shown as a black dotted line. Residues of the L-selectin tail and μ 1A are labeled in red and black, respectively. *B*, model for the AP-1-dependent L-selectin sorting in leukocytes at rest (*left*) and upon stimulation (*right*). P_i , the addition of inorganic phosphate to the L-selectin tail and μ 1A. **Boldface letters** indicate a higher phosphorylation rate of μ 1A in stimulated leukocytes compared with cells at rest. **Boldface arrows** indicate enhanced TGN-to-endosome L-selectin transport by AP-1 due to phosphorylation of μ 1A and thus more activated AP-1 complexes. Phosphorylation of the L-selectin tail on serine 364 in EE causes the transient accumulation of L-selectin in EE, because this L-selectin cannot bind μ 1A, leading to its exocytosis to the plasma membrane (PM).

is influenced by the membrane bilayer. Thus, in a phospholipid bilayer (which mimics the inner leaflet of the plasma membrane), the positively charged N-terminal part of the L-selectin tail interacts with anionic phosphatidylserine lipids at the membrane interface through electrostatic interactions. This prevents calmodulin to interact with the N-terminal part of the L-selectin tail (29) and promotes L-selectin shedding (15). Disruption of the interaction between the positively charged N-terminal part of the L-selectin tail with the phospholipid bilayer is a prerequisite for the binding of calmodulin to the L-selectin tail. Such disruption is brought about by the interaction of the FERM domain of moesin with L-selectin to induce a separation of the cationic N-terminal part of the L-selectin tail from the phospholipids (30). We do not know whether the phospholipid bilayer plays a similar role in intact cells for at least two reasons. First, the lipid bilayer is itself a complex mixture of lipids. Its structure in the plane of the bilayer is that of a microscopic mosaic of regions that differ in composition (*i.e.* domains or clusters) (31). In particular, receptors are found in domains, such as the caveola, that are rich in cholesterol. Second, the cited authors (29, 30) have not taken into account in

their model that calcium interacts with phospholipid bilayers. Such interactions induce changes in lipid dynamics, structure, and affinity of phospholipid-binding proteins (32).

AP complexes are targeted to their respective membrane via binding to specific phosphoinositides, AP-1 to phosphatidylinositol 4-phosphate and AP-2 to phosphatidylinositol 4,5-bisphosphate. There are no data in the literature describing inhibition of a sorting motif/AP complex binding by basic residues located in the vicinity of the sorting motif (23, 25, 26). In fact, phosphatidylinositol 4-phosphate incorporated into liposomes binds AP-1, and the interaction between adaptins and sorting motif-containing proteins is further stabilized by phospholipids during the polymerization of the vesicle coat (33). These results indicate a role of phospholipids and phosphoinositides in the interaction of AP-1 with its binding partners and vesicular assembly.

The μ 2 adaptin of the plasma membrane AP-2 complex collects proteins for clathrin-mediated endocytosis, and many proteins are transported by AP-1 as well as AP-2 CCV (17, 18). μ 2 was not isolated in our screen, despite the fact that it is highly homologous to μ 1A. Therefore, we asked why μ 2 could

not bind L-selectin tail. This is probably explained by the fact that the key amino acids in μ 1A involved in L-selectin tail binding are not found in μ 2. Indeed, only the first glutamic acid residue of the L-selectin binding motif in μ 1A is conserved in μ 2 (Glu-311 > Glu-321, Glu-337 > Ser-347, Glu-340 > Ala-350, Asp-417 > Ile-430). Also, the asparagine residues 318 and 338 and glutamine 419 of μ 1A are not conserved in μ 2, where they are replaced by leucine and glutamic acid residues. These data strongly indicate that the AP-2 complex, which is highly homologous to the AP-1 complex, does not play a role in the regulation of L-selectin cell surface expression.

We tested for L-selectin and AP-1 colocalization *in vivo* by confocal microscopy. Besides the plasma membrane pool of L-selectin, we detected an intracellular pool of L-selectin localized next to AP-1 decorated domains of the TGN. This indicated that AP-1-dependent L-selectin sorting might be responsible for the formation of this putative L-selectin reserve pool.

Murine L-selectin tail contains one unique serine residue at position 364, which is conserved in human L-selectin. It was shown that upon stimulation of lymphoblastoid cell lines with PMA or chemoattractants, the L-selectin tail becomes rapidly phosphorylated (16), and this is probably caused by recruitment of PKC family members to the L-selectin tail (14). However, little is known of the relationship between L-selectin tail phosphorylation and its interaction with binding partners *in vivo*. We found low amounts of μ 1A associated with anti-L-selectin immunoprecipitates in resting Raw 264.7 macrophages. However, this association increased when the cells were stimulated with PMA. This result could be explained in at least two ways, involving phosphorylation of μ 1A and/or L-selectin tail. One phosphorylation step would be the enhanced phosphorylation of μ 1A in response to PMA, because such phosphorylation would augment its interaction with the L-selectin tail. Indeed, the cytoplasmic AP-1 complex is present in a closed conformation, in which the cargo binding domains of μ 1A and σ 1A are blocked by γ 1 and β 1 adaptins. The complex has to undergo a conformational change to an "open" state. This causes the release of the μ 1A C-terminal domain from the large adaptins, enabling its movement toward the membrane and the cytoplasmic sorting domains of membrane proteins. The open conformation is favored by threonine phosphorylation of μ 1A in the flexible linker region connecting the N- and C-terminal domains (34–36). Such a mechanism of activation has also been described for the L-selectin tail-interacting protein moesin. Phosphorylation of moesin by PKC α on a threonine residue located in the C-terminal part unfolds the protein, thus allowing it to interact with the L-selectin tail (37).

The second phosphorylation reaction to explain a PKC-stimulated association between μ 1A and L-selectin tail would be the L-selectin tail phosphorylation. However, μ 1A does not bind the phospho-L-selectin tail peptide, and this is due to the fact that the phosphate group on serine 364 is in close proximity to the negatively charged residues Glu-311 and Glu-340 of μ 1A. In keeping with our results, other investigators showed inhibition of calmodulin binding to L-selectin-p-tail *in vitro* and in intact THP-1 cells (6).

The AP-1 coat is found on the TGN and on early endosomes (EE), and it mediates constitutive protein transport from the EE

to the TGN as well as from the TGN to EE, as exemplified by the mannose 6-phosphate receptors, furin and sortilin (20, 21, 26). Interestingly, besides its role as cargo binding and constitutive regulator of protein transport, the role of AP-1 in regulated secretory pathways has also been shown (38–40). Our data showing that phosphorylation of both μ 1A and the L-selectin tail controls the interaction between these two proteins indicate a regulated, rather than a constitutive, AP-1-dependent secreted pathway for L-selectin. We developed a model of AP-1-dependent L-selectin sorting based on our data and the known functions of AP-1 (Fig. 7B). We propose that stimulation of μ 1A phosphorylation activates AP-1 and enhances AP-1-dependent protein transport from the TGN to EE. Phosphorylation of the L-selectin tail in activated leukocytes would prevent its binding to μ 1A (AP-1) in EE, leading to less incorporation of L-selectin into AP-1 CCV for retrograde EE to TGN transport. Consequently, L-selectin would transiently accumulate in EE, from which it can be readily transported to the plasma membrane through exocytosis (21). Such an indirect function of AP-1 in the regulation of plasma membrane protein transport has been demonstrated by us for both mannose 6-phosphate receptors in mouse fibroblasts (20, 21). Thus large amounts of L-selectin would be exported to the plasma membrane and could be readily clustered on microvilli relevant for leukocyte rolling and adhesion. We are currently investigating this working hypothesis of stimulation-dependent compartment-specific L-selectin tail and μ 1A phosphorylation in leukocytes.

In summary, we have made several novel and important observations regarding L-selectin functions. First, we identified μ 1A as a novel protein interacting with the L-selectin tail. Second, we demonstrated colocalization of an intracellular pool of L-selectin with AP-1. Third, a novel binding motif formed by a triplet of dibasic amino acid motifs in L-selectin (RRXKKXKK) mediates this interaction. Fourth, L-selectin tail phosphorylation inhibits μ 1A binding. From these data we can conclude that AP-1, but not AP-2, is involved in the transport pathway regulating L-selectin secretion to the plasma membrane upon stimulation.

Experimental procedures

Antibodies

The antibodies and their sources were as follows. Anti-L-selectin Ab lam 1–116 (sc-13505) was purchased from Santa Cruz Biotechnology (Heidelberg, Germany); the mouse IgG2a isotype control Ab (UPC-10), was purchased from Sigma-Aldrich (Munich, Germany); the anti- μ 1A Ab (ab 170277) was bought from Abcam (Cambridge, UK), the anti-GST mouse monoclonal Ab (catalog no. 71097-3) was from Novagen/Merck (Darmstadt, Germany).

Chemicals

Protease inhibitor tablets were from Roche Applied Science (Penzberg, Germany), and phosphatase inhibitor mixtures 1 and 2 were purchased from Sigma-Aldrich. Dynabeads Myone streptavidin C1 and MPER buffer were from Invitrogen/Thermo Fisher Scientific (Karlsruhe, Germany), and Protein A-Sepharose TM CL-4B and streptavidin-Sepharose beads

Interaction of L-selectin with μ 1A-adaptin

were purchased from GE Healthcare Life Sciences (Munich, Germany). RPMI 1600 medium was purchased from Gibco/Thermo Fisher Scientific. Fetal bovine serum and octyl- β -glucoside were from Sigma-Aldrich. TAPI-O was from Peptides International (Louisville, KY).

Peptides

Automated solid phase synthesis of peptides using Fmoc (*N*-(9-fluorenyl)methoxycarbonyl)/*t*-Bu chemistry was carried out at the Biochemistry core facility of the Max Planck Institute for Biochemistry (Martinsried, Germany). Characterization and quality control (purity) was done by RP-HPLC. All peptides were desthiobiotinylated, and a spacer (hexanoic acid (Ahx)) between the desthiobiotin group and the first amino acid of the peptides was introduced. The following peptides were used in this study: desthiobiotin-GAhxEGRMKQPDKRDS-KLKRY-OH (scrambled peptide); desthiobiotin-GAhxRRLKKGKKSQERMDDPY-OH (L-selectin tail); desthiobiotin-GAhxRRLKKGKKS_pQERMDDPY-OH (L-selectin-p-tail); desthiobiotin-GAhxRRLKKGKKGKQERMDDPY-OH (L-selectin S364D tail); desthiobiotin-GAhxRALKKGKKSQERMDDPY-OH (L-selectin R357A tail); desthiobiotin-GAhxRRLKAGKKSQERMDDPY-OH (L-selectin K360A tail); desthiobiotin-GAhxRRLKKGKASQERMDDPY-OH (L-selectin K363A tail); desthiobiotin-GAhxRRLKKGKKSQERMNDPY-OH (L-selectin D369N tail); desthiobiotin-GAhxRRLKKGKKSQERMNDPY-OH (L-selectin D370N tail); desthiobiotin-GAhxRRLKKGKKSQERMNNPY-OH (L-selectin D369N, D370N tail).

Cell culture

Raw 264.7 macrophages were grown in 75-cm³ flasks at 37 °C in an atmosphere of 5% CO₂ in RPMI 1600 medium supplemented with 10% FCS, antibiotics, and L-glutamine (20 mM).

Immunoprecipitation and Western blotting

Raw 264.7 macrophages were pretreated with the sheddase inhibitor TAPI-O (10 min, 10 μ g/ml) and then incubated in the absence or presence of PMA (1 μ g/ml) for different time periods. Thereafter, cells were collected by centrifugation (200 \times g, 2 min) and washed with PBS. The tubes were put on ice for 5 min, after which 1 ml of lysis buffer was added. The lysis buffer consisted of 50 mM Tris-HCl, pH 7.4, 30 mM octyl- β -glucoside, 1 mM EDTA, 110 mM NaCl, protease inhibitors, and phosphatase inhibitors. Cell lysates were clarified by centrifugation, and the supernatants were subjected to immunoprecipitation. This was performed by exposure to the anti-L-selectin Ab (1 μ g/ml) or the isotype-matched IgG2a Ab (1 μ g/ml) for 1 h and then to 40 μ l of a 50% slurry of protein A-Sepharose for 1 h. The beads were subsequently collected by centrifugation (200 \times g, 1 min) and washed three times with lysis buffer. The beads were then resuspended in 2 \times concentrated Laemmli sample buffer and boiled under reducing conditions (1 mM DTT) for 5 min. The immunoprecipitated proteins were subjected to electrophoresis on 10% SDS-PAGE and transferred to polyscreen PVDF transfer membranes. The membranes were blocked in TBS supplemented with 0.2% Tween 20 and 3% fat-free skimmed milk and then incubated for 1 h with a primary antibody (1:500 dilution of the rabbit anti- μ 1A Ab (19) or a 1 μ g/ml dilution of

the mouse anti-L-selectin Ab) and thereafter washed three times for 5 min in TBS supplemented with 0.2% Tween. The membranes were subsequently incubated for 1 h with peroxidase-conjugated goat anti-rabbit IgGs (1:10 000) or goat anti-mouse IgGs (1:10 000) in TBS supplemented with 0.2% Tween 20. The blots were washed three times for 5 min with TBS supplemented with 0.2% Tween 20, and antibody binding was visualized by enhanced chemiluminescence (ECL).

Peptide pulldown assays

L-selectin peptides were dissolved in PBS plus 0.01% Tween 20, and then 20 μ l of streptavidin-Sepharose beads were added for 1 h, followed by the addition of 2% BSA for 30 min. The beads were collected by centrifugation (200 \times g, 1 min) and washed once with PBS plus 0.01% Tween 20.

L-selectin tail peptides coupled to streptavidin-Sepharose beads were then incubated in 1 ml of cold buffer composed of 50 mM Tris-HCl, pH 7.4, 0.5% CHAPS, 110 mM NaCl. 50 ng of purified μ 1A expressed as GST-fusion proteins was added. The Eppendorf tubes were rotated for 1 h in the cold room, after which pellets were collected by centrifugation (200 \times g, 1 min). The pellets were washed three times with washing buffer (50 mM Tris-HCl, pH 7.4, 0.5% CHAPS, 300 mM NaCl). The beads were subsequently resuspended in 2 \times concentrated Laemmli sample buffer and boiled under reducing conditions for 5 min. The proteins were subjected to electrophoresis on 10% SDS-PAGE and transferred to polyscreen PVDF transfer membranes, and Western blotting analysis was carried out as described above using a rabbit anti- μ 1A Ab or an anti-GST Ab.

L-selectin peptides coupled to streptavidin-Sepharose beads were also incubated with lysate extracts of Raw 264.7 macrophages. Briefly, the cells were lysed in buffer composed of 50 mM Tris-HCl, pH 7.4, 0.5% CHAPS, 1 mM EDTA, 110 mM NaCl, and protease and phosphatase inhibitors. Lysates were then added for 1 h to L-selectin tail peptides coupled to beads. Thereafter, the beads were collected by low speed centrifugation (200 \times g, 1 min), washed three times with lysis buffer, and finally resuspended in 2 \times concentrated Laemmli sample buffer and boiled under reducing conditions. μ 1A bound to the L-selectin tail was detected by Western blotting analysis as described above.

Proteomics analysis

Raw 264.7 macrophages were stimulated for 1 min with PMA (1 μ g/ml), after which the cells were pelleted and washed with ice-cold PBS. The cells were subsequently lysed with MPER buffer. Clarified lysates were then incubated with Dynabeads Myone streptavidin C1 that had been coupled to either L-selectin tail peptide or scrambled peptide (see above). After a 2-h incubation in the cold room under rotation, the beads were collected with a magnet and washed gently three times with MPER buffer. The beads were then resuspended in 50 mM Tris-HCl, pH 8.0, containing 10 mM biotin to dissociate the biotin analogue desthiobiotin from streptavidin beads. After 1 h in the cold room, supernatants were collected, and proteins were precipitated overnight at -20 °C with 100% acetone. Proteins were pelleted by centrifugation (15,000 \times g, 15 min) in a cold centrifuge. Subsequently, the pellet was air-dried under a fume hood.

20 μ l of 6 M guanidium HCl containing a 10 mM concentration of the reducing agent tris(2-carboxyethyl)phosphine and 40 mM of the alkylation agent 2-chloroacetamine were added to each pellet, and the samples were boiled for 10 min. 180 μ l of LT-Digestion buffer (25 mM Tris-HCl, pH 8.5, containing 10% acetonitrile) was added, followed by 1 μ g of each trypsin and Lys-C endopeptidase. The tubes were left overnight at 37 °C under shaking. Thereafter, 8 μ l of 25% TFA (1% final) was added to stop the reaction. The peptides were bound to triple filter surface (catalog no. SDB-RPS 3X), and after washing the filters three times with 0.2% TFA, proteins were eluted with 60 μ l of buffer containing 80% acetonitrile and 5% ammonium. Peptides were analyzed on an LTQ-Orbitrap analyzer, identified, and quantified using the M_AxQ_Uant software (41).

Purification of GST-fusion proteins

Expression of μ 1A as GST-fusion proteins has been described in detail (42). Briefly, pGEX-5X3 plasmids were transformed into *E. coli* BL21. 1 mM isopropyl β -D-thiogalactopyranoside was added overnight to bacterial cultures grown at 26 °C. Bacteria were harvested, put on ice, washed once with ice-cold PBS, and lysed by the addition of a buffer made up of 50 mM Tris-HCl, pH 7.4, 1% Triton X-100, 110 mM NaCl, and protease inhibitors. This was followed by sonication. The lysates were clarified by centrifugation (10,000 \times g, 15 min), supernatants were collected, and 10% glycerol was added. These supernatants were incubated in the cold room with glutathione-Sepharose beads pre-equilibrated in PBS buffer. After 1 h, the beads were collected by centrifugation (200 \times g, 1 min) and washed three times with ice-cold PBS. The GST-fusion proteins were eluted by adding 200 μ l of buffer containing 50 mM Tris-HCl, pH 8.0, and 10 mM reduced glutathione. Eluted solutions were passed through a G-25 Sephadex column, which was subjected to centrifugation (500 \times g, 1 min), and GST-fusion proteins free of glutathione were collected. Protein concentrations were estimated by spectrometry.

Immunofluorescence confocal microscopy

THP-1 cells stably expressing wild type L-selectin tagged to green fluorescent protein (WT L-selectin-GFP) were generated and cultured as described previously (6). Cultured THP-1 cells were harvested and resuspended in warmed undiluted RPMI at a density of 0.5×10^6 cells/ml. Cells were subsequently seeded onto poly-L-lysine-coated coverslips and allowed to settle for 10 min at 37 °C. RPMI was aspirated, and coverslip-bound cells were fixed in 3% paraformaldehyde for 20 min at room temperature. Cells were washed three times in PBS and then permeabilized in 0.6% saponin for 10 min at room temperature. After washing fixed cells with PBS, cells were blocked in 5% fetal calf serum containing Fc receptor block (Miltenyi Biotec Ltd.) for 1 h at room temperature before staining with 1:300 (v/v) of 0.1 mg/ml TRITC-phalloidin and anti- γ 1 AP-1 mouse monoclonal antibody (250 μ g/ml; BD Biosciences) diluted at 1:75 (v/v) overnight at 4 °C. Cells were washed free from excess phalloidin and antibody, blocked as before, and then stained with secondary goat anti-mouse (Life Technologies) Alexa Fluor 633 (1:400 dilution). Specimens were mounted onto glass slides using mounting medium (Dako). Images were acquired using Leica

SP5 confocal microscope, using a \times 63 objective lens. Scale bar, 7.5 μ m.

Molecular modeling

Molecular docking of the L-selectin tail to μ 1A was conducted using the HEX version 8.0 protein-protein docking program (43). The crystal structure of μ 1A (Protein Data Bank code 4P6Z) and the structural model of 17-residue long L-selectin tail was modeled using the crystal structures of the 8-residue-long L-selectin tail (Protein Data Bank code 2LGF). The remaining 8 amino acid residues of the L-selectin tail were modeled using the loop prediction option of the Prime module within Schrödinger software (44). The L-selectin tail in a wild type as well as mutated forms (S364pS, S364D, D369N, and D370N) were used for docking to facilitate a choice of docking solutions. The μ 1A/L-selectin binding surface interactions were mapped based on shape and electrostatics complementarities as implemented in HEX version 8.0. Default settings for other docking parameters were used. HEX version 8.0 has generated over 100 solutions for each docking run. Besides the HEX energy function, the obtained docking pose of the L-selectin tail in the wild type and mutated forms was evaluated based on its ability to explain the mutagenesis data. In addition, only docking poses on the surface of μ 1A, which is not in an interaction with other subunits within, AP-1 were considered.

Author contributions—K. D. designed the project, performed the research, analyzed the data, and wrote the manuscript; I. G. T. performed the modeling of μ 1A-L-selectin tail interaction, analyzed the data, and wrote part of the manuscript; A. I. performed the colocalization studies; P. S. provided scientific expertise, analyzed the data, and wrote part of the manuscript. All authors read and approved the final manuscript.

Acknowledgments—We thank Yosuke Yoneyama and Shin-Ichiro Takahashi (University of Tokyo) for providing the pGEX-5X3 plasmids encoding for full-length μ 1A or the N-terminal or C-terminal region of μ 1A. Ralph Böttcher, Markus Moser, and Reinhard Fässler (Max Planck Institute for Biochemistry, Martinsried, Germany) are acknowledged for fruitful discussions. Herbert Schiller (Max Planck Institute for Biochemistry, Martinsried, Germany) is acknowledged for help in the identification and quantification of L-selectin tail-interacting proteins using the M_AxQ_Uant software. We thank Stephan Uebel (core facility of the Max Planck Institute for Biochemistry, Martinsried, Germany) for the synthesis of L-selectin tail peptides.

References

- Ley, K., Laudanna, C., Cybulsky, M. I., and Nourshargh, S. (2007) Getting to the site of inflammation: the leukocyte adhesion cascade updated. *Nat. Rev. Immunol.* **7**, 678–689
- von Andrian, U. H., Hasslen, S. R., Nelson, R. D., Erlandsen, S. L., and Butcher, E. C. (1995) A central role for microvillous receptor presentation in leukocyte adhesion under flow. *Cell* **82**, 989–999
- Arbonés, M. L., Ord, D. C., Ley, K., Ratech, H., Maynard-Curry, C., Otten, G., Capon, D. J., and Tedder, T. F. (1994) Lymphocyte homing and leukocyte rolling and migration are impaired in L-selectin deficient mice. *Immunity* **1**, 247–260
- Smalley, D. M., and Ley, K. (2005) L-selectin: mechanism and physiological significance of ectodomain cleavage. *Cell Mol. Med.* **9**, 255–266

Interaction of L-selectin with μ 1A-adaptin

- Marschner, S., Freiberg, B. A., Kupfer, A., Hünig, T., and Finkel, T. H. (1999) Ligation of the CD4 receptor induces activation-independent down-regulation of L-selectin. *Proc. Natl. Acad. Sci. U.S.A.* **96**, 9763–9768
- Rzeniewicz, K., Newe, A., Rey Gallardo, A., Davies, J., Holt, M. R., Patel, A., Charras, G. T., Stramer, B., Molenaar, C., Tedder, T. F., Parsons, M., and Ivetic, A. (2015) L-selectin shedding is activated specifically within trans-migrating pseudopods of monocytes to regulate cell polarity *in vitro*. *Proc. Natl. Acad. Sci. U.S.A.* **112**, E1461–E1470
- Yang, S., Liu, F., Wang, Q. J., Rosenberg, S. A., and Morgan, R. A. (2011) The shedding of CD62L (L-selectin) regulates the acquisition of lytic activity in human tumor reactive T lymphocytes. *PLoS One* **6**, e22560
- Tedder, T. F., Isaacs, C. M., Ernst, T. J., Demetri, G. D., Adler, D. A., and Distche, C. M. (1989) Isolation and chromosomal localization of cDNAs encoding a novel human lymphocyte cell surface molecule, LAM-1: homology with the mouse lymphocyte homing receptor and other human adhesion proteins. *J. Exp. Med.* **170**, 123–133
- Kansas, G. S., Ley, K., Munro, J. M., and Tedder, T. F. (1993) Regulation of leukocyte rolling and adhesion to endothelial venules through the cytoplasmic domain of L-selectin. *J. Exp. Med.* **177**, 833–838
- Kahn, J., Walcheck, B., Migaki, G. I., Jutila, M. A., and Kishimoto, T. K. (1998) Calmodulin regulates L-selectin adhesion molecule expression and function through a protease-dependent mechanism. *Cell* **92**, 809–818
- Ivetic, A., Florey, O., Deka, J., Haskard, D. O., Ager, A., and Ridley, A. J. (2004) The cytoplasmic tail of L-selectin interacts with members of the ezrin-radixin-moesin (ERM) family proteins. *J. Biol. Chem.* **279**, 33263–33272
- Pavalko, F. M., Walker, D. M., Graham, L., Goheen, M., Doerschuk, C. M., and Kansas, G. S. (1995) The cytoplasmic domain of L-selectin interacts with cytoskeletal proteins via α -actinin: receptor positioning in microvilli does not require interaction with α -actinin. *J. Cell Biol.* **129**, 1155–1164
- Xu, T., Chen, L., Shang, X., Cui, L., Luo, J., Chen, C., Ba, X., and Zeng, X. (2008) Critical role of Lck in L-selectin signaling induced by sulfatides engagement. *J. Leukoc. Biol.* **84**, 1192–1201
- Kilian, K., Dervede, J., Mueller, E. C., Bahr, I., and Tauber, R. (2004) The interaction of protein kinase C isozymes α , ι , and θ with the cytoplasmic domain of L-selectin is modulated by phosphorylation of the receptor. *J. Biol. Chem.* **279**, 34472–34480
- Ivetic, A., and Ridley, A. J. (2004) The telling tail of L-selectin. *Biochem. Soc. Trans.* **32**, 1118–1121
- Haribabu, B., Steeber, D. A., Ali, H., Richardson, R. M., Snyderman, R., and Tedder, T. F. (1997) Chemoattractant receptor-induced phosphorylation of L-selectin. *J. Biol. Chem.* **272**, 13961–13965
- Robinson, M. (2004) Adaptable adaptors for coated vesicles. *Trends Cell Biol.* **14**, 167–174
- Bonifacino, J. S. (2014) Adaptor proteins involved in polarized sorting. *J. Cell Biol.* **204**, 7–17
- Zizioli, D., Meyer, C., Guhde, G., Saftig, P., von Figura, K., and Schu, P. (1999) Early embryonic death of mice deficient in γ -adaptin. *J. Biol. Chem.* **274**, 5385–5390
- Meyer, C., Zizioli, D., Lausmann, S., Eskelinen, E. L., Hamann, J., Saftig, P., von Figura, K., and Schu, P. (2000) μ 1A-adaptin-deficient mice: lethality, loss of AP-1 binding and rerouting of mannose 6-phosphate receptors. *EMBO J.* **19**, 2193–2203
- Meyer, C., Eskelinen, E. L., Guruprasad, M. R., von Figura, K., and Schu, P. (2001) μ 1A deficiency induces a profound increase in MPR300/IGF-II receptor internalization rate. *J. Cell Sci.* **114**, 4469–4476
- Owen, D. J., and Evans, P. R. (1998) A structural explanation for the recognition of tyrosine-based endocytotic signals. *Science* **282**, 1327–1332
- Heldwein, E. E., Macia, E., Wang, J., Yin, H. L., Kirchhausen, T., and Harrison, S. C. (2004) Crystal structure of the clathrin adaptor protein 1 core. *Proc. Natl. Acad. Sci. U.S.A.* **101**, 14108–14113
- Griffin, J. D., Spertini, O., Ernst, T. J., Belvin, M. P., Levine, H. B., Kanakura, Y., and Tedder, T. F. (1990) Granulocyte-macrophage colony-stimulating factor and other cytokines regulate surface expression of the leukocyte adhesion molecule-1 on human neutrophils, monocytes, and their precursors. *J. Immunol.* **145**, 576–584
- Traub, L. M. (2009) Tickets to ride: selecting cargo for clathrin-regulated internalization. *Nat. Rev. Mol. Cell Biol.* **10**, 583–596
- Baltes, J., Larsen, J. V., Radhakrishnan, K., Geumann, C., Kratzke, M., Petersen, C. M., and Schu, P. (2014) σ 1B adaptin regulates adipogenesis by mediating the sorting of sortilin in adipose tissue. *J. Cell Sci.* **127**, 3477–3487
- Foote, C., and Nothwehr, S. F. (2006) The clathrin adaptor complex 1 directly binds to a sorting signal in Ste13p to reduce the rate of its trafficking to the late endosome of yeast. *J. Cell Biol.* **173**, 615–626
- Orzech, E., Schlessinger, K., Weiss, A., Okamoto, C. T., and Aroeti, B. (1999) Interactions of the AP-1 Golgi adaptor with the polymer immunoglobulin receptor and their possible role in mediating brefeldin A-sensitive basolateral targeting from the *trans*-Golgi network. *J. Biol. Chem.* **274**, 2201–2215
- Deng, W., Srinivasan, S., Zheng, X., Putkey, J. A., and Li, R. (2011) Interaction of calmodulin with L-selectin at the plasma membrane interface: implication on the regulation of L-selectin shedding. *J. Mol. Biol.* **411**, 220–233
- Deng, W., Cho, S., and Li, R. (2013) FERM domain of moesin desorbs the basic-rich cytoplasmic domain of L-selectin from the anionic membrane surface. *J. Mol. Biol.* **425**, 3549–3562
- Almeida, P. F. F., Pokorny, A., and Hinderliter, A. (2005) Thermodynamics of membrane domains. *Biochim. Biophys. Acta* **1720**, 1–13
- Melcrova, A., Pokorna, S., Pullanchery, S., Kohagen, M., Jurkiewicz, P., Hof, M., Jungwirth, P., and Cremer, P. S., and Cwiklik, L. (2016) The complex nature of calcium cation interactions with phospholipid bilayers. *Nat. Sci. Rep.* 10.1038/srep38035
- Ren, X., Farias, G. G., Canagarajah, B. J., Bonifacino, J. S., and Hurley, J. H. (2013) Structural basis for recruitment and activation of the AP-1 clathrin adaptor complex by Arf1. *Cell* **152**, 755–767
- Conner, S. D., and Schmid, S. L. (2002) Identification of an adaptor-associated kinase, AAK1, as a regulator of clathrin-mediated endocytosis. *J. Cell Biol.* **156**, 921–929
- Ricotta, D., Conner, S. D., Schmid, S. L., von Figura, K., and Honing, S. (2002) Phosphorylation of the AP2 μ subunit by AAK1 mediates high affinity binding to membrane protein sorting signals. *J. Cell Biol.* **156**, 791–795
- Ghosh, P., and Kornfeld, S. (2003) AP-1 binding to sorting signals and release from clathrin-coated vesicles is regulated by phosphorylation. *J. Cell Biol.* **160**, 699–708
- Ng, T., Parsons, M., Hughes, W. E., Monypenny, J., Zicha, D., Gautreau, A., Arpin, M., Gschmeissner, S., Verveer, P. J., Bastiaens, P. I., and Parker, P. J. (2001) Ezrin is a downstream effector of trafficking PKC-integrin complexes involved in the control of cell motility. *EMBO J.* **20**, 2723–2741
- Medigeschi, G. R., Krikunova, M., Radhakrishnan, K., Wenzel, D., Klingauf, J., and Schu, P. (2008) AP-1 membrane-cytoplasm recycling regulated by μ 1A-adaptin. *Traffic* **9**, 121–132
- Radhakrishnan, K., Baltes, J., Creemers, J. W. M., and Schu, P. (2013) *Trans*-Golgi network morphology and sorting is regulated by prolyl-oligopeptidase-like protein PREPL and the AP-1 complex subunit μ 1A. *J. Cell Sci.* **126**, 1155–1163
- Régál, L., Shen, X. M., Selcen, D., Verhille, C., Meulemans, S., Creemers, J. W., and Engel, A. G. (2014) PREPL deficiency with or without cystinuria causes a novel myasthenic syndrome. *Neurology* **82**, 1254–1260
- Cox, J., and Mann, M. (2008) MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat. Biotechnol.* **26**, 1367–1372
- Yoneyama, Y., Matsuo, M., Take, K., Kabuta, T., Chida, K., Hakuno, F., and Takahashi, S. (2013) The AP-1 complex regulates intracellular localization of insulin receptor substrate 1, which is required for insulin-like growth factor I-dependent cell proliferation. *Mol. Cell Biol.* **33**, 1991–2003
- Ghoorah, A. W., Devignes, M. D., Smail-Tabbone, M., and Ritchie, D. W. (2013) Protein docking using case-based reasoning. *Proteins* **81**, 2150–2158
- Prime*, version 3.0 (2011) Schrödinger, LLC, New York