

Assessment of ‘one-step’ versus ‘sequential’ embryo culture conditions through embryonic genome methylation and hydroxymethylation changes

J. Salvaing^{1,2,*}, N. Peynot¹, M. N. Bedhane^{1,3}, S. Veniel¹, E. Pellier^{1,4}, C. Boulesteix¹, N. Beaujean^{1,5}, N. Daniel¹, and V. Duranthon^{1,*}

¹UMR BDR, INRA, ENVA, Université Paris Saclay, 78350 Jouy en Josas, France ²Present address: Laboratoire de Physiologie Cellulaire et Végétale (LPCV), UMR 1417 INRA / UMR 5168 CNRS-CEA-UGA, Bioscience and Biotechnology Institute of Grenoble (BIG), Commissariat à l’Energie Atomique at aux Energies Renouvelables (CEA), 38054 Grenoble Cedex 9, France ³Present address: Jigjiga University, Ethiopia ⁴Present address: Faculté de Médecine, 27 Bd Jean Moulin, 13385 Marseille Cedex C5, France ⁵Present address: INSERM U1208, INRA USC136 I Stem Cell and Brain Research Institute Department of Pluripotent Stem Cells in Mammals, 18 Avenue Doyen Lépine, 69675 Bron, France

*Correspondence address. UMR BDR, INRA, ENVA, Université Paris Saclay, 78350 Jouy en Josas, France.
E-mail: veronique.duranthon@inra.fr (V.D.)/juliette.salvaing@cea.fr (J.S.)

HumReprod 2016;31:2471–2483

The authors would like to apologize for an error in a section of the Materials and methods of the above article. The section titled Recovery of rabbit embryos stated that *in vivo* embryos were collected from oviducts at the 1-cell, 2-cell, 4-cell, 8-cell, 16-cell, morula and blastocyst stages. The time point stated for collection of

in vivo 8-cell embryos was mistakenly stated as 49 hpc (hours post-coitum). The correct time point for collection of *in vivo* 8-cell embryos is 39 hpc.

The authors would like to assure readers that this error does not affect any other content of the article.