

Effect of Light Intensity on Efficiency of Carbon Dioxide and Nitrogen Reduction in *Pisum sativum* L.¹

Received for publication June 6, 1977 and in revised form August 22, 1977

GABOR J. BETHLENFALVAY AND DONALD A. PHILLIPS

Department of Agronomy and Range Science, University of California, Davis, California 95616

ABSTRACT

Photosynthetic efficiency, primary productivity, and N₂ reduction were determined in peas (*Pisum sativum* L. var. Alaska) grown at light intensities ranging from severely limiting to saturating. Plants grown under higher light intensities showed greater carboxylation and light capture potential and higher rates of net C exchange. Uptake of N₂, computed from measured C₂H₂ reduction and H₂ evolution rates, also increased with growth light intensity, while the previously proposed relative efficiency of N₂ fixation, based on these same parameters, declined. The plot of N/C ratios (total nitrogen content/plant dry weight) increased hyperbolically with light intensity, and the plot of N₂/CO₂ uptake ratios (N₂ uptake rate/net CO₂ uptake rate) increased linearly. Both plots extrapolated to the light compensation point. The data indicate that the relative efficiency of N₂ fixation is not necessarily correlated with maximum plant productivity and that evaluation of a plant's capacity to reduce N₂ is related directly to concurrent CO₂ reduction. A measure of whole plant N₂ fixation efficiency based on the N₂/CO₂ uptake ratio is proposed.

Symbiotic N₂ reduction in legumes has long been known to be a function of photosynthesis (21). Both long term (13) and short term (18) CO₂ enrichment studies have suggested that the availability of this molecule limits N₂ fixation in legumes, presumably through an effect on photosynthesis. The importance of suitable light intensity for optimum plant productivity is an elementary fact, which has led to sophisticated considerations of the role of leaf displays in the productivity of plant communities (15). Growth light intensity has been shown to affect symbiotic N₂ reduction (9, 12) and the inhibition of N₂ fixation by combined N (4). Thus, it seems likely that growth light intensity may cause changes in competitive interactions involving legumes in mixed or pure stands by altering the efficiency with which both CO₂ and N₂ are reduced.

This study was undertaken to correlate the rate of NCE² and the amount of C fixed with the rate and amount of N₂ fixation as a function of light intensity, to provide comparisons between different measures for the efficiency of photosynthesis, such as CE (20), NCE (22), and PE (11) with RE which measures N₂ fixation efficiency of nitrogenase (19), and to derive an expression for whole plant N₂ fixation efficiency which incorporates photosynthetic parameters. The effect of growth light intensity

on N₂ fixation may be an important consideration in mixed crop and pasture ecology, where selection of plants, whose efficiency to fix N₂ is not impaired under conditions of partial shade, may be desirable to maximize fixed N input into the soil.

MATERIALS AND METHODS

Growth Conditions. Pea plants were maintained in a growth chamber as described previously (2), but at growth light intensities of 200, 400, 600, or 800 μ Ei. Initial seed weight was 0.22 to 0.26 g. Plants were assayed for N₂ fixation after a 5 hr light period; photosynthetic assays at each light intensity spanned the entire light period.

Carbon Dioxide Fixation. Assimilation of CO₂ by attached leaves was measured in a flow-through gas exchange system with apparatus and data-handling procedures as described by Augustine *et al.* (1). Two plants were selected from uniform stands of six replicates for photosynthetic measurements. Design of the CO₂ assimilation chamber permitted parts of the plant to be inserted in sequential steps without damaging the plant. NCE by the whole plant was determined at the growth light intensity with a CO₂ concentration of 300 μ l/l by sequential insertion of different segments of the shoot. PE was measured on the sixth leaf at the saturating light intensity of 1,200 μ Ei, and the appropriate growth light intensity. PE was calculated as 100 \times the ratio of NCE at growth light intensity/NCE at 1,200 μ Ei. NCE was determined in the fifth leaf at four CO₂ concentrations (50, 100, 150, and 300 μ l/l) and a light intensity of 1,200 μ Ei regardless of growth light intensity. Internal leaf CO₂ concentrations were calculated from physical parameters measured at each of the four external CO₂ concentrations (1). A second order regression line was then computer-plotted through the data points determined by the CER and internal CO₂ concentration values (1). The light compensation point was established as the light intensity intercept of an extrapolated regression line computer-plotted through data points defined by the growth light intensities and the corresponding NCE values. The internal CO₂ compensation point (x intercept) and the slope of the regression line at the compensation point were determined by extrapolation. This slope, an indication of the leaf's capacity to respond to changes in ambient CO₂ concentrations, was used as a measure of carboxylation efficiency as proposed by Tregunna *et al.* (20). Leaf temperatures were maintained at 21 C in all cases. Leaf area was measured with a Lambda Instruments LI-3000 area meter.

Nitrogen Fixation. Production of H₂ and C₂H₂-dependent C₂H₄ by root nodules was determined by techniques and equipment described previously (2). Total N/plant was determined by Kjeldahl analysis (8).

RESULTS

Three measures of photosynthetic efficiency used in this study to evaluate plant response to different light intensities increased with increasing illumination. CE of the fifth leaf, an indicator of

¹ This material is based upon research supported by the National Science Foundation under Grants AER77-07301 and PCM 76-23472. Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the authors and do not necessarily reflect the view of the National Science Foundation.

² Abbreviations: NCE: net carbon exchange; CE: carboxylation efficiency; μ Ei: microeinstein \cdot m⁻² \cdot sec⁻¹; PE: photoefficiency; RE: relative efficiency of nitrogen fixation.

the plant's ability to respond to different CO₂ concentrations, increased linearly with growth light intensity (200, 400, 600 and 800 μEi) when plants were assayed under the same saturating (1,200 μEi) light intensities (Fig. 1). Concurrent measurements of NCE/leaf area in the same leaf (fifth) did not show saturation at the highest growth light intensity used, while the CO₂ compensation point was lowest at 600 μEi (Fig. 2). Photoefficiency was determined to measure the capacity of plants grown under different light intensities to respond to saturating light intensity. Plants grown under low light intensity showed a greater percentage increase in NCE when exposed to saturating light than did plants grown under high light intensity (Fig. 1). Specific activity of NCE (NCE/total plant leaf area) (Fig. 3) and the net CO₂ uptake rate of the whole plant (Fig. 4) reached maximum values near the growth light intensity of 600 μEi. The light compensation point was 5.1 μEi, a value within the range previously established (14) for photorespiring plants.




Fig. 1. Carboxylation efficiency (CE) and photoefficiency (PE) of peas grown at different light intensities. CE was measured on the fifth leaf and PE on the sixth leaf of 24-day-old pea plants. Each point on the CE curve represents the slope at the CO₂ compensation point of a regression line through data points obtained for net C exchange at four CO₂ concentrations. Measurements were made at 1,200 μEi regardless of growth light intensity. The PE curve (extrapolated to the light compensation point) indicates the response of plants grown at lower light intensities to a saturating light intensity (1,200 μEi). PE data were calculated as 100 × NCE at growth light intensity/NCE at 1,200 μEi.




Fig. 2. Effect of growth light intensity on net C exchange rate and internal CO₂ compensation point in peas. Measurements were made on the fifth leaf at the saturating light intensity of 1,200 μEi.




Fig. 3. Correlation of total net C exchange/leaf area with N₂ fixation/nodule wt by pea plants grown at different light intensities. N₂ fixation was computed by the formula (C₂H₂ reduced/3) × RE.




Fig. 4. Net CO₂ uptake by shoots and C₂H₂ reduction or H₂ evolution by root nodules of pea plants grown at different light intensities. Photosynthesis data were determined on two plants by sequential measurement of shoot segments, and root nodule data are means from four plants grown for 24 days at the light intensities indicated.

All plants were at the same stage of development on the basis of leaf numbers.

Both C₂H₂-dependent C₂H₄ production and H₂ evolution increased with increasing light intensity, although at different rates (Fig. 4). As a result, the RE expressed by Schubert and Evans (19) by the formula RE = 1 - H₂ evolved/C₂H₂ reduced, decreased with increasing light intensity (Fig. 5). Analysis of variance showed that differences in RE according to light intensity were highly significant (P ≤ 0.005). Reduction of N₂, however, computed by the formula (C₂H₂ reduced/3) × RE increased linearly with light intensity (Fig. 5). The specific activity of N₂ reduction peaked at 600 μEi, showing a trend similar to the specific activity of NCE (Fig. 3). Ratios of total N to total plant dry wt were comparable in magnitude to values

found in the literature (17) and approached saturation at 800 μEi (Fig. 6). This curve, when extrapolated, intersected the light intensity axis at the light compensation point. The increase in ratio of the rates of N_2 fixation to net CO_2 uptake was linear with light intensity, and the extrapolated x intercept also coincided with the light compensation point (Fig. 6). Nodule and plant dry wt and total N/plant values increased with light intensity (Table I).

DISCUSSION

The data demonstrate that variations in photosynthetic efficiency and plant productivity (Fig. 1 and Table I), which are produced by growth light intensity, are directly correlated with the rate of N_2 fixation (Fig. 5) and N accumulation (Table I). The rate of N_2 fixation and the proposed relative efficiency of N fixation (19) showed opposite trends with variation of growth light intensity (Fig. 5). For these reasons an alternate measure




FIG. 5. Symbiotic N_2 fixation by pea plants grown at different light intensities. Reduction of N_2 was computed from C_2H_2 reduction and H_2 evolution data by the formula: $(\text{C}_2\text{H}_2 \text{ reduced}/3) \times \text{RE}$. Relative efficiency of N_2 fixation (RE), an expression proposed (19) for electron allocation by nitrogenase to reduce N_2 or H^+ , is defined as $\text{RE} = 1 - \text{H}_2 \text{ evolved}/\text{C}_2\text{H}_2 \text{ reduced}$. At the lowest light intensity, H_2 evolution, if present, was not detected by the analytical technique employed. The RE corresponding to the lowest detectable amount of H_2 (at the C_2H_4 level observed) would have been 0.95.




FIG. 6. Whole plant N_2 fixation efficiency in 24-day-old peas grown at different light intensities. Rates of N_2 uptake were calculated from C_2H_2 reduction and H_2 evolution as in Figure 5, and the net CO_2 uptake rate was measured directly (Fig. 4). The weight ratio data, determined as N content/dry wt after subtracting for N and C present in the seed, represent an integration of all N and C metabolism which occurred from the time of planting.

Table I. Leaf area, dry weight and nitrogen content of *Pisum sativum* grown at different light intensities.

Means \pm standard error were calculated from 6 replicates, except for leaf area and nodule weight, which are respectively averages of 2 and 4 replicates. Average dry weight and N content of seeds were respectively 200 ± 5 mg and 7.8 ± 0.4 mg.

Light intensity (μEi)	Leaf area (cm^2)	Dry weight (mg)		Nitrogen content	
		Plant	Nodule	Plant (mg)	%
200	52.2	319 ± 16	10.6 ± 1.1	11.5 ± 0.6	3.6
400	65.3	419 ± 10	22.9 ± 1.6	16.8 ± 0.5	4.0
600	54.5	472 ± 47	32.0 ± 3.8	19.7 ± 1.9	4.2
800	49.7	568 ± 65	47.0 ± 6.9	24.5 ± 3.1	4.3

of whole plant N_2 fixation efficiency was derived on the basis of photosynthetic parameters (Fig. 6).

It has been shown that the light capture and carboxylation mechanisms in leaves develop differently under different growth light intensities (5, 6). In shade plants the light absorption function is dominant, whereas in sun-adapted plants carboxylation takes precedence. This concept is supported by data showing increased CE (Fig. 1) and decreased internal CO_2 compensation point (Fig. 2) with increasing growth light intensity. In addition, low PE (Fig. 1) in plants grown in dim light shows that these plants are capable of a greater increase in photosynthesis when exposed to saturating light intensity than plants grown in bright light. To correlate CO_2 reduction efficiency with N_2 fixation in plants grown under light intensities ranging from severely limiting (200 μEi) to saturating (800 μEi) levels, we therefore measured plant responses to changes in both irradiance levels and CO_2 concentration. The two measures of photosynthetic efficiency used, PE and CE, increased with increasing growth light intensity. NCE, a direct expression of the plant's capacity to assimilate CO_2 , also increased with light intensity whereas the CO_2 compensation point reached a minimum value near 600 μEi (Fig. 2). The above data were obtained on individual leaves at the same stage of development because CE, NCE, and CO_2 compensation point change during ontogeny (2, 3).

Greater photosynthetic productivity/plant at higher light intensities (Table I) was correlated with greater nitrogenase activity/plant as shown by C_2H_2 reduction and H_2 reduction (Fig. 4) and N_2 reduction as computed from these data (Fig. 5). The specific activities of NCE and N_2 fixation (NCE/leaf area and N_2 reduced/nodule mass), however, did not exhibit a linear increase with light intensities (Fig. 3). The maximization of these parameters at 600 to 700 μEi may have resulted from artificially imposed limiting conditions, such as pot size and a uniform watering schedule, but the interdependence of the two processes is underscored by the coincidence of the peaks and the observation that both curves may be extrapolated to the light compensation point.

The decline in RE with increasing growth light intensity (Fig. 5) is not in contradiction to the opposite trend shown by photosynthetic efficiency, for the decline is not an indication of diminished nitrogenase activity. It can be interpreted as a shift in electron allocation (7) from N_2 to H^+ reduction at higher growth light intensities or as increased H_2 uptake by a hydrogenase (10) which may be preferentially expressed under the energy-deficient conditions at the lower growth light intensities. In the latter case hydrogenase activity would lower the amount of H_2 available for measurement, thus giving the appearance of lower H_2 evolution rates. The contrary trends of RE and of actual nitrogen fixation as computed from C_2H_2 reduction and H_2 evolution data (Fig. 5) as well as total N values (Table I) show that this proposed RE is descriptive only of electron allocation to N_2 or H^+ and not of the actual capacity of the symbiosis to fix N_2 . Thus, the N_2/H^+ electron allocation ratio

may be higher (*i.e.* RE greater) under conditions less favorable for photosynthesis than under favorable conditions, yet the latter could permit increased nitrogenase activity and greater N₂ uptake even at lower (less efficient) N₂/H⁺ ratios.

The direct dependence of symbiotic N₂ fixation on photosynthesis suggests the importance of incorporating photosynthetic parameters into any expression of N₂ fixation efficiency in the *Rhizobium*-legume association. Possible expressions of efficiency include the ratio of change in total N during the growth period to change in biomass as a function of growth light intensity (Fig. 6). This plot shows that the N/C ratio approached a maximum value as growth light intensity increased, a possible indication of sink limitation (16) at higher light intensities. A more useful technique for expressing the relation between N₂ fixation and photosynthesis is to plot N₂/CO₂ uptake rates against growth light intensity (Fig. 6). When extrapolated, both lines plotted in Figure 6 intersect the *x* axis at the light compensation point, confirming the previously noted (12) dependence of N₂ fixation on the availability of photosynthetic products. It must be considered, however, that the N₂/CO₂ uptake ratio technique is presumably sensitive to short term fluctuations in environmental parameters and to the time of assay during the light/dark cycle. It is reasonable to assume that a plot of the N₂/CO₂ uptake ratio against any environmental parameter will vary with the direct influence of that parameter on the mechanisms and saturation characteristics of both N₂ and CO₂ uptake. The shape of the plot may also be affected indirectly by an interaction of the two processes. The straight line relationship between the N₂/CO₂ uptake ratio and light intensity (Fig. 6) is therefore considered to be a fortuitous result of the environmental parameters employed in this study.

The advantage of the N₂/CO₂ uptake ratio technique over the simpler dry wt ratio method is that N₂/CO₂ uptake ratio measurements provide for a calculation of short term N₂ and CO₂ reduction ratios which is limited only by the time required for new photosynthate to reach the root nodules. Thus, the effect of environmental parameters on the N₂/CO₂ uptake ratios can be determined allowing the system to achieve an equilibrium in a matter of hours. Such experiments are not feasible using N and C mass increments because the amounts of those elements present at the start of the experimental treatment are difficult to determine precisely. For this reason the N₂/CO₂ uptake ratio under different light intensities is proposed as a measure of whole plant N₂ fixation efficiency. On the whole plant level our proposed measure of N₂ fixation efficiency complements other existing measures such as the "electron-allocation coefficient" of Burns and Hardy (7) or the "relative efficiency of nitrogen fixation" of Schubert and Evans (19), which describe constituent phenomena on the enzymic and bacteroid levels. This proposed measure incorporates both interdependent parameters of the

symbiotic association, which are of interest here: microbial variation in apparent N₂ fixation, and host plant variation in photosynthetic response to varying growth light intensity. These parameters seem to be particularly important as the search continues for *Rhizobium* and legume symbionts which reduce N₂ and CO₂ efficiently under field conditions where competition for light is often significant (15).

LITERATURE CITED

1. AUGUSTINE JJ, MA STEVENS, RW BREIDENBACH, DF PAIGE 1976 Genotypic variation in carboxylation of tomatoes. *Plant Physiol* 57: 325-333
2. BETHLENFALVAY GJ, DA PHILLIPS 1977 Ontogenetic interactions between photosynthesis and symbiotic nitrogen fixation in legumes. *Plant Physiol* 60: 419-421
3. BETHLENFALVAY GJ, DA PHILLIPS 1977 Photosynthesis and symbiotic nitrogen fixation in *Phaseolus vulgaris* L. In *Genetic Engineering for Nitrogen Fixation*. A. Hollaender, ed. Plenum Press, New York, pp 401-408
4. BETHLENFALVAY GJ, DA PHILLIPS 1978 Interactions between N₂ fixation, combined-nitrogen application and photosynthesis in *Pisum sativum* L. *Physiol Plant*. In press
5. BJÖRCKMAN O 1968 Carboxydismutase activity in shade adapted and sun adapted species of higher plants. *Physiol Plant* 21: 1-10
6. BJÖRCKMAN O 1970 Characteristics of the photosynthetic apparatus as revealed by laboratory measurements. In I Setlik, ed. *Prediction and measurement of photosynthetic productivity*. Proc IBP/PPP Technical Meeting, Trebon, pp 267-281
7. BURNS RC, RW HARDY 1975 *Nitrogen Fixation in Bacteria and Higher Plants*. Springer-Verlag, New York, pp 116-122
8. BURRIS RH, PW WILSON 1957 Methods for measurement of nitrogen fixation. *Methods Enzymol* 4: 355-366
9. DIENER T 1950 Über die Bedingungen der Wurzelknöllchenbildung bei *Pisum sativum* L. *Phytopathol Z* 16: 129-170
10. DIXON ROD 1967 Hydrogen uptake and exchange by pea root nodules. *Ann Bot* 31: 179-188
11. GAASTRA P 1962 Photosynthesis of leaves and field crops. *Neth J Agric Sci* 10: (No 5, Special Issue) 311-324
12. GIBSON AH 1975 Recovery and compensation by nodulated legumes to environmental stress. In PS Nutman, ed. *Symbiotic Nitrogen Fixation in Plants*. Internat Biol Programme, Vol 7. Cambridge Univ Press, pp 385-403
13. HARDY RWF, UD HAVELKA 1975 Photosynthate as a major factor limiting nitrogen fixation by field-grown legumes with emphasis on soybeans. In PS Nutman, ed. *Symbiotic Nitrogen Fixation in Plants*. Internat Biol Programme, Vol 7. Cambridge Univ Press, pp 421-439
14. HEATH OVS 1969 *The Physiological Aspects of Photosynthesis*. Stanford Univ Press, Stanford, pp 181-197
15. LOOMIS RS, WA WILLIAMS 1969 Productivity and the morphology of crop stands: patterns with leaves. In RC Dinauer, ed. *Physiological Aspects of Crop Yield*. ASA and CSSA, Madison Wis, pp 27-47
16. MALTHORPE FL, J MOORBY 1969 Vascular transport and its significance in plant growth. *Annu Rev Plant Physiol* 20: 117-138
17. MINCHIN FR, JS PATE 1973 The carbon balance of a legume and the functional economy of its nodules. *J Exp Bot* 24: 259-271
18. PHILLIPS DA, KD NEWELL, SA HASSELL, CE FELLING 1976 The effect of CO₂ enrichment on root nodule development and symbiotic N₂ reduction in *Pisum sativum* L. *Am J Bot* 63: 356-362
19. SCHUBERT KR, HJ EVANS 1976 Hydrogen evolution: a major factor affecting the efficiency of nitrogen fixation in nodulated symbionts. *Proc Nat Acad Sci USA* 73: 1207-1211
20. TREGUNNA EB, G KOTKOV, CD NELSON 1966 Effect of oxygen on the rate of photorespiration in detached tobacco leaves. *Physiol Plant* 19: 723-733
21. WILSON PW 1935 Carbohydrate-Nitrogen Relation in Symbiotic Nitrogen Fixation. Res Bull 129. Agric Exp Sta. Univ Wisconsin, Madison, pp 1-40
22. ZELITCH I 1975 Improving the efficiency of photosynthesis. *Science* 188: 626-633