



# *In vitro* screening for protein tyrosine phosphatase 1B and dipeptidyl peptidase IV inhibitors from selected Nigerian medicinal plants

Yusuf Saidu, Suleiman Alhaji Muhammad, Abdullahi Yahaya Abbas, Andrew Onu, Ibrahim Mohammed Tsado, Luba Muhammad

## ABSTRACT

**Background/Aim:** Protein tyrosine phosphatase 1B (PTP 1B) and dipeptidyl peptidase IV (DPP IV) have been identified as one of the drug targets for the treatment of Type-2 diabetes. This study was designed to screen for PTP 1B and DPP-IV inhibitors from some Nigerian medicinal plants. **Materials and Methods:** PTP 1B and DPP-IV drug discovery kits from Enzo Life Sciences were used to investigate *in vitro* inhibitory effect of crude methanolic extract of 10 plants; *Mangifera indica*, *Moringa oleifera*, *Acacia nilotica*, *Arachis hypogaea*, *Senna nigricans*, *Azadirachta indica*, *Calotropis procera*, *Leptadenia hastata*, *Ziziphus mauritiana*, and *Solanum incanum*. **Results:** The results indicated PTP 1B inhibition by *S. nigricans* ( $68.2 \pm 2.29\%$ ), *A. indica* ( $67.4 \pm 2.80\%$ ), *A. hypogaea* ( $57.2 \pm 2.50\%$ ), *A. nilotica* ( $55.1 \pm 2.19\%$ ), and *M. oleifera* ( $41.2 \pm 1.87\%$ ) were significantly ( $P < 0.05$ ) higher as compared with standard inhibitor, sumarin while that of *L. hastata* ( $18.1 \pm 2.00\%$ ) was significantly lower as compared with sumarin. The PTP 1B inhibition by *M. indica* ( $31.5 \pm 1.90\%$ ) was not significantly ( $P > 0.05$ ) different from that of sumarin. The DPP-IV inhibition by *S. incanum* ( $68.1 \pm 2.71\%$ ) was significantly higher as compared with a known inhibitor, P32/98. *S. nigricans* ( $57.0 \pm 1.91\%$ ), *Z. mauritiana* ( $56.6 \pm 2.01\%$ ), *A. hypogaea* ( $51.0 \pm 1.30\%$ ), *M. indica* ( $44.6 \pm 2.40\%$ ), *C. procera* ( $36.2 \pm 2.00\%$ ), *A. nilotica* ( $35.4 \pm 2.10\%$ ), and *A. indica* ( $33.6 \pm 1.50\%$ ) show significantly ( $P < 0.05$ ) lower inhibitions toward DPP-IV. **Conclusion:** The work demonstrated that these plant materials could serve as sources of lead compounds in the development of anti-diabetic agent(s) targeting PTP 1B and/or DPP-IV.

**KEY WORDS:** Dipeptidyl peptidase IV, inhibition, medicinal plants, protein tyrosine phosphatase 1B, Type-2 diabetes mellitus

Department of  
Biochemistry, Usmanu  
Danfodiyo University,  
Sokoto

**Address for correspondence:**

Suleiman Alhaji Muhammad,  
Department of Biochemistry,  
Usmanu Danfodiyo  
University, Sokoto.  
Phone: +2348063779088.  
E-mail: smbinn98@gmail.  
com

**Received:** August 18, 2016

**Accepted:** December 04, 2016

**Published:** December 22, 2016

## INTRODUCTION

Herbal medicines have gained popularity worldwide due to their natural sources, low-cost, and less toxicity. There are several plant species that have been reported to have antidiabetic effect in diabetic animal models [1-5] and humans [6]. In Nigeria, *Mangifera indica*, *Moringa oleifera*, *Acacia nilotica*, *Arachis hypogaea*, *Senna nigricans*, *Azadirachta indica*, *Calotropis procera*, *Leptadenia hastata*, *Ziziphus mauritiana*, and *Solanum incanum* are being used traditionally for the treatment of diabetes mellitus, but very little is known about the mechanism of actions of antidiabetic activity of these medicinal plants. However, reports show that a number of bioactive ingredients such as flavonoids [7], alkaloids [8], and saponin [9] have been reported to exert antidiabetic activity. Ojiako *et al.* [10] stated that interplay of these bioactive constituents in medicinal plants could be responsible for the hypoglycemic effect. Extracts of *M. indica* [11], *A. hypogaea* [12,13], *A. nilotica* [14], *A. indica* [15],

*L. hastata* [16], *Z. mauritiana* [17], *C. procera* [18], and *M. oleifera* [19] have been reported to possess antidiabetic effects. Recently, efforts are being made toward elucidating the mechanism of actions of some of these medicinal plants and their active constituents.

Diabetes mellitus and its associated complications are the major cause of morbidity and mortality worldwide [20]. In particular, Type-2 diabetes mellitus is the most prevalent form of diabetes accounting for more than 80-90% of the total cases of diabetes [21,22]. Type-2 diabetes mellitus is associated with both macrovascular and microvascular complications that may result in tissue or organ damage. It is estimated that about 415 million people have diabetes in the world and more than 14 million cases in sub-Saharan African [23]. It is expected that this figure would be double in 2040. In Nigeria, there were more than 1.56 million cases of diabetes in 2015 and number of deaths related to adult diabetes (20-79 years) were estimated to be 40,815 [23].

Protein tyrosine phosphatases (PTB) are large family of surface proteins that are central modulators of tyrosine phosphorylation-dependent cellular activities [24,25]. Dipeptidyl peptidase IV (DPP IV) is a proteolytic enzyme that specifically deactivates glucagon-like peptide -1 (GLP-1), an incretin hormone which plays a significant role in the regulation of blood glucose level by stimulating the secretion of insulin, increasing  $\beta$ -cell mass and inhibit the secretion of glucagon [26]. PTP 1B and DPP-IV have been recognized as the best drug target for treatment of Type-2 diabetes mellitus [27,28]. Inhibition of PTP 1B increased the rate of phosphorylation of the insulin receptor and its substrate thereby promoting glucose transporters for the uptake of glucose by insulin sensitive cells while DPP-IV inhibitors maintain the level of active GLP-1 [26,28]. Therefore, medicinal products which contain essentially vast bioactive diversity may serve as potential sources of novel inhibitor(s) of PTP 1B and DPP-IV for the treatment of Type-2 diabetes mellitus. The study was designed to screen medicinal plants with PTP 1B and DPP-IV inhibitory activities.

## MATERIALS AND METHODS

### Chemicals and Reagents

PTP 1B and DPP-IV drug discovery assay kits used were products of Enzo® Life Sciences and other chemicals and reagent used were of analytical grade.

### Plant Materials

A total of 10 plant materials were screened. These included plant extracts studied in our laboratory with *in vivo* hypoglycemic activities and other plant materials that are used by traditional medical practitioners in Northwest Nigeria for the management of diabetes mellitus. Information about antidiabetic plants was sourced by oral interview of traditional medical practitioners, diabetic patients using some of these plants and the general public. In this respect, the following plants were used (Table 1).

The plant materials were collected from farms around Sokoto, Katsina and Kwara States of Nigeria. The plant materials were identified and authenticated by a Taxonomist, Dr. Umar Abdullahi, from Botany Unit, Department of Biological Sciences, Usmanu Danfodiyo University, Sokoto Nigeria. Voucher specimens were deposited at the herbarium of the same institution.

### Sample Preparation

The samples were shade dried and ground to powder using laboratory pestle and mortar, except groundnut seeds. The dried, ground powdered materials were stored in paper bags in desiccators until required. Ten g of the powdered samples were extracted in 100 mL of methanol for 72 h at room temperature, in brown cleaned reagent bottles, with intermittent mixing. At the end of the 72 h, the extracts were filtered using Whatman No. 1 filter paper. The filtrates were concentrated using rotary

evaporator, and the concentrated filtrates were left to dryness in a drying cabinet and the materials obtained were stored in air tight labeled container at 4°C for further analysis. The groundnut seeds were soaked in distilled water for about 3 h and the extract filtered and the dissolved solutes (% w/v) determined. The extracted materials were reconstituted in DMSO at 10 mg/ml and used for the preliminary screening for PTP 1B and DPP-IV inhibitory activities.

### PTP 1B Inhibition Assay

The kit components were thawed on an ice bath with the exception of BIOMOL RED™ that was stored at room temperature. The substrate ('IR5' Insulin receptor  $\beta$ , residues 1142-1153, Py1146) was reconstituted to a concentration of 1.5 mM by assay buffer and distilled H<sub>2</sub>O. The assay buffer, 100 mM MES, pH 6.0 containing 300 mM NaCl, 2mM EDTA, 2 mM DTT and 0.1% NP-40 was diluted with equal volume of distilled H<sub>2</sub>O and maintained on ice. The PTP 1B (human recombinant) was prepared in  $\times 1$  cold assay buffer. Stock of 10 mM of suramin (a known inhibitor) was prepared in assay buffer. The assay mixture was prepared in 96 well plate which contains 10  $\mu$ g per 100  $\mu$ L assay mixture of the sample. The plate reader was read at 620 nm, and all the assay protocol was done in accordance with manufacturer's instructions.

### DPP IV Inhibition Assay

The crude extracts were screened for DPP-IV inhibition at 100  $\mu$ g/mL in a total volume of 100  $\mu$ L using DPP-IV drug discovery assay kits. The inhibitor (P32/98) was diluted in the assay 1 in 10 buffer (50 mM Tris, pH 7.5). The substrate (H-Gly-Pro-pNA) and DPP IV (BML-SE434-9090) were diluted in 1 in 50  $\mu$ L of the assay buffer. The plant samples were reconstituted in 50 mM Tris buffer, pH 7.5, to give 1  $\mu$ g/ $\mu$ L. The assay mixture was prepared in 96 well plates which contains 10  $\mu$ g per 100  $\mu$ L assay mixture of the sample. The assay mixture consists of 15  $\mu$ L of DPP IV (17.3  $\mu$ U/ $\mu$ L) and 50  $\mu$ L of the substrate and was made up to 100  $\mu$ L with the assay buffer while P32/98 was used in the place of extracts as a control. The blank was prepared using the substrate and the buffer only. The plate was read continuously at 405 nm, in a microplate reader at 1 min interval for 10 min. The percentage inhibition of the two enzymes by test extracts was calculated based on the activity in control well as 100% from three independent replicates.

**Table 1: Medicinal plants screened**

Botanical Name	Part used	Voucher number
<i>Mangifera indica</i>	Leaf	UDUS/VS/2011/30
<i>Azadirachta indica</i>	Leaf	UDUS/VS/2011/34
<i>Moringa oleifera</i>	Leaf	UDUS/VS/2011/31
<i>Acacia nilotica</i>	Seed	UDUS/VS/2011/32
<i>Calotropis procera</i>	Leaf	UDUS/VS/2011/28
<i>Leptadenia hastata</i>	Leaf	UDUS/VS/2011/35
<i>Ziziphus mauritiana</i>	Leaf	UDUS/VS/2011/36
<i>Solanum incanum</i>	Fruit	UDUS/VS/2011/22
<i>Senna nigricans</i>	Leaf	UDUS/VS/2011/33
<i>Arachis hypogaea</i>	Whole seed	UDUS/VS/2011/24

## Data Analysis

The values are expressed as mean percentage inhibition  $\pm$  standard deviation. The mean percentage inhibition was analyzed using one-way ANOVA with SPSS (Version 17.0), and  $P < 0.05$  was considered statistically significant.

## RESULTS

The results of the percentage inhibition of PTP 1B and DPP-IV of the crude methanol extract of medicinal plants used in the Northwest Nigeria are presented in Tables 2 and 3, respectively. The result shows that *S. nigricans* and *A. indica* show the highest PTP 1B inhibition of  $68.2 \pm 2.29\%$  and  $67.4 \pm 2.80\%$ , respectively, followed by *A. hypogaea* ( $57.2 \pm 2.58\%$ ), *A. nilotica* ( $55.1 \pm 2.19\%$ ), *M. oleifera* ( $41.2 \pm 1.87\%$ ) which were significantly ( $P < 0.05$ ) higher as compared with sumarin ( $30.1 \pm 2.00\%$ ). The PTP 1B inhibition by *M. indica* ( $31.5 \pm 1.90\%$ ) was not significantly ( $P > 0.05$ ) different as compared with the standard inhibitor, sumarin while *L. hastata* with the least inhibition of  $18.1 \pm 2.00\%$ . *C. procera*, *S. incanum*, and *Z. mauritiana* show no inhibition against PTP 1B activity which could serve as activators.

The results for DPP-IV inhibition indicated that *S. incanum* ( $68.1 \pm 2.71\%$ ) was significantly ( $P < 0.05$ ) higher as compared with a known inhibitor, P32/98 ( $63.1 \pm 2.70\%$ ) while inhibition activity by *S. nigricans* ( $57.0 \pm 1.91\%$ ), *Z. mauritiana* ( $56.6 \pm 2.01\%$ ) *Arachis hypogaea* ( $51.0 \pm 1.30\%$ ), *M. indica* ( $44.6 \pm 2.40\%$ ), *C. procera* ( $36.2 \pm 2.00\%$ ), *A. nilotica* ( $35.4 \pm 2.10\%$ ), and *A. indica* ( $33.6 \pm 1.50\%$ ) were significantly ( $P < 0.05$ ) lower as compared with P32/98. There was no inhibition of DPP-IV activity by *L. hastata* and *M. oleifera* which suggest the plants could act as activators of the enzyme.

## DISCUSSION

The treatment of diabetes mellitus is considered a global challenge and evaluation of plant products with the aim of isolating antidiabetic agents is gaining popularity worldwide due to the presence of several bioactive constituents with minimal side effect. Selective inhibition of PTP 1B and DPP-IV has been suggested as novel therapeutic target for the treatment of Type-2 diabetes mellitus. In this study, inhibitory activities of ten medicinal plants on PTP 1B and DPP-IV were investigated. The result indicated that *S. nigricans*, *A. indica*, *A. hypogaea*, *A. nilotica*, *M. oleifera*, *M. indica*, and *L. hastata* possess significant potentials as sources of lead compounds for the development of PTP 1B inhibitors for the management of Type-2 diabetes mellitus. Similarly, *S. incanum*, *S. nigricans*, *Z. mauritiana*, *A. hypogaea*, *M. indica*, *C. procera*, *A. nilotica* and *A. indica* were active against DPP IV, which may serve as sources of inhibitors of the enzyme in the treatment of Type-2 diabetes mellitus. Natural inhibitors like berberine, an isoquinoline alkaloid has been reported to possess potent antidiabetic properties via inhibition of PTP 1B [29-31] and DPP-IV [32]. Papaverine, a structural analog of berberine which belongs to member of isoquinoline alkaloids have also

**Table 2: Percentage inhibition of crude methanol extract of different medicinal plants against PTP 1B**

Plant materials	% inhibition
<i>Mangifera indica</i>	31.5 $\pm$ 1.90
<i>Azadirachta indica</i>	67.4 $\pm$ 2.80*
<i>Calotropis procera</i>	NI
<i>Acacia nilotica</i>	55.1 $\pm$ 2.19*
<i>Leptadenia hastate</i>	18.1 $\pm$ 2.00*
<i>Solanum incanum</i>	NI
<i>Ziziphus mauritiana</i>	NI
<i>Senna nigricans</i>	68.2 $\pm$ 2.29*
<i>Moringa oleifera</i>	41.2 $\pm$ 1.87*
<i>Arachis hypogaea</i>	57.2 $\pm$ 2.58*
Sumarin	30.12 $\pm$ 2.00

Data are expressed as Mean $\pm$ SD, n=3 replicate, NI- no inhibition, \* $p < 0.05$  when compared with sumarin, standard inhibitor  
PTP 1B: Protein tyrosine phosphatase 1B

**Table 3: Percentage inhibition of crude methanol extract of different medicinal plants against DPP IV**

Plant materials	% inhibition
<i>Mangifera indica</i>	44.6 $\pm$ 2.40*
<i>Azadirachta indica</i>	33.6 $\pm$ 1.50*
<i>Calotropis procera</i>	36.2 $\pm$ 2.00*
<i>Acacia nilotica</i>	35.4 $\pm$ 2.10*
<i>Leptadenia hastate</i>	NI
<i>Solanum incanum</i>	68.1 $\pm$ 2.71*
<i>Ziziphus mauritiana</i>	56.6 $\pm$ 2.01*
<i>Senna nigricans</i>	57.0 $\pm$ 1.91*
<i>Moringa oleifera</i>	NI
<i>Arachis hypogaea</i>	51.0 $\pm$ 1.30*
P32/98	63.1 $\pm$ 2.70

Data are expressed as Mean $\pm$ SD, n=3 replicate, NI- no inhibition, \* $p < 0.05$  when compared with P32/98, standard inhibitor DPP  
IV: Dipeptidyl peptidase IV

been reported to exhibit potent PTP 1B inhibitory activity thereby lowering fasting blood glucose level *in vivo* [33]. Hydroalcoholic extracts of *Terminalia arjuna* and *Commiphora mukul* have been shown to possess significant DPP-IV inhibitory activity [34]. Although hypoglycemic effects of some of these plants screened have been reported, the mechanism of action has not been fully elucidated. It may be interesting to study whether the antidiabetic effect of these plants extracts acts via inhibition of PTP 1B and/or DPP IV activities. Therefore, PTP 1B and DPP-IV inhibitory activities of some of these plants observed in this study indicate that they may serve as potent sources of hypoglycemic agent(s) for the treatment of Type-2 diabetes mellitus. Overexpression of PTP 1B is associated with the development of insulin resistance which could lead to Type-2 diabetes mellitus and obesity [35]. Of all the plants studied, *S. nigricans* had shown to be a better inhibitor of PTP 1B while *S. incanum* had a better effect against DPP-IV. The data reported in this study have shown that *S. nigricans*, *A. nilotica*, *A. hypogaea*, *A. indica*, and *M. oleifera* are better inhibitors of PTP 1B than a known inhibitor, sumarin.

Furthermore, going by the result obtained it may be interesting to further exploit these natural products to investigate *in vivo* hypoglycemic effect, study the kinetics of the two enzymes and possibly isolate and characterize the bioactive component(s)

responsible for the inhibition. *S. nigricans* which have shown to be a better source of inhibitor of PTP 1B and to some extent DPP-IV inhibitors could be very promising sources of lead compound(s) for the treatment of Type-2 diabetes mellitus.

## CONCLUSION

The results of this work indicated that these plants possess either inhibitory activity against PTP 1B and/or DPP-IV. *S. nigricans* possess significant potentials as sources of lead compound(s) in the development of inhibitors for PTP 1B and to some extent DPP-IV than others. It can then be concluded that these plant materials could serve as sources of lead compounds in the development of antidiabetic agent(s) targeting PTP 1B and/or DPP-IV.

## ACKNOWLEDGMENT

This work was supported by research grant from International Foundation of Science (IFS) Grant No F-5020-1.

## REFERENCES

- Trojan-Rodrigues M, Alves TL, Soares GL, Ritter MR. Plants used as antidiabetics in popular medicine in Rio Grande do Sul, Southern Brazil. *J Ethnopharmacol* 2012;139:155-63.
- Nazreen S, Kaur G, Alam MM, Shafi S, Hamid H, Ali M, et al. New flavones with antidiabetic activity from *Callistemon lanceolatus* DC. *Fitoterapia* 2012;83:1623-7.
- Ali S, Igoli J, Clements C, Dima S, Muhammad A, Rashid M-U, et al. Antidiabetic and antimicrobial activities of fractions and compounds isolated from *Berberis brevissima* Jafri and *Berberis parkeriana* schneid. *Bangladesh J Pharmacol* 2013;8:336-42.
- Amran MS, Sultan MZ, Rahman A, Rashid MA. Antidiabetic activity of compounds isolated from the kernel of *Mangifera indica* in alloxan induced diabetic rats. *Dhaka Univ J Pharm Sci* 2013;12:77-81.
- Hua J, Qi J, Yu BY. Iridoid and phenylpropanoid glycosides from *Scrophularia ningpoensis* hemsl. And their  $\alpha$ -glucosidase inhibitory activities. *Fitoterapia* 2014;93:67-73.
- Musabayane CT. The effect of medicinal plants on renal function and blood pressure in diabetes mellitus. *Cardiovasc J Afr* 2012;12:77-81.
- Ma DQ, Jiang ZJ, Xu SQ, Yu X, Hu XM, Pan HY. Effects of flavonoids in *Morus indica* on blood lipids and glucose in hyperlipidemia-diabetic rats. *Chin Herb Med* 2012;4:314-8.
- Prabhakar PK, Doble M. A target based therapeutic approach towards diabetes mellitus using medicinal plants. *Curr Diabetes Rev* 2008;4:291-308.
- Elekofehintia OO, Kamdemb JP, Kadec IJ, Rochab JB, Adanlawod IG. Hypoglycemic, antiperoxidative and antihyperlipidemic effects of saponins from *Solanum anguivi* Lam. Fruits in alloxan-induced diabetic rats. *S Afr J Bot* 2013;88:56-61.
- Ojiako OA, Chikezie PC, Ogbuji AC. Blood glucose level and lipid profile of alloxan-induced hyperglycemic rats treated with single and combinatorial herbal formulations. *J Tradit Complement Med* 2015;6:184-92.
- Mohan CG, Viswanatha GL, Savinay G, Rajendra CE, Halemani PD. 1,2,3,4,6 Penta-O-galloyl- $\beta$ -D-glucose, a bioactivity guided isolated compound from *Mangifera indica* inhibits 11 $\beta$ -HSD-1 and ameliorates high fat diet-induced diabetes in C57BL/6 mice. *Phytomedicine* 2013;20:417-26.
- Billbis LS, Shehu RA, Abubakar MG. Hypoglycemic and hypolipidemic effects of aqueous extract of *Arachis hypogaea* in normal and alloxan-induced diabetic rats. *Phytomedicine* 2002;9:553-5.
- Akter F, Jahan N, Sultana N. Effect of peanut (*Arachis Hypogaea* L.) On fasting blood glucose and Hba1c in alloxan induced diabetic male rats. *J Bangladesh Soc Physiol* 2014;9:48-53.
- Mukundi MJ, Piero NM, Mwaniki NE, Murugi NJ, Daniel AS, Peter GK, et al. Antidiabetic effects of aqueous leaf extracts of *Acacia nilotica* in alloxan induced diabetic mice. *J Diabetes Metab* 2015;6:568.
- Satyanarayana K, Sravanthi K, Shaker IA, Ponnulakshmi R. Molecular approach to identify antidiabetic potential of *Azadirachta indica*. *J Ayurveda Integr Med* 2015;6:165-74.
- Sanda KA, Sandabe UK, Auwal MS, Bulama I, Bashir TM, Sanda FA, et al. Hypoglycemic and antidiabetic profile of the aqueous root extracts of *Leptadenia hastata* in albino rats. *Pak J Biol Sci* 2013;16:190-4.
- Bhatia A, Mishra T. Hypoglycemic activity of *Ziziphus mauritiana* aqueous ethanol seed extract in alloxan-induced diabetic mice. *Pharm Biol* 2010;48:604-10.
- Neto MC, de Vasconcelos CF, Thijan VN, Caldas GF, Araújo AV, Costa-Silva JH, et al. Evaluation of antihyperglycaemic activity of *Calotropis procera* leaves extract on streptozotocin-induced diabetes in wistar rats. *Rev Bras Farmacogn* 2013;23:913-9.
- Al-Malki AL, El Rabey HA. The antidiabetic effect of low doses of *Moringa oleifera* Lam. Seeds on streptozotocin induced diabetes and diabetic nephropathy in male rats. *Biomed Res Int* 2015;2015:381040.
- Zimmet P, Alberti KG, Shaw J. Global and societal implications of the diabetes epidemic. *Nature* 2001;414:782-7.
- Mlinar B, Marc J, Janez A, Pfeifer M. Molecular mechanisms of insulin resistance and associated diseases. *Clin Chim Acta* 2007;375:20-35.
- Dorajoo R, Liu J, Boehm BO. Genetics of Type 2 diabetes and clinical utility. *Genes (Basel)* 2015;6:372-84.
- International Diabetes Federation (IDF). Diabetes Atlas; Available from: <http://www.idf.org/diabetesatlas>. [Last accessed on 2016 Aug 15].
- Tonks NK. Protein tyrosine phosphatases: From genes, to function, to disease. *Nat Rev Mol Cell Biol* 2006;7:833-46.
- Koren S, Fantus IG. Inhibition of the protein tyrosine phosphatase PTP1B: A potential therapy for obesity, insulin resistance and type-2 diabetes mellitus. *Best Pract Res Clin Endocrinol Metab* 2007;21:621-640.
- Wang Y, Li L, Yang M, Liu H, Yang G. Glucagon-like peptide-1 receptor agonist versus insulin in inadequately controlled patients with Type 2 diabetes mellitus: Meta-analysis of clinical trials. *Diabetes Obe Metab* 2011;13:972-82.
- Zhang ZY, Lee SY. PTP1B inhibitors as potential therapeutics in the treatment of Type 2 diabetes and obesity. *Expert Opin Investig Drugs* 2003;12:223-33.
- Singh AK. Dipeptidyl peptidase-4 inhibitors: Novel mechanism of actions. *Indian J Endocrinol Metab* 2014;18:753-9.
- Ko BS, Choi SB, Park SK, Jang JS, Kim YE, Park S. Insulin sensitizing and insulinotropic action of berberine from *Coptidis rhizoma*. *Biol Pharm Bull* 2005;28:1431-7.
- Bustanji Y, Taha MO, Yousef AM, Al-Bakri AG. Berberine potently inhibits protein tyrosine phosphatase 1B: Investigation by docking simulation and experimental validation. *J Enzyme Inhib Med Chem* 2006;21:163-71.
- Chen C, Zhang Y, Huang C. Berberine inhibits PTP1B activity and mimics insulin action. *Biochem Biophys Res Commun* 2010;397:543-7.
- Al-masri IM, Mohammad MK, Taha MO. Inhibition of dipeptidyl peptidase IV (DPP IV) is one of the mechanisms explaining the hypoglycemic effect of berberine. *J Enzyme Inhib Med Chem* 2009;24:1061-6.
- Bustanji Y, Taha MO, Al-Masri IM, Mohammad MK. Docking simulations and *in vitro* assay unveil potent inhibitory action of papaverine against protein tyrosine phosphatase 1B. *Biol Pharm Bull* 2009;32:640-5.
- Borde MK, Suman RK, Mohanty IR, Deshmukh YA. Dipeptidyl peptidase-IV inhibitory activities of medicinal plants: *Terminalia arjuna*, *Commiphora mukul*, *Gymnema sylvest*, *Morinda citrifolia*, *Emblca officinalis*. *Asian J Pharm Clin Res* 2106;9:180-2.
- Panzhinskiy E, Ren J, Nair S. Pharmacological inhibition of protein tyrosine phosphatase 1B: A promising strategy for the treatment of obesity and Type 2 diabetes mellitus. *Curr Med Chem* 2013;20:2609-25.

© EJManager. This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, noncommercial use, distribution and reproduction in any medium, provided the work is properly cited.

Source of Support: Nil, Conflict of Interest: None declared.