

RESEARCH

Open Access

Aspidistra crassifila (Asparagaceae), a new species from Guangxi, China

Chun-Rui Lin¹, Yan Liu^{1*}, Dong-Xin Nong², Yoshiko Kono³ and Ching-I Peng^{3*}

Abstract

Background: *Aspidistra crassifila* Yan Liu & C.-I Peng, a new species of the Asparagaceae from Guangxi Zhuang Autonomous Region, China, is described and illustrated.

Results: The new species is similar to *A. subrotata* Y. Wan & C. C. Huang in the perianth lobes triangular-lanceolate and horizontally spreading, but differs by the perianth campanulate, lobes with appendages at base, stamens 6–8 mm long, filaments enlarged, anthers adnate to perianth tube, connectives extended and upcurved. The chromosome number of the new species was determined to be $2n = 38$, and the karyotype was formulated as $2n = 22m^{2SC} + 4sm + 12st$.

Conclusion: A careful study of the literature, herbarium specimens and living plants, both in the wild and in cultivation in the experimental greenhouse, support the recognition of the new species *Aspidistra crassifila*, which is described herein. *Aspidistra crassifila* is currently known only from Shiwandashan Mountains, which lie in southern Guangxi. A line drawing, color plates and a distribution map are given for the new species to aid in identification.

Keywords: *Aspidistra crassifila*; *Aspidistra subrotata*; Asparagaceae; New species; Chromosome number; Karyotype; Guangxi; China

Background

The genus *Aspidistra* Ker-Gawler (Asparagaceae) was established in 1822 and is comprised of ca. 100 species, more than 60 of which occur in China (Lang et al., 1999; Li, 2004; Tillich, 2005, 2008; Hou et al., 2009; Lin et al., 2009, 2010; Liu et al., 2011). In March 2005, during a field trip to Shiwandashan Mountains in southern Guangxi Zhuang Autonomous Region, China, the second author (Yan Liu) collected and brought back a sterile plant of *Aspidistra* for cultivation, which flowered in Guilin Botanical Garden next spring. We went on another trip to the same locality in Shiwandashan Mountains in January 2007 and were able to collect fruiting materials of this species. Compared with other species of *Aspidistra*, it was recognized as an undescribed species that differs from congeners in its peculiar adnate stamens with extended and upcurved connectives.

Methods

Chromosome preparations

Somatic chromosomes were examined for plants collected from the type collection (*Yan Liu L1380*). Root tips were pretreated in 2 mM 8-hydroxyquinoline at 15–18°C for about 8 h, then fixed overnight in a 3:1 ethanol-acetic acid solution below 4°C. The chromosomes were stained with 2% acetic orcein in 1mol/L hydrochloric acid and observed. Classification of chromosome morphology is based on the position of the centromere, following Levan et al. (1964).

Results and discussion

Taxonomic treatment

Aspidistra crassifila Yan Liu & C.-I Peng, sp. nov.—TYPE: CHINA. Guangxi Zhuang Autonomous Region, Fangcheng (City), Shiwandashan Mountains, alt. 980 m, 10 March 2005. Specimens pressed from plants introduced to Guilin Botanical Garden, Guilin City, Yanshan Township on 11 May 2006, *Yan Liu L1380* (holotype: IBK; isotype: HAST) 粗丝蜘蛛抱蛋 Figures 1, 2.



Species nova *A. subrotatae* Y. Wan & C. C. Huang affinis, sed differt perianthio campanulato (vs. subrotato), lobis basi appendiculatis (vs. non appendiculatis, margine

* Correspondence: gxibly@163.com; bopeng@sinica.edu.tw

¹Guangxi Institute of Botany, Guangxi Zhuangzu Autonomous Region and the Chinese Academy of Sciences, Guilin 541006, P.R. China

³Herbarium (HAST), Biodiversity Research Center, Academia Sinica, Nangang, Taipei 115, Taiwan

Full list of author information is available at the end of the article



reflexis), filamentis dilatatis (vs. non dilatatis), antheris adnatis (vs. versatilibus), connectivis supra antheras productis (vs. non productis).

Description

Herbs perennial, evergreen, rhizomatous. Rhizome creeping, subterete, 8–13 mm thick, covered with scales, nodes dense. Vaginal leaves 4–5, 1–14 cm long, purple-red, enveloping base of petiole, fibrous when withered. Leaves solitary, 2–5 cm apart; petiole stiff, upright, 10–40 cm long, 3–5 mm thick, adaxially sulcate; leaf blade oblong-oblancheolate, 30–60 cm long, 6–12 cm wide, base cuneate, gradually narrowed into petiole, inequilateral, apex acuminate, margin entire. Peduncle purplish red to purplish black, 2.5–6 cm long, with 4–5 bracts, bracts gradually wider from base to top of peduncle; the uppermost bract broadly ovate, purplish red or purplish black, ca. 10 mm long, ca. 12 mm wide, apex subobtusate. Flowers solitary; perianth purplish black, fleshy, campanulate, 4–6 cm in

diam., 8–12 lobed apically; lobes triangular-lanceolate, 15–25 mm long and 5–10 mm wide at base, apex gradually acuminate, horizontally spreading, with appendages at base, tube 8–12 mm long, distal opening 15–20 mm diam.; stamens as many as and opposite to lobes, 6–8 mm long, inserted in the middle of perianth tube, positioned lower than stigma, filaments purplish black, enlarged, 3–4 mm wide at side view, their upper surfaces visible from above (between adaxial surface of perianth tube and margin of stigma), anthers adnate to perianth, pale yellow, oblong, 5–6 mm long, ca. 2 mm wide, connectives extended and upcurved; pistil mushroom-shaped, 2 cm long, style cylindrical, ca. 10 mm long, ca. 4 mm across, purplish red, ovary inconspicuous, stigma enlarged, ca. 10 mm high, 12–15 mm across, upper surface white with purple spots, smooth, lower surface purplish black, irregularly concave and undulate at margin. Berry subglobose, ca. 3.5 cm across, tuberculate.

Additional specimens examined

CHINA. Guangxi Zhuang Autonomous Region, Fangcheng (City), Shiwandashan Mountains, alt. 980 m, 10 March 2005, *Yan Liu L1156* (IBK); same locality, 17 January 2007, *Yan Liu L1425* (IBK).

Chromosome cytology

The chromosome number of *Aspidistra crassifila* was determined to be $2n = 38$ (Figure 3), showing a trimodal variation in chromosome length at mitotic metaphase. Among the 38 chromosomes, the first two were much longer (ca 10.5-10.6 μm) than the rest; the next 16 gradually varied, ca 3.9-7.5 μm ; the remaining 20 chromosomes also gradually varied, ca. 1.9-2.8 μm long. Regardless of the chromosome length, 22 (Nos. 1, 2 and 19-38 in Figure 3B), 4 (Nos. 7, 8 and 17, 18 in Figure 3B), 12 (Nos. 3-6 and 9-16 in Figure 3B) had centromere at the median (m), submedian (sm), and subterminal (st) positions, respectively. Secondary constrictions (SC) were

observed at the proximal regions of the short arms in two submedian chromosomes (arrows in Figure 3A; Nos. 19 and 20 in Figure 3B). Thus, the karyotype formula of *Aspidistra crassifila* is $2n = 38 = 22\text{m}^{2\text{SC}} + 4\text{sm} + 12\text{st}$.

Previously, detailed cytological data were known for 42 species in the genus *Aspidistra* (Bogner and Arnautov, 2004; Li, 2004; Yamashita and Tamura, 2004; Qiao et al., 2008; Hou et al., 2009, Lin et al., 2010; Liu et al., 2011). *Aspidistra crassifila* shared a number of cytological features, namely basic chromosome numbers of $x = 19$, trimodal chromosome complement, the first pair of longer median chromosomes and 10th median SC-chromosomes, in common with those of the *Aspidistra* species previously reported (Lin et al., 2010).

Ecology

On shaded bamboo slopes in seasonal rain forests (monsoon forests).

Distribution

Currently known only from Shiwandashan Mountains, Fangcheng (City), in southern Guangxi Zhuang Autonomous Region, China (Figure 4).

Phenology

Flowering from March to May; fruits maturing in May next year.

Etymology

The specific epithet '*crassifila*' is derived from its enlarged filaments.







Figure 5 *Aspidistra subrotata* Y. Wan & C. C. Huang. **A**, Habit; **B, C**, Flowers; **D, E**, Flower, dissected to show stamens and pistil.

Notes

Aspidistra crassifila (Figures 1, 2) resembles *A. subrotata* Y. Wan & C. C. Huang (Figure 5; Wan and Huang, 1987) in the perianth lobes triangular-lanceolate, horizontally spreading, but differs in its perianth campanulate (vs. subrotate), lobes with appendages at base (vs. without appendages, margin reflexed), filaments enlarged (vs. non enlarged), anthers adnate (vs. versatile) to parianth. Connectives extended and upcurved (vs. not extended). Also the purplish black perianth color is rather rare in *Aspidistra*, known only from *A. renatae* C. Bräuchler (Bräuchler and Ngoc, 2005), *A. nikolai* L.V. Averyanov & H.-J. Tillich (Tillich and Averyanov, 2008), *A. atroviolacea* H.-J. Tillich (Tillich, 2008), and *A. pileata* D. Fang & L. Y. Yu (Fang and Yu, 2002).

Conclusion

A careful study of the literature, herbarium specimens and living plants, both in the wild and in cultivation in the experimental greenhouse, support the recognition of the new species *Aspidistra crassifila*, which is described herein. *Aspidistra crassifila* is currently known only from Shiwandashan Mountains, which lie in southern Guangxi.

Competing interests

The authors declare that they have no competing interests.

Authors' contribution

YL discovered the new species and collected field data; CRL and DXN participated fieldwork and prepared the earlier draft of description; YK performed cytological studies; CIP cultivated the new species to collect morphological and cytological data, and prepared the final manuscript. All authors read and approved the final manuscript.

Acknowledgments

The authors are grateful to Prof. Fa-Nan Wei (IBK) for Latin diagnosis. We also thank Mr. Shun-Qing He (IBK) for the handsome drawing. This study was supported by the Guangxi Natural Science Foundation (2010GXNSFE13004, 2011GXNSFE18001) to Yan Liu; postdoctoral fellowship from Academia Sinica, Taiwan to Yoshiko Kono; and National Geographic Society Grant # 8358-07 (Botanical Exploration of Limestone Karsts of Southern Guangxi, China) to Ching-I Peng.

Author details

¹Guangxi Institute of Botany, Guangxi Zhuangzu Autonomous Region and the Chinese Academy of Sciences, Guilin 541006, P.R. China. ²Guangxi Botanical Garden of Medicinal Plants, Nanning 530023, P.R. China. ³Herbarium (HAST), Biodiversity Research Center, Academia Sinica, Nangang, Taipei 115, Taiwan.

Received: 5 March 2013 Accepted: 11 April 2013

Published: 7 October 2013

References

- Bogner J, Arnautov N (2004) *Aspidistra locii* (Convallariaceae), an unusual new species from Vietnam. *Willdenowia* 34:203–208
- Bräuchler C, Ngoc LH (2005) *Aspidistra renatae* (Ruscaceae), a new species from central Vietnam. *Blumea* 50:527–529
- Fang D, Yu L-Y (2002) Three new species of *Aspidistra* Ker-Gawl. (Liliaceae) from Guangxi, China. *Acta Phytotax Sin* 40:159–163
- Hou M-F, Liu Y, Kono Y, Peng C-I (2009) *Aspidistra dixinensis* (Ruscaceae), a new species from limestone areas in Guangxi. *China Bot Stud* 50:371–378
- Liang K-Y, Li G-Z, Liu Y, Wei Y-G, Wang R-X (1999) Taxonomic and phytogeographic studies on the genus *Aspidistra* Ker-Gawl. (Liliaceae) in China. *Acta Phytotax Sin* 37:468–508
- Levan A, Fredga K, Sandberg AA (1964) Nomenclature for centromeric position on chromosomes. *Hereditas* 52:201–220
- Li G-Z (2004) The Genus *Aspidistra*. Guangxi Science & Technology Publishing House, Nanning
- Lin C-R, Liang Y-Y, Liu Y (2009) *Aspidistra bamaensis* (Ruscaceae), a new species from Guangxi. *China Ann Bot Fenn* 46:416–418
- Lin C-R, Peng C-I, Kono Y, Liu Y (2010) *Aspidistra obconica*, Asparagaceae [Ruscaceae], a new species from limestone areas in Guangxi. *China Bot Stud* 51:263–268
- Liu Y, Kono Y, Lin C-R, Xu W-B, Peng C-I (2011) *Aspidistra erecta* (Asparagaceae), a new species from limestone areas in Guangxi. *China Bot Stud* 52:367–373
- Qiao Q, Zhang C-Q, Ma Y-P, Tian W (2008) Karyotype asymmetry of *Aspidistra* (Convallariaceae, Ruscaceae). *Acta Bot Yunnan* 30:565–569
- Tillich HJ (2005) A key for *Aspidistra* (Ruscaceae), including fifteen new species from Vietnam. *Feddes Repert* 116:313–338
- Tillich HJ (2008) An updated and improved determination key for *Aspidistra* Ker-Gawl. (Ruscaceae, Monocotyledons). *Feddes Repert* 119:449–462
- Tillich HJ, Averyanov LV (2008) Two new species and one new subspecies of *Aspidistra* Ker-Gawl. (Ruscaceae) from Vietnam. *Feddes Repert* 119:37–41
- Wan Y, Huang C-C (1987) New species of the genus *Aspidistra* from Guangxi. *Guihaia* 7:217–224
- Yamashita J, Tamura MN (2004) Phylogenetic analyses and chromosome evolution in Convallariaceae (Ruscaceae sensu lato), with some taxonomic treatments. *J Plant Res* 117:363–370

doi:10.1186/1999-3110-54-43

Cite this article as: Lin et al.: *Aspidistra crassifila* (Asparagaceae), a new species from Guangxi, China. *Botanical Studies* 2013 **54**:43.