



HHS Public Access

Author manuscript

J Occup Environ Med. Author manuscript; available in PMC 2018 May 01.

Published in final edited form as:

J Occup Environ Med. 2017 May ; 59(5): e91–e96. doi:10.1097/JOM.0000000000001004.

Chemical and Physical Exposures in the Emerging US Green Collar Workforce

Charles J. Chen, BS¹, Kevin J. Moore, BA¹, Cristina A. Fernandez, MEd, PhD¹, Kristopher L. Arheart, EdD¹, William G. LeBlanc, PhD¹, Manuel Cifuentes, MD, ScD³, Laura A. McClure, MSPH¹, Sharon L. Christ, PhD⁴, Lora E. Fleming, MD, PhD, MPH, MSc^{1,2}, David J. Lee, PhD¹, and Alberto J. Caban-Martinez, DO, PhD, MPH¹

¹Dept. Public Health Sciences, University of Miami, Miller School of Medicine, Miami, FL

²European Centre for Environment and Human Health and University of Exeter Medical School, Knowledge Spa, Royal Cornwall Hospital, Truro, Cornwall, UK

³Dept. of Public Health, Regis College, Westin, MA

⁴Dept. of Human Development & Family Studies, Purdue University, West Lafayette, IN

Abstract

Objective—“Green collar” workers serve in occupations that directly improve environmental quality and sustainability. This study estimates and compares the prevalence of select physical and chemical exposures among green versus non-green U.S. workers.

Methods—Data from the U.S. 2010 National Health Interview Survey(NHIS) Occupational Health Supplement were linked to the Occupational Information Network(O*NET) Database. We examined four main exposures:1)vapors, gas, dust, fumes(VGDF);2)secondhand tobacco smoke; 3)skin hazards;4)outdoor work.

Results—Green collar workers were significantly more likely to report exposure to VGDF and outdoor work than non-green collar workers(Adjusted Odds Ratio[AOR]=1.25; 95% CI=1.11–1.40; AOR=1.44(1.26–1.63), respectively). Green collar workers were less likely to be exposed to chemicals(AOR=0.80; 0.69–0.92).

Conclusions—Green collar workers appear to be at greater risk for select workplace exposures. As the green industry continues to grow, it is important to identify these occupational hazards in order to maximize worker health.

Corresponding Author: Alberto J. Caban-Martinez, Department of Public Health Sciences, University of Miami, Miller School of Medicine, 1120 N.W. 14th Street, R-669, Clinical Research Building, Office #1025, Miami, Florida 33136, Phone: 305-243-7565, acaban@med.miami.edu.

Author Contributions Statement: CC, KJM, AJCM, DJL, and LEF conceived the study, participated in its design, coordination, performed statistical analyses, and co-drafted the manuscript. WGL and KLA participated in the design of the study, performed statistical analysis and helped with the manuscript draft. SLC, CAF, LAM, and MC read, revised, and approved the final manuscript.

Institution and Ethics Approval and informed consent: This study was approved by the University of Miami Institutional Review Board (IRB #: 20120714).

Disclosure: The authors declare no conflict of interest

Disclaimer: None

Keywords

Green jobs; chemical exposures; physical exposures; National Health Interview Survey; O*NET

INTRODUCTION

“Green” collar workers are individuals employed in businesses whose services and work products directly improve environmental-friendliness and sustainability. Green collar jobs are defined as those that involve the protection of wildlife or ecosystems, the decline of pollution and waste, and/or the reduction of energy usage and carbon emissions.^(1, 2) These jobs span across multiple industries, from the construction of energy efficient buildings and vehicles to the generation of renewable energy power (e.g., biofuels development).^(1, 3) Despite increasing evidence linking climate change to human activity, jobs in environmentally friendly and sustainable industries did not garner significant political support in the US until 2007. The green collar workforce is expected to expand significantly in the coming years with estimates of 40 million jobs in the renewable energy industries by 2030, a significant increase from the 8.5 million jobs in 2008⁽⁴⁾. This new found political and economic support contributed to the increased numbers of Green collar workers and focused attention on the potential occupational hazards and health conditions of this emerging workforce.^(5–7)

Across occupational categories, occupational chemical and physical exposures have been associated with detrimental health consequences. For example, studies have shown that occupational exposures to vapor, dust, and smoke are associated with chronic and non-chronic dry cough and other respiratory symptoms.^(8, 9) Occupational vapor, dust, and smoke exposures have also been linked to complications, such as asthma and chronic obstructive pulmonary disease.⁽¹⁰⁾ Chemical exposures in the workplace have been linked to adverse health effects such as respiratory (e.g., stuffy nose and cough), ocular (e.g., watery, itchy, or burning eyes), and dermal symptoms (e.g., rash and itchy or burning skin).^(11, 12) Some occupational chemical exposures have also been linked to increased risk for cancer (e.g., skin cancer, bladder cancer, and lung cancer).⁽¹³⁾ Studies have also shown that occupational outdoor work is associated with increased sunlight exposure, which in turn has led to an increase risk of skin cancer.⁽¹⁴⁾ Nevertheless, little is known of the specific chemical and physical workplace exposures associated with the emerging green collar workforce.

This study uses nationally representative data to estimate and to compare the prevalence of chemical and physical exposures in green versus non-green collar U.S. workers. Previous studies have shown that 40% of respondents of a similar occupational categorization reported exposure to vapors, gas, dust, or fumes.⁽¹⁵⁾ Shopland et al. analyzed data from the national Current Population Survey (CPS), a survey administered by the Bureau of the Census for the Bureau of Labor Statistics, and found that the percentage of workers who reported a smoke-free workplace policy had increased from 46% in 1993 to 69% in 1999.^(16–18)

The emerging green collar workforce is likely not immune to traditional occupational hazards, and may be exposed to new evolving workplace harmful or protective health and safety factors. Due to advocacy in promoting environmental sustainability possibly extending to their workplaces, we would expect green collar workers to have lower rates of vapors, gas, dust, or fumes, secondhand smoke exposures, and chemical exposures. As for work outdoor exposure, it is evident that a portion of the green collar workforce is dedicated to wildlife preservation and the use of renewable energy resources and construction, which inherently involves more outdoor work than non-green jobs for some green collar workers.^(1–3) Therefore, we hypothesized that green collar workers would report lower rates of exposures to vapors, gas, dust, or fumes, secondhand tobacco smoke, and chemical exposures, but higher rates of work-related outdoor exposure relative to non-green collar workers.

METHODS

Data Sources

2010 National Health Interview Survey (NHIS) Occupational Health

Supplement—The NHIS is a nationally representative survey that collects data on a wide range of health topics. The NHIS is the primary source of information on the health of the civilian non-institutionalized population residing in the U.S since 1957. The cross-sectional household survey is administered annually by the National Center for Health Statistics (NCHS). The NHIS uses multistage, clustered, non-probability sampling techniques. The 2010 NHIS Occupational Supplement provides unique national estimates of the prevalence of common workplace exposures and work-related health conditions. Information on individual job type, employment status, health status, medical conditions, health care utilization and access, and health-related behaviors was collected. ⁽¹⁹⁾ Adult NHIS respondents aged 18 and older who reported being employed in the week prior to survey administration were included in the analyses.

2010 Occupational Information Network online (O*NET), version 19.0—The O*NET is an online occupational database sponsored by the U.S. Department of Labor that is periodically updated with new U.S. job title information. It contains information on over 900 standardized and occupation-specific descriptors over six domains, including: 1) worker characteristics, 2) worker requirements, 3) experience requirements, 4) occupational requirements, 5) workforce characteristics, and 6) occupation-specific information. The O*NET labels each job with Standard Occupational Classification (SOC) codes, which is further classified into green collar occupations based on the activities and technologies of the job requirements (e.g., whether or not they provide green services or produce green goods).⁽²⁰⁾ If a job has at least one “green” task (e.g., whether it provides green services or produces green goods), then it is categorized as a “green collar” job.

NHIS and O*NET Linkage—Linking the publicly 2010 NHIS Occupational Supplement data with the green collar classification in O*NET occurs through the Research Data Center (RDC) at the National Center for Health Statics. While the NHIS publishes publicly a 2-digit condensed occupational and industry code for each employed NHIS survey respondent, the

2-digit public code limits the linkage between the NHIS survey data and the rich job titles and characteristics available in O*NET. After a formal peer-reviewed and secure application process at the RDC, our research team accessed and linked the 4-digit occupational code variable (OCCUPN) in the NHIS (i.e., digits 3 4 5 6) and the 8-digit O*NET SOC code (i.e., 1 2 3 4 5 6 . 7 8). Based on the digit coding and linkage, we created a new variable, “Green Category,” to label NHIS survey respondents as either a Green Collar or non-Green Collar worker. In the case when the O*NET SOC code had a seventh and eighth digit ending in .00, this was considered an exact match with the NHIS data and labeled as green or non-green. However, when the seventh and eighth digit had an extension beyond .00 (such as .01, .02, etc.), we further investigated if each of these detailed occupations were all green, all non-green, or “mixed-green” collar workers. For example, if an O*NET broad occupational group had three different extensions of the seventh and eighth digit codes (e.g., .01, .02, and .03) of which two were classified as green and one was classified as non-green, then the NHIS occupational code was labeled as mixed-green to indicate that the parent job title had mixed jobs. The mixed-green collar workers (n=1,005; 6.8%) were excluded for this analysis.

Dependent Variables

We examined four main outcome variables, reflecting self-reported chemical and physical agent occupational exposures: 1) vapors, gas, dust, or fumes; 2) secondhand tobacco smoke; 3) skin hazards; and 4) outdoor work. Vapors, gas, dust, or fume exposure was measured by the question: “Please tell me if you are/were regularly exposed to vapors, gas, dust or fumes at work twice a week or more?” Secondhand smoke exposure was measured by the question: “During the past 12 months, were you regularly exposed to tobacco smoked from other people at work twice a week or more?” Skin hazard was measured by the question: “During the past 12 months, did you regularly handle or were you in skin contact with chemical products or substances at work twice a week or more?” Outdoor work was measured by asking the respondent, “During the past 12 months, did you regularly work outdoors twice a week or more?” Each response was dichotomized (yes/no) by the NHIS.

Independent Variables

The main independent variable was green collar worker status (“green collar” or “non-green collar”). Self-reported socio-demographic, health characteristics, and job characteristic variables of the NHIS respondent were also included as predictors, including: gender, race (white, black, or other), age, ethnicity (Hispanic or non-Hispanic), educational attainment (greater than high school, high school or GED, or less than high school), health insurance status (insured or uninsured), geographic region (Northeast, Midwest, South, West), body mass index (underweight, normal weight, overweight, obese), number of employees at the place of work (1–9, 10–24, 25–49, 50–99, 100–249, and 250+ employees), employment type (private, government employee, self-employed), and whether the workers had more than one job (yes/no).

Statistical Analyses

Multivariable logistic regression analyses were performed for each of the 4 outcome variables to calculate unadjusted odds ratios (UOR), adjusted odds ratios (AOR), and 95 %

confidence intervals (95% CI). Exposure status regression analyses were adjusted for smoking status, gender, age, ethnicity, education level, health insurance status, geographic region, body mass index, size of company, type of employment, number of jobs. The university institutional review board approved the study protocol. The NHIS data collection involves a complex, multistage design with additional elements of oversampling, clustering, and stratification hence statistical analyses were conducted with SUDAAN 11 (Research Triangle Institute, Research Triangle Park, NC) to account for complex design of NHIS.

RESULTS

Descriptive Information

The socio-demographic and work-related characteristics of the total workforce, including green and non-green collar workers are in Table 1. There was a total of 14,805 workers in the study period of which 2,588 classified as green collar (19%; US population estimate 24,614,939) and 12,217 non-green collar (81%; US population estimate 106,628,031).

Green collar workers were more likely to be male (76% vs. 48% Non-green workers), white race (84% vs. 82%), and be classified as overweight (40% vs. 35%). Fewer green collar workers reported functional limitations (13% vs. 15%), or visual (6% vs. 7%) impairments compared to non-green collar workers. The typical green collar worker was employed in a private company (84% vs. 72%), and worked primarily in that one green collar job as opposed to having a second job (94% vs. 91%).

The prevalence of chemical and physical exposures of green and non-green collar workers is displayed in Table 2. The prevalence of vapors, gas, dust, or fume exposure (32% vs. 23%), secondhand smoke exposure (17% vs. 14%), and working outdoors (34% vs. 22%) was higher in green collar workers than non-green collar workers. In contrast, the prevalence of chemical exposures was reportedly lower in green collar workers than that of non-green collar workers (19% vs. 21%).

Logistic Regression Analyses

In the univariate logistic regression analyses, green collar workers were significantly more likely to be exposed to vapors, gas, dust, or fumes exposure (Unadjusted Odds Ratio, UOR=1.65; 95% CI=1.47–1.85), secondhand smoke exposure (1.27; 95% CI=1.11–1.47), and work outdoors (1.82; 95% CI=1.62–2.04). Green collar workers were less likely to be exposed to chemicals (0.89; 95% CI: 0.78–1.03), although this finding was not statistically significant.

The multivariable logistic regression analyses for chemical and physical exposures of green and non-green collar workers are shown in Table 3. In the multivariate logistic regression, green collar workers were significantly more likely to be exposed to vapors, gas, dust, or fume exposure and working outdoors relative to non-green collar workers (Adjusted Odds Ratio, AOR=1.25; 95% CI=1.11–1.40 and 1.44; 95% CI=1.26–1.63, respectively). However, green collar works were less likely to be exposed to chemicals and skin hazards (0.80; 95% CI=0.69–0.92). There was not a statistically significant difference in secondhand smoke exposure between green and non-green workers (1.06; 95% CI=0.90–1.24).

DISCUSSION

To our knowledge, this is the first study to describe the potential chemical and physical occupational exposures within the emerging U.S. green collar workforce using the uniquely linked 2010 NHIS and O*NET data. Studies have shown that occupational chemical and physical exposures can lead to adverse health outcomes. Our data show that there is a significant difference in self-reported chemical and physical exposures between green and non-green workers. Green collar workers have a greater prevalence in vapors, gas, dust, or fume exposure, secondhand smoke exposure, and outdoor work exposure, whereas non-green workers report a higher prevalence of chemical exposure.

Contrary to our hypothesis, green collar workers have higher rates of vapors, gas, dust, or fume exposures compared to non-green workers. Occupational vapor exposure can lead to adverse health outcomes such as idiopathic pulmonary fibrosis, asthma, and other respiratory symptoms.^(8, 21) These results are unexpected given the green industry's general image to focus on work with non-hazardous materials.⁽²²⁾ Further research is necessary to characterize the vapor exposures in order to determine potential health and wellbeing effects and to develop intervention strategies to reduce workplace exposures.

Although not significant in our multivariable models (1.06; 0.90–1.24), green collar workers may have higher rates of secondhand tobacco smoke exposure compared to non-green workers (32% vs. 24%). Secondhand smoke exposure has been linked to lung cancer, cardiovascular disease, and cerebrovascular disease.^(23–25) Studies have used secondhand smoke serum markers such as cotinine to further quantify the degree of workplace secondhand smoke exposure.⁽²⁶⁾ While the result is unexpected, it is important to note that having a green industry is not necessarily synonymous with a “green” or healthy workplace environment. This may help to explain the discrepancy between the expected hypothesis and the results.

As we hypothesized, occupational outdoor exposure rates were higher among green collar workers when compared to non-green collar workers (1.44; 95% CI=1.26–1.63) after adjusting for other sociodemographic and work characteristics. Among all exposures, the greatest difference between green and non-green workers was the number of workers reporting outdoor work exposure. Increased outdoor exposure may be attributable to the environmentally-friendly and eco-friendly services provided by the green industry such as installation of solar panels or wind turbines. Outdoor work exposure has been linked to increased rates of skin cancer.^(14, 27) Variables such as duration of outdoor exposure and use of sun protection are needed to further characterize outdoor work exposure in the green and non-green workforces.

In contrast to the vapors, gas, dust, or fumes, secondhand smoke, and outdoor exposure rates, non-green collar workers had higher rates of chemical exposure compared to green collar workers. The lower rates of chemical exposure among green collar workers may relate to the workforce's commitment to sustainability and eco-friendliness- non-hazardous, ecofriendly alternatives to chemicals may replace chemicals typically used in non-green occupations. Studies have shown that green collar workers have lower rates of occupational

dermatologic disease, which may account for the lower rates of chemical exposure for green collar workers seen in this study.⁽⁵⁾ However, since it cannot be assumed that “green” chemicals are necessarily less toxic to humans, future studies should better identify the chemical exposures such as the types of chemical, duration of exposure, and frequency of exposure.^(28, 29)

This study is not without limitations. The use of cross-sectional and self-reported data in the NHIS may limit the classification of the exposure or job characteristic given that quantitative or validated individual assessments of the occupational exposures or workplace job tasks/activities are not available in the NHIS.²⁰ The estimates generated by self-reporting survey data can be biased due to the varying subjective assessments performed by the participants. The reasons include low self-confidence, self-biasing and memory recall. Nonetheless self-reported data provides a relatively inexpensive and rapid approach to collect exposure data. The O*NET exposure data are ecological, suggesting that interpretations of these data may be prone to the ecological fallacy.

In addition, we may be over-estimating the true prevalence of green collar workers employed in the U.S. workforce using the NHIS data. While the U.S. economy and workforce expands and contracts throughout the study period, there may be variations in the number of individuals truly employed in green collar occupations. For example, the construction industry has seen an increase in the number of workers and projects in recent years following the global financial crisis; green and sustainable building maybe increasing the number of green collar-related construction jobs. Nonetheless, the BLS used a different sampling frame (i.e. business and government establishments within 325 industries). Furthermore, the BLS measurement of “Green Goods and Services” was different (i.e., consisted of the percentage of the establishment’s revenue related to sale of green goods and services), while the O*NET uses a different mechanism to categorize green and non-green occupations, not industries.

To highlight this importance, we undertook a post-hoc analysis of our univariate logistic regression results, varying the prevalence of green workers in our 2 by 2 exposure-outcome tables to examine the potential influence of misclassification bias. We varied the prevalence from 18% of green collar workers, obtained in our study, down to 6.5%, the prevalence rate noted in the BLS report, and examined the impact this had on odds ratio estimates. Results for two of the outcome measures (vapors, gas, dust, or fumes; and outdoor work) had Odds Ratio estimates similar to the one listed in Table 2 (1.71 to 1.75 and 2.00 to 2.11, respectively). However, odds ratio estimates for secondhand smoke and chemical exposures were variable across the range of green collar prevalence estimates (0.52 to 1.33 and 0.12 to 1.39), respectively. These post-hoc findings highlight the importance of developing a uniform definition of green collar work that can be used both for estimating the size of the workforce and to conduct surveillance on this growing workforce.

Despite these limitations, this preliminary analysis estimating the chemical and physical exposures of green collar workers has several strengths including the large and nationally representative sample of NHIS adult participants, with a snapshot of all U.S. civilian workers. Using uniquely linked and publicly available (NHIS and O*NET data), for the first

time provided a classification scheme of green and non-green collar occupations of the U.S. workforce. Lastly, the time period that NHIS Occupational Health Supplement assessed novel self-reported measures on specific workplace physical and chemical exposures not available in other state- or national level U.S. surveillance systems.

This study documents preliminary findings that the emerging green collar workforce self-reported significantly different chemical and physical occupational exposure rates compared to non-green workers. Green collar workers showed higher prevalence of vapor, gas, dust, or fume exposure, secondhand smoke exposure, and outdoor work exposure compared to non-green collar workers. Previous studies have shown these occupational exposures can be detrimental to worker health. Worker health is directly tied to worker productivity, and these chemical and physical exposures may pose a threat to green collar worker health.⁽³⁰⁾ While exposure rates differ between green and non-green workers, variables such as exposure duration, frequency, and chemical composition are needed to better understand the differences in occupational exposure rates between the two workforces.

Acknowledgments

Funding Support: This work was supported in part by the National Institute for Occupational Safety and Health grant [R03-OH010124]; and by the National Institute for Health Research Health Protection Research Unit (NIHR HPRU) in Environmental Change and Health at the London School of Hygiene and Tropical Medicine in partnership with Public Health England (PHE), and in collaboration with the University of Exeter, University College London, and the Met Office.

References

1. Cleary, J., Kopicki, A. Preparing the Workforce for a “Green Jobs” Economy. Rutgers, NJ: John J Heldrich Center for Workforce Development; 2009.
2. Pinderhughes, R. Race, Poverty & the Environment. 2006. Green collar jobs.
3. Erich, CD., Norton, JJ., Gregory, CM., Rivkin, D., Lewis, P. Greening of the World of Work: Revisiting Occupational Consequences. 2011.
4. Bezdek, RH. Renewable energy and energy efficiency: Economic drivers for the 21st century. American Solar Energy Society; 2007.
5. Moore KJ, Chen C, Lee DJ, LeBlanc WG, Fleming LE, Caban-Martinez AJ. Occupational Skin Conditions in the Emerging US Green Collar Workforce. *Dermatitis*. 2016; 27:155–157. [PubMed: 27172317]
6. Bezdek, RH. American Solar Energy Society. 2009. Green Collar Jobs in the US and Colorado; p. 21
7. Bureau of Labor Statistics. The Economics Daily. 2010. Employment in green goods and services.
8. LeVan TD, Koh W-P, Lee H-P, Koh D, Mimi CY, London SJ. Vapor, Dust, and Smoke Exposure in Relation to Adult-Onset Asthma and Chronic Respiratory Symptoms The Singapore Chinese Health Study. *American journal of epidemiology*. 2006; 163:1118–1128. [PubMed: 16707657]
9. Torén K, Blanc PD. Asthma caused by occupational exposures is common—a systematic analysis of estimates of the population-attributable fraction. *BMC pulmonary medicine*. 2009; 9:7. [PubMed: 19178702]
10. Bergdahl I, Torén K, Eriksson K, et al. Increased mortality in COPD among construction workers exposed to inorganic dust. *The European respiratory journal*. 2004; 23:402. [PubMed: 15065829]
11. Ekenga CC, Parks CG, Sandler DP. Chemical exposures in the workplace and breast cancer risk: A prospective cohort study. *International Journal of Cancer*. 2015
12. Lee SJ, Nam B, Harrison R, Hong O. Acute symptoms associated with chemical exposures and safe work practices among hospital and campus cleaning workers: a pilot study. *American journal of industrial medicine*. 2014; 57:1216–1226. [PubMed: 25223949]

13. Boffetta P, Jourenkova N, Gustavsson P. Cancer risk from occupational and environmental exposure to polycyclic aromatic hydrocarbons. *Cancer Causes & Control*. 1997; 8:444–472. [PubMed: 9498904]
14. Radespiel-Tröger M, Meyer M, Pfahlberg A, Lausen B, Uter W, Gefeller O. Outdoor work and skin cancer incidence: a registry-based study in Bavaria. *International archives of occupational and environmental health*. 2009; 82:357–363. [PubMed: 18649084]
15. Blanc PD, Eisner MD, Balmes JR, Trupin L, Yelin EH, Katz PP. Exposure to vapors, gas, dust, or fumes: assessment by a single survey item compared to a detailed exposure battery and a job exposure matrix. *American journal of industrial medicine*. 2005; 48:110–117. [PubMed: 16032739]
16. Shopland DR, Gerlach KK, Burns DM, Hartman AM, Gibson JT. State-specific trends in smoke-free workplace policy coverage: the current population survey tobacco use supplement, 1993 to 1999. *Journal of Occupational and Environmental Medicine*. 2001; 43:680–686. [PubMed: 11515250]
17. US Department of Health Human Services. The health consequences of involuntary exposure to tobacco smoke: a report of the Surgeon General. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2006. p. 709
18. Arheart KL, Lee DJ, Dietz NA, et al. Declining trends in serum cotinine levels in US worker groups: the power of policy. *Journal of Occupational and Environmental Medicine*. 2008; 50:57–63. [PubMed: 18188082]
19. National Center for Health Statistics. Data File Documentation, National Health Interview Survey, 2010 (machine readable data file and documentation). Hyattsville, Maryland: National Center for Health Statistics, Centers for Disease Control and Prevention; 2011.
20. US Department of Labor. Employment and Training Administration. 2010.
21. Iwai K, Mori T, Yamada N, Yamaguchi M, Hosoda Y. Idiopathic pulmonary fibrosis. Epidemiologic approaches to occupational exposure. *American journal of respiratory and critical care medicine*. 1994; 150:670–675. [PubMed: 8087336]
22. Centers for Disease Control and Prevention. Summary of the Making Green Jobs Safe Workshop. Washington, DC: 2009.
23. Brennan P, Buffler PA, Reynolds P, et al. Secondhand smoke exposure in adulthood and risk of lung cancer among never smokers: a pooled analysis of two large studies. *International Journal of cancer*. 2004; 109:125–131. [PubMed: 14735478]
24. Venn A, Britton J. Exposure to secondhand smoke and biomarkers of cardiovascular disease risk in never-smoking adults. *Circulation*. 2007; 115:990–995. [PubMed: 17296856]
25. Bonita R, Duncan J, Truelsen T, Jackson RT, Beaglehole R. Passive smoking as well as active smoking increases the risk of acute stroke. *Tobacco Control*. 1999; 8:156–160. [PubMed: 10478399]
26. Wortley PM, Caraballo RS, Pederson LL, Pechacek TF. Exposure to secondhand smoke in the workplace: serum cotinine by occupation. *Journal of Occupational and Environmental Medicine*. 2002; 44:503–509. [PubMed: 12085475]
27. Health UDo, Services H. The health consequences of involuntary exposure to tobacco smoke: a report of the Surgeon General. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2006. p. 709
28. Calvert GM, Luckhaupt SE, Sussell A, Dahlhamer JM, Ward BW. The prevalence of selected potentially hazardous workplace exposures in the US: Findings from the 2010 National Health Interview Survey. *American journal of industrial medicine*. 2013; 56:635–646. [PubMed: 22821700]
29. Schulte PA, McKernan LT, Heidel DS, et al. Occupational safety and health, green chemistry, and sustainability: a review of areas of convergence. *Environmental Health*. 2013; 12:1. [PubMed: 23286340]

30. Burton WN, Conti DJ, Chen C-Y, Schultz AB, Edington DW. The role of health risk factors and disease on worker productivity. *Journal of Occupational and Environmental Medicine*. 1999; 41:863–877. [PubMed: 10529942]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1 Socio-demographics and work characteristics of green and non-green collar workers: The National Health Interview Survey, 2010 Occupational Supplement and 2010 O*NET Linkage⁷

Characteristics	Total Worker Population			Green Collar			Non-Green Collar		
	US Estimated Population	N	Percent ² [95% CI]	US Estimated Population	n	Percent ² [95% CI]	US Estimated Population	n	Percent ² [95% CI]
Total	131,296,970	14,805	100.0	24,614,939	2,588	18.7 (18.0–19.4)	106,682,031	12,217	81.2 (80.5–81.9)
Gender									
Male	69,814,890	7,306	53.1 (52.1–54.1)	18,777,269	1,900	76.3 (74.5–78.0)	51,037,621	5,406	47.8 (46.7–48.9)
Female	61,482,080	7,499	46.8 (45.8–47.8)	5,837,670	688	23.7 (21.9–25.4)	55,644,410	6,811	52.1 (51–53.2)
Race									
White	107,663,727	11,224	82 (81–82.9)	20,600,991	2,018	83.7 (81.8–85.4)	87,062,736	9,206	81.6 (80.7–82.5)
Black	15,246,774	2,349	11.6 (10.8–12.3)	2,621,791	365	10.6 (9.1–12.1)	12,624,983	1,984	11.8 (11–12.5)
Other	8,386,469	1,232	6.3 (5.8–6.9)	1,392,157	205	5.7 (4.6–6.6)	6,994,312	1,027	6.6 (5.9–7.1)
Age Group									
18–24	16,733,678	1,559	12.7 (11.9–13.5)	2,260,000	210	9.2 (7.7–10.6)	14,473,678	1,349	13.6 (12.6–14.4)
25–64	109,210,100	12,552	83.1 (82.3–84)	21,453,941	2,279	87.2 (85.4–88.8)	87,756,159	10,273	82.3 (81.3–83.2)
65+	5,353,192	694	4.0 (3.7–4.4)	900,998	99	3.7 (2.8–4.4)	4,452,194	595	4.1 (3.7–4.5)
Ethnicity									
Non-Hispanic	112,465,426	11,865	85.6 (84.8–86.4)	21,076,884	2,110	85.6 (84.1–87.1)	91,388,542	9,755	85.7 (84.8–86.4)
Hispanic	18,831,544	2,940	14.3 (13.5–15.1)	3,538,055	478	14.4 (12.8–15.8)	15,293,489	2,462	14.3 (13.5–15.1)
Educational Level									
>HS	85,961,694	9,479	65.6 (64.5–66.7)	14,887,531	1,537	60.8 (58.2–63.2)	71,074,163	7,942	66.8 (65.6–67.9)
HS	32,367,120	3,574	24.7 (23.8–25.6)	7,176,915	731	29.3 (27–31.5)	25,190,205	2,843	23.7 (22.7–24.6)
<HS	12,585,494	1,713	9.6 (9–10.2)	2,440,156	310	10.0 (8.7–11.1)	10,145,338	1,403	9.5 (8.8–10.2)
Health Insurance									
Not Insured	22,946,866	2,899	17.5 (16.6–18.4)	3,569,671	437	14.5 (12.9–16.2)	19,377,195	2,462	18.2 (17.2–19.2)
Insured	107,795,448	11,863	82.4 (81.5–83.3)	20,934,477	2,142	85.4 (83.7–87)	86,860,971	9,721	81.7 (80.7–82.7)
Regional Location									
Northeast	23,415,050	2,326	17.8 (16.7–18.9)	4,458,522	407	18.1 (16.1–20.1)	18,956,528	1,919	17.7 (16.6–18.9)
Midwest	31,050,486	3,295	23.6 (22.4–24.8)	5,971,088	608	24.2 (22.2–26.2)	25,079,398	2,687	23.5 (22.2–24.7)
South	46,202,248	5,456	35.1 (33.8–36.5)	8,365,611	923	33.9 (31.8–36.1)	37,836,637	4,533	35.4 (34–36.9)
West	30,629,186	3,728	23.3 (22.1–24.5)	5,819,718	650	23.6 (21.6–25.6)	24,809,468	3,078	23.2 (21.9–24.5)
Use Special Equipment									

Characteristics	Total Worker Population			Green Collar			Non-Green Collar		
	US Estimated Population	N	Percent ² [95% CI]	US Estimated Population	<i>J</i> ⁿ	Percent ² [95% CI]	US Estimated Population	<i>J</i> ⁿ	Percent ² [95% CI]
Any Functional Limitations									
Yes	2,262,315	247	1.7 (1.4–1.9)	474,947	45	1.9 (1.3–2.5)	1,787,368	202	1.6 (1.3–1.9)
No	128,929,436	14,543	98.2 (98–98.5)	24,128,685	2,542	98 (97.4–98.6)	104,800,751	12,001	98.3 (98–98.6)
Any Functional Limitations									
Yes	19,175,181	2,188	14.6 (13.9–15.2)	3,262,315	332	13.2 (11.7–14.7)	15,912,866	1,856	14.9 (14.1–15.6)
No	112,121,789	12,617	85.3 (84.7–86)	21,352,624	2,256	86.7 (85.2–88.2)	90,769,165	10,361	85 (84.3–85.8)
Hearing Impairment									
Yes	15,220,235	1,585	11.5 (10.9–12.2)	3,329,911	324	13.5 (11.9–15.1)	11,890,324	1,261	11.1 (10.4–11.8)
No	116,055,331	13,216	88.4 (87.7–89)	21,285,028	2,264	86.4 (84.8–88)	94,770,303	10,952	88.8 (88.1–89.5)
Visual Impairment									
Yes	8,700,869	1,030	6.6 (6.1–7.1)	1,529,915	163	6.2 (5.1–7.3)	7,170,954	867	6.7 (6.1–7.2)
No	122,588,333	13,773	93.3 (92.8–93.8)	23,085,024	2,425	93.7 (92.6–94.8)	99,503,309	11,348	93.2 (92.7–93.8)
Body Mass Index									
Underweight	1,747,585	205	1.3 (1.1–1.6)	184,376	21	0.7 (0.3–1.1)	1,563,209	184	1.5 (1.2–1.7)
Normal Weight	44,199,496	4,969	34.8 (33.8–35.8)	6,925,364	733	28.7 (26.6–30.8)	37,274,132	4,236	36.2 (35–37.3)
Overweight	45,907,356	5,200	36.1 (35.1–37.1)	9,630,198	1,026	39.9 (37.7–42.2)	36,277,158	4,174	35.2 (34.1–36.3)
Obese	35,129,657	3,923	27.6 (26.7–28.5)	7,335,945	739	30.4 (28.2–32.6)	27,793,712	3,184	27 (26–28)
Employees at Work									
1–9 employees	32,784,032	3,735	25.8 (24.9–26.7)	5,311,894	570	22 (20–24)	27,472,138	3,165	26.7 (25.7–27.7)
10–24 employees	18,550,259	2,064	14.6 (13.8–15.4)	3,324,045	348	13.8 (12.1–15.5)	15,226,214	1,716	14.8 (14–15.6)
25–49 employees	14,989,208	1,631	11.8 (11.1–12.5)	2,956,846	296	12.2 (10.5–14)	12,032,362	1,335	11.7 (10.9–12.4)
50–99 employees	13,537,794	1,505	10.6 (10–11.3)	2,381,265	257	9.8 (8.5–11.2)	11,156,529	1,248	10.8 (10.1–11.6)
100–249 employees	15,888,297	1,812	12.5 (11.8–13.1)	3,595,234	377	14.9 (13.3–16.5)	12,293,063	1,435	11.9 (11.2–12.6)
250+ employees	31,017,765	3,528	24.4 (23.5–25.4)	6,502,470	686	27 (25–28.9)	24,515,295	2,842	23.8 (22.8–24.9)
Employment Type									
Private	97,249,605	10,977	74.4 (73.5–75.3)	20,623,010	2,157	83.8 (82.1–85.6)	76,626,595	8,820	72.2 (71.2–73.2)
Government	21,721,619	2,466	16.6 (15.8–17.4)	2,088,164	229	8.4 (7.1–9.8)	19,633,455	2,237	18.5 (17.6–19.3)
Self-Employed	11,684,404	1,285	8.9 (8.3–9.5)	1,870,619	196	7.6 (6.4–8.7)	9,813,785	1,089	9.2 (8.5–9.9)
Incorporated Corp.									
Yes	3,203,863	330	27.5 (24.6–30.4)	720,097	67	38.4 (31.2–45.7)	2,483,766	263	25.4 (22.4–28.4)
No	8,411,963	947	72.4 (69.5–75.3)	1,150,522	129	61.5 (54.2–68.7)	7,261,441	818	74.5 (71.5–77.5)
More than one Job									

Characteristics	Total Worker Population			Green Collar			Non-Green Collar		
	US Estimated Population	N ¹	Percent ² [95% CI]	US Estimated Population	I ¹	Percent ² [95% CI]	US Estimated Population	I ¹	Percent ² [95% CI]
Yes	11,312,689	1,282	8.6 (8–9.2)	1,443,380	168	5.8 (4.8–6.8)	9,869,309	1,114	9.2 (8.6–9.9)
No	119,775,083	13,501	91.3 (90.7–91.9)	23,162,291	2,418	94.1 (93.1–95.1)	96,612,792	11,083	90.7 (90–91.3)
Smoker									
Current	2,783,095	2,818	19.2 (18.4–20.0)	557,846	540	20.5 (18.6–22.4)	2,225,250	2,278	18.9 (18.0–19.8)
Former	2,790,580	2,705	19.2 (18.5–20.0)	594,847	535	21.9 (20.0–23.7)	2,195,733	2,170	18.6 (17.8–19.5)
Never	8,925,572	9,190	61.6 (60.6–62.5)	1,568,056	1,495	57.6 (55.5–59.8)	7,357,516	7,695	62.5 (61.4–63.5)

¹The estimates from this table are based on questions from the National Health Interview Survey; Mixed workers have been excluded

* Green jobs are characterized as having at least one “green” task

¹ Sample size from the National Health Interview Survey for the year 2010;

² Percent (prevalence) estimated from the National Health Interview Survey for the year 2010

Table 2

Prevalence of occupational exposures¹ among green and non-green collar workers: The National Health Interview Survey, 2010 Occupational Supplement and 2010 O*NET Linkage

Characteristics	Total Worker Population			Green Collar			Non-Green Collar		
	US Estimated Population	N ¹	Percent ² [95% CI]	US Estimated Population	n ¹	Percent ² [95% CI]	US Estimated Population	n ¹	Percent ² [95% CI]
Vapor Exposure	31,989,790	3,536	24.4 (23.6–25.3)	7,975,809	846	32.4 (30.4–34.5)	24,013,981	2,690	22.6 (21.6–23.5)
Secondhand Smoke Exposure	18,834,642	2,102	14.3 (13.5–15.1)	4,172,504	446	16.9 (15.2–18.6)	14,662,138	1,656	13.7 (12.9–14.6)
Chemical Exposure	26,683,550	2,876	20.3 (19.5–21.1)	4,599,107	468	18.7 (16.9–20.4)	22,084,443	2,408	20.7 (19.8–21.6)
Work Outdoors	31,396,888	3,419	23.9 (23.0–24.8)	8,268,994	885	33.6 (31.4–35.7)	23,127,894	2,534	21.7 (20.7–22.6)

¹The estimates from this table are based on questions from the National Health Interview Survey that asked respondents "Please tell me if you [fill 2: are/were] regularly exposed to vapors, gas, dust, or fumes at work twice a week or more?"

²Sample size from the National Health Interview Survey for the year 2010;

³Percent (prevalence) estimated from the National Health Interview Survey for the year 2010

Table 3

Multivariable logistic regression analyses predicting occupational exposures in green vs. non-green collar workers: The National Health Interview Survey, 2010 Occupational Supplement and 2010 O*NET Linkage

Independent Variable	Vapor Exposure (n=13,678)	Secondhand Smoke Exposure (n=13,688)	Chemical Exposure (n=13,686)	Work Outdoors (n=13,691)
	AOR (95% CI)	AOR (95% CI)	AOR (95% CI)	AOR (95% CI)
Green Collar (Ref = Non-Green)				
Green	1.25 (1.11–1.40)*	1.06 (0.90–1.24)	0.80 (0.69–0.92)*	1.44 (1.26–1.63)*
Smoker (Ref = Never)				
Current	1.78 (1.57–2.03)*	5.03 (4.39–5.77)*	1.54 (1.34–1.76)*	1.39 (1.22–1.59)*
Former	1.51 (1.34–1.72)*	1.39 (1.18–1.65)*	1.22 (1.05–1.41)*	1.13 (1.00–1.28)*
Gender (Ref = Female)				
Male	2.38 (2.15–2.64)*	1.97 (1.74–2.23)*	1.37 (1.24–1.51)*	4.66 (4.14–5.25)*
Race (Ref = White)				
Black	1.08 (0.92–1.26)	1.41 (1.19–1.67)*	0.79 (0.67–0.93)*	1.03 (0.86–1.24)
Other	0.63 (0.52–0.78)*	0.94 (0.72–1.22)	0.68 (0.53–0.86)*	0.45 (0.36–0.58)*
Age (Ref = 18–24)				
25–64	1.15 (0.96–1.39)	0.58 (0.47–0.71)*	0.80 (0.67–0.96)*	1.10 (0.91–1.33)
65+	0.69 (0.51–0.94)*	0.31 (0.20–0.46)*	0.35 (0.24–0.50)*	0.97 (0.72–1.30)
Ethnicity (Ref = Non-Hispanic)				
Hispanic	0.9 (0.76–1.07)	0.87 (0.71–1.07)	0.89 (0.76–1.04)	0.99 (0.85–1.17)
Education (Ref = HS+)				
High School/GED	1.98 (1.76–2.23)*	1.49 (1.28–1.72)*	1.52 (1.34–1.73)*	1.83 (1.60–2.09)*
Less than High School	2.12 (1.79–2.52)*	1.43 (1.17–1.76)*	1.46 (1.21–1.76)*	2.21 (1.85–2.64)*
Health Insurance Status (Ref = No)				
Yes	0.92 (0.80–1.05)	0.68 (0.58–0.8)*	0.75 (0.66–0.85)*	0.85 (0.74–0.98)*
Geographic Region (Ref = Northeast)				
Midwest	1.20 (1.03–1.40)*	1.18 (0.96–1.45)	1.27 (1.09–1.49)*	1.02 (0.84–1.23)
South	1.14 (1.00–1.32)*	1.47 (1.21–1.78)*	1.12 (0.96–1.30)	1.29 (1.08–1.53)*
West	1.16 (0.98–1.38)	1.21 (0.96–1.51)	1.04 (0.88–1.24)	1.42 (1.19–1.69)*

Independent Variable	Vapor Exposure (n=13,678)	Secondhand Smoke Exposure (n=13,688)	Chemical Exposure (n=13,686)	Work Outdoors (n=13,691)
	AOR (95% CI)	AOR (95% CI)	AOR (95% CI)	AOR (95% CI)
Body Mass Index (Ref = Normal)				
Underweight	0.92 (0.56–1.50)	1.09 (0.65–1.81)	1.26 (0.79–2.00)	0.79 (0.47–1.32)
Overweight	1.17 (1.02–1.33) *	1.20 (1.02–1.42) *	1.11 (0.97–1.27)	1.08 (0.95–1.23)
Obese	1.31 (1.14–1.50) *	1.37 (1.17–1.60) *	1.22 (1.05–1.42) *	1.16 (1.01–1.34) *
Size of Company (Ref = 250+ employees)				
1–9 employees	1.09 (0.94–1.27)	0.89 (0.73–1.09)	1.14 (0.97–1.34)	2.51 (2.11–2.99) *
10–24 employees	0.99 (0.83–1.17)	1.08 (0.89–1.30)	1.09 (0.91–1.30)	2.03 (1.70–2.42) *
25–49 employees	1.12 (0.94–1.34)	1.03 (0.81–1.30)	1.16 (0.97–1.38)	1.85 (1.53–2.24) *
50–99 employees	0.97 (0.79–1.18)	1.17 (0.94–1.44)	0.97 (0.79–1.18)	1.44 (1.16–1.78) *
100–249 employees	1.00 (0.84–1.19)	0.95 (0.76–1.18)	0.83 (0.69–1.01)	1.32 (1.10–1.60) *
Type of Employment (Ref = Government employee)				
Private Employee	1.00 (0.87–1.14)	1.13 (0.93–1.38)	1.15 (0.97–1.35)	0.46 (0.40–0.54)
Self-Employed	1.22 (0.98–1.53)	0.97 (0.70–1.34)	1.39 (1.10–1.76) *	0.80 (0.63–1.02)
More than one job (Ref = No)				
Yes	1.10 (0.91–1.34)	1.20 (0.97–1.49)	1.31 (1.08–1.59)	1.31 (1.10–1.56) *

* Indicates statistically significant ($p < 0.05$)