

ORIGINAL ARTICLE

Open Access



# Antioxidant responses and photosynthetic behaviors of *Kappaphycus alvarezii* and *Kappaphycus striatum* (Rhodophyta, Solieriaceae) during low temperature stress

Hu Li<sup>1,2,3</sup>, Jianguo Liu<sup>1,2\*</sup>, Litao Zhang<sup>1,2</sup> and Tong Pang<sup>1,2</sup>

## Abstract

**Background:** *Kappaphycus* are farmed in tropical countries as raw material for carrageenan, which is widely used in food industry. The sea area available for farming is one limiting factor in the production of seaweeds. Though cultivation is spreading into subtropical regions, the lower seawater temperature is an important problem encountered in subtropical regions for the farming of *Kappaphycus*. This research of physiological response to low temperature stress will be helpful for screening *Kappaphycus* strains for growth in a lower temperature environment.

**Results:** Responses of antioxidant systems and photosystem II (PSII) behaviors in *Kappaphycus alvarezii* and *Kappaphycus striatum* were evaluated during low temperature treatments (23, 20, 17 °C). Compared with the controls at 26 °C, the H<sub>2</sub>O<sub>2</sub> concentrations increased in both species when the thalli were exposed to low temperatures (23, 20, 17 °C), but these increases were much greater in *K. striatum* than in *K. alvarezii* thalli, suggesting that *K. striatum* suffered more oxidative stress. The activities of some important antioxidant enzymes (e.g. superoxide dismutase and ascorbate peroxidase) and the hydroxyl free radical scavenging capacity were substantially higher at 23, 20 and 17 °C than at the control 26 °C in *K. alvarezii*, indicating that the antioxidant system of *K. alvarezii* enhanced its resistance to low temperature. However, no significant increases of antioxidant enzymes activities were observed at 20 and 17 °C in *K. striatum*. In addition, both the maximal efficiency of PSII photochemistry ( $F_v/F_m$ ) and the performance index ( $PI_{ABS}$ ) decreased significantly in *K. striatum* at 23 °C, indicating that the photosynthetic apparatus was damaged at 23 °C. In contrast, no significant decreases of either  $F_v/F_m$  or  $PI_{ABS}$  were observed in *K. alvarezii* at 23 °C.

**Conclusions:** It is concluded that *K. alvarezii* has greater tolerance to low temperature than *K. striatum*.

**Keywords:** Antioxidant systems, *Kappaphycus alvarezii*, *Kappaphycus striatum*, Low temperature stress, Photosystem II

## Background

*Kappaphycus alvarezii* and *Kappaphycus striatum* (Rhodophyta, Solieriaceae), two important carrageenophyte species, are used as the major commercial source of κ-carrageenan, which is widely utilized as a gelling and

stabilizing agent for some food products, including frozen desserts, chocolate-flavoured milk, cottage cheese dressings and soyamilk (Doty 1973; Glenn and Doty 1990; Bixler et al. 2001; Bindu and Levine 2011). The farming of *Kappaphycus* occurs mainly in tropical southeast Asian countries (Ohno et al. 1994), such as the Philippines and Indonesia, serving as a means of livelihood for locals and as a source of revenue for the economy of these countries (Bindu 2011; Bixler and Porse 2011; Ganzon-Fortes et al. 2012). However, although the carrageenan processing technology in

\*Correspondence: jgliu@qdio.ac.cn

<sup>1</sup> Key Laboratory of Experimental Marine Biology, National & Local Joint Engineering Laboratory of Ecological Mariculture, Institute of Oceanology, Chinese Academy of Sciences, 7 Nanhai Road, Qingdao 266071, China  
Full list of author information is available at the end of the article

this industry is mature (Bixler et al. 2001), the supply of carrageenan is not adequate for the global demand. The sea area available for farming is one limiting factor in the production of seaweeds for carrageenan extraction. Currently, cultivation is spreading into subtropical regions (i.e. lower temperature regions for these seaweeds). For example, cultivation during the summer season was introduced in Shikoku Island, Southern Japan (Ohno et al. 1994). The lower seawater temperature is an important problem encountered in subtropical regions, or other low-temperature sea areas, for the farming of *Kappaphycus*.

Photosynthetic processes are exceedingly sensitive to low temperature (Allen and Ort 2001). When plants are exposed to low-temperature stress, photosynthetic enzymes may be degraded and photo-damage may occur, which would lead to decreased photosynthetic activity. The reduced photosynthesis may result in the accumulation of excess energy, which leads to the generation of numerous reactive oxygen species (ROS), such as hydrogen peroxide ( $H_2O_2$ ) and hydroxyl free radical ( $\cdot OH$ ), which are harmful to the plant's photosynthetic apparatus. Collén et al. (1994) found that *Kappaphycus* and *Eucheuma* exposed to abiotic or biotic stresses produced  $H_2O_2$ . The accumulation of  $H_2O_2$  within plant tissues is thought to be a crucial event in initiating adequate antioxidant responses in plants (Gechev et al. 2003). In order to gain a better understanding of the antioxidant response of *Kappaphycus* under low temperature stress, we measured the activities of antioxidant enzymes [e.g. superoxide dismutase (SOD), catalase (CAT) and ascorbate peroxidase (APX)], hydroxyl free radical ( $\cdot OH$ ) scavenging capacity and the level of malondialdehyde (MDA), an important index of oxidative lesions (Barros et al. 2003), in *K. alvarezii* and *K. striatum* thalli under low temperature stress. Moreover, since PSII is the most sensitive part of the photosynthetic apparatus, changes of PSII state may indicate changes in physiological processes in plants (Lim et al. 2014). In recent years, the chlorophyll (Chl) a fluorescence transient (O-J-I-P) has become an important tool in photosynthesis research (Strasser et al. 2004; Strauss et al. 2007), and especially in studying PSII behaviors. The changes of the PSII antenna, the reaction centers, the oxidation–reduction of the plastoquinone (PQ) pool and the electron flow limitations on the acceptor side of PSII, can be reflected by the O-J-I-P transient modulation in higher plants and algae (Stirbet and Govindjee 2012). The effect of low temperature stress on PSII was further assessed in *K. alvarezii* and *K. striatum* thalli by means of the Chl a fluorescence transient.

## Methods

### Algal sample collection and pretreatment

*Kappaphycus alvarezii* (Rhodophyta, Solieriaceae) (reddish brown, Fig. 1a) and *Kappaphycus striatum*

(Rhodophyta, Solieriaceae) (green, Fig. 1b) were harvested in Li'an bay, Hainan Province, China (18°24'N, 110°03'E) (Liu et al. 2009; Pang et al. 2015). Selected healthy thalli of both species were cleaned to remove sludge, miscellaneous algal contaminants and other impurities from the surface of thalli. The algal thalli were acclimatized in a plastic tank (50 × 40 × 40 cm) with 40 L of natural seawater (about 23–24 °C, salinity 33 ‰) collected in Li'an bay (18°24'N, 110°03'E) and irradiated with light (55  $\mu mol photons m^{-2} s^{-1}$ ) in a 12:12 h light:dark (L:D) cycle. They were acclimatized to their new surroundings for a day in the tank before performing the following low temperatures treatments assay.

### Low temperature treatments

Portions of thalli approximately 3 mm in diameter and 1 cm long were cut from the apices of the collected algal thalli. The samples were taken from same parts of the thallus to avoid any difference between individuals. These were all placed in plastic tanks (50 × 40 × 40 cm) with 40 L of natural seawater (about 23–24 °C, salinity 33 ‰) collected in Li'an bay (18°24'N, 110°03'E) and irradiated with a fluorescent lamp at approximately 55  $\mu mol photons m^{-2} s^{-1}$  (Fig. 1c). Sea temperature was adjusted and maintained by an automatic temperature control unit (Guangdong Zhenhua Electrical Appliance Co., LTD). The control temperature was 26 °C, which is the optimum temperature for these two species (Lideman et al. 2013). The samples of two species were exposed to the control temperature (26 °C) and low temperatures 23, 20 and 17 °C for 2 h, respectively. The temperature treatments of *K. alvarezii* and *K. striatum* samples were shown in Fig. 1c.

### Crude extracts and antioxidant system parameters

After treatment at each temperature, algal samples of each species were immediately froze in liquid nitrogen, labeled and stored in a freezer at –80 °C for the following assay (Fariduddin et al. 2014).

The frozen samples (about 2 g) were ground in a mortar in liquid nitrogen and placed in 10-mL centrifuge tubes in ice. The algal powder was mixed with 3 mL of pre-cooled extraction buffer (50 mM phosphate buffer, pH 7.2, 0.1 mM EDTA) and homogenized using an S10 homogenizer (Ningbo Xinzhi Biotechnology Co., Ltd) in an ice-water bath (Nagarani and Kumaraguru 2012). Cell debris was removed by centrifugation at 12,000g for 10 min at 4 °C and the supernatant was used for the following measurements of antioxidant system parameters. Each temperature treatment had three repeats for each species.

The concentration of  $H_2O_2$  in the algal thalli was assessed using a commercially available kit—Hydrogen



**Fig. 1** The photos of *K. alvarezii* and *K. striatum*. **a** *Kappaphycus alvarezii* thalli; **b** *Kappaphycus striatum* thalli; **c** A photo of temperature treatments of *K. alvarezii* and *K. striatum* samples in the tank. Algal samples were irradiated at  $55 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ . The sea water temperature in the tank was adjusted and maintained by an automatic temperature-control bar

Peroxide Assay kit (Nanjing Jiancheng Bioengineering Institute).  $\text{H}_2\text{O}_2$  was bound with molybdenic acid to form a complex, which was measured spectrophotometrically at 405 nm and the concentration of  $\text{H}_2\text{O}_2$  was then calculated (Liu et al. 2013).

Total SOD activity in algal thalli was determined by using a Total Superoxide Dismutase Assay kit (Nanjing Jiancheng Bioengineering Institute) based on measuring the enzyme's ability to inhibit the photochemical reduction of tetrazolium blue. The developed blue color was measured spectrophotometrically at 550 nm (Das et al. 2000). CAT activity was quantified by the method of Malanga et al. (1999). The absorbance decay of  $\text{H}_2\text{O}_2$  was monitored by spectrophotometry at 240 nm, with a molar extinction coefficient of  $\epsilon = 39.4 \text{ mM}^{-1} \text{ cm}^{-1}$ . APX activity was measured as described by Nakano and Asada (1981). Ascorbate consumption was monitored by spectrophotometry at 290 nm, with a molar extinction coefficient of  $\epsilon = 2.8 \text{ mM}^{-1} \text{ cm}^{-1}$ .

Hydroxyl free radical scavenging capacity (HFRSC) was determined by using a commercially available kit—Hydroxyl Free Radical Assay kit (Nanjing Jiancheng Bioengineering Institute) based on the deoxyribose degradation assay described by Halliwell et al. (2006).

To evaluate the extent of oxidative damage in algal cells under low temperature stress, the concentration of MDA, a marker of lipoperoxidation, was measured in the algal thalli by the method of Fraga et al. (1988).

#### Protein determination

Enzyme activities were expressed relative to protein concentrations, which were estimated by the method of Deng et al. (2013) using bovine serum albumin as standard. We employed the dyeing method of Coomassie brilliant blue G-250 to determine the protein concentration of algal samples, measuring the absorbency at the wavelength of 595 nm. Each temperature treatment had three repeats for each species.

#### Chlorophyll (Chl) a fluorescence transient measurement

Fast Chla fluorescence transients were measured with a Plant Efficiency Analyzer (Handy-PEA, Hansatech Instruments Ltd, UK). Each transient obtained from the samples was analyzed according to the JIP-test (Strasser 1978; Strasser et al. 2004; Stirbet and Govindjee 2012; Zhang et al. 2012a). Each temperature treatment had 10 repeats for each species.

The following variables were calculated directly or indirectly.

1.  $\phi_{\text{P}_0}$  (or  $F_v/F_m$ ), the maximum PSII photochemical efficiency, namely the maximum quantum yield of primary photochemistry.  $F_v/F_m = \phi_{\text{P}_0} = \text{TR}_O/\text{ABS}$ , where  $\text{TR}_O$  and ABS denote the trapped and absorbed excitation energy fluxes, respectively.
2.  $\psi_o$ , the probability that a trapped exciton moved an electron into the electron transport chain beyond (plasto)quinone ( $\text{Q}_A^-$ ).  $\psi_o = \text{ET}_O/\text{TR}_O$ , where  $\text{ET}_O$

and  $TR_O$  denote the electron transport and trapped excitation energy fluxes, respectively.

- $\varphi_{Eo}$ , the probability that an absorbed photon moved an electron into the electron transport chain further than  $Q_A^-$ , namely quantum yield of electron transport.  $\varphi_{Eo} = ET_O/ABS$ .
- $PI_{ABS}$ , the performance index on an absorption basis.  $PI_{ABS} = (RC/ABS) \cdot [\varphi_{Po}/(1 - \varphi_{Po})] \cdot [\psi_o/(1 - \psi_o)]$ , where  $RC/ABS$ ,  $\varphi_{Po}/(1 - \varphi_{Po})$  and  $\psi_o/(1 - \psi_o)$  reflect the efficiency of the reaction centers, the light energy absorption efficiency and the accepting efficiency of the electron acceptor, respectively.

### Statistical analysis

Data are presented as mean  $\pm$  SD for at least three replicate measurements (the number (n) of replicates is presented in each figure caption). To determine the statistical significance between experimental groups, the data were examined using a *t* test ( $p < 0.05$ ).


## Results

### Changes in $H_2O_2$ concentrations in *Kappaphycus alvarezii* and *Kappaphycus striatum* thalli

The highest concentrations of  $H_2O_2$  in both *K. alvarezii* and *K. striatum* thalli were observed at 23 °C; they decreased at 20 and 17 °C. A  $H_2O_2$  concentration similar to the control (26 °C) was observed at 17 °C in *K. alvarezii* thalli but *K. striatum* thalli at 17 °C still had a higher  $H_2O_2$  concentration than the control at 26 °C (Fig. 2). In other words, compared with the controls at 26 °C, the  $H_2O_2$  concentration increased in both species when the thalli were exposed to low temperatures (23, 20, 17 °C), but the increase was much greater in *K. striatum* thalli than in *K. alvarezii* thalli. For example, the  $H_2O_2$  concentration at 23 °C was 1.9 times as much as that of the control in *K. alvarezii* thalli, while the  $H_2O_2$  concentration at 23 °C was 3 times as much as that of the control in *K. striatum* thalli. These results suggest that *K. striatum* thalli suffered more serious oxidative stress from  $H_2O_2$  than *K. alvarezii* thalli.

### Antioxidant system responses of *K. alvarezii* and *K. striatum* thalli


Low temperature treatments resulted in distinct responses from the three major antioxidant enzymes (SOD, APX and CAT) in both *K. alvarezii* and *K. striatum* thalli. Compared with the control, SOD activities and APX activities increased significantly in both species when algal thalli were exposed to 23 °C (SOD activities at 23 °C were 2.1 times and 2.8 times as much as that of the controls in *K. alvarezii* thalli and *K. striatum* thalli, respectively; APX activities at 23 °C were 5.5 times and 3.1 times as much as that of the controls in *K. alvarezii*



**Fig. 2** Concentrations of  $H_2O_2$  in thalli of *K. alvarezii* and *K. striatum* exposed to low temperatures (23, 20 and 17 °C) for 2 h, relative to the concentration of the control of each species at 26 °C [ $H_2O_2$  concentration in thalli at the control temperature (26 °C) was  $17.7 \pm 1.3$  mmol  $g^{-1}$  prot for *K. alvarezii* and  $29.2 \pm 2.0$  mmol  $g^{-1}$  prot for *K. striatum*]. Data are means  $\pm$  SD of three replicates ( $n = 3$ )

thalli and *K. striatum* thalli, respectively) (Fig. 3). At even lower temperatures (20 and 17 °C), SOD activities and APX activities of *K. alvarezii* thalli remained elevated (SOD activities were 1.8 times and 1.7 times as much as that of the control, at 20 and 17 °C, respectively; APX activities were 5.3 times and 5.9 times as much as that of the control, at 20 and 17 °C, respectively), whereas those of *K. striatum* thalli had no significant differences at either 20 or 17 °C compared with the control. Although CAT activities did not vary significantly in the two species after exposure to low temperatures (23, 20 and 17 °C) (Fig. 3), *K. alvarezii* thalli maintained higher SOD and APX activities whereas *K. striatum* thalli didn't, under the low temperatures (20 and 17 °C). In addition, compared with the control, hydroxyl free radical scavenging capacity (HFRSC) had significant increases at lowered temperatures (HFRSC were 2.9 times, 2.1 times and 2.1 times as much as the control, at 23, 20 and 17 °C, respectively) in *K. alvarezii* thalli. Hydroxyl free radical scavenging capacity (HFRSC) of *K. striatum* thalli peaked at 23 °C (HFRSC was 2.2 times as much as the control), but dropped to the same level as the control at the lower temperatures, 20 and 17 °C (Fig. 3). Taken together, these results suggest that the antioxidant system of these species increased their activities against the oxidative stress generated by exposure to a low temperature of 23 °C. The antioxidant system of *K. striatum* thalli was unable to resist the lower temperatures (20 and 17 °C) whereas that of *K. alvarezii* thalli still had resistance to the lower temperatures (20 and 17 °C). No significant differences of MDA were found in the two species during exposure to lower temperatures (23 and 20 °C) (Fig. 3). At the lowest temperature of 17 °C, a significant increase of MDA (MDA concentration at 17 °C was 2.2 times as much as






the control) was observed in *K. striatum* thalli, which confirmed the occurrence of oxidative damage; however, no significant increase of MDA was observed in *K. alvarezii* thalli (Fig. 3).

#### Changes in photosynthetic efficiencies (or energy flux ratios)— $\varphi_{Po}$ , $\psi_o$ and $\varphi_{Eo}$

Compared with the control, a significant decrease in the maximum PSII photochemical efficiency ( $F_V/F_m = \varphi_{Po}$ ) was observed in *K. striatum* thalli at 23 °C (the  $F_V/F_m$  value was 0.62 while the control was 0.68) (Fig. 4), indicating that photo-damage had occurred. However, in *K. alvarezii* thalli, there was no significant decrease of  $F_V/F_m$  at that temperature (23 °C). When exposed to 17 °C, both *K. alvarezii* and *K. striatum* thalli had significant decreases in  $F_V/F_m$  compared to the control ( $F_V/F_m$  values were 0.63 and 0.55 at 17 °C in *K. alvarezii* and *K. striatum* thalli, respectively, whereas the respective controls were 0.72 and 0.68). Because the value of  $F_V/F_m$  largely reflected the physiological activity in plants, it was inferred that physiological activity in *K. alvarezii* might



have greater tolerance to low temperature stress than that in *K. striatum*.

With regard to the excitation efficiency for electron transport beyond  $Q_A^-$  ( $\psi_o$ ), both *K. alvarezii* and *K. striatum* thalli only exhibited significant decreases of  $\psi_o$  when the temperature dropped to 17 °C ( $\psi_o$  values were 0.36 and 0.24 in *K. alvarezii* and *K. striatum* thalli, respectively, whereas the respective controls were 0.47 and 0.37) (Fig. 4). At 23 and 20 °C, no significant decreases of  $\psi_o$  were observed, indicating that the electron transport in the PSII electron transport chain was not influenced significantly in either species. Compared with the control, the quantum yield of electron transport ( $\varphi_{Eo}$ ) in *K. striatum* thalli decreased significantly at 20 and 17 °C ( $\varphi_{Eo}$  values were 0.21 and 0.13 at 20 and 17 °C, respectively, while the control was 0.26) (Fig. 4) while a significant decrease of  $\varphi_{Eo}$  was only observed at 17 °C ( $\varphi_{Eo}$  value was 0.23 at 17 °C while the control was 0.34) in *K. alvarezii* thalli. These results suggest that the lowered temperature of 20 °C led to a decreased quantum yield of electron transport in *K. striatum* thalli, whereas *K. alvarezii* thalli maintained a normal capacity for electron transport at that same temperature (20 °C), indicating again that *K. alvarezii* had stronger low temperature tolerance than *K. striatum* with regard to electron transport.

Since the parameters of photosynthetic efficiencies only reflect the energy cascade from light absorption to electron transport and don't indicate the specific PSII component damaged by low temperature stress, the

performance index on an absorption basis ( $PI_{ABS}$ ) and its three variables  $RC/ABS$ ,  $\varphi_{Po}/(1 - \varphi_{Po})$ ,  $\psi_o/(1 - \psi_o)$ , were calculated to probe the specific component of PSII possibly damaged by low temperature stress.

#### Changes in the performance index— $PI_{ABS}$


$PI_{ABS}$  is more sensitive to changes of photosynthetic activity than the maximal photochemical efficiency ( $F_v/F_m$ ). Compared with the control, there were significant decreases of  $PI_{ABS}$  in *K. striatum* thalli when the temperature was below 23 °C ( $PI_{ABS}$  values were 0.33, 0.32 and 0.12 at 23, 20 and 17 °C, respectively, while the control was 0.61), whereas a significant decrease of  $PI_{ABS}$  was observed in *K. alvarezii* thalli only when the temperature was dropped to 17 °C ( $PI_{ABS}$  value was 0.45 at 17 °C while the control was 1.71) (Fig. 5). This result suggests that the 3 °C decrement in temperature from 26 °C resulted in decreased photosynthetic activity of PSII in *K. striatum* thalli, whereas a 9 °C decrement from 26 °C was needed for a negative impact on the PSII activity in *K. alvarezii* thalli.

In order to explore responses of different parts of PSII in the two species during the low temperature stress, three variables  $RC/ABS$ ,  $\varphi_{Po}/(1 - \varphi_{Po})$  and  $\psi_o/(1 - \psi_o)$ , were calculated.  $RC/ABS$ ,  $\varphi_{Po}/(1 - \varphi_{Po})$  and  $\psi_o/(1 - \psi_o)$  reflect the efficiency of reaction centers, the light energy absorption efficiency and the accepting efficiency of the electron acceptor, respectively. When the temperature was dropped to 23 °C,  $RC/ABS$  decreased significantly in *K. striatum* thalli (the ratio  $RC/ABS$  was 0.32 at 23 °C while the control was 0.46), whereas no significant

decrease of  $RC/ABS$  was observed in *K. alvarezii* until the temperature was decreased to 17 °C (the ratio  $RC/ABS$  was 0.45 at 17 °C while the control was 0.72) (Fig. 5). This suggests that reaction centers of PSII in *K. alvarezii* thalli maintained normal efficiency at 23 and 20 °C whereas the efficiency of PSII reaction centers in *K. striatum* thalli had decreased at these temperatures. At 17 °C, PSII reaction centers of both *K. alvarezii* and *K. striatum* were damaged. However, both *K. alvarezii* and *K. striatum* thalli had significant decreases of  $\varphi_{Po}/(1 - \varphi_{Po})$  at 23 °C (the ratios  $\varphi_{Po}/(1 - \varphi_{Po})$  were 2.2 and 1.6 in *K. alvarezii* and *K. striatum* thalli, respectively, while the respective controls were 2.6 and 2.2), indicating that the low temperature reduced the efficiency of the PSII antenna in both species (Fig. 5). In addition, the accepting efficiencies of the PSII electron acceptor ( $\psi_o/(1 - \psi_o)$ ) in both *K. alvarezii* thalli and *K. striatum* thalli were impacted negatively at 17 °C (the ratios  $\psi_o/(1 - \psi_o)$  were 0.58 and 0.32 in *K. alvarezii* and *K. striatum* thalli, respectively, while the respective controls were 0.90 and 0.60), indicating that the acceptor sides of PSII in these two species were damaged (Fig. 5). In summary, in both species, PSII antennae were first damaged by the low temperature stress; damages to the PSII acceptor side was observed only at the lowest temperature (17 °C). In addition, the PSII reaction centers of *K. striatum* had a lower resistance to the low temperature stress than those of *K. alvarezii*.

#### Discussion

PSII contains some sensitive proteins, and the state of PSII is easily affected by low temperature stress (Čajánek et al. 1998; Tang et al. 2007). After exposure to low temperature stress, physiological dysfunctions, including alteration of metabolic processes, increase in ROS and reduction of photosynthetic capacity occur in plants (Allen and Ort 2001). The harsh conditions imposed by various environmental stresses result in an increase of  $H_2O_2$ , which is the main ROS in *K. alvarezii* thalli (Reis et al. 2011; Ling et al. 2015). Moreover, Barros et al. (2006) found that when *K. alvarezii* thalli were subjected to two co-stressors, chilling and high light, the  $H_2O_2$  level increased. In our study, increases of  $H_2O_2$  levels (Fig. 2) were observed during exposure to lowered temperatures (23 and 20 °C) in both *K. alvarezii* and *K. striatum* thalli, suggesting that oxidative stress occurred, but this increases were much greater in *K. striatum* thalli than in *K. alvarezii* thalli, indicating that *K. striatum* suffered more serious oxidative stress from  $H_2O_2$  than *K. alvarezii*. *Kappaphycus alvarezii* at 17 °C had a similar  $H_2O_2$  level as the control at 26°. That might be due to the increase of  $H_2O_2$ —scavenging enzyme (such as APX) activities. Barros et al. (2003) found that *K. alvarezii* released  $H_2O_2$  into the surrounding medium to avoid harmful accumulation



of  $H_2O_2$ . But *K. striatum* at 17 °C still had a higher  $H_2O_2$  level than the control at 26°. Above results indicated that *K. striatum* might suffer greater oxidative damage than *K. alvarezii* during the low temperature.

Some important antioxidant enzymes (SOD, APX and CAT) of *K. alvarezii* increased their activities when *K. alvarezii* thalli is exposed to the cold and high light stress (Barros et al. 2006). In our study, SOD and APX activities were substantially higher at 23 °C than at 26 °C (Fig. 3) in both two species, indicating the antioxidant systems of both species enhanced their antioxidant activities against the oxidative stress caused by the low temperature. However, compared with the control, no significant increases of SOD and APX activities in *K. striatum* thalli were observed at the lower temperatures of 20 and 17 °C, whereas SOD and APX activities of *K. alvarezii* thalli were still higher at these lower temperatures (Fig. 3), indicating that these two antioxidant enzymes of *K. alvarezii* thalli persisted at the lower temperatures (20 and 17 °C) but those of *K. striatum* thalli didn't. However, it is worth noting that no significant difference of CAT activity was observed during any of the low temperature treatments in either of the two species. Because CAT and APX are two important  $H_2O_2$ -scavenging enzymes, why doesn't CAT respond to the increase of  $H_2O_2$ ? Barros et al. (2003) found that APX was an effective system for decomposing lower concentrations of  $H_2O_2$  in *K. alvarezii*, while higher concentrations of  $H_2O_2$  would be more efficiently decomposed by CAT. Ascorbate peroxidase (APX) has a greater affinity for  $H_2O_2$  than CAT. So it is surmised that concentrations of  $H_2O_2$  didn't reach the level at which CAT would play a role during these low temperature treatments. Not only  $H_2O_2$ , but also  $\cdot OH$  (a powerful oxidizing agent) can cross membranes and may oxidize a variety of compounds (Nagarani and Kumaraguru 2012). Hydroxyl free radical scavenging capacity (HFRSC) was congruent with SOD and APX activities for each species (Fig. 3). A significant increase of HFRSC persisted in *K. alvarezii* thalli, whereas no significant increase of HFRSC was observed in *K. striatum* thalli at the lower temperatures (20 and 17 °C), indicating that *K. alvarezii* retained a stronger capacity for eliminating  $\cdot OH$  at 20 and 17 °C but *K. striatum* did not. Besides, the lipoperoxidation damage was only observed in *K. striatum* thalli (MDA, a marker of lipoperoxidation, increased significantly at 17 °C in that species) (Fig. 3). From the different antioxidant responses of these species to low temperature stress, it was inferred that *K. alvarezii* thalli might have greater low temperature resistance than *K. striatum* thalli.

Given that PSII is the most sensitive part of the photosynthetic apparatus, changes of the PSII state can indicate changes in physiological processes during low temperature stress (Lim et al. 2014). The maximum

PSII photochemical efficiency decreased ( $F_v/F_m$  values decreased) in both species as a consequence of low temperature stress, indicating that photo-damage had occurred such that photosynthetic activities decreased in both species.

The energy flux ratios ( $\varphi_{P_0}$ ,  $\psi_0$  and  $\varphi_{E_0}$ ), which were measured by Chl a fluorescence transients, can reflect the energy utilization from light absorption to electron transport and indicate the photosynthetic efficiencies during the energy cascade (Strasser et al. 2004). Li et al. (2014) found that the maximum quantum yield for primary photochemistry ( $\varphi_{P_0}$ ), the efficiency at which a trapped exciton moved an electron into the electron transport chain beyond  $Q_A^-$  ( $\psi_0$ ) and the quantum yield of electron transport ( $\varphi_{E_0}$ ), all decreased sharply under temperature stress. In our results, a significant decrease of  $\varphi_{P_0}$  was observed in *K. striatum* thalli when the temperature dropped to 23 °C, whereas there was no significant decrease of  $\varphi_{P_0}$  at that temperature in *K. alvarezii* thalli (Fig. 4). On the other hand,  $\varphi_{E_0}$  decreased sharply at 20 °C in *K. striatum* thalli while no significant decrease of  $\varphi_{E_0}$  was observed in *K. alvarezii* thalli (Fig. 4), indicating electron transport of *K. alvarezii* PSII wasn't affected at this temperature. These results suggest that *K. alvarezii* thalli had greater low temperature tolerance than *K. striatum* thalli with regard to the PSII energy utilization.

$PI_{ABS}$  is more sensitive to changes in photosynthetic activity than the maximal photochemical efficiency ( $F_v/F_m$ ) and can reflect the states of different photosynthetic components (Strasser et al. 2004). Strauss et al. (2007) found that low soil temperature decreased the value of  $PI_{ABS}$  seriously in soybeans. For red seaweeds, Zhang et al. (2012b) studied the Chl fluorescence characteristics of different strains of *Porphyra yezoensis* under low temperature stress. They found that the strains Sulian and WT had higher photochemical efficiency of PSII than the strain Sutong under low temperature stress, indicating that Sulian and WT might have better tolerance to low temperature. In our study, significant decreases of  $PI_{ABS}$  were observed in *K. striatum* thalli at 23, 20 and 17 °C, whereas a significant decrease of  $PI_{ABS}$  was observed in *K. alvarezii* thalli only at 17 °C. It indicated that the PSII of *K. alvarezii* thalli might have greater low temperature tolerance than the PSII of *K. striatum* thalli. In order to determine the low temperature tolerance of individual photosynthetic components of PSII (reaction centers, antennae and acceptor sides of PSII) for the two species, the values of  $RC/ABS$ ,  $\varphi_{P_0}/(1 - \varphi_{P_0})$ ,  $\psi_0/(1 - \psi_0)$ , were measured (Fig. 5). From our results, PSII antennae were first damaged by the low temperature stress and damage of the PSII acceptor side was observed only at 17 °C in both species.  $RC/ABS$  decreased significantly in *K. striatum* thalli at 23 and 20 °C while no significant difference



of RC/ABS was observed in *K. alvarezii* thalli at these temperatures. This suggests that efficiency of PSII reaction centres in *K. alvarezii* thalli remained at a normal level while the efficiency of PSII reaction centers in *K. striatum* thalli decreased at lower temperatures (23 and 20 °C). It was inferred that PSII reaction centers of *K. striatum* might have weaker resistance to low temperature stress than those of *K. alvarezii*.

## Conclusions

$\kappa$ -Carrageenan-producing *K. alvarezii* and *K. striatum* are important commercial red seaweeds. Strains of *Kappaphycus* could be selected for temperature tolerance by using the Chl fluorescence method, which is non-destructive to the algal tissue and takes only a few secs to record (Zhang et al. 2012b). This study has shown that the excess H<sub>2</sub>O<sub>2</sub> generated by low temperature stress can indirectly damage the PSII apparatus and these two species have somewhat different physiological responses to low temperature stress. By comparing the several important parameters of the Chl fluorescence transient, it is concluded that the PSII of *K. alvarezii* has a stronger low temperature tolerance than *K. striatum*. Given that photosynthesis has been considered to be the physiological process most sensitive to temperature damage and that PSII is the most sensitive part of the photosynthetic apparatus (Wen et al. 2005), it is inferred that *K. alvarezii* thalli might have stronger low temperature tolerance than *K. striatum* thalli as a consequence of PSII characteristics. Our results will be helpful for further studies on the low temperature tolerance of *Kappaphycus* and for screening *Kappaphycus* strains for growth in a lower temperature environment (or higher latitude regions).

## Abbreviations

APX: ascorbate peroxidase; CAT: catalase; Chl: chlorophyll; HFRSC: hydroxyl free radical scavenging capacity; H<sub>2</sub>O<sub>2</sub>: hydrogen peroxide; MDA: malondialdehyde; OEC: oxygen-evolving complex; ·OH: hydroxyl free radical; PQ: plastoquinone; PSII: photosystem II; ROS: reactive oxygen species; SOD: superoxide dismutase.

## Authors' contributions

HL, JGL and TP designed the research; HL performed the experiments and analyzed the data; LTZ and TP analyzed data; HL and JGL wrote the manuscript. All authors read and approved the final manuscript.

## Author details

<sup>1</sup> Key Laboratory of Experimental Marine Biology, National & Local Joint Engineering Laboratory of Ecological Mariculture, Institute of Oceanology, Chinese Academy of Sciences, 7 Nanhai Road, Qingdao 266071, China.

<sup>2</sup> Laboratory for Marine Biology and Biotechnology, Qingdao National Laboratory for Marine Science and Technology, 1 Wenhai Road, Aoshanwei Town, Jimo, Qingdao 266071, China. <sup>3</sup> University of Chinese Academy of Sciences, Beijing 100049, China.

## Acknowledgements

This work was supported by the National Natural Science Foundation of China (41306154), Special Project for Marine Public Welfare Industry (201505033) and

Marine Economy Innovation and Development Fund of Qingdao (2012). The authors would like to thank Zhaoliang Zheng from Lingshui Haotian Company for the provision of *Kappaphycus alvarezii* and *Kappaphycus striatum*, and thank Dr. John van der Meer (Pan-American Marine Biotechnology Association) for his assistance with proofreading.

## Competing interests

The authors declare that they have no competing interests.

Received: 7 May 2016 Accepted: 25 July 2016

Published online: 10 August 2016

## References

- Allen DJ, Ort DR (2001) Impacts of chilling temperatures on photosynthesis in warm-climate plants. *Trends Plant Sci* 6:36–42
- Barros MP, Granbom M, Colepicolo P, Pedersén M (2003) Temporal mismatch between induction of superoxide dismutase and ascorbate peroxidase correlates with high H<sub>2</sub>O<sub>2</sub> concentration in seawater from clofibrate-treated red algae *Kappaphycus alvarezii*. *Arch Biochem Biophys* 420:161–168
- Barros MP, Necchi O, Colepicolo P, Pedersén M (2006) Kinetic study of the plastoquinone pool availability correlated with H<sub>2</sub>O<sub>2</sub> release in seawater and antioxidant responses in the red alga *Kappaphycus alvarezii* exposed to single or combined high light, chilling and chemical stresses. *Biochim Biophys Acta* 1757:1520–1528
- Bindu MS (2011) Empowerment of coastal communities in cultivation and processing of *Kappaphycus alvarezii*—a case study at Vizhinjam village, Kerala, India. *J Appl Phycol* 23:157–163
- Bindu MS, Levine IA (2011) The commercial red seaweed *Kappaphycus alvarezii*—an overview on farming and environment. *J Appl Phycol* 23:789–796
- Bixler HJ, Porse H (2011) A decade of change in the seaweed hydrocolloids industry. *J Appl Phycol* 23:321–335
- Bixler HJ, Johndro K, Falshaw R (2001) Kappa-2 carrageenan: structure and performance of commercial extracts: II. Performance in two simulated dairy applications. *Food Hydrocoll* 15:619–630
- Čajánek M, Štroch M, Lachetová I, Kalina J, Spunda V (1998) Characterization of the photosystem II inactivation of heat-stressed barley leaves as monitored by the various parameters of chlorophyll a fluorescence and delayed fluorescence. *J Photochem Photobiol B—Biol* 47:39–45
- Collén J, Pedersén M, Bornman CH (1994) A stress-induced oxidative burst in *Eucheuma platycladum* (Rhodophyta). *Physiol Plant* 92:417–422
- Das K, Samanta L, Chainy G (2000) A modified spectrophotometric assay of superoxide dismutase using nitrite formation by superoxide radicals. *Indian J Biochem Biophys* 37:201–204
- Deng Y, Wang W, Yu P, Xi Z, Xu L, Li X, He N (2013) Comparison of taurine, GABA, Glu, and Asp as scavengers of malondialdehyde in vitro and in vivo. *Nanoscale Res Lett* 8:1–9
- Doty MS (1973) Farming the red seaweed, *Eucheuma*, for carrageenans. *Micronica* 9:59–73
- Fariduddin Q, Khan TA, Yusuf M (2014) Hydrogen peroxide mediated tolerance to copper stress in the presence of 28-homobrassinolide in *Vigna radiata*. *Acta Physiol Plant* 36:2767–2778
- Fraga CG, Leibovitz BE, Tappel AL (1988) Lipid peroxidation measured as thiobarbituric acid-reactive substances in tissue slices: characterization and comparison with homogenates and microsomes. *Free Radic Biol Med* 4:155–161
- Ganzon-Fortes ET, Trono GC Jr, Villanueva RD, Romero JB, Montañón MNE (2012) 'Endong', a rare variety of the farmed carrageenophyte *Eucheuma denticulatum* (Burman) Collins & Hervey from the Philippines. *J Appl Phycol* 24:1107–1111
- Gechev T, Willekens H, Montagu MV, Eacute DI, Camp WIMV, Toneva V, Minkov I (2003) Different responses of tobacco antioxidant enzymes to light and chilling stress. *J Plant Physiol* 160:509–515
- Glenn EP, Doty MS (1990) Growth of the seaweeds *Kappaphycus alvarezii*, *K. striatum* and *Eucheuma denticulatum* as affected by environment in Hawaii. *Aquaculture* 84:245–255



- Halliwell B, Grootveld M, Gutteridge JMC (2006) Methods for the measurement of hydroxyl radicals in biochemical systems: deoxyribose degradation and aromatic hydroxylation. In: Glick D (ed) *Methods of biochemical analysis*, vol 33. Wiley, New York, pp 59–90
- Li L, Li X, Xu X, Lin L, Zeng F (2014) Effects of high temperature on the chlorophyll a fluorescence of *Alhagi sparsifolia* at the southern Taklamakan Desert. *Acta Physiol Plant* 36:243–249
- Lideman Nishihara GN, Noro T, Terada R (2013) Effect of temperature and light on the photosynthesis as measured by chlorophyll fluorescence of cultured *Eucheuma denticulatum* and *Kappaphycus* sp. (Sumba strain) from Indonesia. *J Appl Phycol* 25:399–406
- Lim PE, Tan J, Phang SM, Nikmatullah A, Hong DD, Sunarpi H, Hurtado AQ (2014) Genetic diversity of *Kappaphycus* Doty and *Eucheuma* J. Agardh (Solieriaceae, Rhodophyta) in Southeast Asia. *J Appl Phycol* 26:1253–1272
- Ling ALM, Yasir S, Matanjun P, Bakar MFA (2015) Effect of different drying techniques on the phytochemical content and antioxidant activity of *Kappaphycus alvarezii*. *J Appl Phycol* 27:1717–1723
- Liu J, Pang T, Wang L, Li J, Lin W (2009) The reasons causing catastrophic death in tropical carrageenan producing seaweeds and their difference in resistance to illness. *Oceanol Limnol Sin* 40:235–241
- Liu L, Liu Y, Cui J, Liu H, Liu Y, Qiao W, Sun H, Yan C (2013) Oxidative stress induces gastric submucosal arteriolar dysfunction in the elderly. *World J Gastroenterol* 19:9439–9446
- Malanga G, Kozak RG, Puntarulo S (1999) N-Acetylcysteine-dependent protection against UV-B damage in two photosynthetic organisms. *Plant Sci* 141:129–137
- Nagarani N, Kumaraguru AK (2012) Chemical characterization, temperature stability, and enzymatic studies on edible marine algae *Kappaphycus alvarezii* (Doty). *J Aquat Food Prod Technol* 21:480–492
- Nakano Y, Asada K (1981) Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in Spinach chloroplasts. *Plant Cell Physiol* 22:867–880
- Ohno M, Largo DB, Ikumoto T (1994) Growth rate, carrageenan yield and gel properties of cultured kappa-carrageenan producing red alga *Kappaphycus alvarezii* (Doty) Doty in the subtropical waters of Shikoku, Japan. *J Appl Phycol* 6:1–5
- Pang T, Liu J, Liu Q, Li H, Li J (2015) Observations on pests and diseases affecting a eucheumatoid farm in China. *J Appl Phycol* 27:1975–1984
- Reis MO, Necchi O, Colepicolo P, Barros MP (2011) Co-stressors chilling and high light increase photooxidative stress in diuron-treated red alga *Kappaphycus alvarezii* but with lower involvement of H<sub>2</sub>O<sub>2</sub>. *Pestic Biochem Phys* 99:7–15
- Stirbet A et al (2012) Chlorophyll a fluorescence induction: a personal perspective of the thermal phase, the J-I-P rise. *Photosynth Res* 113:15–61
- Strasser RJ (1978) The grouping model of plant photosynthesis. In: Akoyunoglou G (ed) *Chloroplast Development*. Elsevier/North Holland Biomedical Press, Amsterdam, pp 513–524
- Strasser RJ, Tsimilli-Michael M, Srivastava A (2004) Analysis of the chlorophyll a fluorescence transient. In: Papageorgiou GC, Govindjee A (eds) *Chlorophyll fluorescence: a signature of photosynthesis, advances in photosynthesis and respiration*. Springer, Dordrecht, pp 321–362
- Strauss AJ, Krüger GH, Strasser RJ, Van Heerden PD (2007) The role of low soil temperature in the inhibition of growth and PSII function during dark chilling in soybean genotypes of contrasting tolerance. *Physiol Plantarum* 19:89–105
- Tang Y, Chen M, Xu Y, Kuang T (2007) Changes in thermostability of photosystem II and leaf lipid composition of rice mutant with deficiency of light-harvesting chlorophyll *a/b* protein complexes. *J Integr Plant Biol* 49:515–522
- Wen X, Qiu N, Lu Q, Lu C (2005) Enhanced thermotolerance of photosystem II in salt-adapted plants of the halophyte *Artemisia anethifolia*. *Planta* 220:486–497
- Zhang L, Zhang Z, Gao H, Meng X, Yang C, Liu J, Meng Q (2012a) The mitochondrial alternative oxidase pathway protects the photosynthetic apparatus against photodamage in *Rumex* K-1 leaves. *BMC Plant Biol* 12:40
- Zhang T, Shen Z, Xu P, Zhu J, Lu Q, Shen Y, Wang Y, Yao C, Li J, Wang Y (2012b) Analysis of photosynthetic pigments and chlorophyll fluorescence characteristics of different strains of *Porphyra yezoensis*. *J Appl Phycol* 24:881–886

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

---

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)

---