



Sodium–Glucose Cotransporter 2 Inhibitors and Diabetic Ketoacidosis: A Case Series From Three Academic Institutions

Diabetes Care 2017;40:e65–e66 | <https://doi.org/10.2337/dc16-2591>

Negin Misaghian-Xanthos,¹
Afreen I. Shariff,² Kavya Mekala,³
Loretta R. Fearrington,¹ Tracy L. Setji,²
Joseph A. Aloji,³ and John B. Buse¹

Sodium–glucose cotransporter 2 inhibitors (SGLT2i) have recently been associated with diabetic ketoacidosis (DKA). Initial case reports were largely among patients using SGLT2i off label in the setting of type 1 diabetes and were unusual because of limited associated hyperglycemia (1). More recently, the problem has been specifically noted in patients with type 2 diabetes (2). Meta-analysis of data from initial randomized controlled trials of SGLT2i suggests little risk of DKA (<0.1%) in patients with type 2 diabetes (3).

To better define the scope of the problem in clinical practice, we conducted a search of three academic health care systems' electronic health records. The study was approved by institutional review boards. Patients were selected from all emergency and inpatient encounters between 1 January 2013 and 30 April 2016 that contained the diagnosis codes for ketosis or acidosis (ICD-9 codes: 250.10, 250.11, 250.12, 250.13, 276.2, 790.6; ICD-10 codes: E08.10, E08.11, E09.10, E09.11, E10.10, E10.11, E13.10, E13.11, E11.65, E11.69, E87.2) and had an SGLT2i noted in a medication list. Individuals who met prespecified criteria for DKA with arterial pH <7.3, venous pH <7.27, serum bicarbonate <18 mEq/L, positive urine or serum ketones

per institutional laboratory cutoffs, or an anion gap >10 were included.

A total of 11,197 individuals had a prescription for SGLT2i documented in a medication list and 39 individuals met the criteria for DKA. Details are provided in Supplementary Table 1. Of the DKA cases, 74% appeared to have type 2 diabetes. Glucose \leq 300 mg/dL was noted at presentation of DKA in 26 of the 39 cases (72%). The mean glucose at presentation was 328 mg/dL, median 266 mg/dL, and range 125–904 mg/dL. With respect to precipitating causes, 49% had nausea/vomiting, although it is unclear if that was a cause, contributor, or consequence of the DKA; 67% had an obvious other precipitating event such as surgery, insulin dose reduction, or decreased oral intake and weight loss.

This is the largest and most comprehensive case series of DKA associated with the use of SGLT2i. Uncertainty about the incidence rate of DKA among SGLT2i-treated patients in our institutions is substantial because we do not have claims data confirming that prescriptions were filled, we likely missed cases that were undiagnosed or incompletely assessed, and some of the cases listed were prescribed SGLT2i by practitioners whose prescribing is not reflected in our

electronic health records. The relatively lower presenting glucose observed in the setting of SGLT2i-associated DKA is likely due to the proposed pathologic mechanisms (4). It is unclear if there are specific characteristics that make some patients more predisposed.

DKA in the setting of SGLT2i therapy is unusual in that it disproportionately affects those with type 2 diabetes and often presents with glucose <300 mg/dL, making recognition challenging. We would recommend a high index of suspicion for DKA in patients taking SGLT2i with unexplained malaise or gastrointestinal symptoms and recommend measuring urine or plasma ketones in that setting. While its occurrence is unusual, we should counsel patients on the signs, symptoms, and possible precipitants of DKA even in the setting of type 2 diabetes when prescribing SGLT2i.

Acknowledgments. The authors thank the data teams at the North Carolina Translational and Clinical Sciences Institute, Duke Clinical and Translational Science Institute, and Wake Forest Clinical and Translational Science Institute for identifying potential cases from the electronic health records.

Funding. The project described was supported at each of the institutions by the National Center

¹University of North Carolina School of Medicine, Chapel Hill, NC

²Duke University School of Medicine, Durham, NC

³Wake Forest School of Medicine, Winston-Salem, NC

Corresponding author: John B. Buse, jbuse@med.unc.edu.

Received 4 December 2016 and accepted 6 March 2017.

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc16-2591/-/DC1>.

N.M.-X., A.I.S., and K.M. contributed equally to this study.

© 2017 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <http://www.diabetesjournals.org/content/license>.

for Advancing Translational Sciences, National Institutes of Health, through Grant Award Numbers UL1-TR-001111, UL1-TR-001420, and UL1-TR-001117.

Duality of Interest. J.B.B.'s institution has received grants and contracts from AstraZeneca, Boehringer Ingelheim, Eli Lilly, Janssen, Sanofi, and Lexicon Pharmaceuticals to support his efforts as investigator and/or advisor. No other potential conflicts of interest relevant to this article were reported.

Author Contributions. All authors participated in study design, analysis, and drafting and approved the final manuscript submitted. N.M.-X., A.I.S., and

K.M. are guarantors of this work and, as such, had full access to all the data at their respective institutions and take responsibility for the integrity of the data and the accuracy of the analysis.

References

1. Peters AL, Buschur EO, Buse JB, Cohan P, Diner JC, Hirsch IB. Euglycemic diabetic ketoacidosis: a potential complication of treatment with sodium–glucose cotransporter 2 inhibition. *Diabetes Care* 2015;38:1687–1693
2. U.S. Food and Drug Administration. SGLT2 inhibitors: drug safety communication - labels

to include warnings about too much acid in the blood and serious urinary tract infections [Internet], 4 December 2015. Available from <http://www.fda.gov/Safety/MedWatch/SafetyInformation/SafetyAlertsforHumanMedicalProducts/ucm475553.htm>. Accessed 17 November 2016

3. Erond N, Desai M, Ways K, Meininger G. Diabetic ketoacidosis and related events in the canagliflozin type 2 diabetes clinical program. *Diabetes Care* 2015;38:1680–1686
4. Rosenstock J, Ferrannini E. Euglycemic diabetic ketoacidosis: a predictable, detectable, and preventable safety concern with SGLT2 inhibitors. *Diabetes Care* 2015;38:1638–1642