



Correction for Khalimonchuk et al., "Formation of the Redox Cofactor Centers during Cox1 Maturation in Yeast Cytochrome Oxidase"

Oleh Khalimonchuk,¹ Megan Bestwick,¹ Brigitte Meunier,² Talina C. Watts,¹ Dennis R. Winge¹

University of Utah Health Sciences Center, Departments of Medicine and Biochemistry, Salt Lake City, Utah, USA¹; Centre de Génétique Moléculaire, CNRS, Gif-sur-Yvette, France²


Volume 30, no. 4, p. 1004–1017, 2010, https://doi.org/10.1128/MCB.00640-09. Page 1007, Fig. 1D: The anti- F_1 control immunoblot was inadvertently duplicated. The correct anti- F_1 blot from the original experiment is shown below. This is the monomeric form of mitochondrial F_1F_0 ATPase (complex V) as a loading control.



Citation Khalimonchuk O, Bestwick M, Meunier B, Watts TC, Winge DR. 2017. Correction for Khalimonchuk et al., "Formation of the redox cofactor centers during Cox1 maturation in yeast cytochrome oxidase." Mol Cell Biol 37:e00127-17. https://doi.org/10.1128/MCB .00127-17.


Copyright © 2017 American Society for Microbiology. All Rights Reserved.

Page 1007, Fig. 1E: The anti- F_1 immunoblot was inadvertently a duplicate of the control blot in Fig. 5A. Therefore, we redid the Fig. 1E study using the original samples from 2009. The resulting images are shown below.




Author Correction Molecular and Cellular Biology

Page 1008, Fig. 2C, lanes 5 and 6: The anti- F_1 immunoblot was a duplicate of the blot in Fig. 1D. The original samples were rerun, and the correct anti-Myc and anti- F_1 blots are shown below.



Page 1011, Fig. 5A, lanes 3 and 4: The anti- F_1 blot was a duplicate of the control in Fig. 1E, as mentioned above. We reran the original samples, and the correct anti-Myc and anti- F_1 blots for Coa1-Myc samples from wild-type (WT) and $cox10\Delta$ cells are shown below.



We regret these errors, but they do not affect the interpretation of the experiments and the conclusions of the study.