

Transcriptional Activity of Baculovirus Very Late Factor 1

Toni-Ann Mistretta¹ and Linda A. Guarino^{1,2*}

*Departments of Biochemistry and Biophysics¹ and Entomology,² Texas A&M University,
College Station, Texas*

Received 20 July 2004/Accepted 8 September 2004

The product of the *vlf-1* (very late factor 1) gene is required for expression of very late genes during the final phase of infection. To determine whether VLF-1 functions as a transcriptional activator, VLF-1 was overexpressed and purified by affinity and cation exchange chromatography. The addition of purified protein to transcription assays containing baculovirus RNA polymerase stimulated transcription of the very late polyhedrin promoter but not the late *39k* promoter. Furthermore, construction and analysis of chimeric templates identified sequences within the polyhedrin promoter that were necessary for enhancement.

Baculoviruses are unusual in having two distinct phases of late gene expression. The standard late genes encode virion structural proteins and are expressed following the onset of DNA replication. The very late genes, which encode proteins required for the formation of occluded virions, are maximally expressed after the formation of budded virus. A viral gene required for the burst of very late gene transcription, called very late expression factor 1 (*vlf-1*), was identified by marker rescue of the temperature-sensitive mutant virus *tsB837* (4). This mutant was viable at the nonpermissive temperature but was defective in the production of occluded virus. The failure of *tsB837* to produce occluded virions was due to a significant reduction in the level of polyhedrin (*polh*) mRNA, which codes for the major occlusion body protein. Transcription and expression of *p10*, another highly expressed very late gene, was also reduced at nonpermissive temperature, although to a lesser extent than polyhedrin.

The specific stimulation of very late promoters by VLF-1 has also been observed in transient expression assays (10, 14). The addition of plasmids encoding VLF-1 stimulated expression from the two very late promoters, *polh* and *p10*, but had little or no effect on two late promoters, *vp39* and *p6.9*. Late and very late promoters differ primarily by the presence or absence of a burst sequence, a sequence downstream of the transcriptional start site that is 90% A+T (3, 5, 7, 9). The burst sequence of a very late promoter is needed in order to respond to stimulation by VLF-1 in transient-expression assays, and gel mobility shift and footprinting assays also indicated that VLF-1 specifically bound to the burst sequence *in vitro* (12).

The mechanism by which VLF-1 stimulates very late genes, however, is unknown. When VLF-1 was originally described, the authors who described it noted sequence similarity with the integrase family of tyrosine recombinases (4). While the homology is intriguing, it lends few clues as to the function of VLF-1. The other members of this family are primarily involved in DNA integration and DNA replication, and none of them are known to act as transcription factors. Furthermore,

all integrases and tyrosine recombinases are characterized by an essential active-site tyrosine which forms a covalent protein-DNA intermediate (1, 6). VLF-1 has this conserved tyrosine but it is not required for VLF-1 to stimulate transcription (14). Thus, it seems unlikely that the mechanism of VLF-1 function in transcription is related to integration or recombination.

To characterize the mechanism of VLF-1 function, we purified VLF-1 by using a baculovirus overexpression system in order to examine its activity in transcription reactions containing purified baculovirus RNA polymerase (RNAPol). A recombinant virus, called AcVLF-1, containing the *vlf-1* gene under the control of the polyhedrin promoter was constructed according to standard procedures (8). A 1,685-bp SpeI-SspI fragment of an AcNPV HindIII-B genomic clone was inserted into the XbaI and SmaI sites of pVL1393. *Spodoptera frugiperda* (Sf9) cells were cotransfected with 5 µg of the resulting transfer vector, pVLF-1, and 2 µg of BakPak viral DNA previously digested with Bsu36I. Individual plaques were further purified, and the correct construction of AcVLF1 was verified by restriction digestion and sequence analysis. Nuclear extracts prepared from AcVLF1-infected cells at 60 h postinfection contained a highly expressed protein compared to extracts infected with wild-type virus (Fig. 1). This protein migrated at the position expected for VLF-1. Crude extracts were precipitated with saturated ammonium sulfate and dialyzed for loading onto a heparin affinity column. VLF-1 was purified by affinity chromatography on a heparin agarose column by using column buffers as previously described for the baculovirus protein PP31 (2). Fractions were monitored by Coomassie blue staining of sodium dodecyl sulfate (SDS)-polyacrylamide gels. Peak fractions were pooled, diluted, and loaded onto a Mono S cation exchange column. Fractions containing purified VLF-1 were pooled and stored in 50 mM HEPES (pH 7.2)–400 mM KCl–50% glycerol–1 mM dithiothreitol–0.1 mM EDTA. The yield of purified VLF-1 ranged from 50 to 200 µg per liter of infected cells. The purified protein was subjected to peptide mass fingerprinting. Masses from 15 tryptic peptides were fit to the masses of predicted VLF-1 peptides, representing 39.3% coverage, which is sufficient to confirm that the purified protein was indeed VLF-1.

To determine whether VLF-1 acts directly at the level of transcription, *in vitro* transcription assays were conducted by

* Corresponding author. Mailing address: Department of Biochemistry, Texas A&M University, 2128 TAMU, College Station, TX 77843-2128. Phone: (979) 845-7556. Fax: (979) 845-9274. E-mail: lguarino@tamu.edu.




FIG. 1. Purification of VLF-1. Baculovirus expression. Nuclear extracts (NE) were prepared at 60 h postinfection from *S. frugiperda* cells infected with wild-type (WT) AcNPV (lane 2) or the recombinant Ac-VLF1 (lane 3). Nuclear extracts were purified on a heparin (Hep) column, and peak fractions (lane 4) were further purified on a Mono S column (S) (lane 5). Each lane contains 2 μ g of protein. Lane 1, crude extract; lane 2, protein eluted from Talon matrix with 100 mM imidazole; lane 3, peak fraction from a Mono S column. The position of VLF-1 is indicated by an arrow on the right, and the positions of relevant molecular markers are indicated on the left.

using purified baculovirus RNA polymerase and VLF-1. Transcription reactions contained equimolar amounts of two templates (Fig. 2A). The plasmid pPolh/CFS is a C-free version of the polyhedrin promoter driving transcription of a 191-nucleotide C-free cassette, while p39kL/CFS is a C-free version of the late *39k* promoter driving the same C-free cassette (11). The transcripts produced from these two templates differ in size; therefore, both templates can be added to the same reaction mixture to determine which is preferred by polymerase under different conditions. In the absence of VLF-1, the late and very late templates produced approximately equal amounts of RNA transcript (Fig. 2B), indicating that RNAPol cannot distinguish between the two types of promoter, as previously shown (11). The addition of VLF-1 resulted in a dramatic increase in the amount of transcript originating from the pPolh/CFS template but did not significantly affect the p39kL/CFS product. Since both templates were added to the same reaction, the preference can best be demonstrated by plotting the ratio of *polh* transcript to *39kL* transcript (Fig. 2C). This ratio increased proportionately with the concentration of VLF-1 added. The highest concentration tested produced a more-than-10-fold increase in transcription from the very late *polh* promoter compared to the level for the late *39k* promoter. Similar results were obtained with His-tagged VLF-1 produced in bacteria (data not shown).

To test whether stimulation of pPolh/CFS was due to the burst sequence, the plasmid pPolh Δ burst/CFS was constructed by inserting a double-stranded oligonucleotide containing the polyhedrin promoter lacking the burst sequence (ATCATGG AGATAATTTAAATGATAACCATCTCGCAAATAAATA AGTATTTT) into the *Sma*I site of pCFS, the parental plasmid for pPolh/CFS (11). In transcription assays involving equimolar amounts of pPolh/CFS and pPolh Δ burst/CFS in the absence of VLF-1, the levels of transcripts from both templates were equivalent. The addition of VLF-1 at a concentration yielding half-maximal stimulation increased the amount of pPolh/CFS




FIG. 2. VLF-1 enhances transcription from the polyhedrin promoter. (A) Schematic diagram of the transcriptional templates pPolh/CFS (polh/CFS) and 39kL/CFS. The open box represents the C-free cassette. The darkly shaded box represents the polyhedrin (*polh*) promoter, the lightly shaded box represents the *39k* late promoter, and the black box represents the burst sequence. (B) Transcription assay. Lane 1, ϕ X174/HinfI marker; lanes 2 to 6, transcripts produced from reaction mixtures containing 0.04 μ M purified RNA polymerase; 0.4 pmol of each template in the absence of VLF-1 (lane 2) or increasing amounts of purified VLF-1 (0.16 to 1.28 μ M in twofold increments [lanes 3 to 6]) were incubated under standard transcription reaction conditions at 30°C for 12 min. The RNA transcripts were resolved on 6% polyacrylamide-8 M urea gel and exposed on PhosphorImager plates. The positions of the transcripts are indicated on the right. The positions of relevant molecular size markers are indicated in nucleotides on the left. (C) Transcripts were quantitated by using ImageQuant software and plotted as the ratio of the *polh* transcript to the *39k* transcript as a function of input VLF-1 concentration.

transcript relative to that of pPolh Δ burst/CFS added to the same tube by a factor of 4.4 ± 0.3 in triplicate reactions (Fig. 3A). This result indicates that the burst sequence is a major determinant of VLF-1 action.




FIG. 3. Transcriptional enhancement by VLF-1 requires a burst sequence. (A) Transcription from the polyhedrin promoter in the presence and absence of a burst sequence. Transcripts were produced from in vitro transcription reaction mixtures containing pPolh/CFS and pPolh Δ Burst/CFS. Transcripts were quantified by using ImageQuant software and calculated as the ratio of pPolh/CFS to pPolh Δ Burst/CFS transcript. (B) Transcription from the chimeric p39kLburst/CFS template is stimulated by VLF-1. Reactions contained equimolar amounts of p39kL/CFS and pp39kLburst/CFS. Transcripts were quantified by using ImageQuant software.

This model also suggests that it should be possible to make a late promoter responsive to VLF-1 by inserting the polyhedrin burst sequence downstream of the transcription start site. To test this hypothesis, a C-free version of the burst sequence (GTGTTTTTGTAAAAGTTTTGTAATAAAAAA AATTATTAATAG) was inserted into the EcoRV site of p39kL, an intermediate in the construction of pp39kL/CFS (11). The two templates p39kL/CFS and pp39kLburst/CFS were assayed in competition reactions (Fig. 3B). In the absence of VLF-1, transcript levels were comparable. The addition of VLF-1 stimulated transcription from pp39kLburst/CFS but had no effect on transcription from p39kL/CFS, as expected. In triplicate reactions, the amount of 39kL/CFS transcript was 2.3 ± 0.06 -fold higher than that of 39kL/CFS. Although this was less than the amount seen with polyhedrin, this result confirms that the burst sequence is important for VLF-1 activity. This lower activity of the synthetic very late promoter could be due to a loss of some secondary structure in the leader region or other unknown sequence features of late promoters compared to very late ones (3).

In vivo, the timing of expression and high level of synthesis of polyhedrin appear to be regulated by the concentration of VLF-1. VLF-1 is a late protein and so is not present during the early stages of infection (4). VLF-1 appears by 15 h postinfection and increases through 24 h postinfection (14). Polyhedrin protein is synthesized at low levels during the late stage of infection but increases significantly thereafter, reaching maximum levels at 36 to 48 h postinfection. Furthermore, the timing of polyhedrin expression can be modified by varying the onset of VLF-1 expression (14). Construction of recombinant viruses with *vlf-1* cloned under the control of a strong late promoter resulted in higher levels of VLF-1 and accelerated expression of polyhedrin. In contrast, production of a recombinant virus with *vlf-1* under the control of a minimal promoter produced lower levels of VLF-1 and reduced expression of polyhedrin. Together, these experiments suggest that polyhedrin expression is regulated by the intracellular concentration of VLF-1 (13). The results presented here are consistent

with these observations, as they show that templates containing a burst sequence were stimulated by VLF-1 in a concentration-dependent manner. Questions still remain regarding the mechanism of VLF-1 function, but the results presented here suggest that it can be studied by using this in vitro system, which is amenable to biochemical analyses.

This research was supported by a grant from the National Science Foundation (MCB 0110925).

REFERENCES

- Esposito, D., and J. J. Scocca. 1997. The integrase family of tyrosine recombinases: evolution of a conserved active site domain. *Nucleic Acids Res.* **25**:3605–3614.
- Guarino, L. A., T. A. Mistretta, and W. Dong. 2002. DNA binding activity of the baculovirus late expression factor PP31. *Virus Res.* **90**:187–195.
- Mans, R. M., and D. Knebel-Morsdorf. 1998. In vitro transcription of p38/polyhedrin hybrid promoters reveals sequences essential for recognition by the baculovirus-induced RNA polymerase and for the strength of very late viral promoters. *J. Virol.* **72**:2991–2998.
- McLachlin, J. R., and L. K. Miller. 1994. Identification and characterization of *vlf-1*, a baculovirus gene involved in very late gene expression. *J. Virol.* **68**:7746–7756.
- Morris, T. D., and L. K. Miller. 1994. Mutational analysis of a baculovirus major late promoter. *Gene* **140**:147–153.
- Nunes-Duby, S. E., H. J. Kwon, R. S. Tirumalai, T. Ellenberger, and A. Landy. 1998. Similarities and differences among 105 members of the Int family of site-specific recombinases. *Nucleic Acids Res.* **26**:391–406.
- Ooi, B. G., C. Rankin, and L. K. Miller. 1989. Downstream sequences augment transcription from the essential initiation site of a baculovirus polyhedrin gene. *J. Mol. Biol.* **210**:721–736.
- O'Reilly, D. R., L. K. Miller, and V. A. Luckow. 1992. *Baculovirus expression vectors: a laboratory manual*. W.H. Freeman & Co., New York, N.Y.
- Rankin, C., B. G. Ooi, and L. K. Miller. 1988. Eight base pairs encompassing the transcriptional start point are the major determinant for baculovirus polyhedrin gene expression. *Gene* **70**:39–49.
- Todd, J. W., A. L. Passarelli, A. Lu, and L. K. Miller. 1996. Factors regulating baculovirus late and very late gene expression in transient-expression assays. *J. Virol.* **70**:2307–2317.
- Xu, B., S. Yoo, and L. A. Guarino. 1995. Differential transcription of baculovirus late and very late promoters: fractionation of nuclear extracts by phosphocellulose chromatography. *J. Virol.* **69**:2912–2917.
- Yang, S., and L. K. Miller. 1999. Activation of baculovirus very late promoters by interaction with very late factor 1. *J. Virol.* **73**:3404–3409.
- Yang, S., and L. K. Miller. 1998. Control of baculovirus polyhedrin gene expression by very late factor 1. *Virology* **248**:131–138.
- Yang, S., and L. K. Miller. 1998. Expression and mutational analysis of the baculovirus very late factor 1 (*vlf-1*) gene. *Virology* **245**:99–109.