



## Correction for Fletcher et al., “Vaccine Potential of the *Neisseria meningitidis* 2086 Lipoprotein”

Leah D. Fletcher, Liesel Bernfield, Vicki Barniak, John E. Farley, Alan Howell, Melissa Knauf, Peggy Ooi, Robert P. Smith, Paige Weise, Mike Wetherell, Xiaoling Xie, Robert Zagursky, Ying Zhang, and Gary W. Zlotnick

Wyeth Vaccines Research, Pearl River, New York 10965

Volume 72, no. 4, p. 2088–2100, 2004, <https://doi.org/10.1128/IAI.72.4.2088-2100.2004>. Page 2089, column 2: Lines 62–64 should read as follows. “. . . Oligonucleotides used for the amplification of these 2086 genes were 5LIP2996 and N2996SPH. The same oligonucleotide PCR primers were used for all five strains. The restriction site BamHI was . . . .”

Page 2091, Table 2: Row 18 labeled “5LIP771” should not be included in the table.

**Citation** Fletcher LD, Bernfield L, Barniak V, Farley JE, Howell A, Knauf M, Ooi P, Smith RP, Weise P, Wetherell M, Xie X, Zagursky R, Zhang Y, Zlotnick GW. 2017. Correction for Fletcher et al., “Vaccine potential of the *Neisseria meningitidis* 2086 lipoprotein.” *Infect Immun* 85:e00212-17. <https://doi.org/10.1128/IAI.00212-17>.

**Copyright** © 2017 American Society for Microbiology. All Rights Reserved.

L.D.F., L.B., V.B., J.E.F., A.H., M.K., P.O., R.P.S., P.W., M.W., X.X., R.Z., and Y.Z. were unreachable for approval of this correction.