



Published in final edited form as:

J Smok Cessat. 2017 June ; 12(2): 88–98. doi:10.1017/jsc.2015.15.

Attitudes and interest in technology-based treatment and the remote monitoring of smoking among adolescents and emerging adults

Erin McClure, Ph.D.^{1,4}, Nathaniel Baker, M.S.², Matthew J. Carpenter, Ph.D.¹, Frank A. Treiber, Ph.D.^{1,3,4}, and Kevin Gray, M.D.¹

¹Department of Psychiatry and Behavioral Sciences, Medical University of South Carolina, Charleston, SC, USA

²Department of Public Health Sciences, Medical University of South Carolina, Charleston, SC, USA

³Department of Nursing, Medical University of South Carolina, Charleston, SC, USA

⁴Technology Applications Center for Healthful Lifestyles(TACHL), South Carolina Center of Economic Excellence, Medical University of South Carolina, Charleston, SC, USA

Abstract

Introduction—Despite the public health relevance of smoking in adolescents and emerging adults, this group remains understudied and underserved. High technology utilization among this group may be harnessed as a tool for better understanding of smoking, yet little is known regarding the acceptability of mobile health (mHealth) integration.

Methods—Participants (ages 14–21) enrolled in a smoking cessation clinical trial provided feedback on their technology utilization, perceptions, and attitudes; and interest in remote monitoring for smoking. Characteristics that predicted greater technology acceptability for smoking treatment were also explored.

Results—Participants (N=87) averaged 19 years old and were mostly male (67%). Technology utilization was high for smart phone ownership (93%), Internet use (98%), and social media use (94%). Despite this, only one-third of participants had ever searched the Internet for cessation tips or counseling (33%). Participants showed interest in mHealth-enabled treatment (48%) and felt that it could be somewhat helpful (83%). Heavier smokers had more favorable attitudes toward technology-based treatment, as did those with smartphones and unlimited data.

Conclusions—Our results demonstrate high technology utilization, favorable attitudes towards technology, and minimal concerns. Technology integration among this population should be

Corresponding author: Erin A. McClure, Addiction Sciences Division, Medical University of South Carolina, 125 Doughty St., Suite 190, MSC 861, Charleston, SC 29425, Phone: 843-792-7192, Fax: 843-792-3982, mccluree@musc.edu.

Conflict of Interest: None declared.

Ethical Standards: The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

pursued, though in a tailored fashion, to accomplish the goal of providing maximally effective, just-in-time interventions.

Keywords

technology; treatment; cessation; survey research; youth tobacco use

Introduction

Cigarette smoking remains the leading cause of preventable death in the United States (US) (Centers for Disease Control and Prevention, 2008) with the majority of adult smokers starting prior to age 18 (Health & Human, 2012; U.S. Department of Health Human Services, 2014). Tobacco use in adolescence reliably predicts being a smoker as an adult (Chassin, Presson, Sherman, & Edwards, 1990), supporting the need for focused research and improved cessation efforts targeting adolescent and emerging adult smokers. Recent data show that current (i.e., past month) use of cigarettes among high school students was approximately 9.2% (Grades 9–12) in the US (Arrazola et al., 2015). Grade-specific estimates of past month cigarette use were shown to be similar (7.2% for 10th and 13.6% for 12th grade students) (Johnston, O'Malley, Meiech, Bachman, & Schulenberg, 2015). Among young adults aged 18–24 years, past month cigarette use is estimated at 18.7% (Jamal et al., 2014). Over half (57%) of adolescent and emerging adult smokers have intentions of quitting (Tworek et al., 2014), and 50–77% have made serious, past-year quit attempts (Bancej, O'Loughlin, Platt, Paradis, & Gervais, 2007; Eaton et al., 2012; Hollis, Polen, Lichtenstein, & Whitlock, 2003; Tworek et al., 2014). However, only 4–6% of unassisted quit attempts among this population are shown to be successful (Centers for Disease Control and Prevention, 2006; Chassin, Presson, Pitts, & Sherman, 2000; W. R. Stanton, McClelland, Elwood, Ferry, & Silva, 1996; Sussman, Lichtman, Ritt, & Pallonen, 1999; Zhu, Sun, Billings, Choi, & Malarcher, 1999), and use of evidence-based treatments and pharmacotherapy is only slightly better (Gray et al., 2011; Gray, Carpenter, Lewis, Klintworth, & Upadhyaya, 2012; Killen et al., 2004; A. Stanton & Grimshaw, 2013; Sussman, Sun, & Dent, 2006). These findings illustrate that young smokers are motivated to quit but do not engage in or with effective cessation support.

Mobile health (mHealth) technology is uniquely suited to address research and treatment gaps within this population, and offers advantages to understand smoking outside of the clinical or research environment in several ways. First, young smokers often face challenges in attending clinic visits, which contributes to study drop-out and missing data. Diminished availability of outcome data leads to inadequately powered trials that continue to constrain the treatment literature (Backinger et al., 2003; Skara & Sussman, 2003; Sussman, 2002). Second, mHealth technology allows for data collection in real-time and in ecologically valid settings, thus providing a more detailed and accurate understanding of smoking. Work in this area began with ecological momentary assessment (EMA), procedures and outcomes of which are now well established in the field (S. Shiffman, 2005; Saul Shiffman, Stone, & Hufford, 2008). Additional innovations now allow for the remote collection and monitoring of carbon monoxide (CO) (Dallery & Glenn, 2005; Hertzberg et al., 2013; Meredith et al., 2014), and the detection of individual puffs through proxies of use, such as arm movements

and respiration (Ali et al.; Raiff, Karata, McClure, Pompili, & Walls, 2014; Sazonov, Lopez-Meyer, & Tiffany, 2013). Remote monitoring offers the opportunity to study health processes at a more granular level, and with the possibility of unobtrusive sensing that may minimize respondent burden and allow for dynamic, interactive approaches. Third, mHealth technology holds the potential to contribute to the delivery, availability, and fidelity of treatment to smokers attempting to quit. Work has been proposed or conducted incorporating mHealth methodology into smoking treatment as a means to engage the individual and provide support in real-time. This has been done through text messaging (Whittaker et al., 2012) and ecological momentary interventions (Heron & Smyth, 2010). Also, work is ongoing to incorporate several features of monitoring and intervention delivery at critical moments in the natural environment (McClernon & Roy Choudhury, 2013). The eventual goal of much of this work is to improve the efficacy and reach of interventions that can be delivered in real-time to improve the likelihood of long-term abstinence.

Adolescents and emerging adults are ideally suited for technology integration into research, and show greater technology utilization compared to other age groups (Lenhert, Ling, Campbell, & Purcell, 2010; Zickuhr, 2011). Among young adults (ages 18–29), 85% are smartphone users, and of these, approximately 15% report that smartphones are their primary means to online access (Pew Research Center, 2015). For those between the ages of 12–17, smartphone use was approximately 47% in 2013 (Madden, Lenhert, Duggan, Cortesi, & Gasser, 2013). Virtually all smartphones are already equipped with features, capabilities, and the necessary computing power to serve as a platform for monitoring technology and intervention delivery.

While high rates of technology utilization among this population may be harnessed as a tool for better understanding of smoking and delivery of treatment, little is known regarding the acceptability and feasibility of mHealth integration to study smoking. Assessing attitudes, interest, and concerns among this target population is critical prior to implementation of mHealth techniques. Identification of characteristics that may predict greater acceptability of mHealth methods and platforms may facilitate the development of acceptable, tailored smoking cessation and relapse prevention tools among sub-groups. Therefore, this study aimed to characterize a broad array of usage, attitudes, and perceptions related to technology-based treatment and the remote monitoring of smoking among adolescent and emerging adult daily cigarette smokers. The survey used in this report was intentionally broad and covered the areas of; the remote assessment of behavior, remote collection of smoking biomarkers (i.e., breath CO), and the remote delivery of treatment for smoking. Specifically, this study aimed to; 1) characterize technology use among this group; 2) assess perceptions, attitudes, and interest in remote monitoring for smoking research and treatment and remote biomarker collection, and 3) determine characteristics that predicted greater acceptability technology for smoking research and treatment.

Methods

Participants

Participants enrolled in a 12-week smoking cessation pharmacotherapy clinical trial (NCT01509547; PI Gray) were approached to complete a questionnaire, typically during the

randomization study visit. Participants eligible for the parent study were daily smokers (5 cigarettes per day over the past 6 months) between the ages of 14–21 years who were interested in making a quit attempt, and had at least one failed quit attempt in their lifetime. Participants were excluded if they had any unstable psychiatric or medical disorder, had any history with suicidal ideation or attempts, were pregnant or breastfeeding, or taking other smoking cessation medications. No additional inclusion or exclusion criteria were implemented for this survey. Administration of the technology questionnaire took place from December 2012 through January 2015 (study recruitment for the parent trial is still ongoing). All procedures were approved by the Institutional Review Board at the Medical University of South Carolina.

Measures

Since we know of no validated surveys to assess smoking-specific technology attitudes, perceptions, and acceptability, a 46 item survey was developed locally. Participants were asked about their use of various forms of technology (mobile phones, Internet, computer, email, social media; 21 items), their interest and concerns regarding the use of technology for the remote monitoring of smoking, remote biomarker collection through breath CO, and treatment delivery (15 items), and the perceived ease of remotely monitoring their smoking and CO (10 items). Among the questions pertaining to interest, concerns, and perceived ease of technology-based treatment, questions and response options were closed-ended. Response options for these items are listed as part of Table 3 below.

Several demographic and smoking-related measures were collected as well. Demographic questions assessed age, gender, education, income, race, and ethnicity. Several smoking measures were also included. A 30-day Timeline Follow-Back (TLFB) (Sobell, Sobell, Leo, & Cancilla, 1988) to assess cigarettes per day was conducted at screening, which has been validated among adolescent smokers (Lewis-Esquerre et al., 2005). Smoking history questions assessed years of regular smoking, age of first cigarette, and number of serious quit attempts. Breath CO and urine cotinine at screening were collected, as well as the modified Fagerström Tolerance Questionnaire (mFTQ) (Prokhorov et al., 2000) and questions to assess the participants' readiness and confidence to quit smoking. Readiness and confidence questions were locally developed and were on a 10-point Likert scale (i.e., "On a scale of 1–10, with 1 being not ready and 10 being extremely ready, how ready are you to quit smoking?").

Statistical Analyses

The survey was administered to 87 participants enrolled in the parent study. Standard descriptive statistics were used to summarize demographic and smoking characteristics. Means and standard deviation are presented for continuous characteristics, while frequency distributions are presented for categorical characteristics. Since this questionnaire constituted an exploratory analysis, possible correlates of favorable technology attitudes and interest were selected from baseline demographic and smoking characteristics as well as technology utilization responses (i.e., gender, race, current and past smoking characteristics, smartphone use and unlimited data). Binary outcome items (yes, no) were analyzed using logistic regression and ordinal outcomes (Not helpful → Very helpful) were analyzed using

ordinal logistic regression. Categorical outcomes that were not ordinal (yes, no, not sure) were analyzed using generalized logistic regression. For all ordinal logit models, the proportional odds assumption was tested and when proportional odds could not be verified, the data were analyzed using generalized logit models. Items with small cell counts (≤ 5) had categories collapsed into logical groups. Results from logistic regression models are presented as odds ratios and associated 95% confidence intervals [OR (95% CI)]. All statistical analyses were performed using the SAS System version 9.3.

Results

Demographic and Smoking Characteristics

Of the 87 participants who completed the questionnaire, the average (SD) age was 18.9 (1.4) years, and the sample was primarily male (58/87; 67%), Caucasian (64/87; 74%), and approximately 68% had graduated from high school (59/87). On average, participants smoked 11.7 (7.6) cigarettes per day, had breath CO readings of 14.2 parts per million (ppm) (8.4) at screening, and urinary cotinine values (n=60) of 1047 ng/ml (619). Nicotine dependence scores averaged 4.4 (1.7) and nearly a quarter of the participants reported substantial nicotine dependence (mFTQ ≥ 6). Participants had been regularly smoking since age 16.2 (1.7) and more than half lived with another smoker (49/87; 56%). Participants were generally motivated to quit smoking, with readiness and confidence scores (on a 10-point scale) averaging 7.7 (1.8) and 7.0 (2.4) respectively.

Technology Utilization

Technology use characteristics are shown in Table 1. As expected for this study sample, technology use was high. Nearly all of the study participants endorsed owning a mobile phone (82/87; 94%) and those who did not own a mobile phone had access to one on a regular basis. All but one participant had the ability to send and receive short message service (SMS) text messages and 93% of participants had smartphones with internet capabilities (81/87). Over half of the study participants reported unlimited data on their mobile phones (45/87; 52%) and the majority reported having yearly contracts (44/87; 51%) and having never changed their mobile phone number (48/87; 55%).

Computer, Internet, email and social media use was also high in this sample. The majority of participants reported using the Internet (85/87; 98%), email (72/87; 83%) and social media (82/87; 94%) on a weekly basis. The most frequently endorsed social media sites used by participants were Facebook (81/87; 93%), Instagram (35/87; 40%) and Twitter (30/87; 35%). Weekly computer use was the least utilized (67/87; 77%), and 65% of participants reported that their mobile phone is the most frequent way that they access the Internet (55/87).

Perceived Ease of Remotely Monitoring Smoking

When participants were asked about the perceived ease of using a remote monitoring technology system to report on their smoking (consisting of remote breath CO monitoring), they were generally favorable in their responses. Responses on the perceived ease of use of remote monitoring technology are shown in Table 2 as median ratings and percentage

distributions of scores for 10-point scale items and percentage distribution for 4-point scale items. Some items were reverse scored and are noted in the table. Specifically, participants responded favorably to being able to carry necessary devices with them on a daily basis and to return study devices. Privacy concerns were relatively low with a median score of 4 (out of 10), while confidentiality concerns were slightly higher (6 out of 10). Participants also endorsed the likelihood of being able to complete remote sessions in a timely fashion, and in a private space.

Attitudes and Interest in Technology for Smoking

Responses regarding attitudes, interest, and concerns with technology for smoking are shown in Table 3. Despite high rates of mobile phone, Internet, email, and computer use, only 33% (29/87) reported that they had ever searched for smoking cessation resources online, and even fewer had ever used health related or self-help applications (apps) on their phones (24/87; 28%). About a quarter of the participants stated that they had no interest in using computer-based smoking cessation counseling (21/87; 24%) and 20% of participants expressed no interest in receiving mobile-phone based cessation counseling (18/87). Nearly half of the sample (42/87; 48%) endorsed being interested in mobile-phone counseling, with far fewer being interested in computer-based counseling (25/87; 29%).

A large percentage of participants felt that mobile phones could be at least somewhat helpful in getting support during a quit attempt (73/87; 84%), and also felt that a quit smoking app may help to motivate them (81/87; 93%). Despite this, about half of the sample still preferred face-to-face counseling exclusively for quitting smoking (44/87; 51%), and most felt that treatment delivered through the Internet would be less effective than in-person treatment (54/87; 62%), though most also reported that Internet-delivered treatment would be more convenient (46/87; 53%). About half of the sample said that they had no concerns regarding technology-based treatment for smoking cessation (43/87; 50%) and remote monitoring of their smoking (52/87; 60%). The most frequent concern for technology-based treatment was that it wouldn't help them to quit (22/87; 25%).

Predictors of Technology Acceptability

Demographic, smoking, and technology characteristics were explored as potential predictors of more favorable acceptability towards technology-based smoking treatment. Several results suggest that smokers with greater nicotine dependence and/or use history were more favorable towards technology integration. First, those with greater dependence (mFTQ scores) were more likely to endorse Internet-delivered treatment as being more effective than in-person treatment (OR=1.35; 95% CI=1.05–1.74; p=0.021). Second, those who had started smoking regularly at a younger age were more likely to have used health-related apps (OR=1.39; 95% CI=1.01–1.91; p=0.043) and were more likely to report computer-based counseling as potentially helpful (OR=1.45; CI=1.04–2.03; p=0.029). Third, smokers with higher CO values (indicative of higher intensity of smoking) were more likely to endorse greater interest in technology-based treatment (OR=1.08; 95% CI=1.01–1.16; p=0.045.); p=0.037). In contrast, those with an earlier age of first cigarette use were less likely to endorse Internet-delivered treatment as being more effective than in-person treatment (OR=0.83; 95% CI=0.69–0.99; p=0.037). Demographically, Caucasian participants were

more likely to endorse that Internet-based treatment would be more convenient compared to non-Caucasian participants ($p=0.025$). Participants who owned smartphones and had unlimited data on their phones were both a) more likely to endorse interest in computer-based cessation (OR=3.33; 95% CI=1.22–9.13; $p=0.019$), more likely to feel that cell phones could be useful when quitting smoking (OR=14.2; 95% CI=2.30–87.8; $p=0.004$) and b) endorse smoking apps as motivating (OR=11.5; 95% CI=1.89–69.9; $p=0.008$).

Discussion

The purpose of this study was to assess technology utilization, perceptions, attitudes, comfort and interest in remote monitoring and technology-based systems for smoking among a treatment-seeking, nicotine dependent sample of adolescents and emerging adults. Exploratory analyses identified potential characteristics that may predict greater acceptability of technology integration. Generally, technology utilization was high for these participants in all forms, which would suggest that they are ideal candidates for technology integration into research and treatment focused on smoking cessation. Despite this, use of technology in the form of apps or Internet searches for information, treatment or tips to quit smoking was low. Participants expressed moderately high interest for technology-based systems for smoking. Results also showed that those with smartphones, unlimited data, greater nicotine dependence and smoking severity viewed technology-based treatment more favorably, with the only exception being for those with a younger age of first cigarette use. It should be noted, however, that predictive analyses were exploratory and significant relationships are interpreted with caution.

These results seem to favor the development and use of mobile-based tools or systems to study and treat smoking. Among this study sample, participants were more interested in mobile-based cessation compared to computer-based programs. This is not surprising given that for many participants, primary access to the Internet was through mobile devices. Also, this study sample showed consistency in mobile phone use and low rates of changing phone numbers. This may suggest that a younger population is less likely to use pay-as-you-go phones that would result in frequent phone number changes, which is a limitation to mobile-based systems. However, it is possible that many study participants may have still been part of a family mobile phone plan, thus contributing to the stability of their mobile access and number. Given that mobile phones are so prevalent among adolescents and young adults, remote monitoring systems that can be incorporated or delivered through mobile platforms are highly desirable, and may help to reduce the burden associated with study participation, data collection, biomarker collection and analysis, and treatment delivery.

This survey study was part of a larger smoking cessation clinical trial (NCT01509547; PI Gray), and as such, participants were motivated to quit smoking and had experienced a failed quit attempt. Even though these participants were treatment-seeking, unfavorable or ambivalent ratings regarding technology-based treatment were still present. For example, 20% and 25% of the sample had no interest in mobile- or computer-based counseling for smoking, respectively. Many more participants said they were “not sure” if they were interested in mobile- (31%) or computer-based counseling (47%), suggesting that this sub-sample is unlikely to engage with technology-based treatment strategies. Additionally, 25%

of the sample felt that technology-based treatment wouldn't help them to quit, which was the most commonly endorsed concern regarding technology-based treatment. These results could have several explanations. First, this may be due to the particular wording of the questions and a lack of concrete examples of the systems being described. Perhaps demonstrating a technology-based system to a user would provide more meaningful measures of acceptability and interest. Second, these data may reflect perceptions that participants have regarding how effective technology-based resources are to quit smoking. Many currently available online and mobile resources are not necessarily evidence-based, which may contribute to perceptions of inefficacy. For example, content analyses of iPhone and Android apps reveal low adherence to evidence-based strategies for quitting smoking (Abrams, Lee Westmaas, Bontemps-Jones, Ramani, & Mellerson, 2013; Abrams, Padmanabhan, Thaweethai, & Phillips, 2011; Bennett et al., 2014), though several apps use strategies to promote behavioral self-monitoring in the form of tracking cigarettes smoked (Bennett et al., 2014). Encouraging adolescents and emerging adults to track and monitor their smoking may be a useful component of a comprehensive intervention or part of in-person treatment, but may not be efficacious independently. It is possible that the self-monitoring of behavior would allow for the collection and use of data specific to the individual that could be used in treatment to encourage and track smoking reduction, understand and avoid triggers, etc. Even in instances where mobile app efficacy is established for smoking cessation among this population, usability and acceptability of these apps will remain a hurdle in their dissemination. It will be essential in the development and evaluation of apps to monitor use and determine which components are most liked and helpful. Also, mobile apps should be developed to be as personalized for the individual as possible, in order to increase efficacy and engagement.

The integration of technology into research and treatment holds great potential as the landscape of novel tobacco products and other substance use changes. Previous work has been done to remotely monitor cigarette smoking through self-report, biochemical verification, and monitoring systems that detect proxies of smoking (Ali et al.; Dallery & Glenn, 2005; Dallery, Raiff, & Grabinski, 2013; Raiff et al., 2014; Sazonov et al., 2013; Saul Shiffman et al., 2008). Technology integration should be pursued to incorporate measures of other tobacco and drug use into remote monitoring systems. This is justified, given that cigarette smoking continues to decline in young smokers (Arrazola et al., 2015; Johnston et al., 2015), while the use of other products are on the rise. For example, use of electronic cigarette (e-cigs) and vaping are consistently on the rise in a younger population (Arrazola et al., 2015; Johnston et al., 2015). For feasibility purposes, remote monitoring and intervention delivery may only be focused on one particular tobacco product, but this may not be sufficient since novel products are gaining popularity at a rapid pace. Research must focus on how best to quantify, monitor, and treat use of novel tobacco products, while potentially incorporating remote methods into this work.

There were several limitations to the current study that should be noted. First, this was a relatively small and homogenous convenience sample of participants that may not generalize widely or be adequately representative. Specifically in terms of motivation to quit smoking, our results cannot necessarily generalize to unmotivated smokers. It will be essential for technology-based treatment systems to attempt to engage unmotivated smokers in order to

increase their motivation and confidence in quitting. It is likely that an unmotivated smoker may be even more ambivalent regarding technology-based treatment than our current sample, but this is an important group of young smokers that must not be overlooked with these treatment strategies. Another limitation is that the questions asked of participants were not validated and only queried interest in mostly hypothetical technology-based systems. The responses, therefore, may not translate to actual use of these systems or compliance with their requirements. Hypothetical acceptability was favorable though, providing justification for the pursuit of technology-based systems for this group.

Adolescent and emerging adult smokers are ideally suited for mHealth integration, and our results reveal that this population has high technology utilization and generally favorable attitudes towards remote monitoring and technology-based systems. The greatest barriers demonstrated in this study were specific to ambivalence towards technology-based systems and the perception that those resources may not be effective. Modifying perceptions regarding lack of efficacy is important to address if these systems are to be used with this target population. We also found some evidence that technology acceptability may vary based on certain characteristics, and this should be carefully considered prior to implementation. Technology integration may need to be tailored to meet smokers where they are in terms of technology use, motivation to quit, and what they perceive as most helpful in their quit attempt. Adolescent and emerging adult smokers tend to be accepting of new technology outlets, and this integration should be pursued to accomplish the goal of providing maximally effective and just-in-time smoking cessation interventions to promote long-term abstinence.

Acknowledgments

Special thanks to the research and medical staff at the Addiction Sciences Division at the Medical University of South Carolina. Specifically, we would like to thank Jessica Hinton for database development, Lori Ann Ueberroth, Casey Johnson, Danielle Paquette, Priscilla Muldrow, Caitlin Morris, Jill Underwood, and Taylor York for the successful execution of the parent trial and this survey component.

Funding: Effort was supported by NIDA grants K01 DA036739 (PI, Erin A. McClure), NIDA grant U01DA031779 (PI, Kevin M. Gray), and pilot funding from the American Cancer Society Institutional Research Grant at the Medical University of South Carolina Hollings Cancer Center (ACS IRG 97-2919-14; PI, Erin A. McClure).

References

- Abroms LC, Lee Westmaas J, Bontemps-Jones J, Ramani R, Mellerson J. A content analysis of popular smartphone apps for smoking cessation. *American Journal of Preventive Medicine*. 2013; 45(6):732–736. DOI: 10.1016/j.amepre.2013.07.008 [PubMed: 24237915]
- Abroms LC, Padmanabhan N, Thaweethai L, Phillips T. iPhone apps for smoking cessation: a content analysis. *American Journal of Preventive Medicine*. 2011; 40(3):279–285. DOI: 10.1016/j.amepre.2010.10.032 [PubMed: 21335258]
- Ali, AA., Hossain, SM., Hovsepian, K., Rahman, M., Plarre, K., Kumar, S. mPuff: Automated detection of cigarette smoking puffs from respiration measurements. Paper presented at the Information Processing in Sensor Networks.; 2012.
- Arrazola RA, Singh T, Corey CG, Husten CG, Neff LJ, Apelberg BJ, Caraballo RS. Tobacco use among middle and high school students - United States, 2011–2014. *MMWR Morbidity and Mortality Weekly Report*. 2015; 64(14):381–385. [PubMed: 25879896]

- Backinger CL, McDonald P, Ossip-Klein DJ, Colby SM, Maule CO, Fagan P, Colwell B. Improving the future of youth smoking cessation. *American Journal of Health Behavior*. 2003; 27(Suppl 2):S170–184. Journal Article. [PubMed: 14521244]
- Bancej C, O’Loughlin J, Platt RW, Paradis G, Gervais A. Smoking cessation attempts among adolescent smokers: a systematic review of prevalence studies. *Tobacco control*. 2007; 16(6):e8.doi: 10.1136/tc.2006.018853 [PubMed: 18048598]
- Bennett ME, Toffey K, Dickerson F, Himelhoch S, Katsafanas E, Savage CLG. A review of Android apps for smoking cessation. *Journal of Smoking Cessation*. 2014; Epub ahead of print (4 February 2014). doi: 10.1017/jsc.2014.1
- Centers for Disease Control and Prevention. Use of cessation methods among smokers aged 16–24 years: United States, 2003. *MMWR Morbidity and Mortality Weekly Report*. 2006; 55(50):1351–1354. [PubMed: 17183227]
- Centers for Disease Control and Prevention. *MMWR. Morbidity and mortality weekly report*. Vol. 57. United States: 2008. Smoking-attributable mortality, years of potential life lost, and productivity losses--United States, 2000–2004; p. 1226–1228.
- Chassin L, Presson CC, Pitts SC, Sherman SJ. The natural history of cigarette smoking from adolescence to adulthood in a midwestern community sample: multiple trajectories and their psychosocial correlates. *Health Psychol*. 2000; 19(3):223–231. [PubMed: 10868766]
- Chassin L, Presson CC, Sherman SJ, Edwards DA. The natural history of cigarette smoking: predicting young-adult smoking outcomes from adolescent smoking patterns. *Health Psychol*. 1990; 9(6): 701–716. [PubMed: 2286181]
- Dallery J, Glenn IM. Effects of an Internet-based voucher reinforcement program for smoking abstinence: A feasibility study. *Journal of Applied Behavior Analysis*. 2005; 38(3):349–357. DOI: 10.1901/jaba.2005.150-04 [PubMed: 16270844]
- Dallery J, Raiff BR, Grabinski MJ. Internet-based contingency management to promote smoking cessation: a randomized controlled study. *Journal of Applied Behavior Analysis*. 2013; 46(4):750–764. DOI: 10.1002/jaba.89 [PubMed: 24114862]
- Eaton DK, Kann L, Kinchen S, Shanklin S, Flint KH, Hawkins J, Wechsler H. Youth risk behavior surveillance - United States, 2011. *MMWR Surveillance Summaries*. 2012; 61(4):1–162.
- Gray KM, Carpenter MJ, Baker NL, Hartwell KJ, Lewis AL, Hiott DW, Upadhyaya HP. Bupropion SR and contingency management for adolescent smoking cessation. *J Subst Abuse Treat*. 2011; 40(1): 77–86. DOI: 10.1016/j.jsat.2010.08.010 [PubMed: 20934835]
- Gray KM, Carpenter MJ, Lewis AL, Klintworth EM, Upadhyaya HP. Varenicline versus bupropion XL for smoking cessation in older adolescents: a randomized, double-blind pilot trial. *Nicotine Tob Res*. 2012; 14(2):234–239. DOI: 10.1093/ntr/ntr130 [PubMed: 21778151]
- Heron KE, Smyth JM. Ecological momentary interventions: incorporating mobile technology into psychosocial and health behaviour treatments. *British journal of health psychology*. 2010; 15(Pt 1): 1–39. DOI: 10.1348/135910709x466063 [PubMed: 19646331]
- Hertzberg JS, Carpenter VL, Kirby AC, Calhoun PS, Moore SD, Dennis MF, Beckham JC. Mobile Contingency Management as an Adjunctive Smoking Cessation Treatment for Smokers With Posttraumatic Stress Disorder. *Nicotine Tob Res*. 2013; 15(11):1934–1938. DOI: 10.1093/ntr/ntt060 [PubMed: 23645606]
- Hollis JF, Polen MR, Lichtenstein E, Whitlock EP. Tobacco use patterns and attitudes among teens being seen for routine primary care. *Am J Health Promot*. 2003; 17(4):231–239. [PubMed: 12640779]
- Jamal A, Agaku IT, O’Connor E, King BA, Kenemer JB, Neff L. Current cigarette smoking among adults--United States, 2005–2013. *MMWR Morbidity and Mortality Weekly Report*. 2014; 63(47): 1108–1112. [PubMed: 25426653]
- Johnston, LD., O’Malley, PM., Meiech, RA., Bachman, JG., Schulenberg, JE. *Monitoring the Future national results on drug use: 1975–2014: Overview, key findings on adolescent drug use*. Ann Arbor, MI: Institute for Social Research, University of Michigan; 2015.
- Killen JD, Robinson TN, Ammerman S, Hayward C, Rogers J, Stone C, Schatzberg AF. Randomized clinical trial of the efficacy of bupropion combined with nicotine patch in the treatment of

- adolescent smokers. *Journal of consulting and clinical psychology*. 2004; 72(4):729–735. DOI: 10.1037/0022-006x.72.4.729 [PubMed: 15301658]
- Lenhart, A., Ling, R., Campbell, S., Purcell, K. Teens and mobile phones. Pew Internet and American Life Project. 2010. Retrived on April 1, 2014 from <http://pewinternet.org/Reports/2010/Teens-and-Mobile-Phones.aspx>
- Lewis-Esquerre JM, Colby SM, Tevyaw TO, Eaton CA, Kahler CW, Monti PM. Validation of the timeline follow-back in the assessment of adolescent smoking. *Drug and Alcohol Dependence*. 2005; 79(1):33–43. DOI: 10.1016/j.drugalcdep.2004.12.007 [PubMed: 15943942]
- Madden, M., Lenhart, A., Duggan, M., Cortesi, S., Gasser, U. Teens and Technology 2013. 2013. Retrieved March 21, 2015, from <http://www.pewinternet.org/2013/03/13/teens-and-technology-2013/>
- McClernon FJ, Roy Choudhury R. I am your smartphone, and I know you are about to smoke: the application of mobile sensing and computing approaches to smoking research and treatment. *Nicotine Tob Res*. 2013; 15(10):1651–1654. DOI: 10.1093/ntr/ntt054 [PubMed: 23703731]
- Meredith SE, Robinson A, Erb P, Spieler CA, Klugman N, Dutta P, Dallery J. A mobile-phone-based breath carbon monoxide meter to detect cigarette smoking. *Nicotine Tob Res*. 2014; 16(6):766–773. DOI: 10.1093/ntr/ntt275 [PubMed: 24470633]
- Pew Research Center. The Smartphone Difference. 2015. Retrieved May 22, 2015, from <http://www.pewinternet.org/2015/04/01/us-smartphone-use-in-2015/>
- Prokhorov AV, De Moor C, Pallonen UE, Hudmon KS, Koehly L, Hu S. Validation of the modified Fagerstrom tolerance questionnaire with salivary cotinine among adolescents. *Addict Behav*. 2000; 25(3):429–433. [PubMed: 10890296]
- Raiff B, Karata Ç, McClure E, Pompili D, Walls T. Laboratory Validation of Inertial Body Sensors to Detect Cigarette Smoking Arm Movements. *Electronics*. 2014; 3(1):87–110. [PubMed: 25553255]
- Sazonov E, Lopez-Meyer P, Tiffany S. A wearable sensor system for monitoring cigarette smoking. *J Stud Alcohol Drugs*. 2013; 74(6):956–964. [PubMed: 24172124]
- Shiffman S. Dynamic influences on smoking relapse process. *Journal of personality*. 2005; 73(6): 1715–1748. DOI: 10.1111/j.0022-3506.2005.00364.x [PubMed: 16274451]
- Shiffman S, Stone AA, Hufford MR. Ecological momentary assessment. *Annual Review of Clinical Psychology*. 2008; 4(Journal Article):1–32. DOI: 10.1146/annurev.clinpsy.3.022806.091415
- Skara S, Sussman S. A review of 25 long-term adolescent tobacco and other drug use prevention program evaluations. *Preventive medicine*. 2003; 37(5):451–474. [PubMed: 14572430]
- Sobell LC, Sobell MB, Leo GI, Cancilla A. Reliability of a timeline method: assessing normal drinkers' reports of recent drinking and a comparative evaluation across several populations. *British journal of addiction*. 1988; 83(4):393–402. [PubMed: 3395719]
- Stanton A, Grimshaw G. Tobacco cessation interventions for young people. *Cochrane Database Syst Rev*. 2013; 8:CD003289.doi: 10.1002/14651858.CD003289.pub5
- Stanton WR, McClelland M, Elwood C, Ferry D, Silva PA. Prevalence, reliability and bias of adolescents' reports of smoking and quitting. *Addiction (Abingdon, England)*. 1996; 91(11):1705–1714.
- Sussman S. Effects of sixty six adolescent tobacco use cessation trials and seventeen prospective studies of self-initiated quitting. *Tobacco induced diseases*. 2002; 1(1):35–81. DOI: 10.1186/1617-9625-1-1-35 [PubMed: 19570247]
- Sussman S, Lichtman K, Ritt A, Pallonen UE. Effects of thirty-four adolescent tobacco use cessation and prevention trials on regular users of tobacco products. *Substance Use and Misuse*. 1999; 34(11):1469–1503. [PubMed: 10468104]
- Sussman S, Sun P, Dent CW. A meta-analysis of teen cigarette smoking cessation. *Health Psychol*. 2006; 25(5):549–557. DOI: 10.1037/0278-6133.25.5.549 [PubMed: 17014271]
- Tworek C, Schauer GL, Wu CC, Malarcher AM, Jackson KJ, Hoffman AC. Youth tobacco cessation: quitting intentions and past-year quit attempts. *American Journal of Preventive Medicine*. 2014; 47(2 Suppl 1):S15–27. DOI: 10.1016/j.amepre.2014.05.009 [PubMed: 25044192]
- U.S. Department of Health and Human Services. Preventing Tobacco Use Among Youth and Young Adults: A Report of the Surgeon General. Atlanta (GA): U.S. Department of Health and Human

Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2012.

U.S. Department of Health Human Services. The Health Consequences of Smoking: 50 Years of Progress. A Report of the Surgeon General. Atlanta (GA): U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2014.

Whittaker R, McRobbie H, Bullen C, Borland R, Rodgers A, Gu Y. Mobile phone-based interventions for smoking cessation. Cochrane database of systematic reviews (Online). 2012; 11:CD006611. Journal Article. doi: 10.1002/14651858.CD006611.pub3

Zhu SH, Sun J, Billings SC, Choi WS, Malarcher A. Predictors of smoking cessation in U.S. adolescents. American Journal of Preventive Medicine. 1999; 16(3):202–207. [PubMed: 10198659]

Zickuhr, K. Generations and their gadgets. Pew Internet and American Life Project. 2011. Retrieved April 5, 2014, from <http://www.pewinternet.org/2011/02/03/generations-and-their-gadgets/>

Table 1

Technology use characteristics.

Mobile phone use	% (N=87)	N (N=87)
Mobile Phone Ownership	94.3	82
Regular Access to a Mobile Phone (do not own)	100	5
Type of Contract		
Pay-as-you-go	10.3	9
Monthly	39.1	34
Yearly	50.6	44
Regular Phone Access > 3 years	78.1	68
Changed Mobile Number (past year)		
Never	55.2	48
1 time	14.9	13
2 times	18.4	16
3 or more times	11.5	10
SMS Text Message Capabilities (send and receive)	98.9	86
Internet Access on Phone - Yes	93.1	81
Unlimited Data - Yes	51.7	45
Uses for Mobile Phone (5 most common listed)		
Text	97.7	85
Phone calls	97.7	85
Social media	85.1	74
Music	85.1	74
Applications (apps)	79.3	69
Email	70.1	61
Computer/Internet/Email Use	% or Mean (N=87)	N or SD (N=87)
Weekly Internet Use - %	97.7	85
Days/Week Internet Use - Mean	6.5	1.3
Sources of Internet Access		
Mobile phone - %	64.7	55
Home computer - %	29.4	25
Other (family/friend cell phone/public library/school) - %	5.9	5
Weekly Computer Use - %	77.0	67
Weekly Email Use - %	82.8	72
Days/Week Email Use - Mean	5.3	2
Weekly Social Media Use - %	94.3	82
Days/Week Social Media Use - Mean	5.7	1.7
Family/Friends on Social Media - %	83.6	19.4
Most Frequently Endorsed Social Media Sites		
Facebook - %	93.1	81
Instagram - %	40.2	35
Twitter - %	34.5	30

Table 2

Perceived ease of using remote monitoring technology.

10-point scale items	Median	1–3 - %	4–6 - %	7–10 - %
Likely to report each cigarette smoked in real time (could not do this → could definitely do this)	7	20	24	56
Accuracy of remembering cigarettes smoked at the end of the day (not accurate → accurate)	7	9	33	57
Ease of carrying 2 devices (CO monitor and phone) (very difficult → very easy)	6	13	41	46
Concerned about privacy (not at all concerned → very concerned)*	4	48	28	24
Concerned about confidentiality (not at all concerned → very concerned)*	6	44	36	21
Would return study devices (would not return → definitely return)	10	1	5	94

4-point scale items	Not at all likely - %	Somewhat likely - %	Moderately likely - %	Definitely likely - %
Able to complete 2–3 remote sessions per day	10	23	33	34
Able to respond immediately to sessions when prompted	19	35	29	17
Carry devices at all times	12	26	26	36
Find a private space to complete sessions	13	25	30	32

Notes:

* indicates reverse scoring for that item on the 1–10 scale

Table 3

Attitudes and interest in technology for smoking.

Previous Tech Use	% (N=87)	N
Used Health or Self-help Apps on Mobile Device - Yes	27.6	24
Used the Internet for Smoking Cessation Counseling, Treatment, Tips - Yes	33.3	29
Interest in Tech		
Interest in Computer-based Smoking Cessation Counseling		
Yes	28.8	25
No	24.1	21
Not sure	47.1	41
Interest in Mobile Phone-based Smoking Cessation Counseling		
Yes	48.3	42
No	20.7	18
Not sure	31.0	27
Smoking Treatment through Mobile Phone or Internet – How Interested?		
Not at all	25.3	22
A little interested	33.3	29
Moderately interested	32.2	28
Very interested	9.2	8
Attitudes and Acceptability		
Computer-based Smoking Cessation – How Helpful?		
Not at all	18.4	16
A little helpful	45.9	40
Moderately helpful	29.9	26
Very helpful	5.8	5
Mobile Phone-based Smoking Cessation – How Helpful?		
Not at all	16.1	14
A little helpful	23.0	20
Moderately helpful	39.1	34
Very helpful	21.8	19
Quit Smoking App – How Motivating?		
Not at all	18.4	16
A little motivating	29.9	26
Moderately motivating	34.5	30
Very motivating	17.2	15
Smoking Cessation Counseling Preference		
Computer only	10.3	9
Face-to-face only	50.6	44
Both computer and face-to-face	27.6	24
No counseling	11.5	10

Previous Tech Use	% (N=87)	N
Internet-delivered treatment vs. In-person Treatment – How effective?		
Less effective	62.1	54
Same	27.6	24
More effective	10.3	9
Internet-delivered treatment vs. In-person Treatment – How convenient?		
Less convenient	23.0	20
Same	24.1	21
More convenient	52.9	46
Comfort with research staff monitoring your smoking through submitted videos		
Sounds very cool	19.5	17
Sounds ok	41.4	36
Don't know	27.6	24
Sounds bad	8.0	7
Sounds awful	3.5	3
Comfort with physician monitoring your smoking through submitted videos		
Sounds very cool	11.5	10
Sounds ok	36.8	32
Don't know	28.7	25
Sounds bad	14.9	13
Sounds awful	8.1	7
Tech Concerns		
Technology-based Treatment Concerns		
Too difficult to access	5.8	5
Too complicated	9.2	8
Too much time	14.9	13
Not confidential enough	12.6	11
Won't help me quit	25.3	22
Might be embarrassing	14.9	13
Other	2.3	2
No concerns	49.4	43
Remote Monitoring Concerns		
Too difficult to access	2.3	2
Too complicated	6.9	6
Too much time	16.1	14
Not confidential enough	18.4	16
Invasion of privacy	11.5	10
Might be embarrassing	13.8	12
Other	1.2	1
No concerns	59.8	52