

SCIENTIFIC REPORTS

OPEN

Corrigendum: A dark green fluorescent protein as an acceptor for measurement of Förster resonance energy transfer

Hideji Murakoshi, Akihiro C. E. Shibata, Yoshihisa Nakahata & Junichi Nabekura

Scientific Reports 5:15334; doi: 10.1038/srep15334; published online 15 October 2015; updated on 07 June 2017

This Article contains typographical errors.

In Table 1 under ‘Förster distance with mEGFP (nm)’, the sREAcH and ShadowG values should read 6.2 and 5.1 respectively.

In addition, in the Materials and Methods section under subheading ‘Plasmid construction’,

‘For construction of mCherry-ERK2 and MEK1 plasmids, the source pCMV5-ratERK2-WT and EYFP-ERK2 plasmids were provided by Natalie Ahn (Addgene plasmid # 40812)²⁹ and by Akihiko Yoshimura, respectively. *mCherry* and *ERK2* or *MEK1* was subcloned into the modified pmEGFP-C1, replacing *mEGFP*.’

should read:

‘For construction of mCherry-ERK2 and MEK1 plasmids, the source pCMV5-ratERK2-WT and EYFP-MEK1 plasmids were provided by Natalie Ahn (Addgene plasmid # 40812)²⁹ and by Akihiko Yoshimura, respectively. *mCherry* and *ERK2* or *MEK1* was subcloned into the modified pmEGFP-C1, replacing *mEGFP*.’



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article’s Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017