

RESEARCH

Open Access



Venoarterial PCO₂-to-arteriovenous oxygen content difference ratio is a poor surrogate for anaerobic metabolism in hemodilution: an experimental study

Arnaldo Dubin*, Gonzalo Ferrara, Vanina Siham Kanoore Edul, Enrique Martins, Héctor Saúl Canales, Carlos Canullán, Gastón Murias, Mario Omar Pozo and Elisa Estenssoro

Abstract

Background: The identification of anaerobic metabolism in critically ill patients is a challenging task. Observational studies have suggested that the ratio of venoarterial PCO₂ (P_{v-a}CO₂) to arteriovenous oxygen content difference (C_{a-v}O₂) might be a good surrogate for respiratory quotient (RQ). Yet P_{v-a}CO₂/C_{a-v}O₂ might be increased by other factors, regardless of anaerobic metabolism. At present, comparisons between P_{v-a}CO₂/C_{a-v}O₂ and RQ have not been performed. We sought to compare these variables during stepwise hemorrhage and hemodilution. Since anemia predictably produces augmented P_{v-a}CO₂ and decreased C_{a-v}O₂, our hypothesis was that P_{v-a}CO₂/C_{a-v}O₂ might be an inadequate surrogate for RQ.

Methods: This is a subanalysis of a previously published study. In anesthetized and mechanically ventilated sheep (*n* = 16), we compared the effects of progressive hemodilution and hemorrhage by means of expired gases analysis.

Results: There were comparable reductions in oxygen consumption and increases in RQ in the last step of hemodilution and hemorrhage. The increase in P_{v-a}CO₂/C_{a-v}O₂ was higher in hemodilution than in hemorrhage (1.9 ± 0.2 to 10.0 ± 0.9 vs. 1.7 ± 0.2 to 2.5 ± 0.1, *P* < 0.0001). The increase in P_{v-a}CO₂ was lower in hemodilution (6 ± 0 to 10 ± 1 vs. 6 ± 0 to 17 ± 1 mmHg, *P* < 0.0001). Venoarterial CO₂ content difference and C_{a-v}O₂ decreased in hemodilution and increased in hemorrhage (2.6 ± 0.3 to 1.2 ± 0.1 vs. 2.8 ± 0.2 to 6.9 ± 0.5, and 3.4 ± 0.3 to 1.0 ± 0.3 vs. 3.6 ± 0.3 to 6.8 ± 0.3 mL/dL, *P* < 0.0001 for both). In hemodilution, P_{v-a}CO₂/C_{a-v}O₂ increased before the fall in oxygen consumption and the increase in RQ. P_{v-a}CO₂/C_{a-v}O₂ was strongly correlated with Hb (*R*² = 0.79, *P* < 0.00001) and moderately with RQ (*R*² = 0.41, *P* < 0.0001). A multiple linear regression model found Hb, RQ, base excess, and mixed venous oxygen saturation and PCO₂ as P_{v-a}CO₂/C_{a-v}O₂ determinants (adjusted *R*² = 0.86, *P* < 0.000001).

Conclusions: In hemodilution, P_{v-a}CO₂/C_{a-v}O₂ was considerably increased, irrespective of the presence of anaerobic metabolism. P_{v-a}CO₂/C_{a-v}O₂ is a complex variable, which depends on several factors. As such, it was a misleading indicator of anaerobic metabolism in hemodilution.

Keywords: Hemodilution, Hemorrhage, Anaerobic metabolism, Oxygen, Carbon dioxide, Respiratory quotient

*Correspondence: arnaldodubin@gmail.com

Cátedra de Farmacología Aplicada, Facultad de Ciencias Médicas,
Universidad Nacional de La Plata, Calle 42 N° 577, 60 y 120, 1900 La Plata,
Argentina

Background

The identification of anaerobic metabolism in critically ill patients can be elusive. Hyperlactatemia, central venous oxygen saturation, or isolated values of oxygen transport and consumption (DO_2 and VO_2) are frequently misleading indicators of tissue hypoxia. In contrast, the acute increase in respiratory quotient (RQ) is an excellent marker of ongoing anaerobic metabolism in exercise [1] and oxygen supply dependency conditions [2, 3]. In both circumstances, there is an excess of carbon dioxide production (VCO_2) compared to VO_2 . This is the result of anaerobic VCO_2 , which arises from the bicarbonate buffering of anaerobically generated protons [1]. The proper measurement of RQ, however, requires analysis of expired gases. This monitoring is not usually available in the critical care setting. Recently, some observational studies have suggested that the ratio of venoarterial PCO_2 ($\text{P}_{\text{v-a}}\text{CO}_2$) to arteriovenous oxygen content difference ($\text{C}_{\text{a-v}}\text{O}_2$) might be a good surrogate for RQ. Accordingly, high $\text{P}_{\text{v-a}}\text{CO}_2/\text{C}_{\text{a-v}}\text{O}_2$ has been associated with hyperlactatemia [4], decreased lactate clearance [5, 6], oxygen supply dependency [7, 8], and worse outcome of critically ill patients [4]. Nevertheless, $\text{P}_{\text{v-a}}\text{CO}_2/\text{C}_{\text{a-v}}\text{O}_2$ might theoretically be increased by several other factors irrespective of the presence of anaerobic metabolism. Moreover, comparisons between $\text{P}_{\text{v-a}}\text{CO}_2/\text{C}_{\text{a-v}}\text{O}_2$ and RQ have not been performed yet.

Given the increasing number of publications about the $\text{P}_{\text{v-a}}\text{CO}_2/\text{C}_{\text{a-v}}\text{O}_2$ and the lack of an adequate validation, further research is needed. This study was derived from a secondary subanalysis of a previous publication that sought to determine the relationship among oxygen transport, microvascular perfusion, and tissue CO_2 in ischemic and anemic hypoxia [9]. The present investigation was focused on the behavior of $\text{P}_{\text{v-a}}\text{CO}_2/\text{C}_{\text{a-v}}\text{O}_2$ and its determinants, as well as its relationship with RQ, during stepwise hemorrhage and hemodilution. Since a progressive hemodilution, which does not compromise aerobic metabolism, will predictably result in increased $\text{P}_{\text{v-a}}\text{CO}_2$ [10] and decreased $\text{C}_{\text{a-v}}\text{O}_2$ [11], our hypothesis was that $\text{P}_{\text{v-a}}\text{CO}_2/\text{C}_{\text{a-v}}\text{O}_2$ might be an inadequate surrogate for RQ in isovolemic anemia.

Methods

Anesthesia and ventilation

Sixteen sheep (20 ± 2 kg, mean \pm SEM) were anesthetized with 30 mg/kg of sodium pentobarbital, intubated, and mechanically ventilated with a Servo Ventilator 900C (Siemens-Elema AB, Solna, Sweden) with a tidal volume of 15 mL/kg, a FiO_2 of 0.21 and a positive end-expiratory pressure of 6 cm H_2O . The initial respiratory rate was set to keep the arterial PCO_2 between 35 and 40 mmHg. This respiratory setting was maintained during the rest

of the experiment. Neuromuscular blockade was performed with pancuronium bromide (0.06 mg/kg). Additional pentobarbital boluses (1 mg/kg) were administered hourly and when clinical signs of inadequate depth of anesthesia were evident. Analgesia was provided by fentanyl as a bolus of 2 $\mu\text{g}/\text{kg}$, followed by 1 $\mu\text{g}/\text{h}/\text{kg}$. These drugs were administered intravenously.

Surgical preparation

A 7.5-French Swan-Ganz standard thermodilution pulmonary artery catheter (Edwards Life Sciences, Irvine, CA, USA) was inserted through an introducer in the right external jugular vein to obtain mixed venous samples; its side port was used to administer fluids and drugs. Catheters were placed in the descending aorta via the left femoral artery to measure blood pressure, perform the bleeding, and obtain blood samples, and in the inferior vena cava to infuse fluids during isovolemic hemodilution.

Measurements and derived calculations

VO_2 , VCO_2 , and RQ were measured by analysis of expired gases (MedGraphics CPX Ultima, Medical Graphics Corporation, St. Paul, MN). VO_2 and VCO_2 were adjusted to body weight.

Arterial and mixed venous PO_2 , PCO_2 , pH, Hb, and O_2 saturation were measured with a blood gas analyzer and a co-oximeter (ABL 5 and OSM 3, Radiometer, Copenhagen, Denmark). $\text{C}_{\text{a-v}}\text{O}_2$ was calculated by standard formulae.

Cardiac output was calculated as VO_2 divided by $\text{C}_{\text{a-v}}\text{O}_2$. DO_2 was calculated as cardiac output multiplied by arterial O_2 content.

We also calculated $\text{P}_{\text{v-a}}\text{CO}_2$ and $\text{P}_{\text{v-a}}\text{CO}_2/\text{C}_{\text{a-v}}\text{O}_2$. According to Fick's principle, venoarterial CO_2 content difference ($\text{C}_{\text{v-a}}\text{CO}_2$) was calculated as VCO_2 divided by cardiac output.

Experimental procedure

Basal measurements were taken after a period of no less than 30 min after systemic VO_2 and VCO_2 became stable. Animals were then assigned to hemodilution ($n = 8$) and hemorrhage ($n = 8$) groups. In the hemodilution group, we performed a stepwise hemodilution through isovolemic exchange of blood with 6% hydroxyethyl starch 130/0.4 in 0.9% NaCl (Voluven, Fresenius Kabi, Bad Homburg, Germany). The amount of blood exchanged to reach desired levels of hematocrit of about 0.15, 0.10, and 0.05 in each step was estimated as previously referred [12]. In the hemorrhage group, consecutive bleedings of 5–10 mL/kg were performed. Similar reductions in systemic VO_2 were pursued in both groups in order to reach comparable degrees of anaerobic metabolism.

Measurements were taken at 30, 60, and 90 min. Blood temperature was kept constant throughout the study with a heating lamp.

At the end of the experiment, the animals were killed with an additional dose of pentobarbital and a KCl bolus.

Data analysis

Data were assessed for normality and expressed as mean ± SEM. Groups were compared with two-way repeated measures of ANOVA. After a $P < 0.05$ for time × group interaction, a post hoc Student’s *t* test with Bonferroni correction was used for pairwise comparisons. Simple linear regression analysis with $P_{v-a}CO_2/C_{a-v}O_2$ as the outcome variable was conducted, and variables showing a P value <0.20 or physiologically plausible were entered in a multiple linear regression model. The final model was tested for the presence of collinearity (VIF test). All analyses were done with Stata statistical software (Stata Corporation, Release 12, College Station, TX, USA).

Results

In both groups, DO_2 fell progressively. In the hemorrhage group, the decrease in DO_2 was primarily related to the reduction in cardiac output from 166 ± 13 to 54 ± 6 mL/min/kg ($P < 0.0001$). In addition, Hb fell from 8.4 ± 0.5 to 6.6 ± 0.4 g/dL ($P < 0.0001$). In hemodilution group, the drop in DO_2 was completely explained by the reduction in Hb from 8.3 ± 0.4 to 1.2 ± 0.1 g/dL ($P < 0.0001$). Cardiac output concurrently increased from 165 ± 16 to 373 ± 41 mL/min/kg ($P < 0.0001$).

In the last stage, there were similar decreases in VO_2 and increases in RQ in both groups. $P_{v-a}CO_2/C_{a-v}O_2$ also increased in the last stage in the hemorrhage group. $P_{v-a}CO_2/C_{a-v}O_2$ increased after the second step in the

hemodilution group, and the increases were higher than in hemorrhage group (Fig. 1).


$P_{v-a}CO_2/C_{a-v}O_2$ was strongly correlated with Hb levels and moderately with RQ (Fig. 2). A similar behavior was observed in hemorrhage group ($R^2 = 0.23, P < 0.002$ and $R^2 = 0.12, P < 0.03$). A multiple linear regression model, developed with data from both groups, found Hb, RQ, base excess, and mixed venous oxygen saturation and PCO_2 as $P_{v-a}CO_2/C_{a-v}O_2$ determinants (adjusted $R^2 = 0.86, P < 0.000001$). Hb was the explanatory variable with the highest independent contribution to the prediction (highest *t* ratio) (Table 1). The model did not exhibit collinearity.

$P_{v-a}CO_2$ increased in the hemorrhage group from the first stage and in hemodilution group only in the last phase. The increases in $P_{v-a}CO_2$ were higher in hemorrhage than in hemodilution, while $C_{v-a}CO_2$ increased in hemorrhage and decreased in hemodilution (Fig. 3). In the hemodilution group, there was a right shift in the relationship between CO_2 pressures and contents (Fig. 4). During reductions in DO_2 , $C_{a-v}O_2$ increased in the hemorrhage group and fell in the hemodilution group (Fig. 3).

Discussion

The main finding of this study was that $P_{v-a}CO_2/C_{a-v}O_2$ failed to properly reflect RQ in hemodilution. It increased before the appearance of the dependency of VO_2 on DO_2 . Its correlation with RQ was moderate, but it showed a strong association with Hb levels. Indeed, $P_{v-a}CO_2/C_{a-v}O_2$ was more explained by Hb levels than by anaerobic metabolism. Changes in the dissociation of CO_2 from Hb mostly account for these results.

Several studies have tried to link $P_{v-a}CO_2/C_{a-v}O_2$ with some events suggestive of anaerobic metabolism such as hyperlactatemia [4], decreased lactate clearance [5, 6],






Fig. 2 Correlation of venoarterial PCO₂ difference-to-arteriovenous O₂ content difference ratio (P_{v-a}CO₂/C_{a-v}O₂) with respiratory quotient (a) and Hb levels (b). The correlation between P_{v-a}CO₂/C_{a-v}O₂ and RQ was statistically significant but moderate. In contrast, P_{v-a}CO₂/C_{a-v}O₂ and Hb levels were strongly correlated

Table 1 Multiple linear regression model for the ratio of venoarterial PCO₂ to arteriovenous oxygen content difference (P_{v-a}CO₂/C_{a-v}O₂)

P _{v-a} CO ₂ /C _{a-v} O ₂	Coefficient	Standard error	t ratio	P value	[95% confidence interval]
Ln hemoglobin (g/dL)	-3.60	0.26	-13.62	<0.000001	-4.13 -3.08
Respiratory quotient	2.43	1.03	2.35	<0.03	0.37 4.49
Base excess (mEq/L)	-0.06	0.03	-2.42	<0.02	-0.12 -0.01
Mixed venous O ₂ saturation (fraction)	0.03	0.01	3.90	<0.0003	0.01 0.04
Mixed venous PCO ₂ (mmHg)	0.15	0.03	4.58	<0.00003	0.08 0.21
Intercept	-0.98	1.72	-0.57	0.57	-4.40 2.44




Fig. 3 Relationship between oxygen transport to venoarterial PCO₂ difference (P_{v-a}CO₂) (a), venoarterial CO₂ content difference (C_{v-a}CO₂) (b), and arteriovenous O₂ content difference (C_{a-v}O₂) (c). Hemodilution produced opposite effects on P_{v-a}CO₂ and C_{v-a}CO₂. C_{v-a}CO₂ decreased in hemodilution and increased in hemorrhage. These changes are the underlying explanation for different behavior of P_{v-a}CO₂/C_{a-v}O₂ in both groups

increased VO₂ in response to fluid challenge [7, 8], and worse outcome [4]. Since RQ was not measured in those studies, it was not clear whether P_{v-a}CO₂/C_{a-v}O₂ effectively reflected the presence of anaerobic metabolism or was only the result of factors that could increase that ratio in the absence of anaerobic metabolism. In fact, P_{v-a}CO₂/

C_{a-v}O₂ is not a straightforward variable. Although related to RQ, it might be hypothetically increased by several factors beyond anaerobic metabolism. Many of the changes in P_{v-a}CO₂/C_{a-v}O₂ might be ascribed to modifications of the CO₂-Hb dissociation curve. Haldane effect, metabolic acidosis, and anemia can increase PCO₂ for




Fig. 4 Changes in the relationship between venoarterial CO₂ pressure and content differences in hemodilution (a) and hemorrhage (b). Hemodilution shifted CO₂Hb dissociation curve to the right

a given CCO₂ [13]. In addition, taking into account the curvilinear characteristics of the dissociation curve, the effects are even greater at higher PCO₂. When the slope of the dissociation curve flattens, substantial increases in P_{v-a}CO₂ may actually represent negligible increases in C_{v-a}CO₂. Therefore, high oxygen venous saturation [14], hyperlactatemia [15], and hemodilution [16] can increase P_{v-a}CO₂ even though C_{v-a}CO₂ remains unchanged.

In line with the previous discussion, our results showed that isovolemic anemia disproportionately increased P_{v-a}CO₂/C_{a-v}O₂, compared to hemorrhage. Furthermore, this ratio was elevated before the beginning of oxygen supply dependency. Progressive hemodilution was associated with opposing effects on P_{v-a}CO₂ and C_{v-a}CO₂: P_{v-a}CO₂ increased and C_{v-a}CO₂ decreased. Previous studies showed that decreasing hemoglobin levels results in widened P_{v-a}CO₂ for a given C_{v-a}CO₂ [16]. In a similar model of progressive hemodilution, the contrasting effects of low Hb levels on P_{v-a}CO₂ and C_{v-a}CO₂ were also noticed [10]. Therefore, increased P_{v-a}CO₂ is a predictable consequence of anemia.

Another expected consequence from hemodilution is the decrease in C_{a-v}O₂ [11]. Increases in oxygen extraction always occur in response to reductions in DO₂, irrespective of the mechanism of oxygen supply limitation. The impact of the increase in oxygen extraction on C_{a-v}O₂, however, depends on cardiac output. According to Fick's principle, C_{a-v}O₂ should widen in conditions of low cardiac output and decreased in states of reduced DO₂ with increased cardiac output, if VO₂ remains constant. Our study also confirmed this assumption.

As a result of the opposite effects of hemodilution on P_{v-a}CO₂ and C_{a-v}O₂, the ratio between both variables markedly augmented in the absence of anaerobic metabolism. The increase in P_{v-a}CO₂/C_{a-v}O₂ was even

higher during the oxygen supply dependency, due to the interplay of the aforementioned factors and the ongoing anaerobic CO₂ production.

Considering the coefficient of determination of the regression ($R^2 = 0.41$), RQ only explains a minor part of the P_{v-a}CO₂/C_{a-v}O₂ variability. As supported by the results of the multiple linear regression model, P_{v-a}CO₂/C_{a-v}O₂ is a complex variable that has several determinants. Although Hb was the main contributor to the prediction of P_{v-a}CO₂/C_{a-v}O₂, it was also influenced by RQ and by the changes in the dissociation of CO₂ from hemoglobin induced by metabolic acidosis and Haldane effect. These effects were magnified at the flattened portion of the CO₂Hb dissociation curve as shown by the impact of mixed venous PCO₂ in the model.

A study has proposed a P_{v-a}CO₂/C_{a-v}O₂ cutoff of 1.4 for the identification of anaerobic metabolism [4]. This suggestion, however, should be carefully interpreted. The development of anaerobic metabolism is identified by acute increases in RQ, not by isolated values [1–3]. Actually, the normal range of RQ is 0.67–1.30 [17] depending also on other factors such as energy source [18] and overfeeding [19]. In our experiments, values of P_{v-a}CO₂/C_{a-v}O₂ during oxygen supply dependency were considerably higher (10.0 ± 2.7 and 2.5 ± 0.4 in hemodilution and hemorrhage groups, respectively).

Our findings do not challenge the value of P_{v-a}CO₂/C_{a-v}O₂ as an outcome predictor of critically ill patients, which was previously described [4]. The composite characteristics of P_{v-a}CO₂/C_{a-v}O₂, however, suggest that the prognostic ability might be mainly related to the interaction of several mechanisms, not only to anaerobic metabolism.

Our study has certain drawbacks. Secondary analyses pose inherent limitations that have been subject to

critiques [20]. In addition, part of our analysis was based on calculations of CCO_2 , not in actual measurements [21]. This last procedure is complex and cumbersome and is not available in our laboratory. Accordingly, we calculated CCO_2 from Fick's principle. We prefer this method, because the different algorithms for computing CCO_2 from blood gases and Hb are frequently misleading and can produce negative $C_{v-a}CO_2$ values. Finally, the experimental model of hemorrhage and hemodilution does not address the applicability of our results to septic conditions.

Conclusions

Hemodilution produced higher increases in $P_{v-a}CO_2/C_{a-v}O_2$, compared to hemorrhage, and this ratio was widened even in the absence of oxygen supply dependency. These findings were related to the effects of anemia on CO_2 Hb dissociation curve and $C_{a-v}O_2$. Our results suggest that $P_{v-a}CO_2/C_{a-v}O_2$ is a multifactorial variable, which results from interactions among anaerobic metabolism, anemia, metabolic acidosis, and Haldane effect. Since it is not an accurate surrogate for RQ, values of $P_{v-a}CO_2/C_{a-v}O_2$ should be cautiously interpreted. Further studies in septic models are needed to confirm the limitations of $P_{v-a}CO_2/C_{a-v}O_2$ in such condition.

Abbreviations

DO_2 : oxygen transport; VO_2 : oxygen consumption; RQ: respiratory quotient; VCO_2 : carbon dioxide production; $P_{v-a}CO_2$: venoarterial PCO_2 ; $C_{a-v}O_2$: arteriovenous oxygen content difference; $P_{v-a}CO_2/C_{a-v}O_2$: ratio of venoarterial PCO_2 to arteriovenous oxygen content difference; $C_{v-a}CO_2$: venoarterial CO_2 content difference.

Authors' contributions

AD, GF, VSKE, EM, HSC, CC, GM, and MOP carried out the animal experiments and participated in the design of the study. AD performed the statistical analysis and drafted the manuscript. EE participated in the study design, statistical analysis, and interpretation of the data. All authors discussed the results and participated in the writing. All authors read and approved the final manuscript.

Acknowledgements

None.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

The data set supporting the conclusions of this article is available from the corresponding author on reasonable request.

Ethics approval

The study was approved by the local Animal Research Committee [0800-009634/11-000]. Care of animals was in accordance with National Institutes of Health (USA).

Funding

This study was supported by the Grant PICT 2010-00495, Agencia Nacional de Promoción Científica y Tecnológica, Argentina.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 24 November 2016 Accepted: 4 June 2017

Published online: 12 June 2017

References

- Wasserman K, Whipp BJ, Koyl SN, Beaver WL. Anaerobic threshold and respiratory gas exchange during exercise. *J Appl Physiol*. 1973;35:236–43.
- Cohen IL, Sheikh FM, Perkins RJ, Feustel PJ, Foster ED. Effect of hemorrhagic shock and reperfusion on the respiratory quotient in swine. *Crit Care Med*. 1995;23:545–52.
- Dubin A, Murias G, Estenssoro E, Canales H, Sottile P, Badie J, Barán M, Rossi S, Laporte M, Pálizas F, Giampieri J, Mediavilla D, Vacca E, Botta D. End-tidal CO_2 pressure determinants during hemorrhagic shock. *Intensive Care Med*. 2000;26:1619–23.
- Mekontso-Dessap A, Castelain V, Anguel N, Bahloul M, Schauvliege F, Richard C, Teboul JL. Combination of venoarterial PCO_2 difference with arteriovenous O_2 content difference to detect anaerobic metabolism in patients. *Intensive Care Med*. 2002;28:272–7.
- Mesquida J, Saludes P, Gruartmoner G, Espinal C, Torrents E, Baigorri F, Artigas A. Central venous-to-arterial carbon dioxide difference combined with arterial-to-venous oxygen content difference is associated with lactate evolution in the hemodynamic resuscitation process in early septic shock. *Crit Care*. 2015;19:126.
- He HW, Liu DW, Long Y, Wang XT. High central venous-to-arterial CO_2 difference/arterial–central venous O_2 difference ratio is associated with poor lactate clearance in septic patients after resuscitation. *J Crit Care*. 2016;31:76–81.
- Monnet X, Julien F, Ait-Hamou N, Lequoy M, Gosset C, Jozwiak M, Persichini R, Anguel N, Richard C, Teboul JL. Lactate and venoarterial carbon dioxide difference/arterial–venous oxygen difference ratio, but not central venous oxygen saturation, predict increase in oxygen consumption in fluid responders. *Crit Care Med*. 2013;41:1412–20.
- Mallat J, Lemyze M, Meddour M, Pepy F, Gasan G, Barrailler S, Durville E, Temime J, Vangrunderbeeck N, Tronchon L, Vallet B, Thevenin D. Ratios of central venous-to-arterial carbon dioxide content or tension to arteriovenous oxygen content are better markers of global anaerobic metabolism than lactate in septic shock patients. *Ann Intensive Care*. 2016;6:10.
- Ferrara G, Kanoore Edul VS, Martins E, Canales HS, Canullán C, Murias G, Pozo MO, Estenssoro E, Ince C, Dubin A. Intestinal and sublingual microcirculation are more severely compromised in hemodilution than in hemorrhage. *J Appl Physiol*. 1985;2016(120):1132–40.
- Dubin A, Estenssoro E, Murias G, Pozo MO, Sottile JP, Barán M, Piacentini E, Canales HS, Etcheverry G. Intramucosal–arterial PCO_2 gradient does not reflect intestinal dysoxia in anemic hypoxia. *J Trauma*. 2004;57:1211–7.
- Laks J, Pilon RN, Klovekorn WP, Anderson W, MacCallum JR, O'Connor NE. Acute hemodilution: its effect of hemodynamics and oxygen transport in anesthetized man. *Ann Surg*. 1974;180:103–9.
- Bourke DL, Smith TC. Estimating allowable hemodilution. *Anesthesiology*. 1974;41:609–12.
- Teboul JL, Scheeren T. Understanding the Haldane effect. *Intensive Care Med*. 2016;43:91–3.
- Jakob SM, Kosonen P, Ruokonen E, Parviainen I, Takala J. The Haldane effect—an alternative explanation for increasing gastric mucosal PCO_2 gradients? *Br J Anaesth*. 1999;83:740–6.
- Sun XG, Hansen JE, Stringer WW, Ting H, Wasserman K. Carbon dioxide pressure–concentration relationship in arterial and mixed venous blood during exercise. *J Appl Physiol*. 1985;2001(90):1798–810.
- Chiara C, Giovannini I, Giuliante F, Vellone M, Ardito F, Tenhunen J, Nuzzo G. Significance of hemoglobin concentration in determining blood CO_2 binding capacity in critical illness. *Respir Physiol Neurobiol*. 2010;172:32–6.

17. McClave SA, Lowen CC, Kleber MJ, McConnell JW, Jung LY, Goldsmith LJ. Clinical use of the respiratory quotient obtained from indirect calorimetry. *J Parenter Enter Nutr.* 2003;27:21–6.
18. MacFie J, Holmfield JH, King RF, Hill GL. Effect of the energy source on changes in energy expenditure and respiratory quotient during total parenteral nutrition. *J Parenter Enter Nutr.* 1983;7:1–5.
19. Hulst JM, van Goudoever JB, Zimmermann LJ, Hop WC, Büller HA, Tibboel D, Joosten KF. Adequate feeding and the usefulness of the respiratory quotient in critically ill children. *Nutrition.* 2005;21:192–8.
20. Mariano ER, Ilfeld BM, Neal JM. “Going fishing”—the practice of reporting secondary outcomes as separate studies. *Reg Anesth Pain Med.* 2007;32:183–5.
21. Van Slyke DD, Neill JM. The determination of gases in blood and other solutions by vacuum extraction and manometric measurement. *J Biol Chem.* 1924;61:523–73.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
