



# Gene regulation during *Drosophila* eggshell patterning

George Pyrowolakis<sup>a</sup>, Ville Veikkolainen<sup>a</sup>, Nir Yakoby<sup>b</sup>, and Stanislav Y. Shvartsman<sup>c</sup>

<sup>a</sup>BIOSS Centre for Biological Signalling Studies and Institute for Biology I, Albert-Ludwigs University of Freiburg, 79104 Freiburg, Germany; <sup>b</sup>Department of Biology and Center for Computational and Integrative Biology, Rutgers University, Camden, NJ 08103; and <sup>c</sup>Lewis-Sigler Institute for Integrative Genomics and Department of Chemical and Biological Engineering, Princeton University, Princeton, NJ 08540

Edited by Neil H. Shubin, The University of Chicago, Chicago, IL, and approved February 10, 2017 (received for review August 8, 2016)

**A common path to the formation of complex 3D structures starts with an epithelial sheet that is patterned by inductive cues that control the spatiotemporal activities of transcription factors. These activities are then interpreted by the *cis*-regulatory regions of the genes involved in cell differentiation and tissue morphogenesis. Although this general strategy has been documented in multiple developmental contexts, the range of experimental models in which each of the steps can be examined in detail and evaluated in its effect on the final structure remains very limited. Studies of the *Drosophila* eggshell patterning provide unique insights into the multiscale mechanisms that connect gene regulation and 3D epithelial morphogenesis. Here we review the current understanding of this system, emphasizing how the recent identification of *cis*-regulatory regions of genes within the eggshell patterning network enables mechanistic analysis of its spatiotemporal dynamics and evolutionary diversification. It appears that *cis*-regulatory changes can account for only some aspects of the morphological diversity of *Drosophila* eggshells, such as the prominent differences in the number of the respiratory dorsal appendages. Other changes, such as the appearance of the respiratory eggshell ridges, are caused by changes in the spatial distribution of inductive signals. Both types of mechanisms are at play in this rapidly evolving system, which provides an excellent model of developmental patterning and morphogenesis.**

enhancer | network | evolution | signal | dynamics

The *Drosophila* eggshell is a proteinaceous structure that houses the future embryo and mediates its interaction with the environment (Fig. 1A). It provides a point for sperm entry and controls the gas exchange needed for embryo respiration. The eggshell is derived from the follicular epithelium, a cell sheet that envelops the germ line cyst, comprising one oocyte and 15 nurse cells (Fig. 1B). This cell sheet is patterned by the combined activities of several signaling pathways, including the epidermal growth factor receptor (EGFR) and bone morphogenetic protein (BMP) pathways (1–3). The EGFR pathway is activated by Gurken (GRK), a TGF $\alpha$ -like ligand that is secreted from the oocyte and generates a dorsoventral gradient of EGFR signaling in the follicle cells (4–6) (Fig. 2A). The BMP pathway is activated by the BMP2/4 homolog Decapentaplegic (DPP), which is distributed in the anteroposterior gradient (7) (Fig. 2B). The joint activities of the EGFR and DPP pathways induce the formation of several eggshell structures, including the respiratory dorsal appendages and the operculum, the region of the eggshell from which the larva hatches when the embryogenesis is completed. Importantly, the follicle cells do not divide during the patterning of the dorsal eggshell structures (8). Consequently, the patterning and morphogenesis of these structures can be studied without the added complexities associated with changing cell numbers.

Changes in the EGFR and DPP signaling profiles can cause dramatic alterations of eggshell morphology. For example, uniform activation of the DPP pathway eliminates the appendages and results in eggshells with greatly expanded operculum (Fig. 1C). At the same time, decreasing the levels of EGFR signaling can result in eggshells with one appendage (Fig. 1D). These and many other important observations were made two decades ago, when eggshell morphology was used as a sensitive readout in

genetic screens that discovered the genes involved in the early steps of body axis specification (9, 10). Genetic studies of eggshell patterning identified numerous molecular components that interpret the GRK and DPP signals in the follicle cells (Fig. 2). Most importantly, it was established that each of the two dorsal appendages is derived from a 2D primordium comprising a patch of cells expressing the Zn-finger transcription factor Broad (BR) and an adjacent L-shaped stripe of cells that express *rhomboid* (RHO), which encodes a ligand-processing enzyme in the *Drosophila* EGFR pathway (11). Cells that express BR and RHO form the top and bottom parts of the future dorsal appendages, respectively (Fig. 2C).

Work over the past decade has identified the transcriptional factors that control these genes and has started to connect them in a regulatory network (2, 12–18). Based on this network, we can predict how the expression patterns of multiple genes will respond to a range of genetic perturbations, including the quantitative changes in the distribution and levels of inductive signals (Fig. 2D). In addition, this network provides a starting point for exploring the evolution of eggshell morphology, which varies greatly across drosophilids. For instance, the eggshell of *Drosophila virilis* has four dorsal appendages (Fig. 1E), whereas the eggshell of *Drosophila willistoni* has a new dorsal structure, the dorsal ridge, that extends from the base of the dorsal appendages toward the posterior of the eggshell (Fig. 1F). What caused these changes in eggshell morphologies? Can they be attributed to changes in the inductive signals or in the regulatory network that interprets these inputs in the follicle cells? Here we review the current knowledge of the eggshell patterning network and the ongoing studies of its evolutionary diversification.

## Patterning Network in *D. melanogaster*

Most of our understanding of tissue patterning by inductive signals is derived from studies of patterns in one dimension, such as the stripes of gene expression in the *Drosophila* blastoderm (19). However, the majority of experimentally observed patterns, including those in the *Drosophila* blastoderm, have a strong 2D component, reflecting their joint control by several inductive signals. Eggshell patterning by the EGFR and DPP pathways provides an excellent model for studying the dynamic emergence of 2D gene expression patterns and features several generally applicable regulatory principles (Fig. 2A and B). These principles include the combined effects of two signaling gradients that act through a network of transcription factors, converging on the enhancers of their target genes (Fig. 2D). Another important aspect of this system, which has been revealed only recently, is that


This paper results from the Arthur M. Sackler Colloquium of the National Academy of Sciences, "Gene Regulatory Networks and Network Models in Development and Evolution," held April 12–14, 2016, at the Arnold and Mabel Beckman Center of the National Academies of Sciences and Engineering in Irvine, CA. The complete program and video recordings of most presentations are available on the NAS website at [www.nasonline.org/Gene\\_Regulatory\\_Networks](http://www.nasonline.org/Gene_Regulatory_Networks).

Author contributions: G.P., V.V., N.Y., and S.Y.S. designed research; G.P., V.V., N.Y., and S.Y.S. performed research; G.P., V.V., N.Y., and S.Y.S. analyzed data; and G.P., V.V., N.Y., and S.Y.S. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

<sup>1</sup>To whom correspondence should be addressed. Email: [stas@princeton.edu](mailto:stas@princeton.edu).




**Fig. 1.** *Drosophila* eggshell, a complex structure derived from an epithelial sheet. (A) Scanning electron microscopy (SEM) image of the eggshell of *D. melanogaster*. The most prominent features are the two dorsal appendages (DAs), the micropyle (Mp), and the operculum (Op). (B) The egg chamber midway through oogenesis stained with phalloidin. (C) SEM image of the eggshell resulting from uniform activation of DPP signaling in the follicle cells. (D) SEM image of the eggshell resulting from reduced EGFR signaling. (E) SEM image of the eggshell from *D. virilis*, a species with four dorsal appendages. (F) SEM image of the eggshell from *D. willistoni*, a species with two dorsal appendages and a dorsal ridge (DR). All SEM images present the dorsal views of the eggshell (anterior to the left).

different features of the emerging 2D patterns are established by processes that are separated in time. In this section, we discuss the network of the inductive cues and transcription factors that govern the 2D pattern of BR, which acts as a master regulator of the dorsal appendage formation.

The dorsoventral gradient of EGFR signaling controls *br* through a feedforward loop, a network motif in which EGFR activates both *br* and its repressor (17, 18, 20, 21). The induction threshold for the repressor is higher than that for *br*, which explains why it is expressed only in cells exposed to intermediate levels of GRK. The activating part of the feedforward loop is mediated by the Iroquois transcription factor Mirror (MIRR), and the repressive part relies on the ETS-domain factor Pointed (PNT) (12, 13, 15, 22). Given the dorsoventral pattern of the EGFR activation by GRK (Fig. 2A), this circuit predicts that the *br* expression domain should look like a horseshoe and extend all the way to the anterior border of the oocyte-associated follicle cells. The real pattern is different, however; the horseshoe is broken at the posterior and pushed away from the anterior border (Fig. 2C). These effects are mediated by two distinct repression events, which are separated both in time and in space. The anterior repression depends on the anterior gradient of DPP signaling (2, 17, 23), which controls *br* through a two-tier cascade that involves repression of BRK, a direct target of DPP in multiple stages of *Drosophila* development (24–26). The posterior repression of *br* relies on the earlier phase of EGFR signaling, when the oocyte nucleus is located at the posterior of the oocyte. At this point of oogenesis, the EGFR pathway is activated in a posterior-to-anterior gradient and represses *br* in the posterior half of the follicular epithelium through the T-box transcription factors Midline (MID) and H15 (Fig. 2D) (16, 27, 28).

The effects of these inductive signals and transcription factors were discovered through their effects on the eggshell morphology and the patterns of BR protein, without direct analysis of the transcriptional regulation of the *br* gene. Important aspects of this regulation have been revealed only recently. Specifically, dissection of the genomic region of *br* revealed the existence of two distinct enhancers, early and late, that combine their activities over time to generate the dynamic expression of *br* in the follicle cells (Fig. 3A) (29). An early enhancer (*brE*) drives uniform expression before stage 10 of oogenesis. At stage 10, this enhancer is repressed in a wide dorsal domain and starts to subside in the rest of the follicular epithelium. At the same time, the late enhancer (*brL*) is activated in two lateral domains within the *brE*-free zone, foreshadowing the formation of the two dorsal appendage primordia (Figs. 2C and 3A).



**Fig. 2.** Inductive signals, target genes, and genetic interactions involved in eggshell patterning. (A) The dorsoventral pattern of EGFR signaling. (B) The anteroposterior pattern of DPP signaling. (C) The fate map for the formation of the respiratory dorsal appendages and the operculum. Each of the appendages is derived from a primordium comprising a 2D domain of cells expressing Broad (BR, blue) and an adjacent line of cells expressing Rhomboid (RHO, red). Cells between the two primordia express the transcription factor Pointed (PNT, orange) and contribute to the formation of the operculum. (D) The network of some of the key interactions involved in eggshell patterning. GRK controls *br* at two different time points: first, when it is distributed in a posterior-to-anterior gradient (P), and then when it is distributed in a dorsoventral (D) gradient (see text for details).





The identification of these enhancers made it possible to determine whether the effects of the key transcription factors controlling *br* are direct and whether they affect the early or the late phases of *br* expression (29). For instance, it was established that MIRR controls both enhancers, but with opposite signs: repressing *brE* and activating *brL* (Fig. 3 B and C). On the other hand, PNT represses *brL*, but does not affect *brE*. The DPP pathway controls only the late phase of *br* expression within the appendage primordia. The net effect of DPP is repressive and is fully dependent on BRK (23). Specifically, the DPP pathway directly represses *brk* in the anterior follicle cells, by acting through a well-characterized direct mechanism (Fig. 3C) (26). In the rest of the follicular epithelium, BRK positively regulates *brL*, likely by repressing an unknown repressor (23).

Established under the joint action of the EGFR and DPP signals, BR represses *rho*, confining its expression to the anterior border of the BR domain. As mentioned above, RHO is a ligand-processing protease within the *Drosophila* EGFR pathway. Remarkably, although neither RHO nor its target (Spitz, SPI) is essential for proper eggshell patterning (15, 18), the BR-dependent repression of *rho* is important. Specifically, low levels of the BR protein resulting from the activity of the *brE* enhancer “protect” the *brL* enhancer from the PNT-mediated repression. In the absence of the early *br* expression, EGFR activation induces premature expression of *rho*, which is normally expressed only after the *brE* enhancer is repressed by GRK. Precocious expression of RHO results in ectopic secretion of SPI and leads to ectopic high levels of EGFR activation. This in turn leads to the PNT-mediated repression of the *brL* enhancer and results in loss of the dorsal appendages (30). Thus, the 2D pattern of BR is shaped by the dynamic interplay of at least two enhancers with very different spatiotemporal activities.

Some of the *cis* and *trans* components in the network establishing the dorsal appendage primordia are shown in Fig. 3 B and C. Based on this network, we can explain the observations made in numerous genetic experiments. For instance, uniform activation of DPP signaling leads to eggshells with no appendages and a large, dorsally located operculum (Fig. 1C). We now understand that this effect is caused by direct repression of BRK, which leads, through a short cascade, to repression of the *brL* enhancer. As another example, uniform activation of the EGFR pathway also results in eggshells with no appendages, but the operculum is now formed around the entire dorsoventral axis (31). This effect is caused by the PNT-dependent repression of *brL*. In both examples, *brL* is repressed, but the mechanisms of repression are different. Note that because all of the effects of the patterning cues ultimately converge on the *brL* enhancer, future studies of the eggshell patterning network should focus on a more complete understanding of this regulatory element. Along with identifying the missing transacting elements, such as a missing repressor that is antagonized by BRK, this requires identifying the *cis*-regulatory sequences that respond to the transcription factors controlled by the EGFR and DPP pathways.

### From *D. melanogaster* to Other Species

Explaining the morphological diversity of eggshells in drosophilids motivated much of the mechanistic studies of the eggshell patterning network (32–38). For example, in contrast to *D. melanogaster*, in which each of the two dorsal appendages is formed from a separate primordium, the four appendages in *D. virilis* arise not from four, but from two separate primordia (Fig. 4A). Similar to *D. melanogaster*, each primordium comprises a 2D domain of *br* bordered by a line of *rho*-expressing cells. Thus, it appears that the same genes are involved in patterning of appendages, but their expression patterns are different (39). What mediates these changes in gene expression? Is it possible to trace them to changes in the inductive signals and the mechanisms by which they are interpreted by the follicle cells? Although work addressing these

questions has just begun, it is already clear that multiple mechanisms are at play in this rapidly evolving system.


Early models of eggshell patterning suggested that quantitative changes in the distribution of the GRK gradient can change the number of dorsal appendages (33, 40, 41). This possibility was conclusively ruled out by an elegant experiment that relied on chimera egg chambers established by pole cell transplantation (42). In this case, egg chambers contained somatic cells from *D. melanogaster* and the germ line cells derived from *D. virilis*. As a result, the follicular epithelium of *D. melanogaster* was patterned by GRK from *D. virilis*. The eggshells formed by these mosaic egg chambers were indistinguishable from the wild-type eggshells of *D. melanogaster*. These results show that, at least in this case, changes in eggshell morphology cannot be explained by changes in the inductive signal alone, pointing to the need to consider alterations in the network that interprets the inductive signal.

As a first step in this direction, Nakamura et al. (42) focused on the regulatory region of *rho* and identified the orthologous enhancer in *D. virilis*. This enhancer recapitulates the endogenous pattern of *rho* expression in *D. virilis* and drives expression in a broken V-shaped pattern. However, when introduced in *D. melanogaster*, this enhancer drives expression in an L-shaped pattern that is very close to the endogenous pattern of *rho* in *D. melanogaster* (Fig. 4B). Thus, changes in the pattern of *rho* expression are caused by changes in *trans* to the regulatory region of *rho*. Consistent with this scenario, the L-shaped activity *rho* enhancer from *D. melanogaster* is altered when this enhancer is introduced to *D. virilis*. As expected, this enhancer is now active in a broken V-shaped pattern (Fig. 4C).

As mentioned in the previous section, one of the main regulators of *rho* is BR, which represses *rho* in cells that form the upper part of the future dorsal appendages (11, 43). At this point of oogenesis, *br* is regulated by the *brL* enhancer. Thus, perhaps one way to alter the expression of *rho* is to alter the activity of the *brL* enhancer. Indeed, the wild-type expression of *rho* in both species can be predicted by the expression pattern of *br*. In both species, *rho* is expressed at the anterior border of the *br* domain (11, 21, 30, 43). If this is correct, then the regulatory region of *brL* should be changed in a way that alters the lateral extent of its activity. In particular, the expression of *brL* should be expanded in the lateral direction, to generate the “handle” characteristic of the wild-type pattern of BR in *D. virilis*. This handle is responsible for defining the anteriormost pair of appendages (39, 44).

To test this possibility, we have begun to identify the regulatory region of *brL* in *D. virilis* and assay its activity in *D. melanogaster*. Our preliminary results demonstrate that the *brL* enhancer from *D. virilis* indeed drives the expression more broadly than the *brL* enhancer from *D. melanogaster* (Fig. 4D'). The most noticeable difference is the expression in a lateral group of cells, resulting in a pattern that begins to resemble the endogenous pattern in *D. virilis* (Fig. 4D–D'). Based on this, we propose that the evolution of the *brL* enhancer is responsible for the diversification of eggshell patterning and morphogenesis. To test this hypothesis, we will have to identify which sequence changes within the *brL* result in the expanded expression of *br* in *D. virilis*. The identification of these changes is ongoing.

Although changes in the spatial distribution of GRK could not explain the differences between the eggshell structures of *D. virilis* and *D. melanogaster*, they are responsible for the diversification of other aspects of eggshell morphogenesis. A clear example of the functional capabilities of changes of inductive signals is provided by studies of the dorsal ridge, a lumen-like structure along the dorsalmost side of the eggshells of several *Drosophila* species, including *D. willistoni* (Fig. 1F) (32, 36). In contrast to *D. melanogaster*, which lacks the dorsal ridge and is patterned by GRK localized around the oocyte nucleus (Fig. 4E), *D. willistoni* is patterned by the GRK profile that is significantly extended toward the posterior end of the follicular epithelium (45) (Fig. 4F). The origin of this dramatic change in the distribution of GRK requires



**Fig. 4.** Two different mechanisms for the evolution of *br* patterning: changes in the regulatory region of *br* and changes in the inductive signals. (A) Schematic representation of the expression domains of *br* and *rho* in *D. virilis*. (B) Summary of the activity of the *rho* enhancer from *D. virilis* in *D. melanogaster*. (C) Summary of the activity of the *rho* enhancer from *D. melanogaster* in *D. virilis*. (D–D'') Comparison of the activities of the *brL* enhancers from *D. melanogaster* (D; *brL-GFP* in green) and *D. virilis* (D'; *brLvir-lacZ*, red), analyzed in *D. melanogaster*. (D'') Merged image including immunostaining for endogenous Br (white nuclear staining) for orientation. The yellow arrow indicates the lateral expansion of *D. virilis* *brL*. (E) Schematic of GRK localization in *D. melanogaster* (lateral view). (F) Schematic of GRK localization in *D. willistoni* (lateral view). (G) Schematics of eggshells in *D. melanogaster*, *D. willistoni*, and *D. melanogaster* patterned by GRK from *D. willistoni*.

further study, but it has been already established that GRK from *D. willistoni* is both necessary for dorsal ridge formation and sufficient for generating a partial dorsal ridge in eggshells of *D. melanogaster* (Fig. 4G). Thus, diversification of eggshell morphogenesis can be caused by the combined effects of changes at very top (GRK) and very bottom (BR) of the patterning network.

### Summary and Outlook

Eggshell morphogenesis in drosophilids provides an excellent opportunity for the detailed analysis of multiple steps that connect inductive signals and transcription factors to 2D patterns of gene expression and 3D structures. The experimental studies of this system have come a long way from the identification of the key components of signaling pathways through their effects on eggshell morphologies to the detailed analysis of gene regulatory sequences and computational models that capture multiple aspects of signaling, transcription, and morphogenesis (30, 46, 47). Much work remains to be done to characterize the regulatory sequences and their control by transcription factors. Furthermore, some of the important players, most notably the activators, are yet to be identified and placed within the existing network.

Recent studies with the enhancer of *br* and data from cross-species analysis strongly suggest that the late enhancer of *br* is one of the main loci for the diversification of eggshell patterning. At this point, the identified regulatory regions for *brL* in both species are still quite large. Shortening these enhancers will enable more efficient exploration of the *cis*-regulatory changes involved in the diversification of eggshell patterning. Once these changes are identified and the activity of minimal enhancers is examined in both species, similar to what has been done for the *rho* enhancer,

their functional effects on eggshell patterning can be tested. This can be done using the recently developed genome editing techniques, by swapping the *brL* regions between the two species. In particular, it will be very interesting to see whether a change in a single enhancer is sufficient to change both the expression pattern of BR and the number of dorsal appendages.

In addition to changes in the *cis*-regulatory region of *br*, which appears to be the key node in the appendage patterning network, it is important to keep in mind the mechanisms that rely on the intracellular modulation of inductive signals. In particular, GRK induces several negative feedback loops that modulate the signals sensed by the enhancers of the genes within the patterning network, changing the 2D patterns of *br* expression and affecting eggshell morphology (15, 48–52). Changing the induction thresholds of these negative feedback regulators provides another degree of flexibility for the evolution of eggshell patterning (18, 28).

One of the most exciting directions for future studies is related to the mechanistic analysis and experimental validation of the computational models that can explain the remarkable morphological diversity of eggshell structures. Current models of eggshell patterning can account for the dynamics of inductive inputs and some of the most important features of the transcriptional network that regulate BR and RHO (20, 21, 28, 30, 37, 53). These models can predict how the spatiotemporal activity of the *brL* enhancer responds to changes in the inductive signals and changes within the transcriptional network, providing a compact summary for a large number of genetic perturbation experiments. At this point, models of eggshell patterning do not directly use the sequence-specific information, but these

capabilities can be added as we acquire more knowledge about the connections between transcription factors and the *brL* sequence (54, 55). In the future, we envision a unified model that accounts for multiple processes, from inductive signals to enhancers, and can generate all of the observed eggshell morphologies by variations of model parameters and sequence variations.

- Berg CA (2005) The *Drosophila* shell game: Patterning genes and morphological change. *Trends Genet* 21(6):346–355.
- Deng WM, Bownes M (1997) Two signalling pathways specify localised expression of the Broad-Complex in *Drosophila* eggshell patterning and morphogenesis. *Development* 124(22):4639–4647.
- Peri F, Roth S (2000) Combined activities of Gurken and decapentaplegic specify dorsal chorion structures of the *Drosophila* egg. *Development* 127(4):841–850.
- Cheung LS, Schüpbach T, Shvartsman SY (2011) Pattern formation by receptor tyrosine kinases: Analysis of the Gurken gradient in *Drosophila* oogenesis. *Curr Opin Genet Dev* 21(6):719–725.
- Schüpbach T (1987) Germ line and soma cooperate during oogenesis to establish the dorsoventral pattern of egg shell and embryo in *Drosophila melanogaster*. *Cell* 49(5):699–707.
- Goentoro LA, et al. (2006) Quantifying the Gurken morphogen gradient in *Drosophila* oogenesis. *Dev Cell* 11(2):263–272.
- Twombly V, et al. (1996) The TGF-beta signaling pathway is essential for *Drosophila* oogenesis. *Development* 122(5):1555–1565.
- Spradling AC (1993) Developmental genetics of oogenesis. The Development of *Drosophila melanogaster* (Cold Spring Harbor Lab Press, Cold Spring Harbor, NY), pp 1–70.
- Nilson LA, Schüpbach T (1999) EGF receptor signaling in *Drosophila* oogenesis. *Curr Top Dev Biol* 44:203–243.
- Pai LM, Barcelo G, Schüpbach T (2000) D-cbl, a negative regulator of the Egfr pathway, is required for dorsoventral patterning in *Drosophila* oogenesis. *Cell* 103(1):51–61.
- Ward EJ, Berg CA (2005) Juxtaposition between two cell types is necessary for dorsal appendage tube formation. *Mech Dev* 122(2):241–255.
- Atkey MR, Lachance JF, Walczak M, Rebello T, Nilson LA (2006) Capicua regulates follicle cell fate in the *Drosophila* ovary through repression of mirror. *Development* 133(11):2115–2123.
- Morimoto AM, et al. (1996) Pointed, an ETS domain transcription factor, negatively regulates the EGF receptor pathway in *Drosophila* oogenesis. *Development* 122(12):3745–3754.
- Goff DJ, Nilson LA, Morisato D (2001) Establishment of dorsal-ventral polarity of the *Drosophila* egg requires capicua action in ovarian follicle cells. *Development* 128(22):4553–4562.
- Boisclair Lachance JF, Fregoso Lomas M, Eleiche A, Bouchard Kerr P, Nilson LA (2009) Graded Egfr activity patterns the *Drosophila* eggshell independently of autocrine feedback. *Development* 136(17):2893–2902.
- Fregoso Lomas M, Hails F, Lachance JF, Nilson LA (2013) Response to the dorsal anterior gradient of EGFR signaling in *Drosophila* oogenesis is prepatterned by earlier posterior EGFR activation. *Cell Reports* 4(4):791–802.
- Yakoby N, Lembong J, Schüpbach T, Shvartsman SY (2008) *Drosophila* eggshell is patterned by sequential action of feedforward and feedback loops. *Development* 135(2):343–351.
- Zartman JJ, Kanodia JS, Cheung LS, Shvartsman SY (2009) Feedback control of the EGFR signaling gradient: Superposition of domain-splitting events in *Drosophila* oogenesis. *Development* 136(17):2903–2911.
- Briscoe J, Small S (2015) Morphogen rules: Design principles of gradient-mediated embryo patterning. *Development* 142(23):3996–4009.
- Lembong J, Yakoby N, Shvartsman SY (2009) Pattern formation by dynamically interacting network motifs. *Proc Natl Acad Sci USA* 106(9):3213–3218.
- Simakov DS, Cheung LS, Pismen LM, Shvartsman SY (2012) EGFR-dependent network interactions that pattern *Drosophila* eggshell appendages. *Development* 139(15):2814–2820.
- Boisclair Lachance JF, et al. (2014) A comparative study of Pointed and Yan expression reveals new complexity to the transcriptional networks downstream of receptor tyrosine kinase signaling. *Dev Biol* 385(2):263–278.
- Charbonnier E, et al. (2015) BMP-dependent gene repression cascade in *Drosophila* eggshell patterning. *Dev Biol* 400(2):258–265.
- Chen Y, Schüpbach T (2006) The role of brinker in eggshell patterning. *Mech Dev* 123(5):395–406.
- Shravage BV, Altmann G, Technau M, Roth S (2007) The role of Dpp and its inhibitors during eggshell patterning in *Drosophila*. *Development* 134(12):2261–2271.
- Hamaratoglu F, Affolter M, Pyrowolakis G (2014) Dpp/BMP signaling in flies: From molecules to biology. *Semin Cell Dev Biol* 32:128–136.
- Fregoso Lomas M, De Vito S, Boisclair Lachance JF, Houde J, Nilson LA (2016) Determination of EGFR signaling output by opposing gradients of BMP and JAK/STAT activity. *Curr Biol* 26(19):2572–2582.
- Zartman JJ, et al. (2011) Pattern formation by a moving morphogen source. *Phys Biol* 8(4):045003.
- Fuchs A, Cheung LS, Charbonnier E, Shvartsman SY, Pyrowolakis G (2012) Transcriptional interpretation of the EGF receptor signaling gradient. *Proc Natl Acad Sci USA* 109(5):1572–1577.
- Cheung LS, Simakov DS, Fuchs A, Pyrowolakis G, Shvartsman SY (2013) Dynamic model for the coordination of two enhancers of broad by EGFR signaling. *Proc Natl Acad Sci USA* 110(44):17939–17944.
- Queenan AM, Ghabrial A, Schüpbach T (1997) Ectopic activation of torpedo/Egfr, a *Drosophila* receptor tyrosine kinase, dorsalizes both the eggshell and the embryo. *Development* 124(19):3871–3880.
- Hinton HE (1981) *Biology of Insect Eggs* (Pergamon Press, Oxford, UK).
- Shvartsman SY, Muratov CB, Lauffenburger DA (2002) Modeling and computational analysis of EGF receptor-mediated cell communication in *Drosophila* oogenesis. *Development* 129(11):2577–2589.
- Nakamura Y, Matsuno K (2003) Species-specific activation of EGF receptor signaling underlies evolutionary diversity in the dorsal appendage number of the genus *Drosophila* eggshells. *Mech Dev* 120(8):897–907.
- Vreede BM, Lynch JA, Roth S, Sucena E (2013) Co-option of a coordinate system defined by the EGFR and Dpp pathways in the evolution of a morphological novelty. *EvoDevo* 4(1):7.
- Niepielko MG, et al. (2014) Chorion patterning: A window into gene regulation and *Drosophila* species' relatedness. *Mol Biol Evol* 31(1):154–164.
- Niepielko MG, Ip K, Kanodia JS, Lun DS, Yakoby N (2012) Evolution of BMP signaling in *Drosophila* oogenesis: A receptor-based mechanism. *Biophys J* 102(8):1722–1730.
- Niepielko MG, Hernáiz-Hernández Y, Yakoby N (2011) BMP signaling dynamics in the follicle cells of multiple *Drosophila* species. *Dev Biol* 354(1):151–159.
- James KE, Berg CA (2003) Temporal comparison of Broad-Complex expression during eggshell-appendage patterning and morphogenesis in two *Drosophila* species with different eggshell-appendage numbers. *Gene Expr Patterns* 3(5):629–634.
- Muratov CB, Shvartsman SY (2003) An asymptotic study of the inductive pattern formation mechanism in *Drosophila* egg development. *Physica D* 186(1–2):93–108.
- Pribly M, Muratov CB, Shvartsman SY (2003) Transitions in the model of epithelial patterning. *Dev Dyn* 226(1):155–159.
- Nakamura Y, et al. (2007) Soma-dependent modulations contribute to divergence of rhomboid expression during evolution of *Drosophila* eggshell morphology. *Development* 134(8):1529–1537.
- Ward EJ, Zhou X, Riddiford LM, Berg CA, Ruohola-Baker H (2006) Border of Notch activity establishes a boundary between the two dorsal appendage tube cell types. *Dev Biol* 297(2):461–470.
- Osterfield M, Schüpbach T, Wieschhaus E, Shvartsman SY (2015) Diversity of epithelial morphogenesis during eggshell formation in drosophilids. *Development* 142(11):1971–1977.
- Niepielko MG, Yakoby N (2014) Evolutionary changes in TGF $\alpha$  distribution underlie morphological diversity in eggshells from *Drosophila* species. *Development* 141(24):4710–4715.
- Osterfield M, Du X, Schüpbach T, Wieschhaus E, Shvartsman SY (2013) Three-dimensional epithelial morphogenesis in the developing *Drosophila* egg. *Dev Cell* 24(4):400–410.
- Fauré A, Vreede BM, Sucena E, Chaouiya C (2014) A discrete model of *Drosophila* eggshell patterning reveals cell-autonomous and juxtacrine effects. *PLoS Comput Biol* 10(3):e1003527.
- Wasserman JD, Freeman M (1998) An autoregulatory cascade of EGF receptor signaling patterns the *Drosophila* egg. *Cell* 95(3):355–364.
- Sapir A, Schweitzer R, Shilo BZ (1998) Sequential activation of the EGF receptor pathway during *Drosophila* oogenesis establishes the dorsoventral axis. *Development* 125(2):191–200.
- Reich A, Sapir A, Shilo B (1999) Sprouty is a general inhibitor of receptor tyrosine kinase signaling. *Development* 126(18):4139–4147.
- Ghigliione C, et al. (1999) The transmembrane molecule kerkon 1 acts in a feedback loop to negatively regulate the activity of the *Drosophila* EGF receptor during oogenesis. *Cell* 96(6):847–856.
- Peri F, Bökel C, Roth S (1999) Local Gurken signaling and dynamic MAPK activation during *Drosophila* oogenesis. *Mech Dev* 81(1–2):75–88.
- Lembong J, Yakoby N, Shvartsman SY (2008) Spatial regulation of BMP signaling by patterned receptor expression. *Tissue Eng Part A* 14(9):1469–1477.
- Samee MA, et al. (2015) A systematic ensemble approach to thermodynamic modeling of gene expression from sequence data. *Cell Syst* 1(6):396–407.
- Samee AH, Sinha S (2013) Evaluating thermodynamic models of enhancer activity on cellular resolution gene expression data. *Methods* 62(1):79–90.