



HHS Public Access

Author manuscript

Environ Int. Author manuscript; available in PMC 2018 January 01.

Published in final edited form as:

Environ Int. 2017 January ; 98: 75–81. doi:10.1016/j.envint.2016.10.003.

Spatial and temporal trends in the mortality burden of air pollution in China: 2004–2012

Miaomiao Liu^a, Yining Huang^a, Zongwei Ma^a, Zhou Jin^a, Xingyu Liu^a, Haikun Wang^a, Yang Liu^b, Jinnan Wang^{a,c}, Matti Jantunen^d, Jun Bi^{d,r,*}, and Patrick L. Kinney^{e,**}

^aState Key Laboratory of Pollution Control and Resource Reuse, School of the Environment, Nanjing University, Nanjing, Jiangsu, China ^bDepartment of Environmental Health, Rollins School of Public Health, Emory University, Atlanta, GA, USA ^cChinese Academy for Environmental Planning, Beijing, China ^dNational Institute for Health and Welfare, Environment and Health Unit, Kuopio, Finland ^eMailman School of Public Health, Columbia University, New York, USA

Abstract

While recent assessments have quantified the burden of air pollution at the national scale in China, air quality managers would benefit from assessments that disaggregate health impacts over regions and over time. We took advantage of a new 10 × 10 km satellite-based PM_{2.5} dataset to analyze spatial and temporal trends of air pollution health impacts in China, from 2004 to 2012. Results showed that national PM_{2.5} related deaths from stroke, ischemic heart disease and lung cancer increased from approximately 800,000 cases in 2004 to over 1.2 million cases in 2012. The health burden exhibited strong spatial variations, with high attributable deaths concentrated in regions including the Beijing–Tianjin Metropolitan Region, Yangtze River Delta, Pearl River Delta, Sichuan Basin, Shandong, Wuhan Metropolitan Region, Changsha–Zhuzhou–Xiangtan, Henan, and Anhui, which have heavy air pollution, high population density, or both. Increasing trends were found in most provinces, but with varied growth rates. While there was some evidence for improving air quality in recent years, this was offset somewhat by the countervailing influences of in–migration together with population growth. We recommend that priority areas for future national air pollution control policies be adjusted to better reflect the spatial hotspots of health burdens. Satellite-based exposure and health impact assessments can be a useful tool for tracking progress on both air quality and population health burden reductions.

Keywords

Air pollution; PM_{2.5}; Health burden; Temporal; Spatial; China

*Correspondence to: J. Bi, School of Environment, Nanjing University, Xianlin Avenue 163#, Nanjing 210023, P. R. China.

**Correspondence to: P. L. Kinney, Rosenfield Building, 722 West 168th Street, Room 1104E, New York, NY 10032, USA.

Role of the funding source

The funder of the study had no role in study design or data collection. ML, JB, PLK and coauthors had full access to all the data in the study. JB and PLK had final responsibility for the decision to submit for publication.

1. Introduction

Over recent decades, rapid economic development has led to worsening air quality in China (Wang et al., 2014; Wu et al., 2012; Zhao et al., 2013a, b), which now ranks as one of the most polluted countries in the world (Van Donkelaar et al., 2010; Verstraeten et al., 2015). As a leading modifiable risk factor for non-communicable diseases, air pollution has attracted considerable interest from both the research and policy communities (Krewski et al., 2009; Laden et al., 2006; Pope et al., 2009; Pope and Dockery, 2006; Shang et al., 2013; West et al., 2016; Yang et al., 2013). However, assessments of health impacts, and their trends over time and space, have been hampered by the limited availability of relevant air monitoring prior to 2013 in China. Initial health impact studies used surrogate measures such as ambient PM₁₀ concentrations measured at fixed monitoring stations to estimate health outcomes attributable to China's air pollution (Cheng et al., 2013; Hou et al., 2012; Matus et al., 2012; Zhang et al., 2008). However, PM₁₀ is a less robust health-related exposure metric than PM_{2.5} (USEPA, 2012), and central-site monitoring data may lead to exposure uncertainty related to spatial variability of PM, population density and demographic characteristics (Steinle et al., 2013).

In recent years, methods have been developed for estimating ground-level PM_{2.5} concentrations based on satellite derived aerosol optical depth (AOD), providing a promising alternative for estimating exposure to outdoor PM_{2.5} and associated health impacts, with more extensive spatial and temporal coverage (Brauer et al., 2012, 2015; Ma et al., 2014; Yao and Lu, 2014). For example, taking advantage of a global PM_{2.5} dataset derived from AOD, the Global Burden of Disease (GBD) project reported that outdoor air pollution in China caused 1.2 million premature deaths and 25 million disability adjusted life years (DALY) losses in 2010 (Yang et al., 2013). A subsequent study reported a lower mortality burden of 0.35–0.50 million premature deaths, based on a different exposure methodology (Chen et al., 2013a). Both of these PM_{2.5}-based health assessments reported results only for a single year and for China as a whole.

Recently, as the Chinese government's demands for air quality management broadened from understanding the scope of the problem to targeting interventions, identification of spatial and temporal trends in health burdens attributable to air pollution is becoming more and more important. A recent study evaluated the temporal trend and spatial distributions of PM₁₀-related health impacts in China based on monitoring data from 2001 to 2011, contributing valuable insights into this question (Cheng et al., 2013). However, because PM₁₀ trends may not track those of PM_{2.5}, spatial and temporal trends derived from PM₁₀ data can be misleading (Cheng et al., 2013; Ma et al., 2016). Therefore, more accurate and refined information of the spatial-temporal characteristics of PM_{2.5} effects are needed to support future policy interventions.

To address this need, we used a new 10 km resolution satellite derived PM_{2.5} dataset in conjunction with fine scale population data to develop novel estimates of PM_{2.5} related health damage in China from 2004 to 2012 at the subnational scale (Burnett et al., 2014; Ma et al., 2016; Yang et al., 2009). The implications of observed temporal trends and spatial distributions of impacts for future policy directions are explored.

2. Methods and data

While several air pollutants are known to have adverse health impacts, we focus here on outdoor PM_{2.5} as the indicator of risk as it is widely regarded as the single best metric of air pollution-related risk to public health (USEPA, 2012). Regarding outcomes, we selected the premature deaths caused by stroke (International Classification of Diseases Revision 10 codes/ICD-10: I60–I69), ischemic heart disease (IHD, ICD-10: I20–I25), and lung cancer (LC, ICD-10: C33–C34) because of strong evidence for causal effects between these outcomes and PM_{2.5} from prior high-quality epidemiological studies and sufficient cause-specific mortality data to estimate outcome-specific effect sizes per unit of exposure (Burnett et al., 2014; NHFPCC, 2005–2013). We excluded chronic obstructive pulmonary disease (COPD) deaths because the mortality data for COPD were not available. Annual cause-specific baseline mortality rates were taken from the China Health Statistical Yearbook.

Population weighted exposure (PWE) to PM_{2.5} was calculated using PM_{2.5} concentrations derived from satellite AOD at 10 km resolution and population distribution maps at 1 km resolution (Ma et al., 2016; Yang et al., 2009). Briefly, we developed and validated a model for estimating PM_{2.5} based on collection 6 AOD retrieved by the US National Aeronautics and Space Administration (NASA) Moderate Resolution Imaging Spectroradiometer (MODIS), meteorology data, land use data and China's ground-monitoring PM_{2.5} data in 2013. Historical daily PM_{2.5} concentrations (2004–2012) were then estimated with this model using historical AOD data as inputs, assuming that the daily relationship between PM_{2.5} and AOD was constant for the same Day of Year (DOY) in each year. This model was calibrated to minimize bias in the monthly and seasonal estimates on PM_{2.5} and has been successfully applied in recent analyses (Chen et al., 2016; Liu et al., 2016b; Ma et al., 2016). The population distribution maps we adopted in this study also have been successfully applied in prior studies (Huang et al., 2014; Liu et al., 2013). In regard to relative risk per unit of exposure for each mortality outcome, the ideal one for the study population should be the results from Chinese cohort studies. However, to our knowledge, there are no published Chinese cohort studies directly examining the impacts of PM_{2.5} on mortality so far. Most of existing Chinese cohort studies used the concentrations of SO₂, NO₂, TSP, or PM₁₀ as exposure surrogates, which were poorer predictors of mortality compared with PM_{2.5} (Cao et al., 2011; Liu et al., 2016a; Zeng et al., 2016; Zhou et al., 2014). In the few published PM_{2.5} cohort studies (Guo et al., 2016; Qian et al., 2016), only the impacts of PM_{2.5} on lung cancer incidence and preterm birth are discussed. Under this circumstance, the integrated exposure response (IER) model developed as part of the GBD study seems to be the most suitable choice for the estimates of mortality burdens attributable to PM_{2.5} in China. Even compared with the RRs estimated based on Chinese TSP cohort and the conversion rate of PM_{2.5}/TSP, the RRs evaluated by the IER model can yield sensible results in the risk analysis over the range of concentrations that prevail in China (Burnett et al., 2014). 5 µg/m³ of annual PM_{2.5} was assumed as the counterfactual theoretical–minimum–risk exposure (Burnett et al., 2014). It should be noted that because the IER model was developed for adults, the age ranges for population and baseline incidence rates used in existing studies using the IER model are usually aged 25 and over or aged 30 and over (Lelieveld et al.,

2015; Silva et al., 2016). However, the data of population and cause-specific baseline incidence rates with detailed age structures are not available in China at such a long time scale and fine spatial resolution. So we used the aggregated all-age data for population and baseline incidence rates in this study. While this might induce some bias, we think the biases are acceptable because most of baseline mortalities from IHD, stroke and lung cancer occur among adults (NHFPPC, 2005–2013). Using these inputs, we computed annual excess deaths attributable to outdoor PM_{2.5} exposure as follows (Cheng et al., 2013; Zhang et al., 2008).

$$ED_{i,j,t} = \left(1 - \frac{1}{RR_{i,j,t}}\right) * I_{i,j,t} * P_{i,j,t} = AF_{i,j,t} * I_{i,j,t} * P_{i,j,t}$$

where $ED_{i,j,t}$ is the excess deaths caused by air pollution for stroke, IHD or LC; $AF_{i,j,t}$ is the attributable fraction (AF), defined as the fraction of the disease burden attributable to PM_{2.5}; $I_{i,j,t}$ is the annual all-age incidence rate of the mortality end point, obtained from China Health Statistical Yearbook (2005–2013) compiled by the Ministry of Health of China according to ICD-10 codes (NHFPPC, 2005–2013); $P_{i,j,t}$ is the permanent all-age population aggregated from population density map at 1 km resolution provided by Chinese Academy of Science (Yang et al., 2009); $RR_{i,j,t}$ is the relative risk of premature mortality due to PM_{2.5} obtained from the integrated exposure response (IER) model (Burnett et al., 2014) and i , j and t are the grid, the prefecture and the year, respectively. Annual excess deaths caused by air pollution were estimated at 10 km resolution and also aggregated for 339 prefectures, 31 provinces and the whole of China, for each year from 2004 to 2012.

3. Results

During the period 2004–2012, over 93% of people in China lived in areas where PM_{2.5} exceeded China's National Air Quality Standard for Grade II of 35 µg/m³ (Fig. 1). Population weighted exposure (PWE) averaged between 67.1 and 76.7 µg/m³ for China as a whole (Table S1). To visualize how regional impacts of PM_{2.5} are distributed across the concentration range, we plot the distribution of population as a function of ambient PM_{2.5} in 2004 (Fig. 1A) and 2012 (Fig. 1B). The distributions were bimodal, with one peak in the range 45–60 µg/m³ and another around 80 µg/m³. Compared with 2004, the population exposure shifted to greater extremes of PM_{2.5} concentrations in 2012.

Total excess premature deaths due to stroke, IHD, and LC attributable to outdoor PM_{2.5} exposure in China increased rapidly from 807 (95% CI: 328–1057) thousand cases in 2004 to 1250 (95% CI: 559–1672) thousand cases in 2012. Specifically, excess deaths due to stroke, IHD, and LC increased from 589 (95% CI: 209–736), 124 (95% CI: 86–194) and 93 (95% CI: 33–127) thousand cases in 2004 to 761 (95% CI: 274–947), 321 (95% CI: 224–498) and 169 (95% CI: 61–227) thousand cases in 2012 (Fig. 2), with the annual average growth rate of 3.25%, 12.63%, and 7.75%, respectively. In 2012, 43.9%, 29.0% and 27.9% of deaths caused by stroke, IHD and LC, respectively, in China were attributed to exposure to outdoor PM_{2.5}.

At the provincial level, the highest ambient PM_{2.5} concentrations were observed in the Beijing–Tianjin Metropolitan Region, Henan, and Shandong, with PWEs of 102.4, 107.0, and 101.1 µg/m³ in 2012, respectively. The Yangtze River Delta and Hunan experienced moderately high PM_{2.5} concentration (~70–90 µg/m³). The provincial variations of population–PM_{2.5} distribution led to significant spatial heterogeneity in air pollution related excess deaths (Fig. S1–S2). Eight provinces - Henan (HeN), Shandong (SD), Hebei (HB), Jiangsu (JS), Sichuan (SC), Guangdong (GD), Anhui (AH) and Hunan (HuN) - contributed over 50% of the national excess deaths. Although the trend over time of excess premature deaths became flat or slightly declined in a few provinces after 2007, increases were seen in most provinces over the nine–year time window (Fig. S1–S2). The annual average growth rate for total excess premature deaths caused by stroke, IHD and LC varied from 0.74% (the lowest in Hainan) to 9.46% (the highest in Beijing).

Maps of mean PM_{2.5}-attributed deaths at 10 km resolution exhibited strong spatial gradients (Fig. 3). As expected, deaths were concentrated in regions including the Beijing–Tianjin Metropolitan Region, Yangtze River Delta, Pearl River Delta, Sichuan Basin, Shandong, Wuhan Metropolitan Region, Changsha–Zhuzhou–Xiangtan, Henan, and Anhui, which had heavy air pollution, high population density, or both. Over time, there was no significant expansion of high–excess deaths centers (Fig. 3A vs. B). In other words, more rapidly increasing excess deaths were mostly located in areas that already experienced high excess deaths in 2004.

4. Discussion

Using a satellite-derived reconstruction of PM_{2.5} concentrations at 10 km resolution across China from 2004 to 2012, we observed significant variations of PM_{2.5} related health burdens over time and across provinces. These new results add useful spatial and temporal dimensions to previous national scale, point-in-time health burden estimates for China (Chen et al., 2013a; Yang et al., 2013). There were several important differences in our analysis as compared to the GBD study. Whereas the GBD study based their estimates on the “registered population”, here we used the higher and more relevant “permanent population”, which captures health impacts related in–migration to more polluted areas. This led to slightly higher estimates here than in the GBD study (Fig. 2). Additional discrepancies in stroke are attributed to the different data sources of the annual incidence rate of mortality. Our results generally confirmed the results found in the 2010 GBD study and added additional spatial and temporal richness.

This study fills an important gap in historical air pollution related health risk assessment in China during the decade preceding the onset of national PM_{2.5} monitoring. Though national excess premature deaths showed increasing trends from 2004 to 2012, some fluctuations were found in individual years after 2007. These fluctuations may reflect the intersection of three interacting trends: air quality change, in–migration induced by urbanization, and population growth. Regarding air quality, due to the combined effects of the global financial crisis starting in September of 2008 and stricter energy conservation and emissions reduction policy implemented in late 2006, national average PM_{2.5} levels experienced a decline from 2007 to 2009 (Lo and Wang, 2013; Ma et al., 2016). Since 2010, PM_{2.5} levels rose slightly

due to the economic recovery but soon showed evidence of further decline due to the onset of new air pollution regulations. It could be concluded that without significant demographic changes, China's efforts on air pollution control would have led to reductions in the associated health damage by 2012. Unfortunately, these efforts were offset by the countervailing influence of in-migration driven by urbanization. The temporal variation of national average PWE (Table S1) illustrates the interaction between air quality change and in-migration on excess deaths. Before 2008, PWE rose at a faster rate ($3.2 \mu\text{g}/\text{m}^3$ per year) than the increase of $\text{PM}_{2.5}$ levels ($1.97 \mu\text{g}/\text{m}^3$ per year), which indicates that during the period of 2004–2008, in-migration patterns caused by rapid urbanization (Fig. S3) showed a strong trend to more polluted urban areas and hence intensified the health damage (Bai et al., 2014; Gong et al., 2012). Since 2008, the intersection between decreased $\text{PM}_{2.5}$ level and in-migration to more polluted areas led to fluctuations of PWE. As for population growth, it also played an important role in the increase of health damage.

We also observed significant spatial variations in PM-related mortality burdens. To better understand the spatial characteristics of health damage, we removed the effects of population size by calculating the attributable fractions (AF) of 31 provinces (Fig. 4). Generally, the provinces with higher AF are burdened with more deaths per thousand attributable to air pollution, holding all other independent factors constant. In the provinces with increasing AF over time, exposure to air pollution is becoming a more and more dangerous risk factor to public health. As shown in Fig. 4, although there were fluctuations for several provinces in individual years, most Chinese provinces showed increased attributable fractions during the period of 2004–2012, which supports the findings from GBD Study 2010 that China has undergone an epidemiological transition from communicable to non-communicable diseases in the past decade (Yang et al., 2013). However, that's not the whole story. There were still ten provinces (Fig. 4) including Hainan (HN), Guizhou (GZ), Guangdong (GD), Guangxi (GX), Fujian (FJ), Yunnan (YN), Shanghai (SH), Hunan (HuN), Jiangxi (JX), and Zhejiang (ZJ) that experienced a decline of AF from 2004 to 2012. All of these provinces are located in southern China, with abundant precipitation and high forest coverage, which is beneficial for the deposition of airborne particles (Guo et al., 2014; Hofman et al., 2014; Li et al., 2014). Also, the lack of winter heating in these southern regions leads to improved air quality and health (Chen et al., 2013b). Lastly, the sea-land breeze may be conducive to atmospheric pollutant dispersion in some coastal regions (Li et al., 2014).

Our findings on spatial characteristics of health burdens have important implications for air pollution control policies in China. In recent years, China's Ministry of Environmental Protection (CMEP) has adopted a "spatial-differentiation" air pollution control strategy that imposes stricter controls in regions where air quality problems are thought to be most severe. In the 12th Five-Year Plan (2011–2015) on Air Pollution Control in Key Regions, thirteen regions were designated as priority areas for air pollution prevention and control in 2012 (Chen et al., 2013a). To examine whether the designated priority areas covered the spatial hotspots of air pollution related health damage, we overlaid our $\text{PM}_{2.5}$ related mortality maps with the boundary map of the priority control regions (Fig. 3). It can be seen that most centers with high excess deaths were located within priority control regions. Compared with non-priority areas, these priority control areas suffered more serious air pollution and higher mortality burdens per squared kilometer (Table S2 and Fig. 3). From 2004 to 2012, air

pollution levels and related mortality burdens in the priority control areas increased before 2008 and then showed slight improvements in recent years (see Table S2). The trends are similar to non-priority control areas, which suggest that the improvements may be attributed to the implementation of nationwide energy conservation and emissions reduction policy in the 11th Five-Year Plan rather than the designation of priority control areas in 2012 (Lo and Wang, 2013; Ma et al., 2016). However, from the historical experience we can expect that the differentiated and stricter emission control strategies in the priority control areas will bring more improvements of air quality and related health in the future compared with other areas. In addition, we found that Henan and Anhui Province are two regions which are suffering serious air pollution and related health damage but haven't yet received enough regulatory attention. From the perspective of health protection, we suggest that in the future, CMEP should expand the scope of the priority areas to include these two regions in the list.

While informative for policy decision making, our analysis has several limitations that should be kept in mind. The first one is related to the satellite-derived $PM_{2.5}$ dataset used in this study. Satellite data are sometimes missing due to the cloud and snow cover, potentially biasing estimates of $PM_{2.5}$ (Ma et al., 2016). To address the problem, we previously carried out a sensitivity analysis for each grid to test how many AOD-derived $PM_{2.5}$ estimations are needed to represent the true mean $PM_{2.5}$ concentrations. We found that a minimum of 6 valid satellite observations in a month or 11 in a season were sufficient to appropriately represent a monthly or seasonal average (Ma et al., 2016). In the current study, we extended the validation to the annual level, which showed that a minimum of 33 data points can predict the annual average $PM_{2.5}$ well ($R^2 = 0.83$, slope = 1.039). Based on this criterion, we found few missing annual average $PM_{2.5}$ estimates, and those were mainly concentrated in western China where the population density is lower than 1 person per square kilometer (Fig. S4). Therefore, missing satellite observations were unlikely to significantly alter the estimates of health burdens in this study. Another limitation relates to spatial scale. While our $PM_{2.5}$ estimates were developed at a relatively fine scale of 10×10 km, this is still too coarse to capture fine patterns of pollution related to proximity of local sources emitting pollution at ground level, such as from traffic, cooking and heating with solid fuels, etc. (Chafe et al., 2014).

Another limitation that affects this study is related to the IER model we adopted. To test the effects of exposure response functions on the results, a sensitivity analysis has been conducted. Specifically, the RRs from IER model were replaced by the RRs projected based on a published Chinese TSP cohort study and the conversion rate of $PM_{2.5}/TSP$ (Cao et al., 2011). Using the RRs in Cao et al. (2011), IHD and stroke mortality decreased by 2–10% and 7–14% while lung cancer mortality increased by 36–43% (Table S3). These differences are mainly caused by the non-linear shape of the IER function (Silva et al., 2016), which indicated that the estimates of mortality burdens are sensible to the exposure response function used in the assessment process. Fortunately, the estimates using the RRs in Cao et al. (2011) were still in the uncertainty range of the estimates using IER model. And the general spatial characteristics of mortality burdens were not significantly altered even though some grid-by-grid differences have been found (Fig. S5).

In addition, we glossed over the source classes responsible for PM_{2.5} related mortality burdens in this study, which may also bias the estimates of mortality burdens. As we know, the coal-dominated energy structure in China determined that a large share of the fine particle pollution were from coal combustion sources. At the same time, however, coal combustion PM_{2.5} posed a IHD risk roughly five times higher than for PM_{2.5} mass in general, on a per microgram/cubic meter PM_{2.5} basis (Thurston et al., 2016). Therefore, the assessment based on PM_{2.5} mass in general in this study may underestimate the mortality burdens. More importantly, the underestimations may vary across different areas within China, smallest for the areas where coal combustion PM_{2.5} accounted for a small fraction and highest for the areas where coal combustion sources contributed the most. This trend could further alter the spatial distributions of mortality burdens within China. Because little is known about the variations of source contributions within China, we cannot directly quantify the spatial effects at this stage. However, some qualitative judgments can be inferred from the observed evidence on the variations of PM_{2.5} constituents because these are largely caused by the variations of source contributions (Thurston et al., 2016). Observations from representative cities across China suggest that compared to the western region, the fraction of the secondary aerosols and elemental carbon is larger in the east of China (Yang et al., 2011). The secondary aerosols especially sulfate particulate are highly related to the coal combustion source; and elemental carbon is a key trace constitute to identify the traffic source, which was shown to be source with the second largest IHD risk (Thurston et al., 2016). If all above is true, the hotspots of health burdens in the east of China identified by our study (Fig. 3) would be further intensified after considering the toxicity of PM_{2.5} from different sources. Certainly, these speculations remain to be tested by future research on source appointment and source-specific toxicity of PM_{2.5}. However, it is clear that clean air efforts in China should focus on key source classes like coal combustion sources and traffic related sources to maximize public health benefits.

5. Conclusions

This study provides a refined estimation of historical health damage attributable to outdoor air pollution in China at 10 × 10 km resolution from 2004 to 2012. To our knowledge, this is the first study to quantify the health consequences of China's PM_{2.5} pollution at both a long time scale and sub-national level. Though some limitations exist, this work contributes valuable insights into the strong spatial and temporal variations of air pollution related health burdens in China. The findings of this study provide important implications for urban planners and policy makers regarding priority areas of national joint air pollution prevention and control in China. Moreover, open publication of our health burden datasets also makes it possible to further explore the determinants of these health impacts.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The study was supported by Natural Sciences Foundation of China (71433007), Jiangsu Science and Technology Support Program (SBE2014070918), Collaborative Innovation Center for Regional Environmental Quality, and

Collaborative Innovation Center for Atmospheric Environment and Equipment. The work of M. Liu was also supported by China Scholarship Council (CSC) under the State Scholarship Fund. The work of P. Kinney was partially supported by NIEHS grant P30ES009089. The work of Y. Liu was partially supported by the NASA Applied Sciences Program (grants NNX11AI53G and NNX14AG01G, Principal Investigator: Liu). We thank Professor Haidong Kan at Fudan University for his valuable assistance in providing the integrated exposure function. The contents of this paper are solely the responsibility of the authors and do not necessarily represent official views of the sponsors. The authors declare that they have no actual or potential competing financial interests.

References

- Bai X, Shi P, Liu Y. Society: realizing China's urban dream. *Nature*. 2014; 509:158–160. [PubMed: 24812683]
- Brauer M, Amann M, Burnett RT, Cohen A, Dentener F, Ezzati M, Henderson SB, Krzyzanowski M, Martin RV, Van Dingenen R, van Donkelaar A, Thurston GD. Exposure assessment for estimation of the global burden of disease attributable to outdoor air pollution. *Environ Sci Technol*. 2012; 46:652–660. [PubMed: 22148428]
- Brauer M, Freedman G, Frostad J, van Donkelaar A, Martin RV, Dentener F, van Dingenen R, Estep K, Amini H, Apte JS, Balakrishnan K, Barregard L, Broday D, Feigin V, Ghosh S, Hopke PK, Knibbs LD, Kokubo Y, Liu Y, Ma S, et al. Ambient air pollution exposure estimation for the global burden of disease 2013. *Environ Sci Technol*. 2015; 50(1):79–88. [PubMed: 26595236]
- Burnett RT, Pope CA, Ezzati M, Olives C, Lim SS, Mehta S, Shin HH, Singh G, Hubbell B, Brauer M, Anderson R, Smith KR, Balmes JR, Bruce NG, Kan HD, Laden F, Prüss-Ustün A, Turner MC, Gapstur SM, Diver WR, Cohen A. An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate matter exposure. *Environ Health Perspect*. 2014; 122:397–403. [PubMed: 24518036]
- Cao J, Yang C, Li J, Chen R, Chen B, Gu D, Kan H. Association between long-term exposure to outdoor air pollution and mortality in China: a cohort study. *J Hazard Mater*. 2011; 186(2):1594–1600. [PubMed: 21194838]
- Chafe ZA, Brauer M, Klimont Z, Dingenen RV, Mehta S, Rao S, Riahi K, Dentener F, Smith KR. Household cooking with solid fuels contributes to ambient PM_{2.5} air pollution and the burden of disease. *Environ Health Perspect*. 2014; 122:1314–1320. [PubMed: 25192243]
- Chen Z, Wang J, Ma G, Zhang Y. China tackles the health effects of air pollution. *Lancet*. 2013a; 382(9909):1959–1960. [PubMed: 24332069]
- Chen Y, Ebenstein A, Greenstone M, Li H. Evidence on the impact of sustained exposure to air pollution on life expectancy from China's Huai River policy. *Proc Natl Acad Sci*. 2013b; 110(32):12936–12941. [PubMed: 23836630]
- Chen, K., Zhou, L., Chen, X., Ma, Z., Liu, Y., Huang, L., Bi, J., Kinney, PL. Urbanization level and vulnerability to heat-related mortality in Jiangsu Province, China. *Environ Health Perspect*. 2016. <http://dx.doi.org/10.1289/EHP243>
- Cheng Z, Jiang J, Fajardo O, Wang S, Hao J. Characteristics and health impacts of particulate matter pollution in China (2001–2011). *Atmos Environ*. 2013; 65:186–194.
- Gong P, Liang S, Carlton EJ, Jiang Q, Wu J, Wang L, Remais JV. Urbanization and health in China. *Lancet*. 2012; 379(9818):843–852. [PubMed: 22386037]
- Guo L, Bao L, She J, Zeng E. Significance of wet deposition to removal of atmospheric particulate matter and polycyclic aromatic hydrocarbons: a case study in Guangzhou, China. *Atmos Environ*. 2014; 83:136–144.
- Guo Y, Zeng H, Zheng R, Li S, Barnett AG, Zhang S, Zou X, Huxley R, Chen W, Williams G. The association between lung cancer incidence and ambient air pollution in China: a spatiotemporal analysis. *Environ Res*. 2016; 144:60–65. [PubMed: 26562043]
- Hofman J, Bartholomeus H, Calders K, Van Wittenberghe S, Wuyts K, Samson R. On the relation between tree crown morphology and particulate matter deposition on urban tree leaves: a ground-based LiDAR approach. *Atmos Environ*. 2014; 99:130–139.
- Hou Q, An X, Wang Y, Tao Y, Sun Z. An assessment of China's PM₁₀-related health economic losses in 2009. *Sci Total Environ*. 2012; 435:61–65. [PubMed: 22846764]

- Huang Y, Jiang D, Zhuang D, Zhu Y, Fu J. An improved approach for modeling spatial distribution of water use profit - a case study in Tuhai Majia Basin, China. *Ecol Indic.* 2014; 36:94–99.
- Krewski D, Jerrett M, Burnett RT, Ma R, Hughes E, Shi Y, Turner MC, Pope CA, Thurston G, Calle EE, Thun MJ, Beckerman B, DeLuca P, Finkelstein N, Ito K, Moore DK, Newbold KB, Ramsay T, Ross Z, Shin H, Tempalski B. Extended follow-up and spatial analysis of the American Cancer Society study linking particulate air pollution and mortality. *Res Rep Health Eff Inst.* 2009; 140:5–114.
- Laden F, Schwartz J, Speizer FE, Dockery DW. Reduction in fine particulate air pollution and mortality: extended follow-up of the Harvard Six Cities study. *Am J Respir Crit Care Med.* 2006; 173(6):667–672. [PubMed: 16424447]
- Lelieveld J, Evans JS, Fnais M, Giannadaki D, Pozzer A. The contribution of outdoor air pollution sources to premature mortality on a global scale. *Nature.* 2015; 525(7569):367–371. [PubMed: 26381985]
- Li L, Qian J, Ou C, Zhou Y, Guo C, Guo Y. Spatial and temporal analysis of Air Pollution Index and its timescale-dependent relationship with meteorological factors in Guangzhou, China, 2001–2011. *Environ Pollut.* 2014; 190:75–81. [PubMed: 24732883]
- Liu M, Wang H, Oda T, Zhao Y, Yang X, Zhang R, Zhang B, Bi J, Chen J. Refined estimate of China's CO₂ emissions in spatiotemporal distributions. *Atmos Chem Phys.* 2013; 13(21):10873–10882.
- Liu W, Huang C, Hu Y, Fu Q, Zou Z, Sun C, Li S, Wang X, Cai J, Pan J, Huang Y, Chang J, Sun Y, Sundell J. Associations of gestational and early life exposures to ambient air pollution with childhood respiratory diseases in Shanghai, China: a retrospective cohort study. *Environ Int.* 2016a; 92:284–293. [PubMed: 27128713]
- Liu C, Yang C, Zhao Y, Ma Z, Bi J, Liu Y, Meng X, Wang Y, Cai J, Kan H, Chen R. Associations between long-term exposure to ambient particulate air pollution and type 2 diabetes prevalence, blood glucose and glycosylated hemoglobin levels in China. *Environ Int.* 2016b; 92:416–421. [PubMed: 27148900]
- Lo K, Wang M. Energy conservation in china's twelfth five-year plan period: continuation or paradigm shift? *Renew Sust Energ Rev.* 2013; 18:499–507.
- Ma Z, Hu X, Huang L, Bi J, Liu Y. Estimating ground-level PM_{2.5} in China using satellite remote sensing. *Environ Sci Technol.* 2014; 48:7436–7444. [PubMed: 24901806]
- Ma Z, Hu X, Sayer AM, Levy R, Zhang Q, Xue Y, Tong S, Bi J, Huang L, Liu Y. Satellite-based spatiotemporal trends in PM_{2.5} concentrations: China, 2004–2013. *Environ Health Perspect.* 2016; 124(2):184–192. [PubMed: 26220256]
- Matus K, Nam KM, Selin NE, Lamsal LN, Reilly JM, Paltsev S. Health damages from air pollution in China. *Glob Environ Chang.* 2012; 22(1):55–66.
- NHFPCC (National Health and Family Planning Commission of China). *China Health Statistical Yearbook 2005–2013.* Peking Union Medical University Press; Beijing: 2005–2013C.
- Pope CA, Dockery DW. Health effects of fine particulate air pollution: lines that connect. *J Air Waste Manage Assoc.* 2006; 56(6):709–742.
- Pope CA, Ezzati M, Dockery DW. Fine-particulate air pollution and life expectancy in the United States. *N Engl J Med.* 2009; 360(4):376–386. [PubMed: 19164188]
- Qian Z, Liang S, Yang S, Trevathan E, Huang Z, Yang R, Wang J, Hu K, Zhang Y, Vaughn M, Shen L, Liu W, Li P, Ward P, Yang L, Zhang W, Chen W, Dong G, Zheng T, Xu S, Zhang B. Ambient air pollution and preterm birth: a prospective birth cohort study in Wuhan, China. *Int J Hyg Environ Health.* 2016; 219(2):195–203. [PubMed: 26639560]
- Shang Y, Sun Z, Cao J, Wang X, Zhong L, Bi X, Li H, Liu W, Zhu T, Huang W. Systematic review of Chinese studies of short-term exposure to air pollution and daily mortality. *Environ Int.* 2013; 54:100–111. [PubMed: 23434817]
- Silva, RA., Adelman, Z., Fry, MM., West, JJ. The impact of individual anthropogenic emissions sectors on the global burden of human mortality due to ambient air pollution. *Environ Health Perspect.* 2016. <http://dx.doi.org/10.1289/EHP177>
- Steinle S, Reis S, Sabel CE. Quantifying human exposure to air pollution –moving from static monitoring to spatio-temporally resolved personal exposure assessment. *Sci Total Environ.* 2013; 443:184–193. [PubMed: 23183229]

- Thurston GD, Burnett RT, Turner MC, Shi Y, Krewski D, Lall R, Ito K, Jerrett M, Gapstur SM, Diver WR, Pope CA III. Ischemic heart disease mortality and long-term exposure to source-related components of US fine particle air pollution. *Environ Health Perspect*. 2016; 124(6):785–794. [PubMed: 26629599]
- USEPA. Provisional Assessment of Recent Studies on Health Effects of Particulate Matter Exposure. U.S Environmental Protection Agency; Washington, DC: 2012. <http://cfpub.epa.gov/ncea/isa/recordisplay.cfm?deid=247132> [accessed Dec 4, 2015]
- Van Donkelaar A, Martin RV, Brauer M, Ralph K, Robert L, Verduzco C, Villeneuve PJ. Global estimates of ambient fine particulate matter concentrations from satellite-based aerosol optical depth: development and application. *Environ Health Perspect*. 2010; 118:847–855. [PubMed: 20519161]
- Verstraeten WW, Neu JL, Williams JE, Bowman KW, Worden JR, Boersma KF. Rapid increases in tropospheric ozone production and export from China. *Nat Geosci*. 2015; 8(9):690–695.
- Wang S, Zhao B, Cai S, Klimont Z, Nielsen CP, Morikawa T, Woo JH, Kim Y, Fu X, Xu J, Hao J, He K. Emission trends and mitigation options for air pollutants in East Asia. *Atmos Chem Phys*. 2014; 14(13):6571–6603.
- West, JJ., Cohen, A., Dentener, F., Brunekreef, B., Zhu, T., Armstrong, B., Bell, ML., Brauer, M., Carmichael, G., Costa, DL., Dockery, DW., Kleeman, M., Krzyzanowski, M., Künzli, N., Liousse, C., Lung, CSC., Martin, RV., Pöschl, U., Pope, CA., Roberts, JM., et al. What we breathe impacts our health: improving understanding of the link between air pollution and health. *Environ Sci Technol*. 2016. <http://dx.doi.org/10.1021/acs.est.5b03827>
- Wu Y, Zhang S, Li M, Ge Y, Shu J, Zhou Y, Xu Y, Hu J, Liu H, Fu X, He K, Hao J. The challenge to NOx emission control for heavy-duty diesel vehicles in China. *Atmos Chem Phys*. 2012; 12(19): 9365–9379.
- Yang X, Huang Y, Dong P, Jiang D, Liu H. An updating system for the gridded population database of China based on remote sensing, GIS and spatial database technologies. *Sensors*. 2009; 9(2):1128–1140. [PubMed: 22399959]
- Yang F, Tan J, Zhao Q, Du Z, He K, Ma Y, Chen G. Characteristics of PM_{2.5} speciation in representative megacities and across China. *Atmos Chem Phys*. 2011; 11(11):5207–5219.
- Yang G, Wang Y, Zeng Y, Gao GF, Liang X, Zhou M, Wan X, Yu S, Jiang Y, Naghavi M, Vos T, Wang H, Lopez AD, Murray CJL. Rapid health transition in China, 1990–2010: findings from the Global Burden of Disease Study 2010. *Lancet*. 2013; 381(9882):1987–2015. [PubMed: 23746901]
- Yao L, Lu N. Spatiotemporal distribution and short-term trends of particulate matter concentration over China, 2006–2010. *Environ Sci Pollut Res*. 2014; 21:9665–9675.
- Zeng X, Vivian E, Mohammed K, Jakhar S, Vaughn M, Huang J, Zelicoff A, Xaverius P, Bai Z, Lin S, Hao Y, Paul G, Morawska L, Wang S, Qian Z, Dong G. Long-term ambient air pollution and lung function impairment in Chinese children from a high air pollution range area: the Seven Northeastern Cities (SNEC) study. *Atmos Environ*. 2016; 138:144–151.
- Zhang M, Song Y, Cai X, Zhou J. Economic assessment of the health effects related to particulate matter pollution in 111 Chinese cities by using economic burden of disease analysis. *J Environ Manag*. 2008; 88(4):947–954.
- Zhao B, Wang S, Liu H, Xu J, Fu K, Klimont Z, Hao J, He K, Cofala J, Amann M. NOx emissions in China: historical trends and future perspectives. *Atmos Chem Phys*. 2013a; 13(19):9869–9897.
- Zhao Y, Zhang J, Nielsen CP. The effects of recent control policies on trends in emissions of anthropogenic atmospheric pollutants and CO₂ in China. *Atmos Chem Phys*. 2013b; 13(2):487–508.
- Zhou M, Liu Y, Wang L, Kuang X, Xu X, Kan H. Particulate air pollution and mortality in a cohort of Chinese men. *Environ Pollut*. 2014; 186:1–6. [PubMed: 24333659]

Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.envint.2016.10.003>.




Fig. 1. National and provincial distribution of population as a function of ambient PM_{2.5} concentration in 2004 (A) and 2012 (B). Red dash vertical lines (in both plots) indicate the annual concentration limits in China's National Air Quality Standard for GradeII. Grey dash vertical lines (in both plots) demarcate boundaries of population quintiles that apportion the PM_{2.5} concentration distribution into 5 bins with equal number of exposed population in 2004.




Fig. 2.

National excess deaths attributable to air pollution from 2004 to 2012. The grey error bars indicate the 95% confidence intervals of excess deaths. The error bars of excess deaths are determined by the 95% confidence intervals of relative risk, which are asymmetrical and in different directions for different outcomes at the higher end of $PM_{2.5}$ concentration.




Fig. 3. Spatial distributions of total excess deaths attributable to PM_{2.5} exposure in 2004 (A) and 2012 (B) (cases per 100 km²). Total deaths here are the sum of deaths caused by lung cancer, IHD and stroke. Black lines show the boundary of priority areas of national joint air pollution prevention and control.




Fig. 4. Averaged attributable fractions of total deaths for 31 provinces of China. Total deaths here are the sum of deaths caused by lung cancer, IHD and stroke.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript