



CIDE gene expression in adipose tissue, liver, and skeletal muscle from obese and lean pigs^{*}

Yue-qin QIU, Xue-fen YANG^{†‡}, Xian-yong MA, Yun-xia XIONG,
 Zhi-mei TIAN, Qiu-li FAN, Li WANG, Zong-yong JIANG

(Ministry of Agriculture Key Laboratory of Animal Nutrition and Feed Science in South China, State Key Laboratory of Livestock and Poultry Breeding, Guangdong Public Laboratory of Animal Breeding and Nutrition, Guangdong Key Laboratory of Animal Breeding and Nutrition, Institute of Animal Science, Guangdong Academy of Agricultural Sciences, Guangzhou 510640, China)

[†]E-mail: yangxuefen@gdaas.cn

Received June 27, 2016; Revision accepted Oct. 5, 2016; Crosschecked May 8, 2017

Abstract: The expression of the cell death-inducing DNA fragmentation factor α -like effector (CIDE) family including *Cidea*, *Cideb*, and *Cidec* was significantly increased in mouse and human models of obesity. However, there was less information on these genes' expression in pigs. Here, we hypothesized that different fat accumulation between lean (Duroc×Landrace×Yorkshire gilts, DLY) and obese (Lantang) pigs was attributed to porcine CIDE-modulating lipid metabolism. Our data showed that *Cidea* and *Cidec* were expressed at a high level in adipose tissue, and at a relatively high level in skeletal muscle, whereas *Cideb* was mainly expressed in the liver in both breeds of pig. Lantang pigs had higher white adipose and skeletal muscle *Cidea* and *Cidec* mRNA abundance, and hepatic and muscle *Cideb* mRNA than DLY pigs. Lipid metabolism-related genes including sterol regulatory element binding protein 1c (*SREBP-1c*), hepatocyte nuclear factor-4 α (*HNF-4 α*), peroxisome proliferator-activated receptor γ coactivator-1 α (*PGC-1 α*), fatty acid synthase (*FASN*), diacylglycerol O-acyltransferase 1 (*DGAT1*), and *DGAT2* showed a higher expression level in adipose tissue from obese pigs than in that from lean pigs. Lantang pigs exhibited higher mRNA abundance for liver *SREBP-1c*, *HNF-4 α* , and *PGC-1 α* , and higher skeletal muscle *SREBP-1c*, *HNF-4 α* , *PGC-1 α* , and *DGAT2* expression, as compared with DLY pigs. However, the *perilipin2* mRNA levels in adipose tissues, liver, and skeletal muscle were significantly lower in obese pigs than in their lean counterparts. Furthermore, plasma non-esterified fatty acid (NEFA), glucose, and triacylglycerol (TAG) levels were greater in obese pigs than in lean pigs. Finally, data from correlation analysis further found that CIDE mRNA expression was positively correlated with back fat thickness (BFT), abdominal fat mass (AFM), and the levels of NEFA, TAG, and glucose in the two breeds. Collectively, these data revealed that the porcine CIDEs possibly modulated lipid metabolism and contributed to the development of fat deposition and obesity in Lantang pigs.

Key words: Cell death-inducing DNA fragmentation factor α -like effector (CIDE); Adipose tissue; Liver; Skeletal muscle; Fat deposition; Lantang pig; DLY pig

<http://dx.doi.org/10.1631/jzus.B1600294>

CLC number: S821.5

[‡] Corresponding author

^{*} Project supported by the National Natural Science Foundation of China (No. 31402086), the China Agriculture Research System (CARS-36), the National Basic Research Program (973) of China (Nos. 2013CB127301 and 2013CB127304), the Guangdong International Science and Technology Cooperation Program (No. 2014A050503049), and the Science and Technology Program of Guangdong Province (No. 2016A020210041), China

ORCID: Xue-fen YANG, <http://orcid.org/0000-0002-6989-7026>

© Zhejiang University and Springer-Verlag Berlin Heidelberg 2017

1 Introduction

Excessive fat deposition affects animal health and production efficiency, and constitutes a health risk to human consumers. Thus, modulation of fat deposition in adipose tissues of pigs is good for both animals and customers (Jiang et al., 2007). Lantang pig is a native breed (obese-type) of South China, whose carcass contains more fat content than hybrid

pigs (Lu *et al.*, 2008; Chen *et al.*, 2010). DLY is the cross breed of three lean-type pigs, Duroc, Landrace, and Yorkshire; the lean percentage of DLY reaches 63%–65% (Lan *et al.*, 2004). Lantang and DLY pigs show an obvious difference in total adipose mass and therefore offer an attractive comparison for studying the mechanism of obesity.

Adipose tissue, liver, and skeletal muscles play important roles in body lipid metabolism in animals (Ahima and Flier, 2000; Hulver *et al.*, 2003; Leonhardt and Langhans, 2004). In pigs, adipose tissue is the central organ for fat synthesis and deposition (O'Hea and Leveille, 1969). Bernlohr *et al.* (2002) demonstrated that fat accumulation in animals depends on levels of triacylglycerol (TAG) synthesis and storage and levels of lipid mobilization and fatty acid oxidation. Excess fatty acids and glucose are converted into TAG after food intake. TAG, a major energy storage form, is stored in lipid droplets, so that other intracellular organelles can avoid lipotoxicity caused by fatty acid (Girousse and Langin, 2012). The lipid droplet is an important subcellular organelle responsible for lipid storage, and the sizes of unilocular lipid droplets reveal the lipid storage capacity and have a positive association with the development of obesity (Bell *et al.*, 2008).

Changes of lipid homeostasis by over-expression or deletion of specific genes often result in obesity. Genetically modified animal models have highlighted that expression of several adipogenic and lipogenic genes, including fatty acid synthase (*FASN*), diacylglycerol *O*-acyltransferase (*DGAT*), sterol regulatory element binding protein 1c (*SREBP-1c*), and peroxisome proliferator-activated receptor γ coactivator-1 α (*PGC-1 α*), played important roles in fatty acid synthesis, TAG synthesis, and lipid storage (Liu *et al.*, 2008; Malaguarnera *et al.*, 2009). Over the past decade, it has been reported that the cell death-inducing DNA fragmentation factor α -like effector (CIDE) family plays an important role in lipid and fat metabolism (Zhou *et al.*, 2003; Gong *et al.*, 2009; Yonezawa *et al.*, 2011). Previous studies have reported that animals with a deficiency in *Cidea*, *Cideb*, or *Cidec* exhibited a typical lean phenotype with high energy expenditure, high levels of plasma TAG and non-esterified fatty acid (NEFA) reduction (Zhou *et al.*, 2003; Li *et al.*, 2007; Nishino *et al.*, 2008), and altered genes' expression in various metabolic and

signaling networks (Li *et al.*, 2010). These studies suggest that the CIDE family plays an important role in TAG synthesis, lipid storage, and the development of obesity (Gong *et al.*, 2009). However, all the studies on CIDEs have been carried out in mice and humans; there was far less information on the expression of these genes in pigs.

In the present study, we evaluated mRNA abundance of the adipose, liver, and skeletal muscle tissues for CIDEs and several lipid metabolism genes in both genetically obese and lean pigs. Our aim was to explain the relationship between porcine CIDE gene expression and lipid accumulation. We think that this basic molecular information might be useful for further investigation of the functions of CIDEs in pig models.

2 Materials and methods

2.1 Animals and sample collection

Eight castrated male Duroc \times Landrace \times Yorkshire pigs of similar liveweight ((20.21 \pm 0.57) kg) as well as eight Lantang pigs, also castrated males of similar liveweight ((16.12 \pm 0.63) kg), were used in this study. All pigs were raised in the pig farm of the animal facilities of the Institute of Animal Science in the Guangdong Academy of Agricultural Sciences, China and were provided feed and watered ad libitum. The diets for the two breeds were made up of different proportions of ingredients because the growth rates of the two breeds were different (Tables 1 and 2). The trials were terminated when the pigs of DLY and Lantang reached 100 and 70 kg body weight, respectively. An ultrasonic instrument (Renco, USA) was used to determine back fat thickness (BFT) at the 1st, 10th, and last ribs before slaughter. Blood sample was obtained from the anterior vena cava using vacuum tubes (using ethylenediaminetetra-acetic acid (EDTA) as anticoagulant) and centrifuged for 5 min at 3000g at 4 °C, and the plasma was separated and stored at -20 °C following slaughter. The left-side carcass was dissected into abdominal fat following the procedure of Walstra and Merkus (1995). Subcutaneous adipose tissue (subcutaneous back fat) and liver and skeletal muscle (longissimus muscle) samples were immediately removed and frozen in liquid nitrogen, and stored at -80 °C.

Table 1 Composition of experimental diets fed to DLY and Lantang pigs

Group	Corn (%)	Soybean meal (%)	Wheat bran (%)	Fish meal (%)	Rice bran (%)	Zeolite powder (%)	Methionine (%)	Calcium hydrophosphate (%)	Salt (%)	Premix*	Total (%)
DLY	67.38	23	4	0	0	0.96	0	0.6	0.06	4	100
Lantang	56.00	18	15	1	5	0.90	0.1	0	0	4	100

*Premix provided the following nutrients per kilogram of diet: vitamin A 1300 IU, vitamin D₃ 150 IU, vitamin E 11 mg, vitamin K₃ 0.5 mg, thiamin 1 mg, riboflavin 2 mg, pyridoxine 1 mg, vitamin B₁₂ 6 µg, niacin 7.5 mg, pantothenic acid 7 mg, biotin 0.05 mg, folacin 0.3 mg, choline chloride 300 mg, Fe 50 mg, Zn 50 mg, Mn 2 mg, Cu 3.5 mg, and I 0.14 mg

Table 2 Nutrient contents of experimental diets fed to DLY and Lantang pigs

Group	Digestive energy (MJ/kg)	Crude protein (%)	Available phosphorus (%)	Lys (%)	Met+cysteine (%)
DLY	13.38	16.0	0.25	0.76	0.58
Lantang	12.04	15.1	0.17	0.72	0.63

2.2 Real-time quantitative PCR analysis

Real-time quantitative polymerase chain reaction (PCR) was done as we described previously (Tian *et al.*, 2016). Briefly total RNA was isolated from subcutaneous adipose and liver tissue samples using TRIzol reagent (Invitrogen Co., Carlsbad, CA, USA) according to the manufacturer's instructions. All complementary DNAs (cDNAs) were synthesized from 1 µg of total RNA using a reverse transcription kit (TaKaRa, Tokyo, Japan) according to the manufacturer's recommendations. Then the synthesized cDNA was diluted (1:10, v/v) and real-time quantitative PCR amplification was performed with SYBR green I (TaKaRa, Tokyo, Japan) and specific primers for pig messenger RNA (mRNA) sequences (Table 3). Conditions for real-time PCR were an initial denaturation at 95 °C for 180 s, followed by 40 cycles at 95 °C for 15 s and 58 °C for 30 s, with a final elongation at 72 °C for 30 s. Each sample for each gene was amplified three times, in three independent wells, in order to have technical replicates. To normalize expression data, we used multiple internal control genes as described by Vandesompele *et al.* (2002). The expression stability was evaluated by the *M* value and pairwise variations of geNorm (Version 3.5; PrimerDesign Ltd., Southampton, Hampshire, UK). We found that *β-actin* had a lower *M* value than glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*), both below 1.5. Thus, *β-actin* was ranked as the most stably expressed gene. Furthermore, the pairwise variations of *β-actin* and *GAPDH* from subcutaneous

adipose and liver tissue samples of Lantang and DLY pigs were below the threshold (0.150) that required the inclusion of an additional normalization gene. Therefore, *β-actin* and *GAPDH* could be used for normalization. In the present study, the expression of the target genes relative to the *β-actin* was analyzed by the $2^{-\Delta\Delta Ct}$ method. $\Delta Ct = Ct(\text{target gene}) - Ct(\beta\text{-actin})$ and $\Delta\Delta Ct = \Delta Ct(\text{Lantang pigs}) - \Delta Ct(\text{DLY pigs})$, where Ct is cycle threshold.

2.3 Analysis of biochemical variables in plasma

Plasma concentrations of high density lipoprotein (HDL), low density lipoprotein (LDL), glucose, cholesterol, NEFA, and TAG were measured using commercial kits purchased from the Nanjing Jiancheng Institute of Bioengineering, China.

2.4 Statistical analysis

Data were presented as mean±standard error of the mean (SEM). Analysis was performed using GraphPad Prism Version 5 (GraphPad Software Inc., San Diego, CA, USA). Significance was predominantly established using a two-tailed Student's *t*-test. $P < 0.05$ was considered a statistically significant difference. Correlation analysis between CIDE mRNA expression levels and apparent index (BFT, abdominal fat mass (AFM), NEFA, TAG, and glucose) was calculated using the Pearson's correlation coefficient of the IBM SPSS Statistics 22 software (IBM Corp., Armonk, NY, USA). $P < 0.05$ and $P < 0.01$ were considered as significant and highly significant correlations, respectively.

Table 3 Primer sequences used in this study

Gene	Sequence (5'→3')	Product size (bp)	GenBank accession
<i>Cidea</i>	Forward: CACCGTGGTAGATACAGAGG	292	NM_001112696.1
	Reverse: GGACAGGAACCGCAACA		
<i>Cideb</i>	Forward: TGGGGACTCTGATGCTGAA	284	NM_001112688.1
	Reverse: CCCGTAGAATGTGGCTTTG		
<i>Cidec</i>	Forward: CGGTGCCTACTCCCTTTCTCT	184	NM_001112689.1
	Reverse: TGGGTCTTTGCCCTTGGT		
<i>Perilipin2</i>	Forward: GCTGGCGACATCTACTCA	250	NM_214200.2
	Reverse: AAGTCCACAACAGAACCCTA		
<i>DGAT1</i>	Forward: AGGACGGACACGACGAT	287	NM_214051.1
	Reverse: GAACGCAGTCACAGCAAAA		
<i>DGAT2</i>	Forward: TCCTGTCTTTCTCGTGC	131	NM_001160080.1
	Reverse: ACCTTTCTTGGGCGTGT		
<i>HNF-4α</i>	Forward: ATCGCCACCATCGTCAA	200	NM_001044571.1
	Reverse: CCTCACCTTTCCACTACCA		
<i>SREBP-1c</i>	Forward: AAGCGGACGGCTCACAA	121	NM_214157.1
	Reverse: GCAAGACGGCGGATTTATT		
<i>PGC-1α</i>	Forward: TCACCACCAAATCCTTAT	295	NM_213963.2
	Reverse: ATTCTTCCCTCTTCAGCCT		
<i>FASN</i>	Forward: CCTGGGAAGAGTGTAAGCA	108	NM_001099930.1
	Reverse: GGAACTCGGACATAGCG		
<i>β-actin</i>	Forward: CATCGTCCACCGCAAAT	210	NC_010445
	Reverse: TGTCACCTTACCGTTCC		

3 Results

3.1 Animal performance

As shown in Table 4, Lantang pigs exhibited significantly greater BFT and AFM than DLY pigs.

3.2 Expression of CIDEs in adipose, liver, and skeletal muscle tissues

As shown in Fig. 1, porcine *Cidea* and *Cidec* mRNAs were highly expressed in white adipose tissue. Both were expressed at a relatively high level in skeletal muscle. However, *Cidea* was not expressed in liver, and *Cidec* mRNA was expressed at a much lower level in liver. *Cideb* mRNA was mainly expressed in porcine liver tissue, at a lower level in adipose tissue, and was not detected in skeletal muscle. Moreover, obese pigs (Lantang) had a significantly higher *Cidea* and *Cidec* mRNA levels in adipose and skeletal muscle tissues, and a higher *Cideb* in adipose and liver than lean breeds (DLY).

3.3 Expression levels of genes responsible for lipid metabolism

The mRNA levels of *PGC-1α* and *SREBP-1c* in adipose tissue from Lantang pigs increased 3.67-fold ($P<0.01$) and 10-fold ($P<0.001$), respectively, in

comparison with DLY pigs. mRNA abundances for *HNF-4α*, *FASN*, *DGAT1*, and *DGAT2* mRNAs were significantly higher in adipose tissue from the Lantang breed than in that from DLY pigs. We observed that Lantang pigs exhibited higher *PGC-1α*, *SREBP-1c*, and *HNF-4α* mRNA expression in liver tissue than the DLY breed ($P<0.05$, $P<0.01$, and $P<0.05$, respectively). *FASN*, *DGAT1*, and *DGAT2* mRNA expression tended to be greater in liver tissue from obese pigs than in that from lean pigs, although at a non-significant level by the Student's *t*-test. In skeletal muscle, the amounts of *PGC-1α*, *SREBP-1c*, and *DGAT2* mRNAs in Lantang pigs were higher than those in DLY pigs. However, the mRNA levels of *perilipin2* in adipose, liver, and skeletal muscle from obese pigs were significantly decreased, as compared with their lean counterparts (Fig. 2).

3.4 Assessment of biochemical variables in different swine breeds

No difference in the level of HDL, LDL, or cholesterol was observed between Lantang and DLY breeds (Figs. 3d–3f). However, plasma NEFA and TAG as well as glucose levels in Lantang pigs were significantly higher than those in DLY breeds (Figs. 3a–3c).

3.5 Relationship between CIDE mRNA expression and BFT, AFM, NEFA, TAG, and glucose

The results of correlation analysis between CIDE mRNA expression and BFT, AFM, NEFA, TAG, and glucose are presented in Table 5. *Cidea* mRNA expression in adipose is positively correlated with BFT, AFM, TAG, and glucose ($P<0.01$) in the two breeds of pig. *Cidea* mRNA expression in skeletal muscle had a positively significant correlation with BFT, AFM, and NEFA ($P<0.05$). The correlations of *Cidec* mRNA expression level in adipose but not in liver or skeletal muscle, and BFT, AFM, NEFA, TAG, and glucose were positive and highly significant ($P<0.01$) in the two breeds. Expression of *Cideb* mRNA in both adipose and liver tissues correlated positively with BFT, AFM, NEFA, TAG, and glucose in two porcine breeds ($P<0.01$).




Fig. 1 CIDE gene expression patterns

Relative mRNA levels of *Cidea*, *Cideb*, and *Cidec* in adipose tissue, liver, and skeletal muscle of Lantang and DLY pigs. Data were expressed as mean±SEM, with $n=8$. * $P<0.05$, ** $P<0.01$, *** $P<0.001$ vs. DLY pigs

Table 4 Growth performance of DLY and Lantang pigs

Group	Initial body weight (kg)	Final body weight (kg)	Average back fat thickness (cm)	Abdominal fat mass (kg)
DLY	20.21±0.57	101.60±2.40	2.10±0.24	0.61±0.06
Lantang	16.12±0.63	68.90±1.83	4.05±0.03*	1.57±0.06*

Data are expressed as mean±SEM, with $n=8$. * $P<0.05$ vs. DLY pigs




Fig. 2 Relative mRNA levels of *SREBP-1c*, *HNF-4a*, *PGC-1a*, *FASN*, *DGAT1*, *DGAT2*, and *perlipin2* in adipose tissue, liver, and skeletal muscle from Lantang and DLY pigs

Data were expressed as mean±SEM, with $n=8$. * $P<0.05$, ** $P<0.01$, *** $P<0.001$ vs. DLY pigs




Fig. 3 Plasma levels of NEFA (a), TAG (b), glucose (c), total cholesterol (d), LDL-cholesterol (e), and HDL-cholesterol (f) in Lantang and DLY pigs

The results show higher plasma levels of NEFA, TAG, and glucose in Lantang pigs than in DLY pigs. Data were expressed as mean \pm SEM, with $n=8$. * $P<0.05$, ** $P<0.01$, *** $P<0.001$. NS: not statistical significant; NEFA: non-esterified fatty acid; TAG: triacylglycerol; LDL: low density lipoproteins; HDL: high density lipoproteins

Table 5 Correlation analysis between CIDE mRNA expression and BFT, AFM, NEFA, TAG, and glucose in adipose, liver, and skeletal muscle

Parameter	<i>Cidea</i> mRNA expression		<i>Cideb</i> mRNA expression		<i>Cidec</i> mRNA expression		
	Adipose	Muscle	Adipose	Liver	Adipose	Liver	Muscle
BFT	0.788**	0.580*	0.707**	0.751**	0.743**	0.120	0.214
AFM	0.818**	0.570*	0.668**	0.614**	0.668**	0.246	0.170
NEFA	0.460	0.514*	0.636**	0.549**	0.618**	-0.265	0.383
TAG	0.685**	0.488	0.665**	0.659**	0.675**	0.123	0.237
Glucose	0.652**	0.385	0.810**	0.583**	0.582**	0.103	0.067

BFT: back fat thickness; AFM: abdominal fat mass; NEFA: non-esterified fatty acid; TAG: triacylglycerol. * $P<0.05$, ** $P<0.01$

4 Discussion

In the present study, we have observed several interesting findings regarding obese Lantang pigs exhibiting higher CIDE gene expression than lean DLY pigs. Some expressions of lipid metabolism genes-related CIDEs also were up-regulated or down-regulated in Lantang pigs, as compared with DLY pigs, and the correlation analysis data found that porcine CIDE mRNA abundances were positively associated with BFT, AFM, NEFA, TAG, and glucose in the two breeds. These observations were considered novel as there were no reports, to our knowledge, comprehensively evaluating the difference

of porcine adipose, liver, or skeletal muscle CIDE mRNA expression between obese and lean pigs, or elucidating the relationship between the roles of CIDEs and fat accumulation.

Earlier studies have reported that *Cidea* was mainly expressed in the heart, and at a lower level in the brain, skeletal muscle, thymus, appendix, lymph nodes, and bone marrow, but neither in normal adult human nor mouse liver tissue (Inohara *et al.*, 1998; Zhou *et al.*, 2003). Our study found that porcine *Cidea* was highly expressed in white adipose and at a relatively high level in skeletal muscle tissue, but was not detected in liver, although Li *et al.* (2009) showed that porcine *Cidea* can be detected in the liver of

Tongcheng and Large White pigs. Danesch *et al.* (1992) demonstrated that *Cidec* was an adipocyte-specific marker gene. Consistently, we found that porcine *Cidec* gene mRNA was expressed at a high level in white adipose tissue, at a relatively high level in skeletal muscle, and at a lower level in liver. It has been reported that *Cideb* mRNA was expressed in many tissues with the highest levels in the liver (Inohara *et al.*, 1998). Our result agreed with the report that porcine *Cideb* was mainly expressed in the liver, but was not detected in skeletal muscle. The differential tissue distribution patterns of the three CIDE genes can imply that they may have different functions.

Previous studies reported that with CIDE deficiency mice showed a typical lean phenotype and a markedly lower adiposity index. Plasma TAG and NEFA levels, and the size of white adipocytes in CIDE-deficient mice were significantly reduced, compared with those in wild-type mice (Li *et al.*, 2007; Nishino *et al.*, 2008; Toh *et al.*, 2008; Gong *et al.*, 2009). In addition, it has been shown that over-expressed *Cidec* can increase lipid droplet size and enhance the accumulation of lipids (Keller *et al.*, 2008), whereas deletion of *Cidea* by RNA interference in human adipocyte stimulated lipolysis (Nordström *et al.*, 2005). In our results, we found that mRNA expressions of *Cidea* and *Cidec* in adipose tissue and skeletal muscle, and *Cideb* in adipose and liver from obese pigs were significantly higher than that from the lean breed. These data indicated that higher mRNA abundance for *Cidea* and *Cidec* in adipose tissue and skeletal muscle, and *Cideb* in liver from Lantang breeds, might play an important role in the development of fat deposition.

The molecular basis for CIDE-modulated lipid metabolism has been elucidated. CIDE deletion in mice resulted in significant reductions of the expression of *SREBP-1c* and its downstream target genes (*ACCI*, *FASN*, *Elovl6*, and *SCD1*) (Abu-Elheiga *et al.*, 2003; Li *et al.*, 2007), which were required for fatty acid synthesis (Shimomura *et al.*, 1999) and TAG synthesis (Horton *et al.*, 2002). Similarly, we found that CIDE-enriched Lantang pigs exhibited greater mRNA abundance for *SREBP-1c* and *FASN*. Therefore, it has been indicated that CIDEs could positively modulate the expression of *SREBP-1c* and its downstream target genes, and lead to increased TAG secretion and lipogenesis (Gong *et al.*, 2009). *PGC-1 α*

and *HNF-4 α* mRNA expression, which were critical to de novo lipogenesis and gluconeogenesis (Herzig *et al.*, 2001) and controlling the expression of *Cidea* and *Cideb* (Hallberg *et al.*, 2008; Chen *et al.*, 2010; Yu *et al.*, 2013), were higher in adipose and liver tissue from obese pigs than in those from their lean counterparts. At the same time, skeletal muscle *PGC-1 α* mRNA abundance from Lantang pigs was increased compared to the DLY pigs. In addition, adipose *DGAT1* and *DGAT2* mRNA expression, which play a central modulation role in animal fat deposition (Nishizuka, 1992), were higher in Lantang pigs than in DLY pigs. Skeletal muscle *DGAT2* mRNA abundance was also increased in DLY pigs. The present data uncovered that the mRNA levels of *perilipin2* in adipose, liver, and skeletal muscle tissue from CIDE-enriched Lantang pigs were lower than those from DLY pigs. Our result was consistent with the report by Singaravelu *et al.* (2013) who demonstrated that *Cideb* over-expression resulted in a significant down-regulation of *perilipin2* protein levels. In addition, a study by Li *et al.* (2012) indicated that *Cideb* and *perilipin2* played opposite roles in modulating TAG accumulation, with *Cideb* as a positive regulator and *perilipin2* as a negative regulator of lipid droplet size in hepatocytes. Collectively, these results revealed that the CIDE gene family modulated fat deposition through controlling the expression of lipid metabolism-related genes.

CIDE-sufficient Lantang pigs also exhibited higher plasma NEFA, TAG, and glucose levels than DLY pigs. Our results agreed with the reports that CIDE deficiency resulted in a reduction of plasma TAG and fatty acid levels (Zhou *et al.*, 2003; Li *et al.*, 2007; Toh *et al.*, 2008). We also found that Lantang pigs had greater BFT and AFM than DLY pigs (Lu *et al.*, 2008). Therefore, there may be a close association between CIDE mRNA expression and fat deposition. The further correlation analysis results uncovered an interesting finding that *Cidea* mRNA expression in adipose was positively correlated with BFT, AFM, TAG, and glucose, and in skeletal muscle was positively correlated with BFT, AFM, and NEFA in the two breeds of pigs. Adipose and liver *Cideb* mRNA abundance and BFT, AFM, TAG, NEFA, and glucose had a positively significant correlation in the two breeds. *Cidec* mRNA in adipose but not in liver or skeletal muscle was positively correlated with BFT,

AFM, TAG, NEFA, and glucose in Lantang and DLY pigs. We made a correlation analysis between porcine CIDE mRNA and fat deposition-related factors. These results further suggested that CIDE genes contributed to fat deposition of the fatty Lantang pigs.

5 Conclusions

In conclusion, our data suggest that the porcine CIDEs possibly modulated lipid metabolism and contributed to the development of fat deposition and obesity.

Compliance with ethics guidelines

Yue-qin QIU, Xue-fen YANG, Xian-yong MA, Yun-xia XIONG, Zhi-mei TIAN, Qiu-li FAN, Li WANG, and Zong-yong JIANG declare that they have no conflict of interest.

All institutional and national guidelines for the care and use of laboratory animals were followed.

References

- Abu-Elheiga, L., Oh, W., Kordari, P., et al., 2003. Acetyl-CoA carboxylase 2 mutant mice are protected against obesity and diabetes induced by high-fat/high-carbohydrate diets. *Proc. Natl. Acad. Sci. USA*, **100**(18):10207-10212. <http://dx.doi.org/10.1073/pnas.1733877100>
- Ahima, R.S., Flier, J.S., 2000. Adipose tissue as an endocrine organ. *Trends Endocrinol. Metab.*, **11**(8):327-332. [http://dx.doi.org/10.1016/S1043-2760\(00\)00301-5](http://dx.doi.org/10.1016/S1043-2760(00)00301-5)
- Bell, M., Wang, H., Chen, H., et al., 2008. Consequences of lipid droplet coat protein downregulation in liver cells: abnormal lipid droplet metabolism and induction of insulin resistance. *Diabetes*, **57**(8):2037-2045. <http://dx.doi.org/10.2337/db07-1383>
- Bernlohr, D.A., Jenkins, A.E., Bennaars, A.A., 2002. Adipose tissue and lipid metabolism. In: Vance, D.E., Vance, J.E. (Eds.), *Biochemistry of Lipids, Lipoproteins and Membranes*, 4th Ed. Elsevier, Amsterdam, p.263-289. [http://dx.doi.org/10.1016/S0167-7306\(02\)36012-5](http://dx.doi.org/10.1016/S0167-7306(02)36012-5)
- Chen, Z.J., Norris, J.Y., Finck, B.N., 2010. Peroxisome proliferator-activated receptor- γ coactivator-1 α (PGC-1 α) stimulates VLDL assembly through activation of cell death-inducing DFFA-like effector B (CideB). *J. Biol. Chem.*, **285**(34):25996-26004. <http://dx.doi.org/10.1074/jbc.M110.141598>
- Chen, Z.M., Qi, X.H., Zhang, H., et al., 2010. Changes of leptin and leptin receptor gene expression in subcutaneous fat and hypothalamus of Lantang and Landrace pigs. *J. Huazhong Agric. Univ.*, **29**(1):67-70 (in Chinese).
- Danesch, U., Hoeck, W., Ringold, G.M., 1992. Cloning and transcriptional regulation of a novel adipocyte-specific gene, FSP27. CAAT-enhancer-binding protein (C/EBP) and C/EBP-like proteins interact with sequences required for differentiation-dependent expression. *J. Biol. Chem.*, **267**(10):7185-7193.
- Girousse, A., Langin, D., 2012. Adipocyte lipases and lipid droplet-associated proteins: insight from transgenic mouse models. *Int. J. Obes. (Lond.)*, **36**(4):581-594. <http://dx.doi.org/10.1038/ijo.2011.113>
- Gong, J., Sun, Z., Li, P., 2009. CIDE proteins and metabolic disorders. *Curr. Opin. Lipidol.*, **20**(2):121-126. <http://dx.doi.org/10.1097/MOL.0b013e328328d0bb>
- Hallberg, M., Morganstein, D.L., Kiskinis, E., et al., 2008. A functional interaction between RIP140 and PGC-1 α regulates the expression of the lipid droplet protein CIDEA. *Mol. Cell. Biol.*, **28**(22):6785-6795. <http://dx.doi.org/10.1128/MCB.00504-08>
- Herzig, S., Long, F., Jhala, U.S., et al., 2001. CREB regulates hepatic gluconeogenesis through the coactivator PGC-1. *Nature*, **413**(6852):179-183. <http://dx.doi.org/10.1038/35093131>
- Horton, J.D., Goldstein, J.L., Brown, M.S., 2002. SREBPs: activators of the complete program of cholesterol and fatty acid synthesis in the liver. *J. Clin. Invest.*, **109**(9):1125-1131. <http://dx.doi.org/10.1172/JCI115593>
- Hulver, M.W., Berggren, J.R., Cortright, R.N., et al., 2003. Skeletal muscle lipid metabolism with obesity. *Am. J. Physiol. Endocrinol. Metab.*, **284**(4):E741-E747. <http://dx.doi.org/10.1152/ajpendo.00514.2002>
- Inohara, N., Koseki, T., Chen, S., et al., 1998. CIDE, a novel family of cell death activators with homology to the 45 kDa subunit of the DNA fragmentation factor. *EMBO J.*, **17**(9):2526-2533. <http://dx.doi.org/10.1093/emboj/17.9.2526>
- Jiang, J.P., Zhou, J., Chen, J., et al., 2007. Effect of chicken egg yolk antibody against adipose tissue plasma membranes on carcass composition and lipogenic hormones and enzymes in pigs. *Livestock Sci.*, **107**(2-3):235-243. <http://dx.doi.org/10.1016/j.livsci.2006.09.020>
- Keller, P., Petrie, J.T., de Rose, P., et al., 2008. Fat-specific protein 27 regulates storage of triacylglycerol. *J. Biol. Chem.*, **283**(21):14355-14365. <http://dx.doi.org/10.1074/jbc.M708323200>
- Lan, L.T., Huang, L.S., Ma, J.W., et al., 2004. Experiment for comparing the performance of Erhualian pig double cross combinations and that of Duroc \times (Landrace \times Large Yorkshire) three-way cross combination. *J. Southwest Univ. Natl.*, **30**(6):741-744.
- Leonhardt, M., Langhans, W., 2004. Fatty acid oxidation and control of food intake. *Physiol. Behav.*, **83**(4):645-651. <http://dx.doi.org/10.1016/j.physbeh.2004.07.033>
- Li, J.Z., Ye, J., Xue, B., et al., 2007. Cideb regulates diet-induced obesity, liver steatosis, and insulin sensitivity by controlling lipogenesis and fatty acid oxidation. *Diabetes*, **56**(10):2523-2532. <http://dx.doi.org/10.2337/db07-0040>
- Li, J.Z., Lei, Y., Wang, Y., et al., 2010. Control of cholesterol biosynthesis, uptake and storage in hepatocytes by Cideb. *Biochim. Biophys. Acta*, **1801**(5):577-586. <http://dx.doi.org/10.1016/j.bbali.2010.01.012>
- Li, X.H., Ye, J., Zhou, L.K., et al., 2012. Opposing roles of cell death-inducing DFF45-like effector B and perilipin 2 in controlling hepatic VLDL lipitation. *J. Lipid Res.*, **53**(9):1877-1889. <http://dx.doi.org/10.1194/jlr.M026591>
- Li, Y.H., Lei, T., Chen, X.D., et al., 2009. Molecular cloning,

- chromosomal location and expression pattern of porcine *CIDEa* and *CIDEc*. *Mol. Biol. Rep.*, **36**(3):575-582.
<http://dx.doi.org/10.1007/s11033-008-9216-5>
- Liu, Y., Millar, J.S., Cromley, D.A., et al., 2008. Knockdown of Acyl-CoA: diacylglycerol acyltransferase 2 with antisense oligonucleotide reduces VLDL TG and ApoB secretion in mice. *Biochim. Biophys. Acta*, **1781**(3):97-104.
<http://dx.doi.org/10.1016/j.bbali.2008.01.001>
- Lu, P., Li, D.F., Yin, J.D., et al., 2008. Flavour differences of cooked longissimus muscle from Chinese indigenous pig breeds and hybrid pig breed (Duroc×Landrace×Large White). *Food Chem.*, **107**(4):1529-1537.
<http://dx.doi.org/10.1016/j.foodchem.2007.10.010>
- Malaguarrera, M., Di Rosa, M., Nicoletti, F., et al., 2009. Molecular mechanisms involved in NAFLD progression. *J. Mol. Med. (Berl.)*, **87**(7):679-695.
<http://dx.doi.org/10.1007/s00109-009-0464-1>
- Nishino, N., Tamori, Y., Tateya, S., et al., 2008. FSP27 contributes to efficient energy storage in murine white adipocytes by promoting the formation of unilocular lipid droplets. *J. Clin. Invest.*, **118**(8):2808-2821.
<http://dx.doi.org/10.1172/JCI34090>
- Nishizuka, Y., 1992. Intracellular signaling by hydrolysis of phospholipids and activation of protein kinase. *Science*, **258**(5082):607-614.
<http://dx.doi.org/10.1126/Science.1411571>
- Nordström, E.A., Rydén, M., Backlund, E.C., et al., 2005. A human-specific role of cell death-inducing DFFA (DNA fragmentation factor- α)-like effector A (CIDEA) in adipocyte lipolysis and obesity. *Diabetes*, **54**(6):1726-1734.
<http://dx.doi.org/10.2337/diabetes.54.6.1726>
- O'Hea, E.K., Leveille, G.A., 1969. Significance of adipose tissue and liver as sites of fatty acid synthesis in the pig and the efficiency of utilization of various substrates for lipogenesis. *J. Nutr.*, **99**(3):338-344.
- Shimomura, I., Bashmakov, Y., Horton, J.D., 1999. Increased levels of nuclear SREBP-1c associated with fatty livers in two mouse models of diabetes mellitus. *J. Biol. Chem.*, **274**(4):30028-30032.
<http://dx.doi.org/10.1074/jbc.274.42.30028>
- Singaravelu, R., Lyn, R.K., Srinivasan, P., et al., 2013. Human serum activates CIDEB-mediated lipid droplet enlargement in hepatoma cells. *Biochem. Biophys. Res. Commun.*, **441**(2):447-452.
<http://dx.doi.org/10.1016/j.bbrc.2013.10.080>
- Tian, Z.M., Ma, X.Y., Yang, X.F., et al., 2016. Influence of low protein diets on gene expression of digestive enzymes and hormone secretion in the gastrointestinal tract of young weaned piglets. *J. Zhejiang Univ.-Sci. B (Biomed. & Biotechnol.)*, **17**(10):742-751.
<http://dx.doi.org/10.1631/jzus.B1600229>
- Toh, S.Y., Gong, J., Du, G., et al., 2008. Up-regulation of mitochondrial activity and acquirement of brown adipose tissue-like property in the white adipose tissue of *Fsp27* deficient mice. *PLoS ONE*, **3**(8):e2890.
<http://dx.doi.org/10.1371/journal.pone.0002890>
- Vandesompele, J., de Preter, K., Pattyn, F., et al., 2002. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol.*, **3**(7):research0034.1.
<http://dx.doi.org/10.1186/gb-2002-3-7-research0034>
- Walstra, P., Merkus, G.S.M., 1995. Procedure for assessment of the lean meat percentage as a consequence of the new EU reference dissection method in pig carcass classification. Report ID-DLO 96.014, Zeist, the Netherlands.
- Yonezawa, T., Kurata, R., Kimura, M., et al., 2011. Which CIDE are you on? Apoptosis and energy metabolism. *Mol. Biosyst.*, **7**(1):91-100.
<http://dx.doi.org/10.1039/c0mb00099j>
- Yu, M., Wang, H., Zhao, J., et al., 2013. Expression of CIDE proteins in clear cell renal cell carcinoma and their prognostic significance. *Mol. Cell. Biochem.*, **378**(1):145-151.
<http://dx.doi.org/10.1007/s11010-013-1605-y>
- Zhou, Z., Yon Toh, S., Chen, Z., et al., 2003. *Cidea*-deficient mice have lean phenotype and are resistant to obesity. *Nat. Genet.*, **35**(1):49-56.
<http://dx.doi.org/10.1038/ng1225>

中文概要

题目: 肥胖型和瘦肉型猪的脂肪、肝脏及骨骼肌组织中 *CIDE* 家族基因表达水平的比较研究

目的: 研究 *CIDE* 家族基因在肥胖型和瘦肉型猪的脂肪、肝脏及肌肉组织中的基因表达水平差异, 并初步探 *CIDE* 家族基因与脂质代谢的关系。

创新点: 首次在肥胖型与瘦肉型猪模型中解释 *CIDE* 家族基因可以调节脂质代谢, 并有助于脂肪沉积及导致肥胖。

方法: 采用荧光定量聚合酶链式反应 (qPCR) 检测肥胖型蓝塘猪和瘦肉型杜长大猪的脂肪、肝脏和骨骼肌中 *CIDE* 家族基因、*SREBP-1c*、*PGC-1 α* 、*HNF-4 α* 、*FASN*、*DGAT1* 和 *DGAT2*、*perlipin 2* 等基因表达水平。采用血浆生化指标试剂盒检测两个品种猪血浆中甘油三酯、葡萄糖、游离脂肪酸及胆固醇的含量。

结论: 肥胖型蓝塘猪脂肪和背最长肌组织中的 *Cidea* 和 *Cidec*, 及肝脏中 *Cidec* 的基因表达量明显高于瘦肉型杜长大猪。在脂肪组织中, 脂质代谢相关的基因 (包括 *SREBP-1c*、*PGC-1 α* 、*HNF-4 α* 、*FASN*、*DGAT1* 和 *DGAT2* 基因) 表达量都是蓝塘猪高于杜长大猪。蓝塘猪肝脏中的 *SREBP-1c*、*HNF-4 α* 和 *PGC-1 α* 基因表达水平显著高于杜长大猪。蓝塘猪背最长肌组织的 *SREBP-1c*、*HNF-4 α* 、*PGC-1 α* 和 *DGAT2* 基因表达量高于杜长大猪。然而, 蓝塘猪的脂肪、肝脏及背最长肌三种组织中的 *perlipin 2* 的表达量显著低于杜长大猪。此外, 蓝塘猪血浆中的甘油三酯、葡萄糖及游离脂肪酸浓度明显高于杜长大猪。通过相关性分析, 我们发现肥胖型和瘦肉型猪不同组织中的 *CIDE* 家族基因表达水平与背部脂肪厚度、腹部脂肪重量、血浆中的甘油三酯、葡萄糖及游离脂肪酸浓度有明显的正向相关性。综上所述, *CIDE* 家族基因可以调节脂质代谢, 并促进脂肪沉积及导致肥胖。

关键词: *CIDE* 家族基因; 脂肪沉积; 脂肪; 肝脏; 骨骼肌; 蓝塘猪; 杜长大猪