



Published in final edited form as:

Cancer Discov. 2017 June ; 7(6): 596–609. doi:10.1158/2159-8290.CD-16-1337.

Prospective comprehensive molecular characterization of lung adenocarcinomas for efficient patient matching to approved and emerging therapies

Emmet J. Jordan^{1,*}, Hyunjae R. Kim^{6,*}, Maria E. Arcila⁶, David Barron³, Debyani Chakravarty⁹, JianJiong Gao⁹, Matthew T. Chang^{2,7}, Andy Ni⁷, Ritika Kundra⁹, Philip Jonsson^{2,7}, Gowtham Jayakumaran⁶, Sizhi Paul Gao², Hannah C. Johnsen², Aphrothiti J. Hanrahan², Ahmet Zehir⁶, Natasha Rekhman⁶, Michelle S. Ginsberg⁸, Bob T. Li⁴, Helena A. Yu⁴, Paul K. Paik⁴, Alexander Drilon⁴, Matthew D. Hellmann⁴, Dalicia N. Reales², Ryma Benayed⁶, Valerie W. Rusch⁵, Mark G. Kris⁴, Jamie E. Chaff⁴, Jose Baselga^{1,2}, Barry S. Taylor^{2,7,9}, Nikolaus Schultz^{7,9}, Charles M. Rudin⁴, David M. Hyman¹, Michael F. Berger^{6,9}, David B. Solit^{1,2,9}, Marc Ladanyi^{2,6}, and Gregory J. Riely⁴

¹Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA

²Human Oncology and Pathogenesis Program, Memorial Sloan Kettering Cancer Center, NY, USA

³Department of Radiation Oncology, Memorial Sloan Kettering Cancer Center, New York, NY, USA

⁴Thoracic Oncology Service, Division of Solid Tumor Oncology, Department of Medicine, Weill Cornell Medical College, Memorial Sloan Kettering Cancer Center, New York, NY, USA

⁵Department of Surgery, Thoracic Oncology Service, Memorial Sloan Kettering Cancer Center, New York, NY, USA

⁶Department of Pathology, Molecular Diagnostics Service, Memorial Sloan Kettering Cancer Center, New York, NY, USA

⁷Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, New York, USA

⁸Department of Radiology, Memorial Sloan Kettering Cancer Center, New York, New York, USA

⁹Marie-Josée and Henry R. Kravis Center for Molecular Oncology, Memorial Sloan Kettering Cancer Center, New York, New York, USA

Abstract

Corresponding Authors: Dr Gregory J. Riely MD, Memorial Sloan Kettering Cancer Center, 1275 York Avenue, New York, NY, 10065, Phone: 646-888-4199, rielyg@mskcc.org.

*These authors contributed equally to this manuscript

Conflicts of interest: DM Hyman is a consultant/advisory board member for Atara, Chugai, CytomX and has received research funding from LOXO, AstraZeneca and PUMA. M Ladanyi is a consultant/advisory board member for NCCN/AstraZeneca and has received research funding from LOXO. GJ Riely has been a consultant for Genentech and Novartis and receives research funding from Novartis, Pfizer, and Ariad. No potential conflicts of interest were reported by other authors.

Tumor genetic testing is standard of care for patients with advanced lung adenocarcinoma but the fraction of patients who derive clinical benefit remains undefined. Here, we report the experience of 860 patients with metastatic lung adenocarcinoma analyzed prospectively for mutations in >300 cancer-associated genes. Potentially actionable genetic events were stratified into one of four levels based upon published clinical or laboratory evidence that the mutation in question confers increased sensitivity to standard or investigational therapies. Overall 37.1% (319/860) of patients received a matched therapy guided by their tumor molecular profile. Excluding alterations associated with standard of care therapy, 14.4% (69/478) received matched therapy with a clinical benefit of 52%. Use of matched therapy was strongly influenced by the level of pre-existent clinical evidence that the mutation identified predicts for drug response. Analysis of genes mutated significantly more often in tumors without known actionable mutations nominated *STK11* and *KEAP1* as possible targetable mitogenic drivers.

Keywords

lung adenocarcinoma; gene sequencing; matched therapy; OncoKB

Introduction

Tumor genetic testing is standard of care for patients with non-small cell lung cancer (NSCLC). Lung adenocarcinomas, which account for approximately 50% of lung cancers, are molecularly sub-classified and their therapy dictated by the presence of distinct molecular alterations including *EGFR* mutations and *ALK* or *ROS1* fusions that confer sensitivity to selective kinase inhibitors (1–4). Additional alterations such as *BRAF V600E*, *RET* fusions, and *ERBB2* amplifications are found in smaller subsets of patients, but when present may also predict for response to targeted inhibitors that are FDA-approved therapies for other tumor types (5–11). In other patients, defined oncogenic drivers such as *KRAS* and *PIK3CA* mutations are detected for which preclinical studies have nominated targeted approaches, but the clinical utility of such therapies has yet to be established (12, 13). As a result of advances in DNA sequencing, the prospective molecular analysis of tumors for mutations in hundreds of cancer-associated genes is now feasible using multiplexed assays that use as input small quantities of formalin fixed paraffin embedded tissue (FFPE). With the goal of optimizing treatment selection in patients with advanced cancer and to address the limitations in sensitivity and breadth of previous prospective clinical testing approaches, we developed the Memorial Sloan Kettering-Integrated Mutation Profiling of Actionable Cancer Targets (MSK-IMPACT) assay, a hybridization capture-based, next-generation sequencing platform (14) for matched tumor: normal sequencing to comprehensively profile somatic alterations in all known cancer genes in solid tumors.

Here, we report our experience with the first 860 patients with recurrent or metastatic lung adenocarcinoma analyzed by MSK-IMPACT, with a focus on defining the fraction of patients for whom such testing influenced treatment selection. Potentially actionable genetic events were stratified into one of four categories based on the level of evidence supporting the utility of the mutation as a predictive biomarker of drug response. Outcome data were collected to determine whether patients were treated with, and benefited from, a therapy

chosen on the basis of a specific molecular event present in their tumor and to determine whether the likelihood of receiving a matched therapy correlated with the level of pre-existent evidence that the particular genomic event correlated with drug response.

Results

Patient Demographics

A total of 915 tumors from 860 patients with recurrent or metastatic lung adenocarcinoma were profiled using MSK-IMPACT during the study period (Table 1). A new biopsy was not required for participation in this study and the median duration between collection of the tumor sample and its use for genomic analysis was 28 days (range 0 to 3274 days, the longer intervals reflecting the testing of older samples resected up to several years previously due to a lack of availability of a more recent biopsy in a minority of patients). In 765 (89%) of the patients tumor tissue 1 year old was utilized, with tissue 30 days old analyzed in 473 (55%). The mean time from receipt of the tumor sample and matched blood sample in the clinical laboratory to the reporting of MSK-IMPACT results was 17 days. Patients had received a median of one (range 0–7) prior systemic treatment prior to MSK-IMPACT testing. At the time of analysis, the median follow up time from diagnosis of metastatic disease was 13 months (1–196 months) and 239 (27.8%) patients had died.

Known mitogenic drivers identified by MSK-IMPACT

Potentially actionable somatic alterations, as defined by the OncoKB classification (15), were identified in 747 patients (86.9%), (Figure 1A, Supplementary Table S1). We next compared the frequency of potential actionable alterations in this prospective cohort to that observed in the retrospective TCGA analysis of untreated lung adenocarcinomas (16), Figure 1B. By definition, all patients in the MSK-IMPACT cohort had recurrent or metastatic disease as compared to 3.9% (9/230) with stage IV disease in the TCGA dataset. Moreover, 37.7% (345/915) of tumors used for MSK-IMPACT analysis were collected following treatment with at least one prior systemic therapy whereas only samples from treatment naïve patients were included in the TCGA cohort. The MSK-IMPACT dataset therefore represents a clinically aggressive cohort distinct from the TCGA set which examined primary resection, treatment naive tumors only. Notably, the MSK-IMPACT cohort included a higher fraction of patients with activating *EGFR* alterations (27% vs 11%; $p < 0.001$) but other molecular subsets showed no significant differences. In addition, 5.5% (47/860) had a *EGFR* T790M mutation, all detected post *EGFR*-TKI therapy, as compared to 0.4% (1/230) in the TCGA dataset ($p < 0.001$). Factors contributing to the higher frequency of *EGFR* mutant patients in this cohort may include a referral bias attributable to the availability of *EGFR* T790M focused clinical trials at our institution during the study period or differences in patient's demographics as 32.2% of the MSK-IMPACT patients were never smokers and 8.4% (72/860 pts) were Asian. Conversely, some oncogenic drivers were present at higher rates in the TCGA cohort than in the MSK-IMPACT cohort, including truncating mutations/deletions in *NFI* (8.3% vs 2%; $p < 0.001$) and *BRAF* mutations (7% vs 3.6%; $p = 0.042$).

Consistent with prior studies, mutations in *EGFR* and *KRAS* were the most commonly identified oncogenic drivers and were with very rare exception mutually exclusive ($p < 0.0001$), (Figure 1C) (17, 18). Oncogenic fusions in *ALK*, *ROS1*, and *RET*, *MET* exon 14 alterations and *BRAF*V600E mutations, all of which predict for significant clinical benefit to targeted inhibitors of these kinases, were identified in 1.7 to 3.8% of patients. Less common actionable drivers were identified in 110 (12.8%) patients (Figure 1A) and included RAS/MAPK pathway lesions such as truncating mutations/deletion of *NFI* (16 pts) and known activating mutations in *NRAS* (10 pts; 9 Q61, 1 G13), *HRAS* (1 pt) and *MAP2K1* (*MEK1*) (E203K; n=1, K57N; n=2, Q56P; n=1, G128V; n=1 and E102_I103 deletion; n=1), (19, 20). Two patients had hotspot *ARAF* mutations at codon 214 (S214Y, S214P) which have been shown to confer sensitivity to sorafenib (21) and two tumors harbored *RAF1* S257L mutations, a previously characterized hotspot (21).

Consistent with prior data, most oncogenic or likely oncogenic PI3K alterations were identified in tumors with a co-occurring higher-level alteration (22, 23). In total, 32/860 (3.7%) tumors had activating mutations in the PI3K/AKT/mTOR pathway as their highest-level actionable alteration including 17 with *PIK3CA* mutations, 6 inactivating *PTEN* alterations, 3 patients each with truncating *TSC1/TSC2* mutations, 2 *AKT1* E17K mutations and one *mTOR* S2215Y mutation. Nine (53%) of the seventeen patients with oncogenic *PIK3CA* mutations as their highest level alteration had a co-occurring *KRAS* mutation. Likely inactivating somatic *BRCA1* (n=3, 0.3%) or *BRCA2* truncating mutations (n=8, 0.9%) were deemed level 2B alterations based upon the FDA approval of the PARP inhibitor olaparib in *BRCA*-mutant ovarian carcinomas. Additionally, one patient had a *CD74-NRG* fusion (24) and a second a *FGFR3-TACC3* fusion (25), as their highest actionable alteration. An activating exon 9 *KIT*E490Q mutation, previously described in thymic carcinoma, was present in one patient (26) and one patient had a *ERBB3*D297H hotspot mutation (27). Although these rare alterations were detected in only a small number of individuals, they highlight the ability of multiplexed sequencing assays to detect rare but potentially actionable drug targets that in aggregate represent key oncogenic drivers in a small but non-trivial 0.5% of lung cancers.

In total, 239 patients had tumors with co-occurring targetable mutations with 46 patients having at least three concurrent level 1–4 alterations (Supplementary Table S2). Fifty two patients had co-occurring level 1–3 mutations of which 25 were patients with co-occurring *EGFR* and *PIK3CA* mutations, Supplementary Fig. S1. Notably, only two *EGFR* mutated patients (out of 214) had a concurrent *KRAS* mutation (level 4). One patient with an *EGFR* L858R mutation had co-occurrent *KRAS* Q61H and Q22K mutations. This specimen was collected prior to *EGFR*-TKI therapy for which the patient received erlotinib for six months until progression. The variant allele frequency of *EGFR* L858R was higher (0.59) as compared to *KRAS* Q61 (0.04) and Q22 (0.03) suggesting subclonal *KRAS*-mutant populations, Supplementary Fig. S2. The second patient had an *EGFR* exon 19 deletion with a *KRAS* Q61R mutation. This tumor was sequenced following disease progression after prior therapy with erlotinib for 26 months. The allele frequency of the *EGFR* mutation was (0.12) with the coexisting *KRAS* Q61R (0.35), Supplementary Fig. S3. *KRAS* mutations at codon 61 are rare in comparison to those located at codon 12 in NSCLC and were seen in 12/235 (0.05%) of *KRAS* mutated tumors in the dataset. Eight patients had a *KRAS* G12

mutation concurrent with a level 2B (*BRCA1/2* or *TSC1/2* loss) alteration. Therefore, multiple potentially targetable lesions with different levels of support sometimes co-existed within tumors, highlighting the challenge of defining therapeutic actionability in the setting of more comprehensive tumor profiling. Despite the frequent identification of two or more targetable driver mutations in individual tumors, no patient in this prospective series was treated on a clinical trial that simultaneously targeted two actionable alterations.

Use of matched therapy

Overall, 37.1% (319/860) of patients received a matched therapy guided by their tumor molecular profile with the likelihood of receiving a matched therapy correlating strongly with the level of evidence that the mutation identified predicts for drug response ($p < 0001$, Figure 2A). Specifically, the majority of patients with level 1 (92%) and 2A (52%) alterations received matched therapy whereas only a minority of patients with level 2B (17%), level 3 (25%), or level 4 (2%) alterations received matched therapy (Figure 2A). In total, 95.3%, 90.9%, and 59.1% of patients with sensitizing *EGFR* mutations, *ALK* fusions, and *ROS1* fusions received matched therapy with clinical benefit documented in 84.8%, 93.3% and 84.6% of patients, respectively (Figure 2B). For patients with level 2A alterations, matched therapy was used for patients with *MET* exon 14 alterations (65.4%), *BRAF*V600E mutations (55.6%), *RET* fusions (53.3%) and amplification of wild-type *MET* (16.7%). Despite the lower use of matched therapy in patients with level 2A mutations, clinical benefit with the matched targeted agent was substantial in such patients with 76.5%, 72.7%, 75%, and 50% of patients with *MET* exon 14 alterations, *RET* fusions, *BRAF*V600E mutations and amplification of wild-type *MET* deriving clinical benefit, respectively. Ongoing active treatment with chemotherapy or immunotherapy was the most common reason that matched therapy was not used in patients with a *RET* (n=7) or *ROS1* fusion (n=6) (Supplementary Fig. S4). There were two patients with *ROS1* fusions (9%) who experienced rapid deterioration and thus did not receive treatment with a *ROS1* kinase inhibitor. Notably, both died after crizotinib had been shown to be active in patients with *ROS1* fusions but prior to its FDA-approval for this indication in March of 2016. One of these patients had local molecular testing for *EGFR* only and had received three lines of systemic therapy prior to referral at which time the *ROS1* fusion was detected by MSK-IMPACT. Unfortunately, the patient died of disease 12 days after referral, before *ROS1*-directed therapy could be initiated. No patients with a presumed inactivating alteration in *TSC1/2* or *BRCA1/2* (level 2B) received matched therapy. Finally, 5/12 (42%) patients with *ERBB2* amplified tumors received matched therapy, of which one demonstrated evidence of clinical benefit (Figure 2B).

For patients with level 3 and 4 alterations, matched therapies were typically offered only within the context of a clinical trial. Most notably, the use of matched therapy was also exceptionally rare in patients with *KRAS* mutant tumors (2/218; 0.9%). 50% of patients with an *ERBB2* mutation received a matched therapy, 40% of whom experienced clinical benefit. Four of 13 patients with a non-V600E *BRAF* alteration (K601E, D594G, T599 duplication, *SND1*-*BRAF* fusion) received matched therapy with none deriving clinical benefit. Three of these patients received single-agent *MEK* or *ERK* inhibitor therapy, while the patient with a *BRAF*-*SND1* fusion received a combination of *BRAF* and *MEK* inhibitors. Two patients

with *AKT1* E17K mutations received matched therapy with one achieving clinical benefit lasting 12 months. No patients with mutations in *MEK1* (*MAP2K1*), *RAF1*, *ARAF*, *FGFR3*, or deletions in *CDKN2A* as their highest actionable alterations received matched therapy. While many of these later patients are still benefiting from chemotherapy or immunotherapy and may receive matched therapy in the future, the lack of clinical trials of targeted agents for these targets and the difficulty of obtaining such drugs for off-indication has been a major impediment to the use of matched therapy in these smaller molecularly defined subsets Supplementary Fig. S5 A–E.

Although patients with level 3 and 4 alterations were only rarely treated on a genotype-matched therapy clinical trial (7%; 26/373); 19% (70/373) did enroll on a therapeutic clinical study (24.6% and 16.6% for level 3 and 4 patients, respectively). In most instances (62.9%; 44/70), these patients enrolled on trials of immunotherapy (Supplementary Fig. S6), with an increasingly larger fraction of patients receiving immunotherapy as a standard treatment towards the end of the study period. The data suggest that the low rate of matched therapy use in patients with level 3 and 4 alterations was not due to a reluctance of such patients to enroll on therapeutic clinical studies but rather attributable to the lack of compelling matched therapy options available for these patients. Notably, there was an inverse trend towards a higher mutational load in patients with level 3 and 4 alterations as compared to those with level 1 and 2 mutations (Supplementary Fig. S7). This was likely attributable to the lower rate of patients who were never smokers in patients with a level 3 (29.2%; 19/65) and level 4 alteration (12.7%; 39/308) as compared to level 1 (62%; 166/269) and 2A (45.1%; 32/71) alterations. As higher mutational load in lung cancer has been associated with a greater likelihood of response to immunotherapy (28), and as matched therapy options were limited in patients with level 3 and 4 alterations, the choice to pursue immunotherapy over a matched therapy in such patients could be considered a rational course of action guided by the clinical sequencing results.

Clinical benefit with EGFR inhibitors in patients with uncommon EGFR mutant alleles

Not all mutations in a cancer gene have similar oncogenic potential or similarly predict for response to a targeted inhibitor. For most cancer genes, clinical response data is often only available for the most commonly mutated alleles and clinical guidelines for the treatment of patients with rare alleles; even in common cancer genes are often based on preclinical drug sensitivities or small case series. Oncogenic *EGFR* alterations known to be predictive of *EGFR* inhibitor sensitivity were identified in 214 (24.9%) patients (Figure 3, Supplementary Table S3). Consistent with prior studies, exon 21 L858R (70 patients) and exon 19 deletions/insertion (113) were the most common variants (85.5% of cases). Less common variants previously shown to activate *EGFR* kinase activity included L861Q (7), E709_ T710delinsD (6 patients), G719A (4) and exon 18–25 kinase domain duplication (EGFR-KDD, 2). Seventeen patients had *EGFR* exon 20 insertions previously shown to confer resistance to *EGFR* inhibitors such as erlotinib and one patient had an exon 20 H773R mutation (29, 30). Notably, one patient with an *EGFR* exon 20 insertion (level 4) received erlotinib without response consistent with prior evidence that such mutations are resistant to this agent (29). Excluding *EGFR* mutations known to confer resistance to erlotinib, nine patients had two or more activating mutations in *EGFR* and all of these patients had either E709A/K or G719X

mutations; alleles previously associated with somatic *EGFR* doublets (31). Overall, 87.3% of patient with sensitizing *EGFR* mutants benefited from the use of an *EGFR* inhibitor but we did observe differences in response as a function of the specific *EGFR* mutant allele present. The rate of clinical benefit was statistically significantly lower in patients with L861Q mutations (43%; $p=0.039$ versus L858R; $p=0.01$ versus exon 19 deletions) or exon 18 deletions (40%; $p=0.02$ versus L858R; $p=0.005$ versus exon 19 deletions). These results are consistent with the lower rate of clinical response of tumors harboring L861Q mutations to first generation TKI therapy reported in prior studies (32, 33) and support the clinical evaluation of afatinib and osimertinib in such patients, as these agents have demonstrated greater potency against this allele (34). Notably, the patient with a L861Q mutation who had the longest duration of clinical benefit received dacomitinib, a second generation TKI. Consistent with published data (35), both patients with *EGFR*-kinase domain duplication (*EGFR*-KDD) treated with erlotinib and afatinib, respectively, derived clinical benefit.

Unknown Mitogenic Driver (UMD) Set

In total, 103 patients had tumors for which no level 1–4 alteration was identified and these tumors were therefore designated as unknown mitogenic driver (UMD). To ensure that UMD samples were not enriched for low-purity samples, we generated estimates of their purity and looked for any sequencing reads supporting hotspot mutations. This analysis confirmed that this set of UMD samples was not significantly enriched with low tumor content samples (estimated purity by FACETS analysis of 12–92%, mean 35% versus 38%; range 4–95% for samples with Level 1–4 alterations), Figure 4A. We then sought to compare the frequency of alterations in these samples to those with a known mitogenic driver with the goal of nominating additional oncogenic drivers as candidates for future drug development. In comparison to tumors harboring a level 1 to 4 alteration, alterations in *TP53*, *STK11*, *KEAP1*, *KMT2D*, and *PDGFRA* were all significantly more common ($p<0.05$) in the UMD cohort (Figure 4B). These findings corroborate the enrichment of *TP53* and *KEAP1* in mitogenic driver negative samples in the TCGA cohort (16). In contrast to the TCGA set which identified enrichment of *RIT1* mutations in the oncogene-negative set, no *RIT1* mutations were identified within the UMD set. Further analysis identified a number of genes that were statistically enriched in the UMD cohort versus samples with level 1–4 mitogenic drivers when subdividing these patients according to smoking history (Figure 4B). For example, chromatin modifying genes such as *KMT2C*, *SETD2* and *CREBBP* and genes involved in homologous recombination (*MRE11A*, *BRCA2*) were more commonly identified ($p<0.05$) in patients with a never/former light smoking history within the UMD cohort as compared to level 1–4 patient samples. Patients with a former/current heavy smoking history in the UMD cohort had higher rates of *STK11* and *TP53* alterations as compared to those who had a level 1–4 driver. Co-occurring mutations in *TP53* and *STK11* are known to be synergistic in tumorigenesis (36), but represented only 27% (18/68) of patients with *TP53* or *STK11* mutations in the former/current heavy smoking UMD subset. While *ERBB4* alterations were identified in the UMD subset, none were hotspot alterations or mutations previously demonstrated to confer sensitivity to HER kinase inhibitors. Furthermore, all *BRCA2* mutations were somatic missense mutations of uncertain significance and thus further laboratory and clinical studies will be needed to clarify the significance of these findings.

A potentially targeted approach was utilized in two of the UMD patients; off label azacitidine, a DNA methyltransferase inhibitor, in a patient with a *KDM5C* frameshift mutation (6 months, stable disease), everolimus for an *mTOR* L2383F mutation (1 month; no benefit). In the latter case, the *mTOR* mutation was a novel missense variant of unknown significance. One patient in the UMD cohort without an *EGFR* mutation or amplification had clinical benefit from erlotinib treatment. A tumor biopsy after disease progression revealed an uncharacterized *ERRF1* (*ERBB* receptor feedback inhibitor 1, a.k.a. *MIG6*) A143D missense mutation as the sole non-synonymous mutation. *ERRF1* loss of function causes hyperactivation of *EGFR* and persistent mitogen-activated protein kinase (MAPK) signaling, with tumors in mice responsive to gefitinib (37) and it has been shown to accelerate initiation and progression of *EGFR*-mutated lung adenocarcinoma in mice (38). *ERRF1* mutations were identified in six (0.7%) patients, four of whom had higher level alterations (*EGFR* exon 19 deletion (2), *KRAS* G12 (2)).

As real-time functional validation of all non-recurrent somatic missense mutations of unknown significance is not currently feasible, we used *in silico* modeling to identify other potentially functional missense variants in the UMD cohort. *KEAP1* (kelch-like ECH-associated-protein-1), a negative regulator of *Nrf2* (nuclear factor erythroid 2-related factor 2), was altered in 34% (35/103) of the UMD cases (10 truncations and 25 missense variants) with *Nrf2* mutations noted in 2.9% (3/103). Using 3D structure-based computational analysis of the *KEAP1* protein structure, we found that many of these *KEAP1* mutations cluster in the Kelch domain, which interacts with *Nrf2* (Figure 4C). *KEAP1* mutations can induce increased *Nrf2* accumulation resulting in chemoresistance through induced expression of cellular antioxidants and xenobiotic detoxification enzymes (39). Targeting *Nrf2* with inhibitors such as luteolin and brusatol may enhance chemotherapy sensitization (40, 41), but trials testing this hypothesis have yet to be conducted in NSCLC.

Discussion

Rapid advances in sequencing methodology have made it feasible to prospectively profile increasing numbers of cancer-associated genes using the small quantities of FFPE-derived DNA that are typically available as part of the routine clinical care of patients with advanced cancer. However, the fraction of patients who derive clinical benefit from molecular characterization remains undefined. Here, we report the prospective clinical experience with MSK-IMPACT testing in the first 860 patients with lung adenocarcinoma with a focus on defining the fraction of patients who received a matched therapy and derived clinical benefit from such treatment.

Overall, 37.1% of lung adenocarcinoma patients who had undergone MSK-IMPACT testing received a matched therapy with the likelihood of matched treatment correlating strongly with the level of evidence that the mutation identified predicts for drug response. For patients with *EGFR* mutations and *ALK* or *ROS1*-fusions (level 1 alterations), 93% received the corresponding matched therapy with 85.8% deriving clinical benefit. Alterations in these genes are now recognized by the FDA as predictive biomarkers of drug response. Of the patients for whom a treatment other than the corresponding matched therapy was chosen, 85% (17/20) remain clinically stable on chemotherapy or immunotherapy. With longer

followup, we anticipate that most if not all of these patients will ultimately receive matched targeted therapy.

The described patients with *ROS1* fusions identified too late in their clinical course to receive matched therapy illustrate the potential importance of early broad molecular testing that includes genes beyond those recognized as biomarkers by the FDA, in particular those mutations categorized as level 2A, defined here as standard of care biomarkers for FDA-approved drugs in lung cancer patients based on currently accepted practice guidelines such as those issued by the National Comprehensive Cancer Network. We found that the proportion of patients with level 2A alterations who received a matched therapy was significantly lower ($p < 0.0001$) than patients with level 1 alterations (93% vs 52%) but the rate of clinical benefit (76%) was compelling in those who were treated with appropriate matched therapy. This lower use of matched therapy in patients with level 2A alterations was likely attributable to limited access to the corresponding matched therapies, in particular the lack of access to several of the agents outside the context of a clinical trial during at least a portion of the study period. As an example, the clinical efficacy of *MET* inhibitors in patients with exon 14 *MET* alterations was first reported in August 2015 (42), and the availability of *MET* inhibitors for patients with activating alterations in *MET* was limited outside of clinical trials prior to this date. A substantial fraction of patients with lung adenocarcinoma are still not screened for alterations in *MET*, *RET*, and *BRAF*. Notably, 39% (28/71) of patients with level 2A alterations were screened after at least one previous systemic therapy and identification of such alterations by MSK-IMPACT testing was more likely to occur in the setting of symptomatic advanced disease, which may have led at least some patients to have been deemed poor candidates for clinical studies of the corresponding matched therapy. As 11.3% (8/71) of the Level 2A patients had rapid clinical deterioration and never received treatment with the appropriate matched therapy, broader screening for such alterations in lung cancer patients at the time of diagnosis may result in improved outcomes.

In contrast to the high uptake of matched therapies in the standard-of-care setting (level 1 and 2A alterations), only 7.6% (31/407) of patients with level 2B-4 alterations received a matched therapy. These results are consistent with a prior study that assessed the use of matched therapy in the investigational setting (43). The low frequency of matched therapy treatment in patients with level 2B-4 alterations was likely not due to a lack of interest by such patients or their physicians in participating in clinical trials as 17.9% (73/407) of patients with level 2B-4 alterations were enrolled on a therapeutic study; 70% (51/73) employed immunotherapy/other investigational compounds with 30% informed by the patients mutational profile. This low frequency of matched therapy use in patients with level 2B-4 alterations is likely attributable to the lack of compelling matched therapy studies for such alterations and the inability of patients to access matched therapies outside the context of a clinical trial and/or the reluctance of patients/physicians to pursue compassionate use of such treatments. The results were particularly striking for patients with *ERBB2* amplifications and inactivating mutations of *BRCA1/2* and *TSC1/2*, which are standard-of-care predictive biomarkers of response to *HER2*, *PARP*, and *mTOR* inhibitors respectively in other cancer types. Based upon this early experience, we have sought to open basket

clinical trials that would allow for treatment of patients with such alterations in the context of a clinical study (NCT02201212, NCT02675829).

With the use of a broader sequencing panel, we often identified multiple potentially actionable targets co-existing within individual tumors. In total, 239 (27.8%) patients had two or more actionable mutations but not a single such patient received a combination of matched therapies (Supplementary Table S2). While responses to single agent targeted therapy are often dramatic in patients with lung adenocarcinoma, intrinsic and acquired resistance continue to be major hurdles in achieving the promise of a precision medicine approach. Given the various signaling pathways involved in oncogenesis and their interdependence through cross-talk signaling and feedback mechanisms, the use of combinations of targeted agents could prevent or delay the emergence of drug resistant clones. However, the complexity imposed by drug-drug interactions, the potential for increased toxicity and the need to identify an optimal dose and timing schedule (sequential or co-administration) require that each potential combination be explored in the context of a clinical trial prior to broach use. This complexity along with the significant logistical and financial challenges in targeting more than one potentially actionable alteration suggest to us that novel clinical trial designs will be needed to achieve progress with combination strategies in patients with lung cancer (44, 45).

Finally, in 13.1% of cases, we did not identify any Level 1–4 alterations. While broader molecular testing such as whole exome, genome, or transcriptome sequencing may have identified potentially actionable fusions or other alterations in some of these tumors, the majority did harbor mutations in cancer genes that have in laboratory models been shown to contribute to lung cancer pathogenesis such as alterations in the tumor suppressor genes *TP53* and *STK11*. While certain targeted agents have been proposed as rational treatments in the setting of several of the genes mutated in the UMD cohort, our review of the literature suggests that such mutations were not compelling biomarkers of drug response in lung adenocarcinoma in these cases. As an example, mTOR is a kinase downstream of *STK11/LKB1* and hence mTOR inhibitors have been proposed as a potential therapeutic approach in patients with *STK11* mutations (46). However, no patients with *STK11/LKB1* alterations received an mTOR inhibitor in our cohort. A phase 2 trial of the mTOR inhibitor everolimus in patients with solid malignancies that harbor *TSC1/2*, *NF1/2* or *STK11* mutations was recently initiated (NCT02352844), but we were unable to find any trials testing the utility of *STK11* as a predictive biomarker of response to mTOR inhibition in lung cancer.

In summary, we find that 37.1% of lung cancer patients who underwent MSK-IMPACT testing received a matched therapy based on their mutational profile. Of these patients, 78.1% (249 patients) derived clinical benefit. Excluding standard of care therapy (*EGFR* mutations, *ALK* and *ROS1* fusions), 14.4% (69/478) of patients with a level 2–4 alteration as their highest actionable target received matched therapy with 52% (36/69) exhibiting clinical benefit. The use of matched therapy was strongly influenced by the level of clinical evidence that the mutation identified predicts for drug response. Our data suggest that the use of matched therapies is limited by a lack of access to FDA-approved drugs in patients with level 2 alterations and by the lack of compelling clinical trials of investigational agents in patients with level 3 and 4 mutations.

Methods

Patient Selection

All patients had recurrent or metastatic lung adenocarcinoma and were referred for genomic testing from January 2014 to March 2016. Clinical data were collected within the context of a prospective clinical trial (ClinicalTrials.gov, NCT01775072), under an Institutional Review Board approved protocol allowing genomic testing on patients' tumors. Informed consent was obtained from all participating patients. This study was conducted in accordance with the Declaration of Helsinki.

Genomic sequencing

Tumor and germline DNA were processed to generate bar-coded libraries and subjected to exon capture using custom-designed probes (14). The average sequence coverage across all tumors was 615X, providing high sensitivity to detect mutations at low allele frequencies in heterogeneous or low purity specimens. Matched normal DNA, available for 97% of samples, was analyzed simultaneously to identify and filter out germline single nucleotide polymorphisms (SNPs). Genomic analysis was performed using the MSK-IMPACT assay, a clinical test approved by the New York State Department of Health designed to detect mutations, copy number alterations and select fusions involving 341 (version 1) or 410 (version 2) cancer-associated genes cancer-associated genes, Supplementary Table S4 (14). Genomic analysis was performed using assay version-1 (341 genes) for 296 samples and version-2 (410 genes) for 619 samples.

Analysis

We stratified potentially actionable genetic events into one of four levels based on published clinical or laboratory evidence that the mutation in question confers increased sensitivity to standard or investigational therapies. An interactive compendium of the mutations deemed actionable is available at the OncoKB website (15). Level 1 alterations included mutations and fusions that are FDA-approved biomarkers in patients with lung cancer (sensitizing *EGFR* mutations, *ALK* or *ROS1*-fusions), whereas level 2A events were alterations that were deemed to be standard of care biomarkers for FDA-approved drugs in lung cancer patients based on currently accepted practice guidelines such as those issued by the National Comprehensive Cancer Network (NCCN). Level 2B alterations included those that are FDA-approved biomarkers in another cancer indication (e.g. *ERBB2* amplification) but not in patients with lung cancer. Level 3 included alterations for which compelling clinical evidence links the biomarker to drug response in patients but use of the biomarker is not currently a standard-of-care in any cancer type (e.g. *ERBB2* mutation). Finally, level 4 alterations were those in which compelling preclinical data associates the biomarker with drug response (e.g. *NFI* loss). Patients with two or more level 1–4 oncogenic drivers were grouped with the highest-level actionable driver.

Clinical records for all patients were reviewed to determine whether the patient received a matched targeted therapy or immunotherapy and whether the patient was enrolled on a therapeutic clinical trial. Clinical trials were designated as either matched therapy based on the assigned oncogenic mutational profile, immunotherapy, or “other” if it did not meet the

aforementioned criteria. Patients were deemed to have derived clinical benefit if there was a reduction in tumor size on imaging and documented symptom improvement or stable disease on two consecutive imaging scans 30 days apart with symptom improvement. All clinical and genomic data are available in electronic form through the cBioPortal for Cancer Genomics (47, 48).

Samples without a potentially actionable mitogenic driver mutation (UMD: unknown mitogenic driver) were subsequently analyzed for the presence of sequencing reads with non-reference bases at mutational hotspots (27). To guard against false negative results due to insufficient tumor content, the purity of the UMD samples was also estimated by allelic copy number analysis using FACETS (49). Cochran-Armitage tests were used to assess the trend in the probability of receiving matched therapy and immunotherapy across the level of evidence categories, followed by Fisher's exact tests for pairwise comparisons. The Cuzick trend test was used to assess the trend in the number of mutations as a function of the level of evidence category. Fisher's exact tests were used to compare the rates of mutation for each gene between MSK-IMPACT and TCGA patients. All statistical tests were two-sided and a p value <0.05 considered significant.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Sources of funding: Marie-Josée and Henry R. Kravis Center for Molecular Oncology (CMO)

References

1. Shaw AT, Kim DW, Nakagawa K, Seto T, Crino L, Ahn MJ, et al. Crizotinib versus chemotherapy in advanced ALK-positive lung cancer. *The New England journal of medicine*. 2013; 368(25):2385–94. [PubMed: 23724913]
2. Solomon BJ, Mok T, Kim DW, Wu YL, Nakagawa K, Mekhail T, et al. First-line crizotinib versus chemotherapy in ALK-positive lung cancer. *The New England journal of medicine*. 2014; 371(23):2167–77. [PubMed: 25470694]
3. Rosell R, Carcereny E, Gervais R, Vergnenegre A, Massuti B, Felip E, et al. Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EURTAC): a multicentre, open-label, randomised phase 3 trial. *The Lancet Oncology*. 2012; 13(3):239–46. [PubMed: 22285168]
4. Bergethon K, Shaw AT, Ou SH, Katayama R, Lovly CM, McDonald NT, et al. ROS1 rearrangements define a unique molecular class of lung cancers. *Journal of clinical oncology: official journal of the American Society of Clinical Oncology*. 2012; 30(8):863–70. [PubMed: 22215748]
5. Stephens P, Hunter C, Bignell G, Edkins S, Davies H, Teague J, et al. Lung cancer: intragenic ERBB2 kinase mutations in tumours. *Nature*. 2004; 431(7008):525–6.
6. Mazieres J, Barlesi F, Filleron T, Besse B, Monnet I, Beau-Faller M, et al. Lung cancer patients with HER2 mutations treated with chemotherapy and HER2-targeted drugs: results from the European EUHER2 cohort. *Annals of oncology: official journal of the European Society for Medical Oncology/ESMO*. 2016; 27(2):281–6.
7. Tissot C, Couraud S, Tanguy R, Bringuier PP, Girard N, Souquet PJ. Clinical characteristics and outcome of patients with lung cancer harboring BRAF mutations. *Lung cancer (Amsterdam, Netherlands)*. 2016; 91:23–8.

8. Kris MG, Camidge DR, Giaccone G, Hida T, Li BT, O'Connell J, et al. Targeting HER2 aberrations as actionable drivers in lung cancers: phase II trial of the pan-HER tyrosine kinase inhibitor dacomitinib in patients with HER2-mutant or amplified tumors. *Annals of oncology: official journal of the European Society for Medical Oncology/ESMO*. 2015; 26(7):1421–7.
9. Drilon A, Wang L, Hasanovic A, Suehara Y, Lipson D, Stephens P, et al. Response to Cabozantinib in patients with RET fusion-positive lung adenocarcinomas. *Cancer discovery*. 2013; 3(6):630–5. [PubMed: 23533264]
10. Planchard D, Kim TM, Mazieres J, Quoix E, Riely G, Barlesi F, et al. Dabrafenib in patients with BRAF-positive advanced non-small-cell lung cancer: a single-arm, multicentre, open-label, phase 2 trial. *The Lancet Oncology*. 2016
11. Drilon A, Rekhtman N, Arcila M, Wang L, Ni A, Albano M, et al. Cabozantinib in patients with advanced RET-rearranged non-small-cell lung cancer: an open-label, single-centre, phase 2, single-arm trial. *The Lancet Oncology*. 2016
12. Janne PA, Shaw AT, Pereira JR, Jeannin G, Vansteenkiste J, Barrios C, et al. Selumetinib plus docetaxel for KRAS-mutant advanced non-small-cell lung cancer: a randomised, multicentre, placebo-controlled, phase 2 study. *The Lancet Oncology*. 2013; 14(1):38–47. [PubMed: 23200175]
13. Vansteenkiste JF, Canon JL, Braud FD, Grossi F, De Pas T, Gray JE, et al. Safety and Efficacy of Buparlisib (BKM120) in Patients with PI3K Pathway-Activated Non-Small Cell Lung Cancer: Results from the Phase II BASALT-1 Study. *Journal of thoracic oncology: official publication of the International Association for the Study of Lung Cancer*. 2015; 10(9):1319–27.
14. Cheng DT, Mitchell TN, Zehir A, Shah RH, Benayed R, Syed A, et al. Memorial Sloan Kettering-Integrated Mutation Profiling of Actionable Cancer Targets (MSK-IMPACT): A Hybridization Capture-Based Next-Generation Sequencing Clinical Assay for Solid Tumor Molecular Oncology. *The Journal of molecular diagnostics: JMD*. 2015; 17(3):251–64. [PubMed: 25801821]
15. OncoKB.org. Precision Oncology Knowledge Base [Internet]. 2016. Available from <http://www.OncoKB.org>
16. Comprehensive molecular profiling of lung adenocarcinoma. *Nature*. 2014; 511(7511):543–50. [PubMed: 25079552]
17. Kris MG, Johnson BE, Berry LD, Kwiatkowski DJ, Iafrate AJ, Wistuba II, et al. Using multiplexed assays of oncogenic drivers in lung cancers to select targeted drugs. *Jama*. 2014; 311(19):1998–2006. [PubMed: 24846037]
18. Barlesi F, Mazieres J, Merlio JP, Debievre D, Mosser J, Lena H, et al. Routine molecular profiling of patients with advanced non-small-cell lung cancer: results of a 1-year nationwide programme of the French Cooperative Thoracic Intergroup (IFCT). *Lancet*. 2016; 387(10026):1415–26. [PubMed: 26777916]
19. Ohashi K, Sequist LV, Arcila ME, Lovly CM, Chen X, Rudin CM, et al. Characteristics of lung cancers harboring NRAS mutations. *Clinical cancer research: an official journal of the American Association for Cancer Research*. 2013; 19(9):2584–91. [PubMed: 23515407]
20. Arcila ME, Drilon A, Sylvester BE, Lovly CM, Borsu L, Reva B, et al. MAP2K1 (MEK1) Mutations Define a Distinct Subset of Lung Adenocarcinoma Associated with Smoking. *Clinical cancer research: an official journal of the American Association for Cancer Research*. 2015; 21(8):1935–43. [PubMed: 25351745]
21. Imielinski M, Greulich H, Kaplan B, Araujo L, Amann J, Horn L, et al. Oncogenic and sorafenib-sensitive ARAF mutations in lung adenocarcinoma. *The Journal of clinical investigation*. 2014; 124(4):1582–6. [PubMed: 24569458]
22. Chaft JE, Arcila ME, Paik PK, Lau C, Riely GJ, Pietanza MC, et al. Coexistence of PIK3CA and other oncogene mutations in lung adenocarcinoma-rationale for comprehensive mutation profiling. *Molecular cancer therapeutics*. 2012; 11(2):485–91. [PubMed: 22135231]
23. Eng J, Woo KM, Sima CS, Plodkowski A, Hellmann MD, Chaft JE, et al. Impact of Concurrent PIK3CA Mutations on Response to EGFR Tyrosine Kinase Inhibition in EGFR-Mutant Lung Cancers and on Prognosis in Oncogene-Driven Lung Adenocarcinomas. *Journal of thoracic oncology: official publication of the International Association for the Study of Lung Cancer*. 2015; 10(12):1713–9.

24. Fernandez-Cuesta L, Plenker D, Osada H, Sun R, Menon R, Leenders F, et al. CD74-NRG1 fusions in lung adenocarcinoma. *Cancer discovery*. 2014; 4(4):415–22. [PubMed: 24469108]
25. Capelletti M, Dodge ME, Ercan D, Hammerman PS, Park SI, Kim J, et al. Identification of recurrent FGFR3-TACC3 fusion oncogenes from lung adenocarcinoma. *Clinical cancer research: an official journal of the American Association for Cancer Research*. 2014; 20(24):6551–8. [PubMed: 25294908]
26. Schirosi L, Nannini N, Nicoli D, Cavazza A, Valli R, Buti S, et al. Activating c-KIT mutations in a subset of thymic carcinoma and response to different c-KIT inhibitors. *Annals of oncology: official journal of the European Society for Medical Oncology/ESMO*. 2012; 23(9):2409–14.
27. Chang MT, Asthana S, Gao SP, Lee BH, Chapman JS, Kandoth C. Identifying recurrent mutations in cancer reveals widespread lineage diversity and mutational specificity. 2016; 34(2):155–63.
28. Rizvi NA, Hellmann MD, Snyder A, Kvistborg P, Makarov V, Havel JJ, et al. Cancer immunology. Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. *Science (New York, NY)*. 2015; 348(6230):124–8.
29. Yasuda H, Kobayashi S, Costa DB. EGFR exon 20 insertion mutations in non-small-cell lung cancer: preclinical data and clinical implications. *The Lancet Oncology*. 2012; 13(1):e23–31. [PubMed: 21764376]
30. Arcila ME, Nafa K, Chaffaer JE, Rekhtman N, Lau C, Reva BA, et al. EGFR exon 20 insertion mutations in lung adenocarcinomas: prevalence, molecular heterogeneity, and clinicopathologic characteristics. *Molecular cancer therapeutics*. 2013; 12(2):220–9. [PubMed: 23371856]
31. Chen Z, Feng J, Saldivar JS, Gu D, Bockholt A, Sommer SS. EGFR somatic doublets in lung cancer are frequent and generally arise from a pair of driver mutations uncommonly seen as singlet mutations: one-third of doublets occur at five pairs of amino acids. *Oncogene*. 2008; 27(31):4336–43. [PubMed: 18372921]
32. Locatelli-Sanchez M, Couraud S, Arpin D, Riou R, Bringuier PP, Souquet PJ. Routine EGFR molecular analysis in non-small-cell lung cancer patients is feasible: exons 18–21 sequencing results of 753 patients and subsequent clinical outcomes. *Lung*. 2013; 191(5):491–9. [PubMed: 23749122]
33. Lee VH, Tin VP, Choy TS, Lam KO, Choi CW, Chung LP, et al. Association of exon 19 and 21 EGFR mutation patterns with treatment outcome after first-line tyrosine kinase inhibitor in metastatic non-small-cell lung cancer. *Journal of thoracic oncology: official publication of the International Association for the Study of Lung Cancer*. 2013; 8(9):1148–55.
34. Banno E, Togashi Y, Nakamura Y, Chiba M, Kobayashi Y, Hayashi H, et al. Sensitivities to various EGFR-TKIs of uncommon EGFR mutations L861Q and S768I: what is the optimal EGFR-TKI? *Cancer science*. 2016
35. Gallant JN, Sheehan JH, Shaver TM, Bailey M, Lipson D, Chandramohan R, et al. EGFR Kinase Domain Duplication (EGFR-KDD) Is a Novel Oncogenic Driver in Lung Cancer That Is Clinically Responsive to Afatinib. *Cancer discovery*. 2015; 5(11):1155–63. [PubMed: 26286086]
36. Wei C, Amos CI, Stephens LC, Campos I, Deng JM, Behringer RR, et al. Mutation of Lkb1 and p53 genes exert a cooperative effect on tumorigenesis. *Cancer research*. 2005; 65(24):11297–303. [PubMed: 16357136]
37. Ferby I, Reschke M, Kudlacek O, Knyazev P, Pante G, Amann K, et al. Mig6 is a negative regulator of EGF receptor-mediated skin morphogenesis and tumor formation. *Nature medicine*. 2006; 12(5):568–73.
38. Maity TK, Venugopalan A, Linnoila I, Cultraro CM, Giannakou A, Nemati R, et al. Loss of MIG6 Accelerates Initiation and Progression of Mutant Epidermal Growth Factor Receptor-Driven Lung Adenocarcinoma. *Cancer discovery*. 2015; 5(5):534–49. [PubMed: 25735773]
39. Singh A, Misra V, Thimmulappa RK, Lee H, Ames S, Hoque MO, et al. Dysfunctional KEAP1-NRF2 interaction in non-small-cell lung cancer. *PLoS medicine*. 2006; 3(10):e420. [PubMed: 17020408]
40. Chian S, Thapa R, Chi Z, Wang XJ, Tang X. Luteolin inhibits the Nrf2 signaling pathway and tumor growth in vivo. *Biochemical and biophysical research communications*. 2014; 447(4):602–8. [PubMed: 24747074]

41. Ren D, Villeneuve NF, Jiang T, Wu T, Lau A, Toppin HA, et al. Brutal enhances the efficacy of chemotherapy by inhibiting the Nrf2-mediated defense mechanism. *Proceedings of the National Academy of Sciences of the United States of America*. 2011; 108(4):1433–8. [PubMed: 21205897]
42. Paik PK, Drilon A, Fan PD, Yu H, Rekhtman N, Ginsberg MS, et al. Response to MET inhibitors in patients with stage IV lung adenocarcinomas harboring MET mutations causing exon 14 skipping. *Cancer discovery*. 2015; 5(8):842–9. [PubMed: 25971939]
43. Meric-Bernstam F, Brusco L, Shaw K, Horombe C, Kopetz S, Davies MA, et al. Feasibility of Large-Scale Genomic Testing to Facilitate Enrollment Onto Genomically Matched Clinical Trials. *Journal of clinical oncology: official journal of the American Society of Clinical Oncology*. 2015; 33(25):2753–62. [PubMed: 26014291]
44. Yap TA, Omlin A, de Bono JS. Development of therapeutic combinations targeting major cancer signaling pathways. *Journal of clinical oncology: official journal of the American Society of Clinical Oncology*. 2013; 31(12):1592–605. [PubMed: 23509311]
45. Kummar S, Chen HX, Wright J, Holbeck S, Millin MD, Tomaszewski J, et al. Utilizing targeted cancer therapeutic agents in combination: novel approaches and urgent requirements. *Nature reviews Drug discovery*. 2010; 9(11):843–56. [PubMed: 21031001]
46. Klumpen HJ, Queiroz KC, Spek CA, van Noesel CJ, Brink HC, de Leng WW, et al. mTOR inhibitor treatment of pancreatic cancer in a patient With Peutz-Jeghers syndrome. *Journal of clinical oncology: official journal of the American Society of Clinical Oncology*. 2011; 29(6):e150–3. [PubMed: 21189378]
47. Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer discovery*. 2012; 2(5):401–4. [PubMed: 22588877]
48. Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, et al. Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Science signaling*. 2013; 6(269):p11. [PubMed: 23550210]
49. Shen R, Seshan VE. FACETS: allele-specific copy number and clonal heterogeneity analysis tool for high-throughput DNA sequencing. *Nucleic acids research*. 2016

Significance

An increasing number of therapies that target molecular alterations required for tumor maintenance and progression have demonstrated clinical activity in patients with lung adenocarcinoma. The data reported here suggests that broader, early testing for molecular alterations that have not yet been recognized as standard of care predictive biomarkers of drug response could accelerate the development of targeted agents for rare mutational events and could result in improved clinical outcomes.




Figure 1. Potentially actionable oncogenic drivers identified by MSK-IMPACT testing
A. Spectrum of oncogenic drivers assigned to 860 patients with lung adenocarcinoma identified by MSK-IMPACT. **B.** Comparison of selected gene alteration frequencies in the MSK-IMPACT and TCGA cohorts. **C.** Oncoprint of select gene alterations identified by MSK-IMPACT in patients with a level 1–4 alterations or those with no actionable mutation (Unknown Mitogenic Driver).





Figure 2.

Figure 2A. Use of matched therapy (MT) correlates strongly with the level of evidence that the mutation identified predicts for drug response.

A. Use of Matched Therapy (MT), immunotherapy (IT) and clinical trial participation in patients with level 1–4 alterations or in the unknown mitogenic driver (UMD) cohorts. **B:** Use of matched therapy and immunotherapy and clinical benefit from matched therapy in patients whose tumors harbored alterations in select level 1–4 genes.



EGFR Mutations	No. pts	Receive TKI	Clinical Benefit TKI	Chemotherapy	Immunotherapy	Clinical Trial	Median time on TKI (months)
Exon21							
L858R	70	66 (96%)	56* (89%)	49 (67%)	5 (7%)	23 (33%)	9 (1-47)
L861Q	7	7 (100%)	3 (43%)	5 (71%)	3 (43%)	2 (29%)	4.5 (1-33)
Exon 19							
Exon 19 deletion	112	109 (97%)	100* (94%)	55 (49%)	9 (8%)	46 (41%)	10 (2-85)
Exon 19 insertion	1	1 (100%)	1 (100%)	1 (100%)	1 (100%)	0 (0%)	3 (ongoing)
L747P	3	3 (100%)	2 (67%)	3 (100%)	1 (33%)	0 (0%)	7 (6-15)
Exon 18							
E709_T710delinsD	6	5 (67%)	2(40%)	4 (50%)	2 (33%)	2 (33%)	2 (1.5-22)
G719A	4	4 (100%)	2 (50%)	4 (100%)	2 (50%)	1 (25%)	5 (2-10)
E709K+G719S	2	1 (50%)	1 (100%)	2 (100%)	0 (0%)	1 (50%)	12
E709K+G719A	1	1 (100%)	0(0%)	1 (100%)	0 (0%)	0 (0%)	2 (erlotinib)**
E709A+G719S	1	1 (100%)	1 (100%)	1 (100%)	0 (0%)	0 (0%)	7 (ongoing)
Other							
G719C+S768I	2	2 (100%)	2 (100%)	2 (100%)	1 (50%)	1 (50%)	11.5 (3-20)
G719S+S768I	1	1 (100%)	1 (100%)	0 (0%)	0 (0%)	0 (0%)	23 (ongoing)
G719A+L861Q	1	1 (100%)	1 (100%)	1 (100%)	0 (0%)	0 (0%)	5 (ongoing)
G719A+S768I	1	1 (100%)	1 (100%)	1 (100%)	0 (0%)	1 (100%)	12
EGFR-KDD	2	2 (100%)	2 (100%)	3 (100%)	1 (50%)	1 (50%)	8 (both pts ongoing)
Exon 20 Insertion	17	1 (6%)	0 (0%)	16 (94%)	6 (38%)	4 (25%)	1.5
Exon 20 H773R	1	1 (100%)	1 (100%)	1 (100%)	0 (0%)	1 (100%)	6

L858R *3 patients await imaging
 Exon 19 deletion* 3 patients await imaging
 ** E709K+G719A: currently receiving Afatinib (2 months ongoing)

Figure 3. Use of matched therapy and clinical benefit in patients with known activating mutations in EGFR

(Top) Frequency of known activating and resistance mutations in EGFR identified by MSK-IMPACT. (Bottom) Use of matched therapy, chemotherapy, immunotherapy, clinical trial enrollment and clinical benefit from matched therapy as a function of the specific EGFR mutation identified in the patient’s tumor.




Figure 4. Potential driver alterations in the unknown mitogenic driver (UMD) cohort
A. Estimated purity analysis by FACETS in samples with Level 1–4 alteration to the UMD sample set. **B.** Oncoprint of the most common gene alterations in 103 patients with no actionable Level 1–4 driver mutations with a comparative frequency of select recurrently altered genes in the UMD cohort level 1–4 samples according to smoking history. P values were calculated using Fisher’s exact test **C.** Distribution of missense mutations in KEAP1 detected by MSK-IMPACT (upper panel); x-axis represents amino acid positions and y-axis represents number of samples mutated. The PFAM domains were also displayed as context. Protein structure analysis revealing that KEAP1 missense mutations identified in lung adenocarcinoma patients (the ones with side chains displayed) clustered the interaction interface with Nrf2 (nuclear factor erythroid 2-related factor 2). Nrf2 peptide is colored in green.

Table 1

Clinical characteristics of the 860 patients (915 samples) with metastatic or recurrent lung adenocarcinoma profiled by MSK-IMPACT

Characteristics	N=860 (%)	Samples (n=915)
Age		
18–50	122 (14.2)	134
51–75	615 (71.5)	652
>75	123 (14.3)	129
Sex		
Men	354 (41.2)	369
Women	506 (58.8)	546
Smoking status		
Never	277 (32.2)	302
Former light (≤ 15 pack year)	153 (17.8)	167
Former heavy (>15 pack year)	420 (48.8)	436
Current heavy	10 (1.2)	10
Sites of Tissue for MSK-IMPACT		
Lung		420
Lymph node		169
Pleura/Pleural fluid		110
Other		81 [*]
Liver		59
Brain		47
Bone		29

* Other: Soft tissue n=36 (chest wall mass; n=11, epidural tumor; n=7, gluteal mass; n=1, groin; n=1, iliac mass n=1, ischiorectal mass; n=1, L2 soft tissue mass; n=1, L5 soft tissue mass; n=1, pararenal mass; n=1, paraspinal mass; n=3, paratracheal mass; n=1, soft tissue mass scapula; n=2, pelvic mass; n=2, T1 soft tissue mass; n=1, T12 soft tissue mass; n=1, retroperitoneum; n=1), Adrenal; n=16, skin/subcutaneous nodule; n=12, Pericardium/pericardial fluid; n=6, ascites/omentum; n=4, pancreas; n=2, breast; n=1, colon; n=1, spleen; n=1, uterus; n=1, diaphragm; n=1.