



# HHS Public Access

Author manuscript

*Rev Environ Health*. Author manuscript; available in PMC 2018 March 01.

Published in final edited form as:

*Rev Environ Health*. 2017 March 01; 32(1-2): 65–72. doi:10.1515/reveh-2016-0041.

## Impact of nutrition on pollutant toxicity: an update with new insights into epigenetic regulation

Jessie B Hoffman<sup>1,2</sup>, Michael C Petriello<sup>1,3</sup>, and Bernhard Hennig<sup>1,3,\*</sup>

<sup>1</sup>Superfund Research Center, University of Kentucky, Lexington, KY 40536

<sup>2</sup>Department of Pharmacology and Nutritional Sciences, College of Medicine, University of Kentucky, Lexington, KY 40536

<sup>3</sup>Department of Animal and Food Sciences, College of Agriculture, Food and Environment, University of Kentucky, Lexington, KY 40536

### Abstract

Exposure to environmental pollutants is a global health problem and is associated with the development of many chronic diseases including cardiovascular disease, diabetes, and metabolic syndrome. There is a growing body of evidence that nutrition can both positively and negatively modulate the toxic effects of pollutant exposure. Diets high in pro-inflammatory fats, such as linoleic acid, can exacerbate pollutant toxicity while diets rich in bioactive and anti-inflammatory food components, including omega-3 fatty acids and polyphenols, can attenuate toxicant-associated inflammation. Previously, researchers have elucidated direct mechanisms of nutritional modulation including alteration of NF- $\kappa$ B signaling, but recently increased focus has been given to the ways in which nutrition and pollutants affect epigenetics. Nutrition has been demonstrated to modulate epigenetic markers that have been linked either to increased disease risks or to protection against diseases. Overnutrition (i.e. obesity) and undernutrition (i.e. famine) have been observed to alter prenatal epigenetic tags that may increase the risk of offspring developing disease later in life. Conversely, bioactive food components, including curcumin, have been shown to alter epigenetic markers that suppress activation of NF- $\kappa$ B, thus reducing inflammatory responses. Exposure to pollutants also alters epigenetic markers and may contribute to inflammation and disease. It has been demonstrated that pollutants, via epigenetic modulations, can increase activation of NF- $\kappa$ B and upregulate miRNAs associated with inflammation, cardiac injury, and oxidative damage. Importantly, recent evidence suggests that nutritional components, including EGCG, can protect against pollutant-induced inflammation through epigenetic regulation of pro-inflammatory target genes of NF- $\kappa$ B. Further research is needed to better understand how nutrition can modulate pollutant toxicity through epigenetic regulation. Further research is needed to better understand how nutrition can modulate pollutant toxicity through epigenetic regulation. Therefore, the objective of this review is to elucidate the current evidence linking epigenetic changes to pollutant-induced diseases and how this regulation may be modulated by nutrients allowing for the development of future personalized lifestyle interventions.

\*Correspondence should be directed to: Bernhard Hennig, 900 S. Limestone Street, Superfund Research Center, University of Kentucky, Lexington, KY 40536, USA. Tel.: +1 859-218-1343; fax: +1 859-257-1811; bhennig@uky.edu.

The authors declare that there are no competing financial interests.

## Keywords

Nutrition; environmental pollutants; epigenetics; anti-inflammatory nutrients; antioxidant response

---

## I. Introduction

The levels of environmental pollutants continue to rise worldwide, despite our increasing knowledge base of the negative health effects of pollutant exposure. There is increasing evidence that exposure to pollutants including persistent organic pollutants (POPs), heavy metals, and air pollution, can contribute to the progression of chronic diseases such as cardiovascular disease, diabetes, and cancer. Currently, the strongest evidence linking POPs with inflammatory diseases is related to pollutant-induced diabetes (1, 2). Multiple meta-analyses show positive associations of exposures to POPs such as polychlorinated biphenyls (PCBs) and pesticides with increased risk of developing type 2 diabetes (1, 2). The health effects of pollutant exposure not only poses a problem for the individual, but also contributes to the global burden of disease.

The mechanisms by which many of these pollutants contribute to disease involves an increase in cellular oxidative stress and subsequent inflammatory responses. For example, coplanar polychlorinated biphenyls (PCBs), a class of POPs, increase downstream inflammatory responses via binding to the aryl hydrocarbon receptor (AhR) which can upregulate the transcription of cytochrome p450 (Cyp1a1) (3, 4). Cyp1a1 is primarily involved in xenobiotic detoxification, however in the presence of certain PCBs, results in the production of reactive oxygen species (ROS) via an uncoupling mechanism (5). This production of ROS yields an increase in oxidative stress due to the imbalance in the cellular redox status (4). This pro-oxidative cellular status contributes to a state of chronic inflammation, which is a hallmark of many diseases including atherosclerosis, diabetes, and other metabolic disorders (6).

The most common routes of exposure to these environmental pollutants are through inhalation of airborne pollutants as well as ingestion of contaminated foods (7). POPs, including PCBs and organochloride pesticides, are lipophilic and therefore readily accumulate in the tissue of living animals (8). While production of many of these pollutants has been banned, their resistance to degradation and stability in the environment has allowed for their bioaccumulation throughout the ecosystem (8). Additionally, in humans, accumulation of these lipophilic pollutants over the course of many years can lead to activation of inflammatory pathways contributing to the development of numerous chronic inflammatory diseases (9, 10). Obesity is linked to higher pollutant body burden, but there is not a consensus if obesity makes a person more or less susceptible to pollutant-induced disease. Interestingly, there is some experimental evidence that points to the sequestration of pollutants in large adipose depots as actually being protective (11–13). Lipophilic pollutants may accumulate within adipocytes and lipid droplets which limits their availability to other target cell types. Importantly, it appears that obese individuals undergoing weight loss either through bariatric surgery or lifestyle intervention may be a highly susceptible population to pollutant-induced inflammation (11–13). Because overweight and obese individuals are not

only capable of accumulating greater levels of pollutants but are already at an increased risk for developing chronic inflammatory diseases, dietary interventions that provide a means of combatting pollutant toxicity and chronic disease risk factors are of great importance. (8, 14, 15).

## II. Nutrition as a modulator of pollutant toxicity

### Poor nutrition can exacerbate the toxicity of pollutants

It is well established that exposure to environmental pollutants can increase risk of developing chronic inflammatory diseases including cardiovascular disease, diabetes, and metabolic syndrome (6–9). Additionally, evidence implicates that individuals with a less nutritious or healthful diet (i.e. rich in processed foods and low in fruits and vegetables) are also at a greater risk of developing these same chronic diseases (16). Common to both pollutant exposure and poor nutrition is the activation of pro-inflammatory molecular proteins, including NF- $\kappa$ B, a key player in the development of inflammatory responses (17). Because of the overlap between pathways that mediate the negative effects of poor nutrition and environmental pollutants, there is an increasing body of experimental evidence demonstrating that certain nutritional components or overall nutritional status can exacerbate the negative health effects of environmental pollutant exposure.

Obesity is a global health problem and is a risk factor for the development of numerous inflammatory diseases (18). The rise in obesity is often attributed to overconsumption of processed, nutritionally poor foods (i.e. high energy density and low nutrient density) and foods high in pro-inflammatory fats (18). It has been consistently demonstrated that high fat diets, particularly those with greater proportions of fatty acids coming from saturated, trans, and linoleic fatty acids can increase inflammation and risk of cardiovascular disease as well as other metabolic complications (19, 20). In addition to overconsumption of nutritionally poor foods, emerging evidence now links exposure to environmental pollutants with increased adiposity and weight gain (21). Pollutants that elicit this physiological effect are termed “obesogens” and are believed to contribute to obesity and metabolic disorder through promotion of chronic imbalances in lipid storage mechanisms and adipocyte hyperplasia (21). Current evidence suggests that pollutant exposure can intensify the detrimental effects of a pro-inflammatory high-fat diet (22). For example, non-alcoholic fatty liver disease (NAFLD), commonly seen in developed countries as a hepatic indicator of obesity and metabolic syndrome, may be worsened by pollutant exposure. It has been demonstrated that the pollutant PCB153 exacerbates diet-induced obesity and NAFLD in mice through alterations in adipokines and disruptions in hepatic lipid metabolism (23). Additionally, high-fat diets appear to worsen arsenic-induced liver fibrosis and inflammation (24) while diets high in saturated fat appear to increase the toxicity and carcinogenic effects of persistent organic pollutants (POPs) (25). Furthermore, the flame retardant hexobromocyclododecane, was observed to enhance high-fat diet-induced obesity, hepatic steatosis, and impair glucose and lipid homeostasis in mice (26).

Cardiovascular diseases (CVDs) are the leading cause of death worldwide, and it is estimated that approximately 90% of CVDs can be prevented (27). The primary modifiable risk factors contributing to the development of CVDs include poor dietary habits, physical

Author Manuscript

Author Manuscript

Author Manuscript

inactivity, and smoking (27). Similarly, to the development of obesity, diets containing certain fatty acid profiles may promote the development of CVDs. For example, consumption of pro-inflammatory fatty acids such as linoleic and trans-fatty acids can contribute to endothelial cell dysfunction, a beginning stage of atherosclerosis (19, 20). Additionally, it has been demonstrated, primarily in experimental models, that exposure to environmental pollutants can cause endothelial cell dysfunction and chronic inflammation and may contribute to the development of CVDs (3, 28, 29). Furthermore, there is increasing experimental evidence that poor nutrition and pollutant exposure can interact and synergistically increase CVD risk (30). For example, it has recently been shown that trimethylamine N-oxide (TMAO), produced from metabolism of dietary carnitine or phosphatidyl choline, is a strong clinical biomarker of cardiovascular disease.(31–35) Interestingly, a recent preclinical paper from our group found that dioxin-like pollutant exposure increases the hepatic expression of the enzyme flavin-containing monooxygenase (FMO3) which is critical for production of TMAO. As such, this increase in hepatic expression of FMO3 resulted in increases in circulating TMAO (36). While this study was conducted in mice and the transcriptional regulation of FMO3 could differ in humans, this finding is important in that it provides new evidence on how certain nutritional components and pollutants may interact to contribute to a greater risk of cardiovascular disease.

It is not surprising that pollutant exposure exhibits detrimental combinatory effects with sub-optimal nutrition since pollutants and nutritional components can target the same molecular pathways, for example, the proinflammatory NF-kB pathway (4). However, this same idea can be utilized from the perspective of nutrition as a protective modulator of pollutant toxicity and may be more meaningful for those already exposed to high levels of pollutants and exhibiting signs of chronic inflammatory diseases.

### **Healthful nutrition can protect against the health effects of pollutant exposure**

Author Manuscript

Author Manuscript

It is well understood that a nutritious diet, rich in foods containing bioactive food components such as polyphenols and anti-inflammatory fatty acids, can be protective against chronic inflammation, diabetes, metabolic syndrome, and cardiovascular disease (16). Diets rich in these bioactive food components often contain greater levels of antioxidant and anti-inflammatory compounds and thus have the capacity to attenuate the inflammatory and oxidative properties of pollutant exposure. This paradigm has been demonstrated consistently throughout the literature (37–43). For example, we have shown that consumption of polyphenol rich green tea can decrease oxidative stress in response to PCB 126 exposure and does so via upregulation of antioxidant enzymatic pathways (38). Additionally, it has been observed that epigallocatechin gallate (EGCG), a major polyphenol in green tea, can attenuate arsenic-induced cardiovascular inflammation and toxicity (42). Resveratrol, a polyphenol found abundantly in fruits and other plants, has been demonstrated to protect against PCB-induced neuronal cell death as well as impairment in adipocyte glucose homeostasis (44, 45)

Omega-3 polyunsaturated fatty acids including eicosapentanoic (EPA) and docosahexenoic acid (DHA), found abundantly in fatty fish and fish oil, exhibit potent anti-inflammatory properties (46, 47). Indeed, individuals consuming a diet abundant in omega-3 fatty acids,

such as the well-studied Mediterranean diet, are documented as having lower incidences of cardiovascular disease, diabetes, and metabolic syndrome (48). It is not surprising then, that experimental evidence shows that omega-3 fatty acids can be protective against pollutant induced inflammation and toxicity. As such, we have demonstrated that omega-3 fatty acids can protect against PCB-induced vascular endothelial cell dysfunction (37) and others have shown that these fatty acids can protect against pollutant associated atherosclerotic plaque formation (49). Additionally, omega-3 fatty acids appear to be protective against air pollutant-associated oxidative stress and cardiovascular complications through increasing the activity of endogenous antioxidants and decreasing lipoperoxidation (40, 43).

In addition to consuming a diet rich in polyphenols and anti-inflammatory components, choosing to live an active lifestyle may also protect against pollutant toxicity. For example, a recent study by our laboratory observed that exercise is capable of reducing inflammation and cardiovascular disease risk associated with PCB exposure (50). Furthermore, other groups have demonstrated that exercise can attenuate PCB-associated alterations in the mouse gut microbiome (51). This is an important finding because the gut microbiome has recently been observed to regulate numerous health conditions including obesity, cardiovascular disease, immune function, and neurological disorders (52–54). Importantly, the gut microbiome is incredibly plastic and can change rapidly in response to nutrition, exercise, and even pollutant exposure (52–54). Utilizing nutrition and exercise to combat pollutant-induced alterations in the gut microbiome that may contribute to disease may allow for important advances in therapeutic interventions and is a critical area of research that needs to be further explored.

Because many of these environmental pollutants are lipophilic and accumulate within the adipose tissue of exposed individuals, examining nutritional means by which to promote excretion and reduction of body burden are an important facet as well. It has been noted that individuals consuming a vegetarian or vegan diet trended towards lower amounts of organochloride body burden (55). However, it remains unclear whether this observation was due to less exposure to contaminated foods (e.g. contaminated meats, fish, or dairy) or whether a plant-based diets contain nutritional components that can promote excretion of these pollutants. While further research is still warranted to examine nutritional means by which to reduce pollutant body burden, some studies have indicated promising opportunities for therapeutic interventions in exposed individuals (56–58). Specifically, it has been observed that consumption of Olestra, a dietary fat substitute, is effective at enhancing the excretion rate in individuals exposed to high concentrations of PCBs and is believed to do so through interference with the enterohepatic circulation of these pollutants (58–62). Because Olestra is no longer in production due to decreased popularity associated with adverse gastrointestinal side effects, there is a need for research examining other dietary components that may alter enterohepatic circulation and/or enhance pollutant excretion. As discussed, it is well documented that nutrition and lifestyle choices can both positively and negatively regulate pollutant toxicity. Recently, the field of epigenetics has become important in furthering our understanding how environment and lifestyle choices differentially influence gene expression and may alter susceptibility to disease on an individual level.

### III. Epigenetics as an emerging area of pollutant toxicity research and avenue for future intervention

#### Epigenetics overview

Epigenetics is a term given to describe all heritable changes in gene expression that do not involve changes within the genetic code (63). One of the most commonly studied means of epigenetic modifications is DNA methylation, which involves the addition of a methyl group to the 5 position of a cytosine. This modification most often occurs on a cytosine-guanine dinucleotide (CpG). CpG dinucleotides tend to be clustered in certain regions, termed CpG islands, and are often located in the promoter regions of protein-coding genes (63, 64). DNA methylation is involved in the regulation of numerous cellular processes, ranging from chromosome stability, to X-chromosome inactivation, to gene transcription. Generally, hypermethylation of CpG islands within the promoter regions results in gene silencing. Another common type of epigenetic alteration includes histone modifications. Histones are nuclear proteins that aid in the condensation of DNA into structural units called nucleosomes. Histones can undergo posttranslational modifications such as acetylation, methylation, and phosphorylation, which affect the interactions of histones with the DNA and associated nuclear proteins (63). Through these histone modifications, alterations in chromatin structure and subsequently gene expression are often observed (63, 64). A third major type of epigenetic modification involves microRNAs (miRNAs). miRNAs are small, single stranded, non-coding RNA molecules that when fully mature, lead to the degradation of target mRNA by direct binding. This downregulation of target mRNAs can ultimately lead to decreased protein production. miRNAs regulate numerous biological processes and it has been observed that alterations in miRNA expression is associated with the development of diseases such as cardiovascular disease, metabolic disorders and cancer (63–65).

Environment and lifestyle play critical roles in contributing to the epigenetic alterations discussed above (65). Studies on monozygotic twin pairs have demonstrated that throughout the aging process, identical twins develop differences in their epigenetic status despite their common genetic make-up (63). These observations and the knowledge that differences in epigenetic modifications can play a role in disease, contributed to increasing interest in examining the ways and extent to which our environment influences our epigenetic code.

#### Epigenetics, environmental exposures, and nutrition

It has been consistently demonstrated that exposure to environmental pollutants can influence epigenetic modifications (65–69). Specifically, exposure to persistent organic pollutants can alter the epigenetic code through several mechanisms. For example, we have demonstrated that exposure to coplanar PCBs, including PCB 77 and PCB126, can induce vascular endothelial cell dysfunction and inflammation through alterations in histone modifications of the p65 subunit of NF- $\kappa$ B that result in its increased activation (66). Additionally, we have shown that the commercial PCB mixture Aroclor 1260 can upregulate miRNAs associated with cardiac injury and inflammatory pathways in primary human endothelial cells (69). In humans, it has been observed that exposure to polycyclic aromatic hydrocarbons (PAH) results in alterations in specific miRNAs associated with oxidative DNA damage and lipid peroxidation in coke oven workers (67). Moreover, it was noted that

exposure to air pollution was associated with expression of miRNAs that may mediate detrimental biological responses and health effects. (68)

One common theme with many pollutant-associated epigenetic alterations is that they induce modifications that contribute to pro-inflammatory and oxidative pathways (65, 70). Interestingly, nutrition and positive lifestyle choices can also contribute to epigenetic modifications that may combat these detrimental effects by increasing antioxidant and anti-inflammatory responses (71, 72). Polyphenolic compounds found in many fruits and vegetables exhibit significant antioxidant and anti-inflammatory capacities and thus are good candidates for studies examining epigenetic means of protection against pollutant toxicity. For example, curcumin, a component found in turmeric that exhibits anti-inflammatory properties, has been observed to reduce histone acetylation of the NF- $\kappa$ B coactivator p300 histone acetyltransferase and thus suppress activation of NF- $\kappa$ B (73). Similar results were observed by our group with the tea catechin EGCG (74). Furthermore, it has been reported that numerous polyphenols can activate SIRT1, a member of the sirtuin family of histone deacetylases (HDACs). When activated, SIRT1 can participate in the deacetylation of transcription factors such as NF- $\kappa$ B, forkhead box class O (FoxO) and p53, in turn modulating cellular pathways involved in inflammation, metabolism, aging, and several other conditions (71).

In addition to healthful nutrition, physical exercise also contributes to epigenetic modifications that may provide protection against metabolic, cardiovascular, and other diseases. For example, one study found that acute exercise resulted in changes in the methylation of the promoter region of genes involved in energy and glucose homeostasis including pyruvate dehydrogenase kinase isoenzyme 4 (PDK4), peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC1 $\alpha$ ), and peroxisome proliferator-activated receptor delta (PPAR $\delta$ ) within skeletal muscle (75). Furthermore, physical exercise was observed to inhibit the miRNA miR-33 (76, 77). Cellular effects of inhibition of miR-33 involve upregulation of AMPK and subsequent increases in  $\beta$ -oxidation, which may aid in balancing factors related to metabolic syndrome by increasing the breakdown of fatty acids from adipose stores (77). The role of  $\beta$ -oxidation in metabolic syndrome is complex and is influenced both by dietary components and exercise leading to an increase in fatty acid oxidation and ultimately decreased obesity and metabolic syndrome (78–80). Understanding the role of miRNAs in lipid homeostasis may provide potential therapeutic strategies for reducing metabolic syndrome.

While epigenetics can be modulated by healthful nutrition and positive lifestyle choices, there is also evidence that epigenetics can be modified by poor nutrition. Epidemiological data are very helpful at examining epigenetic alterations, as there appears to be inheritance to specific DNA methylation patterns over generations. For example, it has been shown that individuals who were prenatally exposed to famine during The Dutch Famine (1940s) had reduced DNA methylation of the insulin-like growth factor II (IGF2) gene six decades later, compared to their siblings who were unexposed to the famine (81). Also, it has been demonstrated that parental obesity can alter DNA methylation patterns in imprinted genes of their offspring, compared to offspring of non-obese parents and this is believed to increase the risk for the development of obesity and metabolic diseases later in life (82). While there

is still much to be studied about the effects of these alterations related to epigenetic modifications, it is hypothesized that in both cases of undernutrition and over-nutrition, the acquired epigenetic alterations may be passed down to the subsequent generation and may increase susceptibility to development of chronic diseases throughout life (81–83).

The epigenetic alterations in response to nutrition and lifestyle choices have been well documented; however, studies examining the epigenetic effects of nutritional components on the modulation of pollutant toxicity are limited. A recent study conducted by our group showed that EGCG, the most abundant polyphenol found in green tea, can prevent against PCB126 induced endothelial cell inflammation through epigenetic alterations of proinflammatory target genes of NF- $\kappa$ B (74). This study was important in that it was one of the first studies to indicate the epigenetic role of healthful nutrition in the modulation of pollutant toxicity. Because of the gaps in the literature regarding epigenetic regulation of pollutant toxicity via nutritional and lifestyle interventions, there is a great need for further research to formulate a better understanding of this topic.

#### IV. Conclusions and future directions

Environmental pollutant exposure is associated with numerous health complications ranging from cardiovascular disease, diabetes, and metabolic syndrome. There is evidence that pollutant toxicity can be modulated by nutrition and lifestyle choices. Western diets, abundant in processed foods, excess caloric content and pro-inflammatory fatty acids, may contribute to the development of obesity, cardiovascular diseases, and can enhance pollutant toxicity. Conversely, diets rich in bioactive food components such as polyphenols and omega 3 fatty acids (i.e. the Mediterranean diet), are associated with reduced risk of inflammatory diseases and can attenuate the negative health effects of pollutant exposure. Recently, epigenetics has become a focus of both nutrition and pollutant research and may provide a deeper mechanistic insight into how these two environmental factors can interact. It has been shown that both over and undernutrition can contribute to epigenetic modifications that are associated with an increased risk of disease. Additionally, exposure to pollutants can alter epigenetic markers that may also contribute to the development of health complications. Importantly, we have shown that bioactive food components (e.g. EGCG) can induce epigenetic alterations that attenuate inflammatory responses associated with PCB exposure and thus may protect against the development atherosclerosis. Epigenetic marks differ for every individual and therefore future studies may allow us to understand how certain foods may be utilized to alter specific epigenetic tags to reduce the risk of pollutant-associated disease. Future studies also need to focus on metabolites of both pollutants and nutrients and the complex interplay of epigenetic regulation of chronic diseases. Fostering an understanding at this detailed level could help in developing personalized nutritional interventions that could epigenetically favor good health and thus reduce the risk of disease.

#### Acknowledgments

This work was supported by the National Institute of Environmental Health Sciences at the National Institutes of Health [P42ES007380], and the University of Kentucky Agricultural Experiment Station.



## References

1. Tang M, Chen K, Yang F, Liu W. Exposure to organochlorine pollutants and type 2 diabetes: a systematic review and meta-analysis. *PLoS one*. 2014; 9(10):e85556. [PubMed: 25329153]
2. Taylor KW, Novak RF, Anderson HA, Birnbaum LS, Blystone C, Devito M, et al. Evaluation of the association between persistent organic pollutants (POPs) and diabetes in epidemiological studies: a national toxicology program workshop review. *Environmental health perspectives*. 2013; 121(7):774–83. [PubMed: 23651634]
3. Lim EJ, Majkova Z, Xu S, Bachas L, Arzuaga X, Smart E, et al. Coplanar polychlorinated biphenyl-induced CYP1A1 is regulated through caveolae signaling in vascular endothelial cells. *Chemico-biological interactions*. 2008; 176(2–3):71–8. [PubMed: 18786521]
4. Majkova Z, Smart E, Toborek M, Hennig B. Up-regulation of endothelial monocyte chemoattractant protein-1 by coplanar PCB77 is caveolin-1-dependent. *Toxicology and applied pharmacology*. 2009; 237(1):1–7. [PubMed: 19265715]
5. Schlezinger JJ, Struntz WD, Goldstone JV, Stegeman JJ. Uncoupling of cytochrome P450 1A and stimulation of reactive oxygen species production by co-planar polychlorinated biphenyl congeners. *Aquatic toxicology*. 2006; 77(4):422–32. [PubMed: 16500718]
6. Kuehn BM. Environmental pollutants tied to atherosclerosis. *Jama*. 2011; 306(19):2081. [PubMed: 22089709]
7. Carpenter DO. Polychlorinated biphenyls (PCBs): routes of exposure and effects on human health. *Reviews on environmental health*. 2006; 21(1):1–23. [PubMed: 16700427]
8. Dirinck E, Jorens PG, Covaci A, Geens T, Roosens L, Neels H, et al. Obesity and persistent organic pollutants: possible obesogenic effect of organochlorine pesticides and polychlorinated biphenyls. *Obesity*. 2011; 19(4):709–14. [PubMed: 20559302]
9. Goncharov A, Pavuk M, Foushee HR, Carpenter DO, Anniston Environmental Health Research C. Blood pressure in relation to concentrations of PCB congeners and chlorinated pesticides. *Environmental health perspectives*. 2011; 119(3):319–25. [PubMed: 21362590]
10. Sergeev AV, Carpenter DO. Hospitalization rates for coronary heart disease in relation to residence near areas contaminated with persistent organic pollutants and other pollutants. *Environmental health perspectives*. 2005; 113(6):756–61. [PubMed: 15929900]
11. Baker NA, Karounos M, English V, Fang J, Wei Y, Stromberg A, et al. Coplanar polychlorinated biphenyls impair glucose homeostasis in lean C57BL/6 mice and mitigate beneficial effects of weight loss on glucose homeostasis in obese mice. *Environmental health perspectives*. 2013; 121(1):105–10. [PubMed: 23099484]
12. Dirinck E, Dirtu AC, Jorens PG, Malarvannan G, Covaci A, Van Gaal LF. Pivotal Role for the Visceral Fat Compartment in the Release of Persistent Organic Pollutants During Weight Loss. *The Journal of clinical endocrinology and metabolism*. 2015; 100(12):4463–71. [PubMed: 26469381]
13. Kim MJ, Marchand P, Henegar C, Antignac JP, Alili R, Poitou C, et al. Fate and complex pathogenic effects of dioxins and polychlorinated biphenyls in obese subjects before and after drastic weight loss. *Environmental health perspectives*. 2011; 119(3):377–83. [PubMed: 21156398]
14. Uemura H, Arisawa K, Hiyoshi M, Kitayama A, Takami H, Sawachika F, et al. Prevalence of metabolic syndrome associated with body burden levels of dioxin and related compounds among Japan's general population. *Environmental health perspectives*. 2009; 117(4):568–73. [PubMed: 19440495]
15. La Merrill M, Emond C, Kim MJ, Antignac JP, Le Bizec B, Clement K, et al. Toxicological function of adipose tissue: focus on persistent organic pollutants. *Environmental health perspectives*. 2013; 121(2):162–9. [PubMed: 23221922]
16. Chiuvè SE, Sampson L, Willett WC. The association between a nutritional quality index and risk of chronic disease. *American journal of preventive medicine*. 2011; 40(5):505–13. [PubMed: 21496749]

17. Wang L, Reiterer G, Toborek M, Hennig B. Changing ratios of omega-6 to omega-3 fatty acids can differentially modulate polychlorinated biphenyl toxicity in endothelial cells. *Chemico-biological interactions*. 2008; 172(1):27–38. [PubMed: 18155686]
18. Kopelman PG. Obesity as a medical problem. *Nature*. 2000; 404(6778):635–43. [PubMed: 10766250]
19. Baum SJ, Kris-Etherton PM, Willett WC, Lichtenstein AH, Rudel LL, Maki KC, et al. Fatty acids in cardiovascular health and disease: a comprehensive update. *Journal of clinical lipidology*. 2012; 6(3):216–34. [PubMed: 22658146]
20. Kuipers RS, de Graaf DJ, Luxwolda MF, Muskiet MH, Dijck-Brouwer DA, Muskiet FA. Saturated fat, carbohydrates and cardiovascular disease. *The Netherlands journal of medicine*. 2011; 69(9): 372–8. [PubMed: 21978979]
21. Grun F. Obesogens. *Current opinion in endocrinology, diabetes, and obesity*. 2010; 17(5):453–9.
22. Hennig B, Slim R, Toborek M, Robertson LW. Linoleic acid amplifies polychlorinated biphenyl-mediated dysfunction of endothelial cells. *Journal of biochemical and molecular toxicology*. 1999; 13(2):83–91. [PubMed: 9890193]
23. Wahlang B, Falkner KC, Gregory B, Ansert D, Young D, Conklin DJ, et al. Polychlorinated biphenyl 153 is a diet-dependent obesogen that worsens nonalcoholic fatty liver disease in male C57BL6/J mice. *The Journal of nutritional biochemistry*. 2013; 24(9):1587–95. [PubMed: 23618531]
24. Wu J, Liu J, Waalkes MP, Cheng ML, Li L, Li CX, et al. High dietary fat exacerbates arsenic-induced liver fibrosis in mice. *Experimental biology and medicine*. 2008; 233(3):377–84. [PubMed: 18296743]
25. Harris DL, Washington MK, Hood DB, Roberts LJ 2nd, Ramesh A. Dietary fat-influenced development of colon neoplasia in Apc Min mice exposed to benzo(a)pyrene. *Toxicologic pathology*. 2009; 37(7):938–46. [PubMed: 19841130]
26. Yanagisawa R, Koike E, Win-Shwe TT, Yamamoto M, Takano H. Impaired lipid and glucose homeostasis in hexabromocyclododecane-exposed mice fed a high-fat diet. *Environmental health perspectives*. 2014; 122(3):277–83. [PubMed: 24398136]
27. McGill HC Jr, McMahan CA, Gidding SS. Preventing heart disease in the 21st century: implications of the Pathobiological Determinants of Atherosclerosis in Youth (PDAY) study. *Circulation*. 2008; 117(9):1216–27. [PubMed: 18316498]
28. Perkins JT, Petriello MC, Newsome BJ, Hennig B. Polychlorinated biphenyls and links to cardiovascular disease. *Environmental science and pollution research international*. 2016; 23(3): 2160–72. [PubMed: 25877901]
29. Petriello MC, Han SG, Newsome BJ, Hennig B. PCB 126 toxicity is modulated by cross-talk between caveolae and Nrf2 signaling. *Toxicology and applied pharmacology*. 2014; 277(2):192–9. [PubMed: 24709675]
30. Petriello MC, Newsome B, Hennig B. Influence of nutrition in PCB-induced vascular inflammation. *Environmental science and pollution research international*. 2014; 21(10):6410–8. [PubMed: 23417440]
31. Senthong V, Li XS, Hudec T, Coughlin J, Wu Y, Levison B, et al. Plasma Trimethylamine N-Oxide, a Gut Microbe-Generated Phosphatidylcholine Metabolite, Is Associated With Atherosclerotic Burden. *Journal of the American College of Cardiology*. 2016; 67(22):2620–8. [PubMed: 27256833]
32. Bennett BJ, de Aguiar Vallim TQ, Wang Z, Shih DM, Meng Y, Gregory J, et al. Trimethylamine-N-oxide, a metabolite associated with atherosclerosis, exhibits complex genetic and dietary regulation. *Cell metabolism*. 2013; 17(1):49–60. [PubMed: 23312283]
33. Brown JM, Hazen SL. The gut microbial endocrine organ: bacterially derived signals driving cardiometabolic diseases. *Annual review of medicine*. 2015; 66:343–59.
34. Seldin MM, Meng Y, Qi H, Zhu W, Wang Z, Hazen SL, et al. Trimethylamine N-Oxide Promotes Vascular Inflammation Through Signaling of Mitogen-Activated Protein Kinase and Nuclear Factor-kappaB. *Journal of the American Heart Association*. 2016; 5(2)
35. Senthong V, Wang Z, Li XS, Fan Y, Wu Y, Tang WH, et al. Intestinal Microbiota-Generated Metabolite Trimethylamine-N-Oxide and 5-Year Mortality Risk in Stable Coronary Artery

- Disease: The Contributory Role of Intestinal Microbiota in a COURAGE-Like Patient Cohort. *Journal of the American Heart Association*. 2016; 5(6)
36. Petriello MC, Hoffman JB, Sunkara M, Wahlang B, Perkins JT, Morris AJ, et al. Dioxin-like pollutants increase hepatic flavin containing monooxygenase (FMO3) expression to promote synthesis of the pro-atherogenic nutrient biomarker trimethylamine N-oxide from dietary precursors. *The Journal of nutritional biochemistry*. 2016; 33:145–53. [PubMed: 27155921]
  37. Majkova Z, Layne J, Sunkara M, Morris AJ, Toborek M, Hennig B. Omega-3 fatty acid oxidation products prevent vascular endothelial cell activation by coplanar polychlorinated biphenyls. *Toxicology and applied pharmacology*. 2011; 251(1):41–9. [PubMed: 21130106]
  38. Newsome BJ, Petriello MC, Han SG, Murphy MO, Eske KE, Sunkara M, et al. Green tea diet decreases PCB 126-induced oxidative stress in mice by up-regulating antioxidant enzymes. *The Journal of nutritional biochemistry*. 2014; 25(2):126–35. [PubMed: 24378064]
  39. Petriello MC, Newsome BJ, Dziubla TD, Hilt JZ, Bhattacharyya D, Hennig B. Modulation of persistent organic pollutant toxicity through nutritional intervention: emerging opportunities in biomedicine and environmental remediation. *The Science of the total environment*. 2014; 491–492:11–6.
  40. Romieu I, Garcia-Esteban R, Sunyer J, Rios C, Alcaraz-Zubeldia M, Velasco SR, et al. The effect of supplementation with omega-3 polyunsaturated fatty acids on markers of oxidative stress in elderly exposed to PM(2.5). *Environmental health perspectives*. 2008; 116(9):1237–42. [PubMed: 18795169]
  41. Slim R, Toborek M, Robertson LW, Hennig B. Antioxidant protection against PCB-mediated endothelial cell activation. *Toxicological sciences : an official journal of the Society of Toxicology*. 1999; 52(2):232–9. [PubMed: 10630576]
  42. Sun TL, Liu Z, Qi ZJ, Huang YP, Gao XQ, Zhang YY. (-)-Epigallocatechin-3-gallate (EGCG) attenuates arsenic-induced cardiotoxicity in rats. *Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association*. 2016; 93:102–10. [PubMed: 27170490]
  43. Tong H, Rappold AG, Diaz-Sanchez D, Steck SE, Berntsen J, Cascio WE, et al. Omega-3 fatty acid supplementation appears to attenuate particulate air pollution-induced cardiac effects and lipid changes in healthy middle-aged adults. *Environmental health perspectives*. 2012; 120(7):952–7. [PubMed: 22514211]
  44. Guida N, Laudati G, Anzilotti S, Secondo A, Montuori P, Di Renzo G, et al. Resveratrol via sirtuin-1 downregulates RE1-silencing transcription factor (REST) expression preventing PCB-95-induced neuronal cell death. *Toxicology and applied pharmacology*. 2015; 288(3):387–98. [PubMed: 26307266]
  45. Baker NA, English V, Sunkara M, Morris AJ, Pearson KJ, Cassis LA. Resveratrol protects against polychlorinated biphenyl-mediated impairment of glucose homeostasis in adipocytes. *The Journal of nutritional biochemistry*. 2013; 24(12):2168–74. [PubMed: 24231106]
  46. Siriwardhana N, Kalupahana NS, Moustaid-Moussa N. Health benefits of n-3 polyunsaturated fatty acids: eicosapentaenoic acid and docosahexaenoic acid. *Advances in food and nutrition research*. 2012; 65:211–22. [PubMed: 22361189]
  47. Swanson D, Block R, Mousa SA. Omega-3 fatty acids EPA and DHA: health benefits throughout life. *Advances in nutrition*. 2012; 3(1):1–7. [PubMed: 22332096]
  48. Sofi F, Cesari F, Abbate R, Gensini GF, Casini A. Adherence to Mediterranean diet and health status: meta-analysis. *Bmj*. 2008; 337:a1344. [PubMed: 18786971]
  49. Turunen AW, Jula A, Suominen AL, Mannisto S, Marniemi J, Kiviranta H, et al. Fish consumption, omega-3 fatty acids, and environmental contaminants in relation to low-grade inflammation and early atherosclerosis. *Environmental research*. 2013; 120:43–54. [PubMed: 23089109]
  50. Murphy MO, Petriello MC, Han SG, Sunkara M, Morris AJ, Esser K, et al. Exercise protects against PCB-induced inflammation and associated cardiovascular risk factors. *Environmental science and pollution research international*. 2016; 23(3):2201–11. [PubMed: 25586614]
  51. Choi JJ, Eum SY, Rampersaud E, Daunert S, Abreu MT, Toborek M. Exercise attenuates PCB-induced changes in the mouse gut microbiome. *Environmental health perspectives*. 2013; 121(6):725–30. [PubMed: 23632211]

52. Round JL, Mazmanian SK. The gut microbiota shapes intestinal immune responses during health and disease. *Nature reviews Immunology*. 2009; 9(5):313–23.
53. Kinross JM, Darzi AW, Nicholson JK. Gut microbiome-host interactions in health and disease. *Genome medicine*. 2011; 3(3):14. [PubMed: 21392406]
54. Tremaroli V, Backhed F. Functional interactions between the gut microbiota and host metabolism. *Nature*. 2012; 489(7415):242–9. [PubMed: 22972297]
55. Arguin H, Sanchez M, Bray GA, Lovejoy JC, Peters JC, Jandacek RJ, et al. Impact of adopting a vegan diet or an olestra supplementation on plasma organochlorine concentrations: results from two pilot studies. *The British journal of nutrition*. 2010; 103(10):1433–41. [PubMed: 20030906]
56. Jandacek RJ, Genuis SJ. An assessment of the intestinal lumen as a site for intervention in reducing body burdens of organochlorine compounds. *The Scientific World Journal*. 2013; 2013:205621. [PubMed: 23476122]
57. Jandacek RJ, Tso P. Enterohepatic circulation of organochlorine compounds: a site for nutritional intervention. *The Journal of nutritional biochemistry*. 2007; 18(3):163–7. [PubMed: 17296488]
58. Jandacek RJ, Heubi JE, Buckley DD, Khoury JC, Turner WE, Sjodin A, et al. Reduction of the body burden of PCBs and DDE by dietary intervention in a randomized trial. *The Journal of nutritional biochemistry*. 2014; 25(4):483–8. [PubMed: 24629911]
59. Jandacek RJ, Rider T, Keller ER, Tso P. The effect of olestra on the absorption, excretion and storage of 2,2',5,5' tetrachlorobiphenyl; 3,3',4,4' tetrachlorobiphenyl; and perfluorooctanoic acid. *Environment international*. 2010; 36(8):880–3. [PubMed: 19616301]
60. Jandacek RJ, Anderson N, Liu M, Zheng S, Yang Q, Tso P. Effects of yo-yo diet, caloric restriction, and olestra on tissue distribution of hexachlorobenzene. *American journal of physiology Gastrointestinal and liver physiology*. 2005; 288(2):G292–9. [PubMed: 15513954]
61. Jandacek RJ. Intervention to reduce PCBs: learnings from a controlled study of Anniston residents. *Environmental science and pollution research international*. 2016; 23(3):2022–6. [PubMed: 25721531]
62. Moser GA, McLachlan MS. A non-absorbable dietary fat substitute enhances elimination of persistent lipophilic contaminants in humans. *Chemosphere*. 1999; 39(9):1513–21. [PubMed: 10481251]
63. Egger G, Liang G, Aparicio A, Jones PA. Epigenetics in human disease and prospects for epigenetic therapy. *Nature*. 2004; 429(6990):457–63. [PubMed: 15164071]
64. Chuang JC, Jones PA. Epigenetics and microRNAs. *Pediatric research*. 2007; 61(5):24R–9R. Pt 2.
65. Vrijens K, Bollati V, Nawrot TS. MicroRNAs as potential signatures of environmental exposure or effect: a systematic review. *Environmental health perspectives*. 2015; 123(5):399–411. [PubMed: 25616258]
66. Liu D, Perkins JT, Petriello MC, Hennig B. Exposure to coplanar PCBs induces endothelial cell inflammation through epigenetic regulation of NF-kappaB subunit p65. *Toxicology and applied pharmacology*. 2015; 289(3):457–65. [PubMed: 26519613]
67. Deng Q, Dai X, Guo H, Huang S, Kuang D, Feng J, et al. Polycyclic aromatic hydrocarbons-associated microRNAs and their interactions with the environment: influences on oxidative DNA damage and lipid peroxidation in coke oven workers. *Environmental science & technology*. 2014; 48(7):4120–8. [PubMed: 24555497]
68. Hou L, Barupal J, Zhang W, Zheng Y, Liu L, Zhang X, et al. Particulate Air Pollution Exposure and Expression of Viral and Human MicroRNAs in Blood: The Beijing Truck Driver Air Pollution Study. *Environmental health perspectives*. 2016; 124(3):344–50. [PubMed: 26068961]
69. Wahlang B, Petriello MC, Perkins JT, Shen S, Hennig B. Polychlorinated biphenyl exposure alters the expression profile of microRNAs associated with vascular diseases. *Toxicology in vitro : an international journal published in association with BIBRA*. 2016; 35:180–7. [PubMed: 27288564]
70. Rozek LS, Dolinoy DC, Sartor MA, Omenn GS. Epigenetics: relevance and implications for public health. *Annual review of public health*. 2014; 35:105–22.
71. Ayissi VB, Ebrahimi A, Schluesener H. Epigenetic effects of natural polyphenols: a focus on SIRT1-mediated mechanisms. *Molecular nutrition & food research*. 2014; 58(1):22–32. [PubMed: 23881751]

72. Ross SA, Davis CD. The emerging role of microRNAs and nutrition in modulating health and disease. *Annual review of nutrition*. 2014; 34:305–36.
73. Yun JM, Jialal I, Devaraj S. Epigenetic regulation of high glucose-induced proinflammatory cytokine production in monocytes by curcumin. *The Journal of nutritional biochemistry*. 2011; 22(5):450–8. [PubMed: 20655188]
74. Liu D, Perkins JT, Hennig B. EGCG prevents PCB-126-induced endothelial cell inflammation via epigenetic modifications of NF-kappaB target genes in human endothelial cells. *The Journal of nutritional biochemistry*. 2016; 28:164–70. [PubMed: 26878794]
75. Barres R, Yan J, Egan B, Treebak JT, Rasmussen M, Fritz T, et al. Acute exercise remodels promoter methylation in human skeletal muscle. *Cell metabolism*. 2012; 15(3):405–11. [PubMed: 22405075]
76. Davalos A, Goedeke L, Smibert P, Ramirez CM, Warriar NP, Andreo U, et al. miR-33a/b contribute to the regulation of fatty acid metabolism and insulin signaling. *Proceedings of the National Academy of Sciences of the United States of America*. 2011; 108(22):9232–7. [PubMed: 21576456]
77. Pareja-Galeano H, Sanchis-Gomar F, Garcia-Gimenez JL. Physical exercise and epigenetic modulation: elucidating intricate mechanisms. *Sports medicine*. 2014; 44(4):429–36. [PubMed: 24399634]
78. Yang Z, Cappello T, Wang L. Emerging role of microRNAs in lipid metabolism. *Acta pharmaceutica Sinica B*. 2015; 5(2):145–50. [PubMed: 26579440]
79. Monsalve FA, Pyarasani RD, Delgado-Lopez F, Moore-Carrasco R. Peroxisome proliferator-activated receptor targets for the treatment of metabolic diseases. *Mediators of inflammation*. 2013; 2013:549627. [PubMed: 23781121]
80. Huang CW, Chien YS, Chen YJ, Ajuwon KM, Mersmann HM, Ding ST. Role of n-3 Polyunsaturated Fatty Acids in Ameliorating the Obesity-Induced Metabolic Syndrome in Animal Models and Humans. *International journal of molecular sciences*. 2016; 17(10)
81. Heijmans BT, Tobi EW, Stein AD, Putter H, Blauw GJ, Susser ES, et al. Persistent epigenetic differences associated with prenatal exposure to famine in humans. *Proceedings of the National Academy of Sciences of the United States of America*. 2008; 105(44):17046–9. [PubMed: 18955703]
82. Soubry A, Murphy SK, Wang F, Huang Z, Vidal AC, Fuemmeler BF, et al. Newborns of obese parents have altered DNA methylation patterns at imprinted genes. *Int J Obes (Lond)*. 2015; 39(4):650–7. [PubMed: 24158121]
83. Vickers MH. Early life nutrition, epigenetics and programming of later life disease. *Nutrients*. 2014; 6(6):2165–78. [PubMed: 24892374]