



Correction to “Detection of Active Mammalian GH31 α -Glucosidases in Health and Disease Using In-Class, Broad-Spectrum Activity-Based Probes”

Jianbing Jiang, Chi-Lin Kuo, Liang Wu, Christian Franke, Wouter W. Kallemeijn,^{id} Bogdan I. Florea, Eline van Meel, Gijsbert A. van der Marel, Jeroen D. C. Codée,^{id} Rolf G. Boot, Gideon J. Davies,^{id} Herman S. Overkleeft,^{*id} and Johannes M. F. G. Aerts^{*}

ACS Cent. Sci. 2016, 2, 351–358. DOI: [10.1021/acscentsci.6b00057](https://doi.org/10.1021/acscentsci.6b00057)

In this paper we report an IC_{50} value for 1,6-*epi*-cyclophellitol (compound 7, JJB307) as a human lysosomal α -glucosidase (GAA) inhibitor of 54.1 ± 4.9 nM *in vitro* and 97.6 ± 14.5 nM *in situ* (Figure 3a). We re-evaluated these data and now find that compound 7 is actually a low micromolar GAA inhibitor with an IC_{50} of 14.6 ± 1.6 μ M *in vitro* and >50 μ M *in situ*.

Received: April 28, 2017

Published: May 18, 2017

