

Published in final edited form as:

Paediatr Perinat Epidemiol. 2016 September ; 30(5): 511–520. doi:10.1111/ppe.12307.

Associations of infant subcutaneous fat mass with total and abdominal fat mass at school-age. The Generation R Study

Susana Santos^{a,b}, Romy Gaillard^{b,c,d}, Andreia Oliveira^{a,e}, Henrique Barros^{a,e}, Marieke Abrahamse-Berkeveld^f, Eline M van der Beek^f, Albert Hofman^d, and Vincent WV Jaddoe^{b,c,d}

^aEPI-Unit, Institute of Public Health, University of Porto, Porto, Portugal ^bThe Generation R Study Group, Erasmus MC, University Medical Center, Rotterdam, The Netherlands ^cDepartment of Pediatrics, Erasmus MC, University Medical Center, Rotterdam, The Netherlands ^dDepartment of Epidemiology, Erasmus MC, University Medical Center, Rotterdam, The Netherlands ^eDepartment of Clinical Epidemiology, Predictive Medicine and Public Health, University of Porto Medical School, Porto, Portugal ^fNutricia Research, Danone Nutricia Early Life Nutrition, Utrecht, The Netherlands

Abstract

Background—Skinfold thickness enables the measurement of overall and regional subcutaneous fatness in infancy and may be associated with total and abdominal body fat in later childhood. We examined the associations of subcutaneous fat in infancy with total and abdominal fat at school-age.

Methods—In a population-based prospective cohort study among 821 children, we calculated total subcutaneous fat (sum of biceps, triceps, suprailiacal and subscapular skinfold thicknesses) and central-to-total subcutaneous fat ratio (sum of suprailiacal and subscapular skinfold thicknesses/total subcutaneous fat) at 1.5 and 24 months. At 6 years, we measured fat mass index (total fat/height³), central-to-total fat ratio (trunk fat/total fat) and android-to-gynoid fat ratio (android fat/gynoid fat) by dual-energy X-ray absorptiometry and preperitoneal fat mass area by abdominal ultrasound.

This is the peer reviewed version of the following article: Santos, S., Gaillard, R., Oliveira, A., Barros, H., Abrahamse-Berkeveld, M., van der Beek, E. M., Hofman, A. and Jaddoe, V. W. V. (2016), Associations of Infant Subcutaneous Fat Mass with Total and Abdominal Fat Mass at School-Age: The Generation R Study. *Paediatr. Perinat. Epidemiol.*, 30: 511–520. doi:10.1111/ppe.12307, which has been published in final form at <http://onlinelibrary.wiley.com/doi/10.1111/ppe.12307/abstract>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

Corresponding author: Vincent WV Jaddoe, MD, PhD. The Generation R Study Group (Na29-15). Erasmus MC, University Medical Center, PO Box 2040, 3000 CA Rotterdam, The Netherlands (v.jaddoe@erasmusmc.nl). Phone: +31 (0) 10 7043405, Fax: +31 (0) 10 7044645.

Conflict of Interest Statement

No conflict of interest was declared.

Author Contributions

SS, RG and VVWJ designed the research project; AH and VVWJ were involved in the design and planning of the study and data collection; SS and RG conducted the analyses; SS, RG and VVWJ wrote the paper; AO, HB, MAB, EMvdB and AH critically reviewed the manuscript; SS, RG and VVWJ had primary responsibility for the final content. All authors read and approved the final manuscript.

Results—Central-to-total subcutaneous fat ratio at 1.5 months was positively associated with fat mass index and central-to-total fat ratio at 6 years, whereas both total and central-to-total subcutaneous fat ratio at 24 months were positively associated with all childhood adiposity measures ($p < 0.05$). A 1-standard-deviation scores higher total subcutaneous fat at 24 months was associated with an increased risk of childhood overweight (Odds Ratio 1.70 [95% Confidence Interval 1.36, 2.12]). These associations were weaker than those for body mass index and stronger among girls than boys.

Conclusions—Subcutaneous fat in infancy is positively associated with total and abdominal fat at school-age. Our results also suggest that skinfold thicknesses add little value to estimate later body fat, as compared to body mass index.

Keywords

infancy; subcutaneous fat; tracking; childhood; abdominal fat

Infancy seems to be a critical period for the development of obesity.¹ An accumulating body of evidence has suggested that nutrition and growth in infancy are related to the risk of obesity in later life.^{2, 3} Also, infant growth patterns seem to be related not only to body mass index, but also to an adverse body fat distribution.^{4, 5} Several studies have shown that compared to body mass index, body fat distribution plays a greater role in the development of obesity-related complications, such as cardiovascular disease and type 2 diabetes.⁶ Although previous studies have suggested that central fat mass tracks moderately from mid-childhood into adulthood,^{7, 8, 9, 10} it is not known whether body fat distribution measures in infancy are associated with similar measures in later childhood.

Skinfold thickness is a valid measurement of subcutaneous fat mass that enables assessment of overall and regional fatness in infancy.¹¹ Previously, we reported the tracking of subcutaneous fat mass measured by skinfold thickness during the first 2 years of life.¹² Assessing the associations of these specific fat mass measures during infancy with fat mass measures during childhood helps to further understand the stability of body fat across childhood. Also, skinfold thickness measurements in infancy may be associated with an adverse body fat pattern in later childhood.

Therefore, we examined, in a population-based prospective cohort study among 821 children, the associations of infant subcutaneous fat mass measures with total and abdominal fat mass measures and with the risk of overweight at school-age.

Methods

Study design

This study was embedded in the Generation R Study, a population-based prospective cohort study from early pregnancy onwards among 9778 mothers and their children living in Rotterdam, the Netherlands.¹³ The study was approved by the local Medical Ethical Committee. Written informed consent was obtained from all mothers. Additional detailed assessments of fetal and postnatal growth and development were conducted in a subgroup of Dutch mothers and their children from late pregnancy onwards. Of all approached women,

80% agreed to participate. Of the total of 1205 singleton children participating in the subgroup study, 965 children had body mass index or skinfold thicknesses measured at the age of 1.5 or 24 months. Of the group of 965 children, 821 children had follow-up measurements at the age of 6 years (Flow chart is given in Supplemental Figure S1).

Body fat measurements during infancy

We measured weight to the nearest gram in naked infants at the age of 1.5 months by using an electronic infant scale and at 24 months by using a mechanical personal scale (SECA, Almere, The Netherlands). Body length at the age of 1.5 months was measured in supine position to the nearest millimeter by using a neonatometer and body height at 24 months was measured in standing position by using a Harpenden stadiometer (Holtain Limited, Dyfed, UK). Body mass index (kg/m^2) was calculated.

We measured skinfold thicknesses at the ages of 1.5 and 24 months on the left side of the body at the biceps, triceps, suprailiacal and subscapular area by using a skinfold caliper (Slim Guide, Creative Health Products) according to standard procedures.¹² We calculated total subcutaneous fat mass from the sum of all four skinfold thicknesses, and central subcutaneous fat mass from the sum of suprailiacal and subscapular skinfold thicknesses.^{14, 15} To create total subcutaneous fat mass independent of length or height and central subcutaneous fat mass independent of total subcutaneous fat mass, we estimated the optimal adjustment by log-log regression analyses.¹⁶ Details of these regressions are given in the Supplemental Methods. Total subcutaneous fat mass was only weakly correlated with length or height, and was not adjusted for it whereas a central-to-total subcutaneous fat mass ratio was calculated as central divided by total subcutaneous fat mass.

Body fat measurements at school-age

Measurements were performed in a dedicated research center by a well-trained staff.⁴ We measured height to the nearest millimeter by using a Harpenden stadiometer (Holtain Limited, Dyfed, UK) and weight to the nearest gram by using a mechanical personal scale (SECA, Almere, The Netherlands) in standing position without shoes and heavy clothing. We calculated body mass index (kg/m^2), and defined overweight and obesity as described by Cole et al.¹⁷

We measured total and regional body fat mass using a DXA scanner (iDXA, GE-Lunar, 2008, Madison, WI, USA, enCORE software v.12.6), according to standard procedures.⁴ Previous studies have validated DXA against computed tomography for body fat assessment.^{18, 19, 20} We divided total fat mass by height³ in order to obtain a fat mass index uncorrelated with height, as confirmed by a log-log regression analysis.^{16, 21} We assessed central fat mass as fat mass of the trunk, and divided by total fat mass in order to obtain a central-to-total fat mass ratio effectively uncorrelated with total fat mass. We calculated the ratio of android and gynoid fat mass, which reflects the relation between fat mass in the abdomen (android) and hip (gynoid) regions.

Abdominal preperitoneal fat mass, as a proxy for visceral fat mass, was measured by abdominal ultrasound (GE LOGIQ E9, Milwaukee, WI, USA), as described previously.²² Briefly, a linear (L12-5 MHz) transducer was placed perpendicular to the skin surface on the

median upper abdomen.²³ We scanned longitudinally from the xiphoid process to the navel along the midline (linea alba). Preperitoneal fat mass area was measured as the area of 2 cm length along the midline starting from the reference point in direction of the navel.

Covariates

Information on maternal age, educational level, parity, pre-pregnancy weight and smoking habits during pregnancy was assessed using self-reported questionnaires during pregnancy. We measured maternal height at enrolment, and calculated pre-pregnancy body mass index (kg/m^2). Information about child's sex, gestational age and weight at birth was obtained from medical records. Information about breast feeding duration, timing of introduction of solid foods and average television watching time at 6 years old was obtained by questionnaires.

Statistical analysis

We examined differences between boys and girls for maternal and child's characteristics with Student's t-tests and Mann-Whitney tests for normally and non-normally distributed variables, respectively and with χ^2 -test for dichotomous variables. First, we used Pearson's or Spearman's rank correlation coefficients to estimate correlations of subcutaneous fat mass measures at 1.5 or 24 months with total and abdominal fat mass measures at 6 years. We, subsequently, assessed the associations of infant subcutaneous fat mass measures with childhood total and abdominal fat mass measures using linear regression models and with the risk of childhood overweight using logistic regression models. These regression models were adjusted for maternal age, educational level, parity, pre-pregnancy body mass index and smoking habits during pregnancy, and child's gestational age-adjusted birthweight SDS, breast feeding duration, timing of introduction of solid foods, and TV watching time. We included covariates in the models when they changed the effect estimates substantially ($>10\%$), or when they were strongly associated with body fat mass in our or previous studies. Since we observed statistically significant interactions between infant subcutaneous fat mass measures and child's sex in the association with childhood total and abdominal fat mass measures, we performed all analyses for the total group and for boys and girls separately. No significant interactions were observed with body mass index categories at 1.5 or 24 months. We constructed SDS [(observed value - mean)/SD] for all continuous body fat measures at each age to take into account the expected changes in body composition and fat distribution across ages and also to enable comparisons in effect size for different exposure and outcome measures. Missing values in covariates (ranging from 0 to 15%) were multiple-imputed, by using Markov chain Monte Carlo approach. Five imputed datasets were created and analyzed together. We performed statistical analyses using the Statistical Package of Social Sciences version 21.0 for Windows (SPSS Inc, Chicago, IL, USA).

Results

Subject characteristics

Tables 1 and 2 show the subject characteristics. Boys had higher body mass index, whereas girls had higher central-to-total subcutaneous fat mass ratio at 1.5 months and a higher total subcutaneous fat mass and central-to-total subcutaneous fat mass ratio at 24 months

($p < 0.05$). At 6 years, girls had higher fat mass index, central-to-total fat mass ratio, and abdominal preperitoneal fat mass area than boys ($p < 0.05$). Supplemental Table S2 gives all subcutaneous fat mass measures at 1.5 and 24 months. Non-response analyses showed that as compared to children who did not participate in the follow-up studies, those who did participate were born with a higher weight and gestational age at birth and were breastfed for a longer period ($p < 0.05$). (Supplemental Table S3).

Body fat from infancy to childhood

The crude correlations of infant subcutaneous fat mass measures at 1.5 or 24 months with total and abdominal fat mass measures at 6 years old for the total group and by sex are shown in Supplemental Tables S4 and S5, respectively.

A 1-standard-deviation scores (SDS) higher body mass index at 1.5 months was only associated with a 0.16 [95% Confidence Interval (CI) 0.09, 0.24] SDS increase in body mass index at 6 years (Table 3). A 1-SDS higher central-to-total subcutaneous fat mass ratio at 1.5 months was associated with a 0.12 [95% CI 0.05, 0.19] SDS increase in fat mass index and a 0.11 [95% CI 0.04, 0.18] SDS increase in central-to-total fat mass ratio at 6 years. No other associations of fat mass at 1.5 months with fat outcomes at 6 years were observed. Also, no sex differences were observed.

A 1-SDS higher body mass index at 24 months was associated with an increase of 0.48 [95% CI 0.42, 0.54] SDS in body mass index, 0.33 [95% CI 0.26, 0.40] SDS in fat mass index, 0.18 [95% CI 0.11, 0.26] SDS in central-to-total fat mass ratio, and 0.16 [95% CI 0.08, 0.23] SDS in android-to-gynoid fat mass ratio (Table 4). A 1-SDS higher total subcutaneous fat mass at 24 months was associated with an increase of 0.30 [95% CI 0.23, 0.37] SDS in body mass index, 0.36 [95% CI 0.29, 0.43] SDS in fat mass index, 0.28 [95% CI 0.20, 0.35] SDS in central-to-total fat mass ratio, 0.23 [95% CI 0.16, 0.31] SDS in android-to-gynoid fat mass ratio and 0.19 [95% CI 0.12, 0.26] SDS in preperitoneal fat mass area. The associations of central-to-total subcutaneous fat mass ratio at 24 months with fat mass measures at 6 years were also significant but weaker. The associations tended to be stronger among girls than among boys.

Infant body fat and risk of childhood overweight

A 1-SDS higher body mass index at 1.5 months was associated with an increased risk of overweight at 6 years (Odds Ratio (OR) 1.40 [95% CI 1.08, 1.81]) (Figure 1), whereas a 1-SDS higher central-to-total subcutaneous fat mass ratio at 1.5 months was associated with an increased risk of overweight at 6 years among girls only (OR 1.61 [95% CI 1.09, 2.38]). A 1-SDS higher body mass index and total subcutaneous fat mass at 24 months were associated with increased risks of childhood overweight (ORs 2.76 [95% CI 2.07, 3.69] and 1.70 [95% CI 1.36, 2.12]), respectively). We did not observe associations for central-to-total subcutaneous fat mass ratio at 24 months with the risk of childhood overweight. Stronger associations were present among girls than among boys.

Comments

We observed that infant subcutaneous fat mass measures, calculated from skinfold thickness, are associated with total and abdominal fat mass at school-age. The effect estimates were stronger for body mass index and total subcutaneous fat mass than for central-to-total subcutaneous fat mass ratio, and for girls compared to boys. Also, the effect estimates were stronger for 24 months than for 1.5 months fat mass measures.

Interpretation of main findings

Body mass index tends to track from infancy onwards. A previous review among 21 studies has shown that large body size (weight or body mass index) in 0- to 4-year-old children was related to large body size at primary school age.²⁴ A meta-regression analysis among 48 cohort studies has shown a high degree of tracking for body mass index among children until 10 years of age. The tracking estimates gradually decreased over follow-up time and were not influenced by body mass index at first measurement or sex.²⁵ In our study, we observed a moderate tracking of body mass index from infancy to childhood. Stronger associations were observed from 24 months onwards than from 1.5 months onwards, in line with the stronger tracking reported in the meta-regression for shorter follow-up periods. We did not observe differences in results when we used ponderal index (weight/height³) at 1.5 months (data not shown). We observed slightly stronger effect estimates among girls compared to boys.

Body mass index provides limited information about body fat distribution. Android-to-gynoid fat mass ratio as well as preperitoneal fat mass have previously been associated with an adverse cardiovascular risk profile in childhood and adulthood, independently of body mass index.^{26, 27} Previous studies have suggested that total and central fat mass track moderately from childhood into adulthood,^{7, 8, 9, 10, 28, 29, 30} although with a lower tracking as compared to body mass index. Since we used different body fat mass measures in infancy and at school-age, we could not directly estimate tracking coefficients. However, we observed that total subcutaneous fat mass at 24 months was positively associated with fat mass index at school-age, suggesting tracking of total fat mass from infancy into school-age. Higher total and central-to-total subcutaneous fat mass ratio in infancy were associated with higher central-to-total and android-to-gynoid fat mass ratios measured by DXA and preperitoneal fat mass area measured by abdominal ultrasound at 6 years old. Our results suggest that besides higher body mass index, higher total subcutaneous fat mass and central-to-total subcutaneous fat mass ratio in infancy relate to an adverse body fat profile at school-age.

We observed stronger effect estimates for infant total subcutaneous fat mass than for central-to-total subcutaneous fat mass ratio, in the associations with similar measures at school-age, though both showed weaker effect estimates compared to the tracking of body mass index from infancy to childhood. The latter may be explained by the differences in methodology to assess the fat mass components in infancy and childhood. In infancy, skinfold thickness measurements reflect the subcutaneous depots only, whereas in childhood DXA comprise both subcutaneous and intra-abdominal depots.¹¹ However, during the first 4 months of life approximately 90% of body fat is located subcutaneously,³¹ and preperitoneal fat mass

seems to increase only from the second year of life onwards.³² Finally, skinfold thickness measurements may be more liable to measurement error than body mass index,^{33, 34} which may also lead to an underestimation of the effect estimates for subcutaneous fat measures from infancy onwards. We also observed stronger effect estimates among girls than boys. Sex-specific adiposity differences have been attributed to sex hormones. During infancy, testosterone concentrations increase during the first week of life before decreasing around 6 months of life among boys, whereas oestrogen increases shortly after birth and remains raised until 2-3 years among girls.³⁵ The stronger associations that we observed among girls may be partly a result of a more stable body fat development during infancy driven by less hormonal fluctuations. Further studies are needed to explore the mechanisms underlying the observed sex differences in body fat development. The stronger associations observed for 24 months than for 1.5 months could be due to the shorter interval between 24 months and 6 years old, as previously reported for body mass index.²⁵ Also, 1.5 months might be more reflective of fetal growth patterns which seemed to be less associated, as compared to postnatal growth, with later abdominal adiposity in our previous study.⁵

Whether skinfold thickness measurements during infancy are useful in clinical practice is not known. Our results suggest that compared to body mass index, subcutaneous fat mass measures in infancy add little value to estimate total or abdominal fat mass at school-age. Given the challenges of obtaining precise and reliable skinfold thickness measurements, the additional clinical value of skinfold thicknesses compared to body mass index may be only limited.

Methodological considerations

Major strengths of this study are the population-based prospective design with detailed infant and childhood body fat measurements available. Of the 965 singleton children with information on body fat mass measures at the age of 1.5 or 24 months, 85% (821) participated in the adiposity follow-up study at 6 years old. The non-response could lead to biased effect estimates if the associations of interest would differ between children included and not included in the analyses. Children included in the analyses were born with a higher weight and gestational age and were breastfed for a longer period compared to those not included. It is difficult to speculate if these differences might have influenced our effect estimates. However, this seems unlikely since children that did not participate in the follow-up studies did not differ from those who did participate regarding infant body mass index and subcutaneous fat mass measures. Our sample was ethnically homogeneous, which may limit the generalizability of our results to other ethnic groups.²⁵ We used skinfold thickness which is a valid measurement of subcutaneous fat in children but in extremely overweight children the measurement error is larger.¹⁵ The inter- and intra-observer measurement error is also larger as compared to other anthropometric measurements.^{33, 34} We did not have available skinfold thickness measurements from the lower limbs such as thigh or calf, which could have improved our estimates of subcutaneous fat mass. We obtained detailed measures of adiposity at school-age by using DXA that quantifies body fat content with high precision and abdominal ultrasound which is a valid method to assess abdominal preperitoneal fat mass area.²³ Finally, although we adjusted for a large number of potential confounders,

residual confounding in the observed associations might still occur, as in any observational study.

Conclusion

Subcutaneous fat mass measures in infancy are positively associated with total and abdominal fat mass at school-age. The effect estimates were stronger for body mass index and total subcutaneous fat mass than for central-to-total subcutaneous fat mass ratio, and for girls compared to boys. Our results suggest that skinfold thickness measurements in infancy add little value to estimate body fat in later childhood, as compared to body mass index.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We gratefully acknowledge the contribution of general practitioners, hospitals, midwives, and pharmacies in Rotterdam.

Funding

The Generation R Study is made possible by financial support from Erasmus MC, University Medical Center, Rotterdam, Erasmus University Rotterdam, Netherlands Organization for Health Research and Development (ZonMw), Netherlands Organisation for Scientific Research (NWO), Ministry of Health, Welfare and Sport and Ministry of Youth and Families. Research leading to these results has received funding from an unrestricted grant from Nutricia Research and the European Union's Seventh Framework Programme (FP7/2007-2013), project EarlyNutrition under grant agreement n°289346. SS received a grant from the Portuguese Foundation for Science and Technology (SFRH/BD/81123/2011). VJ received an additional grant from the Netherlands Organization for Health Research and Development (VIDI 016.136.361) and Consolidator Grant from the European Research Council (ERC-2014-CoG-648916).


References

1. Gillman MW. The first months of life: a critical period for development of obesity. *The American Journal of Clinical Nutrition*. 2008; 87:1587–1589. [PubMed: 18541543]
2. McCarthy A, Hughes R, Tilling K, Davies D, Smith GD, Ben-Shlomo Y. Birth weight; postnatal, infant, and childhood growth; and obesity in young adulthood: evidence from the Barry Caerphilly Growth Study. *The American Journal of Clinical Nutrition*. 2007; 86:907–913. [PubMed: 17921364]
3. Taal HR, Vd Heijden AJ, Steegers EA, Hofman A, Jaddoe VW. Small and large size for gestational age at birth, infant growth, and childhood overweight. *Obesity (Silver Spring)*. 2013; 21:1261–1268. [PubMed: 23666877]
4. Gishti O, Gaillard R, Manniesing R, Abrahamse-Berkeveld M, van der Beek EM, Heppe DH, et al. Fetal and infant growth patterns associated with total and abdominal fat distribution in school-age children. *The Journal of Clinical Endocrinology and Metabolism*. 2014; 99:2557–2566. [PubMed: 24712569]
5. Durmus B, Mook-Kanamori DO, Holzhauser S, Hofman A, van der Beek EM, Boehm G, et al. Growth in foetal life and infancy is associated with abdominal adiposity at the age of 2 years: the generation R study. *Clinical Endocrinology*. 2010; 72:633–640. [PubMed: 19769622]
6. Despres JP. Body fat distribution and risk of cardiovascular disease: an update. *Circulation*. 2012; 126:1301–1313. [PubMed: 22949540]
7. Chrzanowska M, Suder A, Kruszelnicki P. Tracking and risk of abdominal obesity in the adolescence period in children aged 7-15. The Cracow Longitudinal Growth Study. *American Journal of Human Biology*. 2012; 24:62–67. [PubMed: 22131191]

8. Freitas D, Beunen G, Maia J, Claessens A, Thomis M, Marques A, et al. Tracking of fatness during childhood, adolescence and young adulthood: a 7-year follow-up study in Madeira Island, Portugal. *Annals of Human Biology*. 2012; 39:59–67. [PubMed: 22148930]
9. Monyeki KD, Kemper HC, Makgae PJ. Development and tracking of central patterns of subcutaneous fat of rural South African youth: Ellisras longitudinal study. *BMC Pediatrics*. 2009; 9:74. [PubMed: 20003226]
10. Psarra G, Nassis GP, Sidossis LS. Short-term predictors of abdominal obesity in children. *European Journal of Public Health*. 2006; 16:520–525. [PubMed: 16230317]
11. Wells JC, Fewtrell MS. Measuring body composition. *Archives of Disease in Childhood*. 2006; 91:612–617. [PubMed: 16790722]
12. Ay L, Hokken-Koelega AC, Mook-Kanamori DO, Hofman A, Moll HA, Mackenbach JP, et al. Tracking and determinants of subcutaneous fat mass in early childhood: the Generation R Study. *International Journal of Obesity (Lond)*. 2008; 32:1050–1059.
13. Jaddoe VW, van Duijn CM, Franco OH, van der Heijden AJ, van Iizendoorn MH, de Jongste JC, et al. The Generation R Study: design and cohort update 2012. *European Journal of Epidemiology*. 2012; 27:739–756. [PubMed: 23086283]
14. Ketel IJ, Volman MN, Seidell JC, Stehouwer CD, Twisk JW, Lambalk CB. Superiority of skinfold measurements and waist over waist-to-hip ratio for determination of body fat distribution in a population-based cohort of Caucasian Dutch adults. *European Journal of Endocrinology*. 2007; 156:655–661. [PubMed: 17535865]
15. Freedman DS, Wang J, Ogden CL, Thornton JC, Mei Z, Pierson RN, et al. The prediction of body fatness by BMI and skinfold thicknesses among children and adolescents. *Annals of Human Biology*. 2007; 34:183–194. [PubMed: 17558589]
16. Wells JC, Cole TJ, ALSPAC study team. Adjustment of fat-free mass and fat mass for height in children aged 8 y. *International Journal of Obesity and Related Metabolic Disorders*. 2002; 26:947–952. [PubMed: 12080448]
17. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*. 2000; 320:1240–1243. [PubMed: 10797032]
18. Kaul S, Rothney MP, Peters DM, Wacker WK, Davis CE, Shapiro MD, et al. Dual-energy X-ray absorptiometry for quantification of visceral fat. *Obesity (Silver Spring)*. 2012; 20:1313–1318. [PubMed: 22282048]
19. Shuster A, Patlas M, Pinthus JH, Mourtzakis M. The clinical importance of visceral adiposity: a critical review of methods for visceral adipose tissue analysis. *The British Journal of Radiology*. 2012; 85:1–10.
20. Helba M, Binkovitz LA. Pediatric body composition analysis with dual-energy X-ray absorptiometry. *Pediatric Radiology*. 2009; 39:647–656. [PubMed: 19415261]
21. VanItallie TB, Yang MU, Heymsfield SB, Funk RC, Boileau RA. Height-normalized indices of the body's fat-free mass and fat mass: potentially useful indicators of nutritional status. *The American Journal of Clinical Nutrition*. 1990; 52:953–959. [PubMed: 2239792]
22. Mook-Kanamori DO, Holzhauser S, Hollestein LM, Durmus B, Manniesing R, Koek M, et al. Abdominal fat in children measured by ultrasound and computed tomography. *Ultrasound in Medicine and Biology*. 2009; 35:1938–1946. [PubMed: 19800165]
23. Suzuki R, Watanabe S, Hirai Y, Akiyama K, Nishide T, Matsushima Y, et al. Abdominal wall fat index, estimated by ultrasonography, for assessment of the ratio of visceral fat to subcutaneous fat in the abdomen. *The American Journal of Medicine*. 1993; 95:309–314. [PubMed: 8368228]
24. Stocks T, Renders CM, Bulk-Bunschoten AM, Hirasings RA, van Buuren S, Seidell JC. Body size and growth in 0- to 4-year-old children and the relation to body size in primary school age. *Obesity reviews : an official journal of the International Association for the Study of Obesity*. 2011; 12:637–652. [PubMed: 21426479]
25. Bayer O, Kruger H, von Kries R, Toschke AM. Factors associated with tracking of BMI: a meta-regression analysis on BMI tracking. *Obesity (Silver Spring)*. 2011; 19:1069–1076. [PubMed: 20948517]

26. Fox CS, Massaro JM, Hoffmann U, Pou KM, Maurovich-Horvat P, Liu CY, et al. Abdominal visceral and subcutaneous adipose tissue compartments: association with metabolic risk factors in the Framingham Heart Study. *Circulation*. 2007; 116:39–48. [PubMed: 17576866]
27. Gishti O, Gaillard R, Durmus B, Abrahamse M, van der Beek EM, Hofman A, et al. Body mass index, total and abdominal fat distribution and cardiovascular risk factors in school-age children. *Pediatric Research*. 2015; 77:710–718. [PubMed: 25665058]
28. Wright CM, Emmett PM, Ness AR, Reilly JJ, Sherriff A. Tracking of obesity and body fatness through mid-childhood. *Archives of Disease in Childhood*. 2010; 95:612–617. [PubMed: 20522467]
29. Toselli S, Brasili P, Di Michele R. Tracking of weight status and body fatness in Italian children. *Eating and Weight Disorders*. 2013; 18:383–388. [PubMed: 24097345]
30. Vink EE, van Coeverden SC, van Mil EG, Felijs BA, van Leerdam FJ, Delemarre-van de Waal HA. Changes and tracking of fat mass in pubertal girls. *Obesity (Silver Spring)*. 2010; 18:1247–1251. [PubMed: 19875991]
31. Olhager E, Flincke E, Hannerstad U, Forsum E. Studies on human body composition during the first 4 months of life using magnetic resonance imaging and isotope dilution. *Pediatric Research*. 2003; 54:906–912. [PubMed: 12930921]
32. Holzhauser S, Zwijssen RM, Jaddoe VW, Boehm G, Moll HA, Mulder PG, et al. Sonographic assessment of abdominal fat distribution in infancy. *European Journal of Epidemiology*. 2009; 24:521–529. [PubMed: 19639387]
33. WHO Multicentre Growth Reference Study Group. Reliability of anthropometric measurements in the WHO Multicentre Growth Reference Study. *Acta Paediatrica Supplement*. 2006; 450:38–46.
34. Moreno LA, Joyanes M, Mesana MI, Gonzalez-Gross M, Gil CM, Sarria A, et al. Harmonization of anthropometric measurements for a multicenter nutrition survey in Spanish adolescents. *Nutrition*. 2003; 19:481–486. [PubMed: 12781845]
35. Grumbach MM. A window of opportunity: the diagnosis of gonadotropin deficiency in the male infant. *The Journal of Clinical Endocrinology and Metabolism*. 2005; 90:3122–3127. [PubMed: 15728198]

A. 1.5 months



B. 24 months




Figure 1. Associations of subcutaneous fat mass measures at 1.5 or 24 months with risk of overweight at 6 years old^{a-c}

^aValues are odds ratios (95% Confidence Interval) on a log scale and represent the risk of overweight at 6 years per 1-standard-deviation scores increase in body mass index and subcutaneous fat mass measures at 1.5 or 24 months. Body mass index = weight/height². Total subcutaneous fat mass = biceps + triceps + suprailiacal + subscapular skinfold thicknesses. Central-to-total subcutaneous fat mass ratio = (suprailiacal + subscapular skinfold thicknesses)/total subcutaneous fat mass. SDS, standard-deviation scores.

^bModels are adjusted for maternal age, educational level, parity, pre-pregnancy body mass index, smoking habits during pregnancy, and child's gestational age-adjusted birthweight standard-deviation scores, breast feeding duration, timing of introduction of solid foods, and TV watching time.

^cP-value for interaction of child's sex with total subcutaneous fat mass and central-to-total subcutaneous fat mass ratio at 1.5 months = 0.05. P-value for interaction of child's sex with body mass index at 24 months <0.05.

Table 1Characteristics of study participants^a

	Total group (n = 821)	Boys (n = 412)	Girls (n = 409)	P-value
Maternal characteristics				
Age (years), mean (SD)	32.0 (3.9)	31.8 (3.9)	32.1 (3.8)	0.346
Highest completed education, n (%)				
Primary school	10 (1.2)	3 (0.7)	7 (1.7)	0.428
Secondary school	265 (32.3)	135 (32.8)	130 (31.8)	
Higher education	546 (66.5)	274 (66.5)	272 (66.5)	
Parity, n (%) primiparae	522 (63.6)	262 (63.6)	260 (63.6)	0.995
Pre-pregnancy body mass index (kg/m ²), mean (SD)	23.5 (4.0)	23.4 (4.0)	23.7 (4.0)	0.208
Smoking habits during pregnancy, n (%) yes	175 (21.3)	87 (21.1)	88 (21.5)	0.776
Child's characteristics				
Sex, %		412 (50.2)	409 (49.8)	
Birthweight (g), mean (SD)	3533 (522)	3588 (503)	3477 (536)	0.002
Gestational age at birth (weeks), median (95% range)	40.3 (36.3-42.4)	40.3 (36.5-42.4)	40.3 (36.0-42.4)	0.710
Breast feeding duration (months), mean (SD)	4.8 (3.8)	4.7 (3.7)	4.9 (3.9)	0.442
Introduction of solid foods, n (%)				
<3 months	47 (5.7)	23 (5.6)	24 (5.9)	0.885
3 to 6 months	631 (76.9)	316 (76.7)	315 (77.0)	
>6 months	143 (17.4)	73 (17.7)	70 (17.1)	
TV watching time, n (%) 2 hours/day	76 (9.3)	45 (10.9)	31 (7.6)	0.131

^aValues are expressed as means (SD), medians (95% range) or numbers of subjects (%). The values represent the pooled results after multiple imputation. Observed data are given in Supplemental Table S1. SD, standard deviation.

Table 2

Body fat at 1.5 and 24 months by skinfold thicknesses and at 6 years old by dual-energy X-ray absorptiometry and abdominal ultrasound^a

	Total group	Boys	Girls	P-value
1.5 months	n = 742	n = 372	n = 370	
Age (months), mean (SD)	1.6 (0.4)	1.6 (0.4)	1.6 (0.5)	0.340
Body mass index (kg/m ²), mean (SD)	15.1 (1.4)	15.3 (1.5)	15.0 (1.3)	0.001
Total subcutaneous fat mass (mm), mean (SD)	23.9 (7.1)	23.8 (7.0)	24.0 (7.1)	0.705
Central-to-total subcutaneous fat mass ratio, mean (SD)	0.50 (0.05)	0.49 (0.05)	0.51 (0.05)	<0.001
24 months	n = 746	n = 381	n = 365	
Age (months), mean (SD)	25.2 (1.1)	25.3 (1.1)	25.2 (1.1)	0.408
Body mass index (kg/m ²), mean (SD)	15.9 (1.3)	16.0 (1.3)	15.9 (1.3)	0.145
Total subcutaneous fat mass (mm), mean (SD)	27.4 (7.5)	26.7 (7.2)	28.1 (7.7)	0.012
Central-to-total subcutaneous fat mass ratio, mean (SD)	0.43 (0.06)	0.42 (0.06)	0.44 (0.06)	0.001
6 years	n = 821	n = 412	n = 409	
Age (years), mean (SD)	6.0 (0.3)	6.0 (0.3)	6.0 (0.2)	0.450
Body mass index (kg/m ²), mean (SD)	15.9 (1.4)	15.9 (1.3)	15.9 (1.5)	0.955
Overweight and obese (IOTF), n (%)	88 (10.7)	34 (8.3)	54 (13.2)	0.023
Fat mass index (kg/m ³), mean (SD)	3.2 (0.8)	2.9 (0.7)	3.4 (0.9)	<0.001
Central-to-total fat mass ratio, mean (SD)	0.33 (0.04)	0.32 (0.03)	0.34 (0.04)	<0.001
Android-to-gynoid fat mass ratio, mean (SD)	0.24 (0.05)	0.24 (0.05)	0.25 (0.06)	0.089
Preperitoneal fat mass area (cm ²), median (95% range)	0.4 (0.2-0.9)	0.3 (0.2-0.7)	0.4 (0.2-1.0)	<0.001

^aValues are expressed as means (SD), medians (95% range) or numbers of subjects (valid %). Body mass index = weight/height². Total subcutaneous fat mass = biceps + triceps + suprailiacal + subscapular skinfold thicknesses. Central-to-total subcutaneous fat mass ratio = (suprailiacal + subscapular skinfold thicknesses)/total subcutaneous fat mass. Fat mass index = total fat mass/height³. Central-to-total fat mass ratio = trunk fat mass/total fat mass. Android-to-gynoid fat mass ratio = android fat mass/gynoid fat mass. IOTF, International Obesity Task Force. SD, standard deviation.

Table 3

Associations of subcutaneous fat mass measures at 1.5 months with total and abdominal fat mass at 6 years old^{a-c}

Fat mass measures at 1.5 months	Fat mass measures at 6 years in standard-deviation scores Difference [95% Confidence Interval]				
	Body mass index	Fat mass index	Central-to-total fat mass ratio	Android-to-gynoid fat mass ratio	Preperitoneal fat mass area
Total group					
Body mass index	0.16 [0.09,0.24]**	0.05 [-0.03,0.12]	-0.03 [-0.11,0.05]	0.04 [-0.04,0.12]	0.05 [-0.03,0.13]
Total subcutaneous fat mass	0.06 [-0.02,0.13]	0.05 [-0.02,0.13]	0.01 [-0.07,0.09]	0.01 [-0.07,0.09]	0.05 [-0.02,0.12]
Central-to-total subcutaneous fat mass ratio	0.04 [-0.02,0.12]	0.12 [0.05,0.19]**	0.11 [0.04,0.18]**	0.07 [0.00,0.14]	0.07 [-0.01,0.14]
Boys					
Body mass index	0.25 [0.14,0.36]**	0.21 [0.09,0.33]**	0.10 [-0.02,0.22]	0.13 [0.02,0.25]*	0.20 [0.07,0.33]**
Total subcutaneous fat mass	0.12 [0.01,0.23]*	0.07 [-0.04,0.19]	0.09 [-0.03,0.20]	0.03 [-0.08,0.14]	0.14 [0.02,0.26]*
Central-to-total subcutaneous fat mass ratio	0.03 [-0.08,0.13]	0.07 [-0.03,0.18]	0.06 [-0.05,0.16]	0.07 [-0.04,0.17]	0.01 [-0.11,0.13]
Girls					
Body mass index	0.08 [-0.02,0.18]	0.02 [-0.08,0.12]	-0.06 [-0.16,0.05]	-0.03 [-0.14,0.08]	0.02 [-0.09,0.12]
Total subcutaneous fat mass	-0.01 [-0.11,0.09]	0.01 [-0.10,0.11]	-0.08 [-0.19,0.03]	-0.02 [-0.13,0.09]	-0.03 [-0.13,0.06]
Central-to-total subcutaneous fat mass ratio	0.08 [-0.01,0.18]	0.10 [0.01,0.20]*	0.09 [-0.01,0.19]	0.06 [-0.04,0.17]	0.06 [-0.03,0.16]

^a Values are standardized regression coefficients [95% confidence interval] and represent the difference in standard-deviation scores for fat mass measures at 6 years per 1-standard-deviation scores increase in body mass index and subcutaneous fat mass measures at 1.5 months. Body mass index = weight/height². Total subcutaneous fat mass = biceps + triceps + suprailiacal + subscapular skinfold thicknesses. Central-to-total subcutaneous fat mass ratio = (suprailiacal + subscapular skinfold thicknesses)/total subcutaneous fat mass. Fat mass index = total fat mass/height³. Central-to-total fat mass ratio = trunk fat mass/total fat mass. Android-to-gynoid fat mass ratio = android fat mass/gynoid fat mass.

^b Models are adjusted for maternal age, educational level, parity, pre-pregnancy body mass index, smoking habits during pregnancy, and child's gestational age-adjusted birthweight standard-deviation scores, breast feeding duration, timing of introduction of solid foods, and TV watching time.

^c P-value for interaction of child's sex with fat mass measures at 1.5 months >0.05.

* P-value<0.05

** P-value<0.01.

Table 4

Associations of subcutaneous fat mass measures at 24 months with total and abdominal fat mass at 6 years old
a-b

Fat mass measures at 24 months	Fat mass measures at 6 years in standard-deviation scores Difference [95% Confidence Interval]				
	Body mass index	Fat mass index	Central-to-total fat mass ratio	Android-to-gynoid fat mass ratio	Preperitoneal fat mass area
Total group					
Body mass index	0.48 [0.42,0.54] ^{**†}	0.33 [0.26,0.40] ^{**†}	0.18 [0.11,0.26] ^{**†}	0.16 [0.08,0.23] ^{**†}	0.07 [0.00,0.14] [†]
Total subcutaneous fat mass	0.30 [0.23,0.37] ^{**}	0.36 [0.29,0.43] ^{**†}	0.28 [0.20,0.35] ^{**}	0.23 [0.16,0.31] ^{**†}	0.19 [0.12,0.26] ^{**}
Central-to-total subcutaneous fat mass ratio	0.09 [0.01,0.16] ^{*†}	0.14 [0.06,0.22] ^{**†}	0.12 [0.04,0.20] ^{**†}	0.10 [0.03,0.18] ^{**†}	0.10 [0.02,0.17] [*]
Boys					
Body mass index	0.46 [0.36,0.55] ^{**}	0.35 [0.25,0.45] ^{**}	0.20 [0.09,0.30] ^{**}	0.10 [-0.01,0.21]	0.01 [-0.12,0.13]
Total subcutaneous fat mass	0.26 [0.15,0.36] ^{**}	0.33 [0.22,0.43] ^{**}	0.22 [0.11,0.32] ^{**}	0.12 [0.01,0.23] [*]	0.14 [0.01,0.27] [*]
Central-to-total subcutaneous fat mass ratio	-0.01 [-0.12,0.10]	-0.02 [-0.13,0.10]	0.00 [-0.11,0.11]	0.01 [-0.10,0.12]	0.11 [-0.02,0.24]
Girls					
Body mass index	0.50 [0.41,0.59] ^{**}	0.43 [0.33,0.52] ^{**}	0.25 [0.14,0.36] ^{**}	0.22 [0.11,0.32] ^{**}	0.15 [0.05,0.25] ^{**}
Total subcutaneous fat mass	0.34 [0.24,0.44] ^{**}	0.38 [0.28,0.47] ^{**}	0.30 [0.20,0.41] ^{**}	0.31 [0.20,0.41] ^{**}	0.21 [0.12,0.29] ^{**}
Central-to-total subcutaneous fat mass ratio	0.15 [0.05,0.26] ^{**}	0.18 [0.07,0.28] ^{**}	0.14 [0.04,0.25] ^{**}	0.14 [0.03,0.25] [*]	0.05 [-0.05,0.15]

^aValues are standardized regression coefficients [95% confidence interval] and represent the difference in standard-deviation scores for fat mass measures at 6 years per 1-standard-deviation scores increase in body mass index and subcutaneous fat mass measures at 24 months. Body mass index = weight/height². Total subcutaneous fat mass = biceps + triceps + suprailiacal + subscapular skinfold thicknesses. Central-to-total subcutaneous fat mass ratio = (suprailiacal + subscapular skinfold thicknesses)/total subcutaneous fat mass. Fat mass index = total fat mass/height³. Central-to-total fat mass ratio = trunk fat mass/total fat mass. Android-to-gynoid fat mass ratio = android fat mass/gynoid fat mass.

^bModels are adjusted for maternal age, educational level, parity, pre-pregnancy body mass index, smoking habits during pregnancy, and child's gestational age-adjusted birthweight standard-deviation scores, breast feeding duration, timing of introduction of solid foods, and TV watching time.

[†]P-value for interaction of child's sex with fat mass measures at 24 months < 0.05.

* P-value < 0.05;

** P-value < 0.01.