

Host Genome Influence on Gut Microbial Composition and Microbial Prediction of Complex Traits in Pigs

Amelia Camarinha-Silva,^{*,1} Maria Maushammer,^{*,1} Robin Wellmann,^{*} Marius Vital,[†] Siegfried Preuss,^{*} and Jörn Bennewitz^{*,2}

^{*}Institute of Animal Science, University of Hohenheim, 70599 Stuttgart, Germany and [†]Microbial Interactions and Processes Research Group, Helmholtz Centre for Infection Research, 38124 Braunschweig, Germany

ABSTRACT The aim of the present study was to analyze the interplay between gastrointestinal tract (GIT) microbiota, host genetics, and complex traits in pigs using extended quantitative-genetic methods. The study design consisted of 207 pigs that were housed and slaughtered under standardized conditions, and phenotyped for daily gain, feed intake, and feed conversion rate. The pigs were genotyped with a standard 60 K SNP chip. The GIT microbiota composition was analyzed by 16S rRNA gene amplicon sequencing technology. Eight from 49 investigated bacteria genera showed a significant narrow sense host heritability, ranging from 0.32 to 0.57. Microbial mixed linear models were applied to estimate the microbiota variance for each complex trait. The fraction of phenotypic variance explained by the microbial variance was 0.28, 0.21, and 0.16 for daily gain, feed conversion, and feed intake, respectively. The SNP data and the microbiota composition were used to predict the complex traits using genomic best linear unbiased prediction (G-BLUP) and microbial best linear unbiased prediction (M-BLUP) methods, respectively. The prediction accuracies of G-BLUP were 0.35, 0.23, and 0.20 for daily gain, feed conversion, and feed intake, respectively. The corresponding prediction accuracies of M-BLUP were 0.41, 0.33, and 0.33. Thus, in addition to SNP data, microbiota abundances are an informative source of complex trait predictions. Since the pig is a well-suited animal for modeling the human digestive tract, M-BLUP, in addition to G-BLUP, might be beneficial for predicting human predispositions to some diseases, and, consequently, for preventative and personalized medicine.

KEYWORDS GenPred; shared data resource; complex traits; genomic selection; gut microbial composition; microbial prediction; pig

HOST–MICROBIOTA interactions have received considerable attention in human studies in recent years, and it has been shown repeatedly that host genetics, as well as the environment, affect gut microbiota composition (Spor *et al.* 2011). In order to unravel host genetic effects, the use of a model organism that is kept in a stable environment with minimum environmental variations is needed (Zhao *et al.* 2013). Kostic *et al.* (2013) described different model organisms, such as mice, zebrafish, the fruit fly *Drosophila*, and the bobtail squid, as important sources of information on host–microbiota homeostasis. The pig can be used as a model for human-related research, as the human and porcine

physiology, metabolism, and gastrointestinal tract (GIT) microbiota are similar (Heinritz *et al.* 2013). It has already been used as an animal model for microbiota-associated diseases such as *Helicobacter pylori* infections, necrotizing enterocolitis disease, obesity, and diabetes, and to formulate dietary strategies for overcoming obesity and other metabolic syndromes (Heinritz *et al.* 2013). Moreover, pigs are not only suitable model organisms for human-related research but also some of the most important livestock species used for meat production worldwide. Breeding is frequently carried out using genome-wide SNP data for the prediction of selection candidate breeding values (Knol *et al.* 2016)—a technique that is recognized as a form of genomic selection (Meuwissen *et al.* 2001). The underlying assumption is that each SNP affects complex traits of interest only marginally, but modeling all SNPs jointly in a prediction equation is attributed with remarkably high levels of prediction accuracy. Genomic prediction is also utilized in plant breeding (Jannink *et al.* 2010), and has been proposed as a tool for predicting complex genetic predispositions in humans

Copyright © 2017 by the Genetics Society of America
doi: <https://doi.org/10.1534/genetics.117.200782>

Manuscript received February 2, 2017; accepted for publication April 27, 2017; published Early Online May 2, 2017.

Supplemental material is available online at www.genetics.org/lookup/suppl/doi:10.1534/genetics.117.200782/-/DC1.

¹These authors contributed equally to this work.

²Corresponding author: Institute of Animal Science, University of Hohenheim, Garbenstrasse 17, 70599 Stuttgart, Germany. E-mail: j.bennewitz@uni-hohenheim.de

(de los Campos *et al.* 2010). Inspired by the success of genomic prediction methods, Ross *et al.* (2013) developed a metagenomic prediction approach. These authors used human and cattle datasets with massive parallel sequencing data to form metagenomic relationship matrices, which in turn were used to predict complex traits with considerable accuracy. This study clearly highlights the potential to consider alternative high-dimensional host-related explanatory variables beyond SNP markers for prediction purposes, and, more generally, opportunities to apply quantitative-genetic methods in holistic analyses of data on microbiota, host genetics, and complex traits.

Currently, there is a lack of research in pigs regarding the genetic influences on gut microbial community, and the effect of this community on complex host traits. Besides studies on the influence of nutrition and medication on the microbiota, one study has approached the question of how the early-life pig gut microbiota impacts host phenotypes (Mach *et al.* 2015).

In this study, standard quantitative-genetic methods were extended and applied to analyze the interrelationship between pig GIT microbiota compositions, complex traits, and pig genomes. The specific aims were (i) to characterize GIT microbiota for pigs of a mature age, (ii) to analyze the effects of host genetics on GIT microbial composition, (iii) to investigate the role of GIT microbial composition on key host complex traits, and (iv) to evaluate genomic as well as microbial predictions of complex host traits.

Materials and Methods

Sample collection

The animal experiments were performed in accordance with German Animal Welfare legislation. All procedures regarding animal handling and treatment were approved by the University of Hohenheim Committee of Animal Welfare under authorization number S411/14TZ. The pigs belonged to the Piétrain breed. This is an important sire line breed (Stratz *et al.* 2014) for which genomic selection is practiced (Wellmann *et al.* 2013). Housing, slaughtering, and the recording of phenotypes of the pigs was performed under standardized conditions at one experimental farm. Performance testing started with a weight of 30 kg and ended with a weight of 105 kg. Animal feed intake (FI) and daily gain (DG) values were recorded during performance testing. The feed conversion (FC) value was calculated as a ratio of the consumed feed and weight gain occurring during performance testing. See Supplemental Material, Table S1 for descriptive statistics of these traits. By reaching 105 kg, the pigs were slaughtered with an average slaughter age of 188 (± 14) days, and an average performance testing duration of 100 (± 11) days. In total, colon and blood samples from 207 Piétrain sows were collected on 14 slaughter days. Blood samples were taken directly during the slaughtering and stored on ice. After opening of the abdomen, colon samples were collected from

the mid-colon and also stored on ice. For long-term storage, blood samples were kept at -20° and colon samples at -80° .

Illumina amplicon sequencing

Colon digesta samples were thawed on ice and homogenized, and 250 mg of each sample was used to extract DNA using the FastDNA SPIN Kit for Soil (MP Biomedicals, Solon, OH) according to the manufacturer's instructions. PCR targeting of the V1-2 region of the 16S rRNA gene was carried out as described in Camarinha-Silva *et al.* (2014). Based on a previous work of our group (Burbach *et al.* 2016), this 16S region and DNA extraction method was giving the best coverage of the microbial community. Amplicons were purified and normalized using a SequalPrep Normalization Kit (Invitrogen, Carlsbad, CA), and were pooled and sequenced with 250 bp paired-end sequencing chemistry applied on an Illumina MiSeq platform.

Bioinformatic processing of sequences was done according to Camarinha-Silva *et al.* (2014) with some modifications. Raw reads were assembled (Cole *et al.* 2014) and subsequently aligned using MOTHUR (gotoh algorithm with the SILVA reference database) prior to preclustering of the sequences with two mismatches (diffs = 2). Low abundance operational taxonomic units (OTUs), if present in <5 samples in relative abundances <0.01%, were removed. Finally, $40,379 \pm 1149$ sequences were obtained per sample, comprising a total of 2714 OTUs that were taxonomically assigned using the naïve Bayesian RDP classifier (Wang *et al.* 2007). The OTUs were then evaluated against the RDP database using Seqmatch function, which belongs to the RDP database. Note that no differences were observed in the microbiota regarding the day of DNA extraction, which was performed on 5 consecutive days by the same person.

Genotyping

In total, 207 German Piétrain sows were genotyped using an Illumina PorcineSNP60 BeadChip (Ramos *et al.* 2009). Genotypes from individuals were filtered with respect to call rates (removal of SNPs with a call rate of <95%), minor allele frequencies (exclusion of SNP with a minor allele frequency of <5%), significant deviations from Hardy-Weinberg equilibrium ($P < 0.0001$), and SNPs on the Y-chromosome were removed. The call rate across all animals was ≥ 0.994 . After quality control measures were performed, 51,970 SNPs remained for further analysis.

Statistical analysis

Microbial community: A multivariate dataset comprising the relative abundance of each phylotype across each sample was analyzed using v.6.1.6, PRIMER-E (Plymouth Marine Laboratory, UK; Clarke and Warwick 2001). The Bray-Curtis coefficient (Bray and Curtis 1957) was used to create a sample-similarity matrix, and microbial community structures were explored via nonmetric multidimensional scaling (MDS) (Clarke and Warwick 2001). Statistical comparisons between *a priori* defined groups (*e.g.*, weight, age, DG, and FC) were drawn

using analysis of similarity (ANOSIM) based on 999 permutations, and were considered significantly different at a P -value of < 0.05 . Species responsible for observed differences were identified based on similarity percentages (SIMPER) (Clarke and Warwick 2001). Tax4fun (Aßhauer *et al.* 2015) was used to predict the functional capabilities of microbial communities detected in colons based on the 16S rRNA sequencing data.

Genetic parameters of bacterial genera: Genetic parameters were estimated for bacterial genera rather than for OTUs, because the latter would result in numerous statistical analysis, and, thus, in a strong multiple testing problem. An analysis of variance was performed by fitting a univariate genomic mixed linear model to test for significant effects. The age and weight measured at the test station and the slaughter weight were included as fixed covariables in the model, and the slaughter day (SD) was considered as a random effect in the model. The observation vector included the relative abundance of one genus. Only genera with abundance values exceeding 0.1% were considered. By backward elimination, nonsignificant effects (significance level for the elimination of $\alpha = 0.05$) were removed from the model. An univariate analysis was performed to estimate the heritability of each genus. Statistical analyses were performed using the ASReml package available through R (Butler *et al.* 2009). The mixed linear model is written as follows:

$$y = X_b \mathbf{b} + Z_{SD} \mathbf{SD} + \mathbf{a} + \mathbf{e}, \quad (1)$$

where \mathbf{y} is the observation vector, which includes the abundance of one genus, \mathbf{b} is the vector of fixed effects (described above), \mathbf{SD} is the vector with random slaughter day effects with variance σ_{SD}^2 , \mathbf{a} is a vector with random animal genetic effects, X_b and Z_{SD} are the corresponding design matrices, and \mathbf{e} is the residual term with residual variance σ_e^2 . Note that a random pen effect as well as a random maternal effect were not significant, and thus were not included in this model. The distribution of the random animal effect is $\mathbf{a} \sim N(0, G\sigma_A^2)$, with G being the genomic relationship matrix, and σ_A^2 being the additive genetic variance. The G matrix was estimated using SNP genotypes following VanRaden (2008) as

$$G = \frac{(Z - 2Q)(Z - 2Q)^T}{\sum_m 2p_m(1 - p_m)},$$

where Z is the gene content matrix with entries 0, 1, or 2 for each SNP and each animal, and matrix Q contains the frequency p_m of each SNP m . Narrow-sense heritability was estimated as $h^2 = \sigma_A^2 / \sigma_P^2$, with $\sigma_P^2 = \sigma_{SD}^2 + \sigma_A^2 + \sigma_e^2$. The p -values of the heritability estimates were calculated by conducting a likelihood-ratio test on random animal effects. The null-hypothesis, that the variance of the random effect is 0, was rejected if twice the difference in the log-likelihoods of the full model, and the reduced model without the random

effect was larger than the 0.95-quantile of a χ^2 -distribution with 1 d.f. We chose a significance level of P -value = 0.05. A total of 49 genera was analyzed, which resulted in a multiple testing problem. In order to judge how many false positives were among the significant results, we applied the false discovery rate (FDR) technique. We calculated for each test an FDR q -value using the software QVALUE (Storey and Tibshirani 2003). The FDR q -value of the significant genera with the lowest test statistic (P -value ≈ 0.05) provided an estimate of the proportion of false positives among the significant outcomes. Note that model (1) is a mixed linear model, which assumes normality of the data. The relative abundance of the genera were, in general, not normally distributed, and sometimes peaked at zero. However, due to the small data set we did not apply generalized linear mixed models.

Genetic and microbial parameters of host traits: First, explanatory variables for host traits FC, DG and FI were estimated via backward elimination, using ages and weights measured upon test station arrival and slaughter weights as fixed covariables, and SD and pen as random effects. Nonsignificant effects of $\alpha = 0.05$ were excluded. To estimate genetic variance components and narrow sense heritabilities of the host traits, the following model was applied

$$y = X_b \mathbf{b} + Z_{SD} \mathbf{SD} + Z_{pen} \mathbf{pen} + \mathbf{a} + \mathbf{e}, \quad (2)$$

where \mathbf{y} is a vector of observations (FC, DG, or FI), \mathbf{b} is a vector of fixed effects (*i.e.*, for FC and FI weights measured upon test station arrival, and for DG slaughter weights), and \mathbf{SD} and \mathbf{pen} are vectors of random slaughter day, and pen effects, respectively, with variance components σ_{SD}^2 and σ_{pen}^2 . X_b , Z_{SD} , Z_{pen} are corresponding design matrices, and \mathbf{e} denotes the residual term. The distribution of the random animal effect \mathbf{a} is the same as described in model (1). The heritability was estimated in the same manner as shown in model (1) for the bacteria genera. The P -value of the additive genetic variance was calculated by performing a likelihood ratio test of the animals' random effects, as described for the random effect in model (1).

The microbial variance component was estimated by applying the following univariate microbial mixed linear model (fitted in ASReml R):

$$y = X_b \mathbf{b} + Z_{SD} \mathbf{SD} + Z_{pen} \mathbf{pen} + \mathbf{m} + \mathbf{e}, \quad (3)$$

where the model parameters are as described in model (2) except vector \mathbf{m} , which contains the random effect of the animal microbiota for each individual with $\mathbf{m} \sim N(0, M\sigma_m^2)$, where σ_m^2 is the microbial variance. The microbial relationship matrix M was calculated as follows: we have $M = 1/NXX^T$, with matrix X (dimension $n \times N$ where n is the number of animals and N is the number of OTUs), constructed from matrix P (dimension $n \times N$). The elements P_{ik} are the relative abundance of OTU k in animal i (plus 0.01). Following this, the elements in X are




Figure 1 Nonmetric multidimensional scaling (nMDS) plot illustrating similarities in the global bacterial community structure of pig colon digesta. The pigs are colonized with higher abundances of Firmicutes (●) and Bacteroidetes (●). While a two-dimensional (2D) stress value of 0.2 denotes some stress on the plot, this is considered acceptable since 207 samples are ordinated together.

$$X_{ik} = \frac{\log P_{ik} - \overline{\log P_{\cdot k}}}{sd(\log P_{\cdot k})}$$

Thus, the off-diagonals in **M** are calculated as $M_{ij} = 1/N \sum_{k=1}^N X_{ik} X_{jk}$, and the diagonals as

$$M_{ii} = \frac{1}{N} \sum_{k=1}^N X_{ik}^2 = \frac{1}{N} \sum_{k=1}^N \frac{(\log P_{ik} - \overline{\log P_{\cdot k}})^2}{var(\log P_{\cdot k})}$$

Note, that it might happen that the diagonals are >1 . This is the case if

$$|\log P_{ik} - \overline{\log P_{\cdot k}}| > sd(\log P_{\cdot k}),$$

i.e., $\log P_{ik} < \overline{\log P_{\cdot k}} - sd(\log P_{\cdot k})$ or $\log P_{ik} > \overline{\log P_{\cdot k}} + sd(\log P_{\cdot k})$,

which means that the abundances of the OTUs in animal *i* deviate strongly from the average values, *e.g.*, if OTUs are missing in the animal that are present in most other animals, and if OTUs are present that are rare in other animals. Note also that the microbial relationships are affected by some errors remaining in the OTU data. More research is needed regarding the effect of data screening on the precision of the microbial relationship estimation.

The fraction of the phenotypic variance explained by the microbial variance was calculated as $m^2 = \sigma_m^2 / \sigma_p^2$, where $\sigma_p^2 = \sigma_m^2 + \sigma_{SD}^2 + \sigma_{pen}^2 + \sigma_e^2$ is the phenotypic variance. This fraction was termed microbiability by Difford *et al.* (2016). The *P*-value of the microbial variance was calculated by performing a likelihood ratio test of the animals' random microbiota effects, as described for the random animal effect in model (1).

Genomic and microbial prediction: To predict host traits FC, DG, and FI using genomic and microbiota data, G-BLUP and




Figure 2 Relative abundance of the most abundant genera of the Bacteroidetes (first heatmap row) and Firmicutes groups (second heatmap row). Groups with an asterisk (*) are significantly different ($P < 0.05$).

M-BLUP models were applied, respectively. For the G-BLUP predictions (VanRaden 2008), model (2) was applied with previously estimated variance components. In a similar vein, for the M-BLUP predictions, model (3) was applied. For both types of predictions, a repeated cross validation was performed with 10,000 iterations, where 80% of the individuals were sampled randomly without replacement to train the prediction model (reference population). The pig trait phenotypes (FC, DG, and FI) were predicted from the remaining 20% of the pigs based on the results of the reference population analysis (validation population). The accuracy of a prediction was defined as the correlation between predicted and observed trait phenotypes in the validation population. The mean correlation was calculated as the mean of 10,000 correlation estimates. Confidence intervals of correlations were estimated as 2.5 and 97.5% quantiles of the 10,000 ordered correlation estimates.

Effects of single OTUs on complex host traits: To identify the drivers of prediction accuracy levels, the marginal effects of OTUs on phenotypic traits (*i.e.*, single OTU effects not captured by the remaining OTUs) were estimated from the solutions of the M-BLUP model. To do this, an adapted version of the back solving method proposed by Strandén and Garrick (2009) was used, which is described in the supplementary information (File S1).

Table 1 Estimated heritability (h^2) and P -value for the relative abundances of bacterial genera

Bacteria	h^2	SE	P -value ^a
<i>Alloprevotella</i>	0.34	0.16	0.01
<i>Blautia</i>	0.33	0.14	<0.01
<i>Catenibacterium</i>	0.39	0.16	0.01
<i>Lactobacillus</i>	0.34	0.16	0.02
Uncultured <i>Spirochaetales</i>	0.52	0.15	<0.01
Uncultured <i>Spirochaetes</i>	0.32	0.17	0.01
Uncultured <i>Succinivibrionaceae</i>	0.57	0.14	<0.01
Uncultured <i>Veillonellaceae</i>	0.33	0.15	0.01

^a All p -values showed a FDR < 0.12.

Data availability

The authors state that all data necessary for confirming the conclusions presented in the article are fully represented within the tables and figures. Sequences are available at the European Nucleotide Archive (ENA) under accession number PRJEB18070 (<http://www.ebi.ac.uk/ena/data/view/PRJEB18070>). File S2 and File S3 contain genotypes and phenotypes for each individual, respectively. File S4 contains relative abundances at OTU level.

Results

Microbial community characterization and heritability of gut microbiota compositions

The main phyla that account for higher abundances were found to be Firmicutes (54%), Bacteroidetes (42%), Proteobacteria (2%), and Spirochaetes (1%). The most abundant families found in colons were Ruminococcaceae (24%), Prevotellaceae (21%), Porphyromonadaceae (8%), Clostridiaceae 1 (6%), Rikenellaceae (6%), and Lachnospiraceae (5%), with all other families found to be present at average levels of <5%. The microbial community harboring the different pigs showed a similarity of 35% between animals. Overall, the predicted KEGG pathways of higher abundance were found to be related to pathways associated with carbohydrate metabolism, such as starch and sucrose pyruvate, fructose and mannose metabolism; amino acid metabolism; environmental information processing such as membrane transport and signal transduction; genetic information processing such as replication and repair; and translation (Figure S1).

In Figure 1, a nonmetric multidimensional scaling plot is provided to illustrate similarities in the global bacterial community structure of pig colon digesta. The stress value associated with this plot measures the difficulty involved in compressing the samples relationship into two dimensions. It was 0.22, indicating some stress, but we consider this as acceptable, because larger data sets (as in our case) general result in larger stress values. A significant difference was observed between two groups of animals in this plot: one colonized at higher levels with Firmicutes (F), and another colonized with Bacteroidetes (B) ($R = 0.339$, P -value = 0.001). The groups showed an average dissimilarity value of 68%,

Table 2 Estimated microbiability (m^2) and heritability (h^2) with SE and P -values for DG, FC, and feed intake

Trait	m^2	SE	P -value	h^2	SE	P -value
DG	0.28	0.13	0.01	0.42	0.14	<0.01
FC	0.21	0.14	0.01	0.19	0.13	0.08
FI	0.16	0.10	0.03	0.11	0.11	0.22

$m^2 = \sigma_m^2 / \sigma_p^2$, as defined by Difford *et al.* (2016).

where the average similarity level between all samples in group B was found to be 33%, and that for group F was 38%. Microorganisms contributing to this separation belonged to the genera *Clostridium sensu stricto* ($B = 20.7\%$, $F = 12.8\%$), *Lactobacillus* ($B = 1.3\%$, $F = 5.2\%$), *Prevotella* ($B = 5.3\%$, $F = 2.7\%$), and *Clostridium XI* ($B = 1.9\%$, $F = 5.8\%$) (Figure 2). Additional groups of microorganisms presenting significant differences between both groups are shown in Figure 2. The reasons for the separation into these two groups could not be identified and might also be due to different conditions at birth and weaning period, which were unknown to us. Despite the different colon colonization patterns found, no effect of the Firmicutes/Bacteroidetes ratio was observed for DG, FC, or FI (Figure S2). Eight genera generated significant heritability estimates (P -value < 0.05), which are shown in Table 1. The FDR of these significant results was <0.11. Heritability estimates of all 49 bacterial genera are shown in Table S2.

Heritability and microbiability of host traits and prediction results

The heritability and microbiability of the DG, FC, and FI traits are shown in Table 2. The heritability estimates are likely slightly underestimated, due to the use of SNP chip data instead of pedigree data. However, they are within a typical range for complex pig traits measured under standardized conditions. The heritability and microbiability estimates were significant with the exception of the heritabilities of FC and FI. For these traits, the microbiability estimates were higher than the heritability estimates. The microbial relationship matrix **M**, and genomic relationship matrix **G**, underlying these calculations are shown as heatmaps in Figure S3 and Figure S4, respectively. The mean of the diagonal values of **G** was 0.99, and the values ranged from 0.85 to 1.19. The mean of the diagonal values of **M** was 0.995, and the values ranged from 0.66 to 1.97.

The results of the genomic and microbial predictions are shown in Table 3. The microbial prediction generated an accuracy of 0.41 for DG, and 0.33 for FC and FI. These prediction accuracies were higher than those obtained from genomic predictions. In addition, confidence intervals showed that the accuracies were significantly >0 (as was twice the case for genomic predictions).

A plot of marginal OTU effects is shown in Figure 3. Some outlier effects were detected (Table 4), but none of the marginal OTU effects showed substantial effects. For DG, the outliers were assigned to uncultured *Veillonellaceae*, uncultured *Prevotellaceae* and uncultured *Proteobacteria*. For FC two OTUs with outlier effects were detected, which were

Table 3 Accuracy of microbial (r_m) and genomic predictions (r_g) of DG, FC, and FI, with C.I.

Trait	Microbial Prediction		Genomic Prediction	
	r_m	97.5% CI	r_g	97.5% CI
DG	0.41	0.18:0.62	0.35	0.08:0.58
FC	0.33	0.07:0.54	0.23	-0.04:0.48
FI	0.33	0.15:0.51	0.20	-0.08:0.46

assigned to uncultured *Bacteroidales* and uncultured *Clostridiales*. One outlier of OTU effects was found for FI, and was assigned to uncultured *Clostridiales*.

Discussion

In this study, we analyzed the microbial composition of pig colon samples. The general pattern of microbiota composition is in agreement with reports on colon (Looft *et al.* 2014; Kim and Isaacson 2015). No interrelationships were found between the Firmicutes-Bacteroidetes ratio and DG, FC, or FI, as previously shown by Mach *et al.* (2015). In former studies conducted in mice and humans, this ratio was considered as an important marker for obesity (Ley *et al.* 2005, 2006; Turnbaugh *et al.* 2006). However, other studies showed contradictory results, or even no evidence of a possible effect in human obesity (Duncan *et al.* 2008; Schwartz *et al.* 2010; Jumpertz *et al.* 2011).

The host genetic variance on the GIT microbiota composition was substantial for some bacterial genera as denoted by the heritability estimates (Table 1). This result is in agreement with earlier findings of Estellè *et al.* (2014) and O'Connor *et al.* (2014), who reported similar heritabilities for *Blautia* and *Lactobacillus* in a French Large White pig population, and in a segregated mouse population, respectively. The underlying mechanism of this host genetic determination remains largely unknown thus far. In general, host genetics can influence microbiota compositions through differences in immunoglobulin and antibacterial molecules secreted into gut lumen (Wen *et al.* 2008; Vijay-Kumar *et al.* 2010; Shulzhenko *et al.* 2011), owing to differences in mucosal gut structures (Sommer *et al.* 2014; Wlodarska *et al.* 2014) and bile acid metabolism (Ryan *et al.* 2014). Genome-wide association studies (GWAS) may help identify host genes affecting microbiota compositions, and, thus, derive and substantiate novel hypotheses on the genetic mechanism underlying the heritability of microbiota compositions. However, this involves the use of large datasets, and was therefore not possible in this study.

Microbiability as first defined by Difford *et al.* (2016) allows for a holistic view of the influence of microbiota on host traits. For all three investigated traits, microbiability levels were found to be significant, and, for FC and FI, microbiability estimates were higher than the heritability estimates (Table 3). This points to a strong effect of GIT microbiota compositions on these traits. As microbiota are partly under the control of host genes, from an animal breeder's perspective they can be




Figure 3 OTU marginal effects (absolute values) of each OTU for each trait. Outliers are marked in red.

viewed as host traits. This highlights the possibility of breeding for optimized microbiota to indirectly improve complex host traits. Indeed, from the heritabilities shown in Table 1, selection responses can be expected for at least eight (from a total of 49) bacterial genera. This targeted breeding strategy might be especially beneficial for important and so-called hard-to-measure traits, for which precise data collection is restricted to few individuals. Examples include the utilization of certain nutrients in monogastric animals (Beck *et al.* 2016), and greenhouse gas emissions in ruminants (Hayes *et al.* 2013; Roehe *et al.* 2016).

The microbial prediction method M-BLUP is closely related to the well-known G-BLUP model, which is widely used in animal breeding. The key difference is that relationships between individuals are modeled based on relative microbiota abundances at the OTU level, for which calculations are straightforward. The M-BLUP predictions outperformed the G-BLUP predictions in terms of prediction accuracy levels (Table 3). This further underscored the importance of microbiota compositions for trait variability. Thus, it seems that, in addition to SNP data, microbiota abundances are an informative source of complex trait prediction data for pigs, which again may be of special interest for hard-to-measure traits. To identify drivers of prediction accuracy levels, we estimate the marginal effects of a single OTU. Since only few outliers were detected (Table 4), it can be tentatively concluded that many of the OTUs explain a small fraction of the trait variability and that such traits, like DG, FC, and FI, are not only polygenic in nature (Wellmann *et al.* 2013), but also highly polymicrobial determined in this pig population. However, this must be investigated further because this putative polymicrobial trait determination might, at least in part, also be attributable to M-BLUP model assumptions.

Table 4 Outliers of marginal OTU effect estimates (\hat{u}) with designations and average abundances for DG, FC, and FI

OTU	Trait	Designation	Average Abundance	\hat{u}
1284	DG	<i>Unc. Veillonellaceae</i>	0.017	-0.009
1459	DG	<i>Unc. Prevotellaceae</i>	<0.001	-0.008
4300	DG	<i>Unc. Proteobacteria</i>	0.002	-0.008
901	FC	<i>Unc. Bacteroidales</i>	0.029	-0.006
2525	FC	<i>Unc. Clostridiales</i>	0.008	0.008
2812	FI	<i>Unc. Clostridiales</i>	0.007	0.006

One limit of the microbial prediction model pertains to the fact that the GIT microbial composition for pigs is itself not constant, but changes from birth to adulthood (Kim *et al.* 2011; Pajarillo *et al.* 2014; Mach *et al.* 2015), and is, of course, affected by environmental conditions. In the current study, these effects were minimized by housing pigs under standardized conditions, and by collecting microbiota data on pigs of the same age, which may have additionally contributed to the high level of microbial prediction accuracy achieved. However, it remains to be determined whether GIT microbial compositions at a juvenile stage can be used as predictors of complex host traits measured at a mature age. Further, it is likely that microbial compositions collected from different locations of the GIT, or from feces, will show various complex trait predictive capacities.

The two microbiota pig breeding strategies described above (*i.e.*, breeding for an optimized microbiome and applying microbial prediction) define the microbiota composition differently. For the former, it is treated as a quantitative host trait, whereas, for the latter, it is used as an explanatory variable for prediction purposes. Detailed investigations show that not all microbiota genera are heritable host traits, and not all microbiota OTUs are equally important for predictions (Figure 3). Hence, a comparatively detailed analysis of these two components is desirable, but larger datasets must be used. In addition to GWAS of microbiota compositions and complex host traits, this interplay may be analyzed through structural equation models as introduced to the field of livestock genetics by Gianola and Sorensen (2004). In a quantitative genetic setting, structural equation models allow for the separation of direct and indirect genetic effects shaping genetic relationships among traits. Direct genetic effects result from linkage disequilibrium between genes affecting traits or from pleiotropic effects. However, when a causal relationship between two traits exists, genes directly affecting only one trait may also affect the second trait indirectly via the causal relationship between the traits. Methods for identifying causal structures (Valente *et al.* 2010) would help simultaneously identify which microbiota bacteria present host genetic variance and the impact of these bacteria on complex host trait variations.

This study was conducted using pig samples. Since the pig is an animal well-suited for modeling the human digestive tract, our results may have implications for predicting human predisposition to disease, and consequently for preventative

and personalized medicine. For genetically determined traits such as type-2 diabetes, G-BLUP has already been proven to be useful (de los Campos *et al.* 2013). This study extends the scope of predictions toward using microbiota data, which might be of special interest for traits where it is known that the microbiome plays an important role, *e.g.*, metabolic diseases and obesity (Karlsson *et al.* 2013; Le Chatelier *et al.* 2013). Applications of M-BLUP with appropriate microbiota data may help quantify the risks of suffering from such diseases, and may further the development of personalized preventative strategies.

Methods that combine highly dimensional and correlated predictors (G-BLUP and M-BLUP) with cumulative prediction power will have to be developed in future studies. For this purpose, the present data set is too small.

Acknowledgments

We thank Dr. Jörg Heinkel and the animal care staff at the experimental farm of the Landesanstalt für Schweinezucht in Boxberg, Germany, for assisting with animal sampling procedures. The authors declare no conflicts of interest.

Literature Cited

- Aßhauer, K. P., B. Wemheuer, R. Daniel, and P. Meinicke, 2015 Tax4Fun: predicting functional profiles from metagenomic 16S rRNA data. *Bioinformatics* 31: 2882–2884.
- Beck, P., H. P. Piepho, M. Rodehutschord, and J. Bennewitz, 2016 Inferring relationships between Phosphorus utilization, feed per gain, and bodyweight gain in an F2 cross of Japanese quail using recursive models. *Poult. Sci.* 95: 764–773.
- Bray, J. R., and J. T. Curtis, 1957 An ordination of the upland forest communities of Southern Wisconsin. *Ecol. Monogr.* 27: 325–349.
- Burbach, K., J. Seifert, D. H. Pieper, and A. Camarinha-Silva, 2016 Evaluation of DNA extraction kits and phylogenetic diversity of the porcine gastrointestinal tract based on Illumina sequencing of two hypervariable regions. *MicrobiologyOpen* 5: 70–82.
- Butler, D., B. R. Cullis, A. R. Gilmour, and B. J. Gogel, 2009 *ASREML-R, Reference Manual Version 3*. Queensland Department of Primary Industries and Fisheries, Brisbane.
- Camarinha-Silva, A., R. Jáuregui, D. Chaves-Moreno, A. P. Oxley, F. Schaumburg *et al.*, 2014 Comparing the anterior nare bacterial community of two discrete human populations using Illumina amplicon sequencing. *Environ. Microbiol.* 16: 2939–2952.
- Clarke, K. R., and R. M. Warwick, 2001 *Change in Marine Communities: An Approach to Statistical Analysis and Interpretation*, Ed. 1. E Ltd, Plymouth, UK.
- Cole, J. R., Q. Wang, J. A. Fish, B. Chai, D. M. McGarrell *et al.*, 2014 Ribosomal database project: data and tools for high throughput rRNA analysis. *Nucleic Acids Res.* 42: D633–D642.
- de los Campos, G., D. Gianola, and D. B. Allison, 2010 Predicting genetic predisposition in humans: the promise of whole-genome markers. *Nat. Rev. Genet.* 11: 880–886.
- de los Campos, G., A. L. Vazquez, R. Fernando, Y. C. Klimentidis, and D. Sorensen, 2013 Prediction of complex human traits using the genomic best linear unbiased predictor. *PLoS Genet.* 9: e1003608.
- Difford, G. F., J. Lassen, and P. Løvendahl, 2016 Genes and microbes, the next step in dairy cattle breeding. In *Proceedings, EAAP—67th Annual Meeting, Belfast*.

- Duncan, S. H., G. E. Loble, G. Holtrop, J. Ince, A. M. Johnstone *et al.*, 2008 Human colonic microbiota associated with diet, obesity and weight loss. *Int. J. Obes.* 32: 1720–1724.
- Estellé, J., N. Mach, Y. Ramayo-Caldas, F. Levenez, G. Lecomnier *et al.*, 2014 The influence of host's genetics on the gut microbiota composition in pigs and its links with immunity traits. In *Proceedings, 10th World Congress of Genetics Applied to Livestock Production*, Vancouver.
- Gianola, D., and D. Sorensen, 2004 Quantitative genetic models for describing simultaneous and recursive relationships between phenotypes. *Genetics* 167: 1407–1424.
- Hayes, B. J., H. A. Lewin, and M. E. Goddard, 2013 The future of livestock breeding: genomic selection for efficiency, reduced emissions intensity, and adaptation. *Trends Genet.* 29: 206–214.
- Heinritz, S. N., R. Mosenthin, and E. Weiss, 2013 Use of pigs as a potential model for research into dietary modulation of the human gut microbiota. *Nutr. Res. Rev.* 26: 191–209.
- Jannink, J. L., A. J. Lorenz, and H. Iwata, 2010 Genomic selection in plant breeding: from theory to practice. *Brief. Funct. Genomics* 9: 166–177.
- Jumpertz, R., D. S. Le, P. J. Turnbaugh, C. Trinidad, C. Bogardus *et al.*, 2011 Energy-balance studies reveal associations between gut microbes, caloric load, and nutrient absorption in humans. *Am. J. Clin. Nutr.* 94: 58–65.
- Karlsson, F. H., V. Tremaroli, I. Nookaew, G. Bergstrom, C. J. Behre *et al.*, 2013 Gut metagenome in European women with normal, impaired and diabetic glucose control. *Nature* 498: 99–103.
- Kim, H. B., and R. E. Isaacson, 2015 The pig gut microbial diversity: Understanding the pig gut microbial ecology through the next generation high throughput sequencing. *Vet. Microbiol.* 177: 242–251.
- Kim, H. B., K. Borewicz, B. A. White, R. S. Singer, S. Sreevatsan *et al.*, 2011 Longitudinal investigation of the age-related bacterial diversity in the feces of commercial pigs. *Vet. Microbiol.* 153: 124–133.
- Knol, E. F., B. Nielsen, and P. W. Knap, 2016 Genomic selection in commercial pig breeding. *Anim. Front.* 6: 15.
- Kostic, A. D., M. R. Howitt, and W. S. Garrett, 2013 Exploring host – microbiota interactions in animal models and humans. *Genes Dev.* 27: 701–718.
- Le Chatelier, E., T. Nielsen, J. Qin, E. Prifti, F. Hildebrand *et al.*, 2013 Richness of human gut microbiome correlates with metabolic markers. *Nature* 500: 541–546.
- Ley, R. E., F. Backhed, P. Turnbaugh, C. A. Lozupone, R. D. Knight *et al.*, 2005 Obesity alters gut microbial ecology. *Proc. Natl. Acad. Sci. USA* 102: 11070–11075.
- Ley, R. E., P. J. Turnbaugh, S. Klein, and J. I. Gordon, 2006 Microbial Ecology: Human gut microbes associated with obesity. *Nat* 444: 1022–1023.
- Looft, T., H. K. Allen, B. L. Cantarel, U. Y. Levine, D. O. Bayles *et al.*, 2014 Bacteria, phages and pigs: the effects of in-feed antibiotics on the microbiome at different gut locations. *ISME J.* 8: 1566–1576.
- Mach, N., M. Berri, J. Estellé, F. Levenez, G. Lecomnier *et al.*, 2015 Early-life establishment of the swine gut microbiome and impact on host phenotypes. *Environ. Microbiol. Rep.* 7: 554–569.
- Meuwissen, T. H. E., B. J. Hayes, and M. E. Goddard, 2001 Prediction of total genetic value using genome-wide dense marker maps. *Genetics* 157: 1819–1829.
- O'Connor, A., P. M. Quizon, J. E. Albright, F. T. Lin, and B. J. Bennett, 2014 Responsiveness of cardiometabolic-related microbiota to diet is influenced by host genetics. *Mamm. Genome* 25: 583–599.
- Pajarillo, E. A. B., J. P. Chae, M. P. Balolong, H. B. Kim, K. S. Seo *et al.*, 2014 Pyrosequencing-based analysis of fecal microbial communities in three purebred pig lines. *J. Microbiol.* 52: 646–651.
- Ramos, A. M., R. P. M. A. Crooijmans, N. A. Affara, A. J. Amaral, A. L. Archibald *et al.*, 2009 Design of a high density SNP genotyping assay in the pig using SNPs identified and characterized by next generation sequencing technology. *PLoS One* 4: e6524.
- Roehe, R., R. J. Dewhurst, C. A. Duthie, J. A. Rooke, N. McKain *et al.*, 2016 Bovine host genetic variation influences rumen microbial methane production with best selection criterion for low methane emitting and efficiently feed converting hosts based on metagenomic gene abundance. *PLoS Genet.* 12: e1005846.
- Ross, E. M., P. J. Moate, L. C. Marett, B. G. Cocks, and B. J. Hayes, 2013 Metagenomic predictions: from microbiome to complex health and environmental phenotypes in humans and cattle. *PLoS One* 8: e73056.
- Ryan, K. K., V. Tremaroli, C. Clemmensen, P. Kovatcheva-Datchary, A. Myronovych *et al.*, 2014 FXR is a molecular target for the effects of vertical sleeve gastrectomy. *Nature* 509: 183–188.
- Schwartz, D., K. Taras, K. Schäfer, S. Beijer, N. A. Bos *et al.*, 2010 Microbiota and SCFA in lean and overweight healthy subjects. *Obesity (Silver Spring)* 18: 190–195.
- Shulzhenko, N., A. Morgun, W. Hsiao, M. Battle, M. Yao *et al.*, 2011 Crosstalk between B lymphocytes, microbiota and the intestinal epithelium governs immunity vs. metabolism in the gut. *Nat. Med.* 17: 1585–1593.
- Sommer, F., N. Adam, M. E. V. Johansson, L. Xia, G. C. Hansson *et al.*, 2014 Altered mucus glycosylation in core 1 O-glycan-deficient mice affects microbiota composition and intestinal architecture. *PLoS One* 9: e85254.
- Spor, A., O. Koren, and R. Ley, 2011 Unravelling the effects of the environment and host genotype on the gut microbiome. *Nat. Rev. Microbiol.* 9: 279–290.
- Storey, J. D., and R. Tibshirani, 2003 Statistical significance for genomewide studies. *Proc. Natl. Acad. Sci. USA* 100: 9440–9445.
- Strandén, I., and D. J. Garrick, 2009 Technical note: derivation of equivalent computing algorithms for genomic predictions and reliabilities of animal merit. *J. Dairy Sci.* 92: 2971–2975.
- Stratz, P., K. Wimmers, T. H. E. Meuwissen, and J. Bennewitz, 2014 Investigations on the pattern of linkage disequilibrium and selection signatures in the genomes of German Piétrain pigs. *J. Anim. Breed. Genet.* 131: 473–482.
- Turnbaugh, P. J., R. E. Ley, M. A. Mahowald, V. Magrini, E. R. Mardis *et al.*, 2006 An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature* 444: 1027–1031.
- Valente, B. D., G. J. M. Rosa, G. De Los Campos, D. Gianola, and M. A. Silva, 2010 Searching for recursive causal structures in multivariate quantitative genetics mixed models. *Genetics* 185: 633–644.
- VanRaden, P. M., 2008 Efficient methods to compute genomic predictions. *J. Dairy Sci.* 91: 4414–4423.
- Vijay-Kumar, M., J. D. Aitken, F. A. Carvalho, T. C. Cullender, S. Mwangi *et al.*, 2010 Metabolic syndrome and altered gut microbiota in mice lacking toll-like receptor 5. *Science* 344: 228–231.
- Wang, Q., G. M. Garrity, J. M. Tiedje, and J. R. Cole, 2007 Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.* 73: 5261–5267.
- Wellmann, R., S. Preuß, E. Tholen, J. Heinkel, K. Wimmers *et al.*, 2013 Genomic selection using low density marker panels with application to a sire line in pigs. *Genet. Sel. Evol.* 45: 28.
- Wen, L., R. E. Ley, P. Y. Volchkov, P. B. Stranges, L. Avanesyan *et al.*, 2008 Innate immunity and intestinal microbiota in the development of Type 1 diabetes. *Nature* 455: 1109–1113.
- Wlodarska, M., C. A. Thaiss, R. Nowarski, J. Henao-Mejia, J. P. Zhang *et al.*, 2014 NLRP6 inflammasome orchestrates the colonic host-microbial interface by regulating goblet cell mucus secretion. *Cell* 156: 1045–1059.
- Zhao, L., G. Wang, P. Siegel, C. He, H. Wang *et al.*, 2013 Quantitative genetic background of the host influences gut microbiomes in chickens. *Sci. Rep.* 3: 1–6.

Communicating editor: M. P. L. Calus