



Measuring the effect of Western diet on liver tissue architecture by FLIM autofluorescence and harmonic generation microscopy: erratum

SUMAN RANJIT,¹ ALEXANDER DVORNIKOV,¹ EVGENIA DOBRINSKIKH,²
XIAOXIN WANG,² YUHUAN LUO,² MOSHE LEVI,² AND ENRICO GRATTON^{1,*}

¹Laboratory for Fluorescence Dynamics, Department of Biomedical Engineering, University of California, Irvine, CA, USA

²Departments of Medicine, University of Colorado-Anschutz Medical Campus, Aurora, Colorado, USA
*egratton22@gmail.com

Abstract: We correct a formula in our recent paper: [Biomed. Opt. Express **8**, 3143 (2017)].

©2017 Optical Society of America

OCIS codes: (170.2520) Fluorescence microscopy; (100.0100) Image processing; (170.3650) Lifetime-based sensing.

Formula 2 of this manuscript should be

$$\text{Fraction of bound NADH} = \text{distance}(\text{intercept to blue}) / \text{distance}(\text{green to blue}) \quad (2)$$

Instead of the formula in the manuscript

$$\text{Fraction of free NADH} = \text{distance}(\text{intercept to blue}) / \text{distance}(\text{green to blue}) \quad (2)$$