



Published in final edited form as:

*Nat Commun.* ; 6: 8139. doi:10.1038/ncomms9139.

## An *Arabidopsis* PWI and RRM motif-containing protein is critical for pre-mRNA splicing and ABA responses

Xiangqiang Zhan<sup>1</sup>, Bilian Qian<sup>2</sup>, Fengqiu Cao<sup>1</sup>, Wenwu Wu<sup>1</sup>, Lan Yang<sup>1</sup>, Qingmei Guan<sup>2</sup>, Xianbin Gu<sup>2</sup>, Pengcheng Wang<sup>3</sup>, Temiloluwa A. Okusolubo<sup>2</sup>, Stephanie L. Dunn<sup>2</sup>, Jian-Kang Zhu<sup>1,3</sup>, and Jianhua Zhu<sup>2</sup>

<sup>1</sup>Shanghai Center for Plant Stress Biology, Shanghai Institutes of Biological Sciences, Chinese Academy of Sciences, Shanghai 200032, China

<sup>2</sup>Department of Plant Science and Landscape Architecture, University of Maryland, 2121 Plant Sciences Building, College Park, Maryland 20742, USA

<sup>3</sup>Department of Horticulture and Landscape Architecture, Purdue University, West Lafayette, Indiana 47907, USA

### Abstract

The phytohormone abscisic acid (ABA) is important for growth, development and stress responses in plants. Recent research has identified ABA receptors and signalling components that regulate seed germination and stomatal closure. However, proteins that regulate ABA signalling remain poorly understood. Here we use a forward-genetic screen to identify *rbm25-1* and *rbm25-2*, two *Arabidopsis* mutants with increased sensitivity to growth inhibition by ABA. Using RNA-seq, we found that RBM25 controls the splicing of many pre-mRNAs. The protein phosphatase 2C *HAB1*, a critical component in ABA signalling, shows a dramatic defect in pre-mRNA splicing in *rbm25* mutants. Ectopic expression of a *HAB1* complementary DNA derived from wild-type mRNAs partially suppresses the *rbm25-2* mutant phenotype. We suggest that RNA splicing is of particular importance for plant response to ABA and that the splicing factor RBM25 has a critical role in this response.

---

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

Correspondence and requests for materials should be addressed to X.Z. (xqzhan@sibs.ac.cn) or to J.Z. (jhzhu@umd.edu).

#### Author contributions

X.Z., B.Q., F.C., J.Z. and J.-K.Z. designed the research. X.Z. performed the following experiments: isolation and characterization of the *roa1* mutants, mapping and gene complementation, subcellular localization, and isolation of total RNA for RNA-seq experiments and validation of RNA-seq data. J.Z. and Q.G. performed the water loss, heat stress responses, genetic analysis experiments, seed germination and contributed to experiments on responses of seedlings in ABA-containing media. B.Q., X.G., T.A.O and S.L.D. analysed responses of *roa1* mutants to ethylene and auxin. B.Q. performed the multiple sequence alignment and construction of phylogenetic tree of ROA1 and its orthologues. P.W. performed ROA1 phosphorylation profiling experiment. W.W., J.Z., B.Q. and L.Y. analysed the RNA-seq data. X.Z., J.Z. and B.Q. analysed the results and prepared the figures and tables for the article. J.Z., X.Z., and J.-K.Z. wrote the article.

#### Additional information

**Accession codes:** The RNA-seq data have been deposited in NCBI Gene Expression Omnibus (GEO) database under the accession code GSE66737.

**Supplementary Information** accompanies this paper at <http://www.nature.com/naturecommunications>

**Competing financial interests:** The authors declare no competing financial interests.

The phytohormone abscisic acid (ABA) is critical for many processes in the life cycle of plants, such as the promotion of seed maturation, establishment and maintenance of seed dormancy, regulation of stomatal aperture and adaptation to environmental stresses<sup>1–3</sup>. On exposure to environmental stresses, plants rapidly increase their ABA content as a result of increased ABA synthesis, decreased ABA catabolism, decreased formation of inactive ABA conjugates, or a combination of these factors<sup>4</sup>. The *de novo* biosynthesis of ABA is through the cleavage of a C<sub>40</sub> carotenoid precursor, followed by the conversion of the intermediate xanthoxin to ABA via ABA aldehyde<sup>2</sup>. The cleavage step catalysed by a 9-*cis* epoxy-carotenoid dioxygenase (NCED) has generally been considered the rate-limiting step in the pathway<sup>4,5</sup>. Wang *et al.*<sup>6</sup> found that *CED1* (for NCED defective 1), which is a putative  $\alpha/\beta$  hydrolase domain-containing protein and is allelic to the *BODYGUARD* gene that is essential for cuticle biogenesis, plays an important role in regulating the expression of *NCED3* and genes encoding other signalling components downstream of ABA biosynthesis.

Following ABA biosynthesis, multiple signal transduction pathways, which amplify the primary signal produced when ABA binds to its receptors, are required to control the ABA-regulated adaptive responses. These adaptive responses include stomatal closure (which allows the plant to conserve water), the accumulation of compatible osmolytes such as proline<sup>7</sup>, and the differential expression of a wide array of stress-responsive genes<sup>8,9</sup>. The core ABA signal transduction pathway is mediated by receptors in the PYR/PYL/RCAR family, which, on binding to ABA, inhibit the type 2C protein phosphatases (PP2Cs) such as ABI1, ABI2 and HAB1 (refs<sup>3,10–12</sup>). The PP2Cs are negative regulators of the ABA signalling pathway<sup>13,14</sup>. Inhibition of the PP2Cs leads to the activation of protein kinases in the sucrose non-fermenting related kinase 2 (SnRK2) family. In the absence of ABA, the PP2Cs can directly inactivate SnRK2 kinases by dephosphorylating a critical amino acid residue in their kinase activation loop<sup>15,16</sup>. These kinases are required for the subsequent regulation of the activities of downstream effectors in ABA signalling.

Since the three SnRK2 kinases, SnRK2.2, SnRK2.3 and SnRK2.6, control virtually all of the ABA responses in plants, the substrate proteins of the SnRK2s function as the downstream effectors of ABA responses<sup>17</sup>. The best known of the substrates include several transcription factors and ion channels<sup>18–21</sup>. The basic leucine zipper family transcription factors such as ABI5 and other ABA response factors or ABA-responsive element-binding factors (AREBs) control the transcription of ABA-responsive genes<sup>18,20</sup>. The ion channels such as the slow anion channel SLAC1 and the potassium channel KAT1 are critical for guard cell regulation by ABA<sup>19,21–24</sup>. Recent phosphoproteomics studies identified dozens of additional substrate proteins of the SnRK2s including several proteins with potential roles in RNA processing and proteins involved in chloroplast function<sup>25,26</sup>. These newly identified substrate proteins are potential effectors of ABA signalling, although their functions in ABA responses are largely unknown.

To identify cellular factors important for ABA responses, particularly ABA responses in leaf development, we conducted a genetic screen in *Arabidopsis thaliana* for mutants with defective leaf greening response to ABA treatment. We report here that *roa1* (*regulator of ABA response*) mutants are hypersensitive to ABA (HAB) as indicated by impaired leaf greening in response to ABA. The *ROA1/RBM25* locus defines a splicing factor required for

the splicing of transcripts of many genes including those involved in ABA signal transduction pathways. Our work has revealed a critical role of pre-mRNA splicing in ABA responses in plants.

## Results

### *roa1* mutants are hypersensitive to ABA

Identification of a large number of mutants defective in seed germination or guard cell movements has led to the discovery of genes involved in ABA biosynthesis and its downstream signalling events<sup>3</sup>. High levels of ABA inhibit plant growth and development and cause leaves to become pale or yellow<sup>27,28</sup>. The cellular factors controlling these processes are poorly understood. To identify critical regulators for ABA responses at the seedling growth stage, we screened an *A. thaliana* transfer DNA (T-DNA) insertion population with known integration sites (the SALK collection from ABRC) for mutants with increased sensitivity to exogenous ABA in the growth medium in terms of root growth and leaf yellowing. A similar screen was performed with an in-house generated T-DNA insertional population in a *TOUCH 4* promoter:luciferase reporter line<sup>29</sup> (*pTCH4:LUC*). These mutants were designated as *roa*. The *roa1-1* (SALK\_064472 from the SALK collection) and *roa1-2* (from the *pTCH4:LUC* population) mutants were chosen for in-depth characterization (Supplementary Fig. 1a). To compare *roa1-1* and *roa1-2* side-by-side, *roa1-2* was backcrossed with Col-0 and *roa1-2* allele without the *pTCH4:LUC* transgene was obtained. In the absence of exogenous ABA, *roa1-1* and *roa1-2* seedlings do not exhibit obvious growth defects (Fig. 1a). However, after being transferred to media supplemented with 5  $\mu$ M ABA, the newly emerged leaves in *roa1-1* and *roa1-2* seedlings are yellow, whereas all of the leaves in the wild-type seedlings are green (Fig. 1a). The mutants also had reduced primary root elongation in response to 5  $\mu$ M ABA (Fig. 1a,b). There is substantially less accumulation of chlorophylls in ABA-treated *roa1-1* and *roa1-2* leaves (Fig. 1a,c). Genetic analysis showed that both *roa1-1* and *roa1-2* mutations are recessive and are caused by mutations in a single nuclear gene (Supplementary Data 1). Pair-wise crosses between *roa1-1* and *roa1-2* suggested that the *roa1-1* and *roa1-2* mutations are allelic to each other (Fig. 1d).

To investigate whether the *roa1-1* and *roa1-2* mutations alter plant responses to other plant hormones or abiotic stress, we examined the responses of *roa1-1* and *roa1-2* plants to heat stress and two additional hormones. Sensitivity of the soil-grown *roa1-1* and *roa1-2* plants at both vegetative and reproductive stages to heat stress is not altered when they are compared with the wild-type plants (Supplementary Fig. 1b). We used the precursor of ethylene biosynthesis, 1-amino-cyclopropane-1-carboxylic acid (ACC), to examine the effect of ethylene on hypocotyl elongation of dark grown *roa1-1* and *roa1-2* seedlings. Hypocotyl elongation of *roa1-1*, *roa1-2* and wild-type seedlings was equally inhibited by exogenous application of ACC (Fig. 1e; Supplementary Fig. 1c). We further examined effects of auxin on hypocotyl and root elongation of *roa1-1* and *roa1-2* seedlings. *roa1-1* and *roa1-2* seedlings essentially responded to the same extent to indole-3-acetic acid in the growth media (Fig. 1f; Supplementary Fig. 1d,e). Therefore, while ROA1 may be involved in the

response to stresses and hormones other than ABA, we find no evidence of its involvement in the response to auxin, ethylene or heat.

### **ROA1 encodes a PWI and RRM motif-containing protein**

Although *roa1-1/SALK\_064472* was annotated as having a T-DNA insertion in the *At1g60200* gene, we could not find any T-DNA insertion in this gene. We carried out thermal asymmetric interlaced PCR (TAIL-PCR) analysis and identified a T-DNA insertion downstream of *At1g60200* in *roa1-1* (Fig. 2a). Although *roa1-2* also came from a T-DNA mutant population, there is no T-DNA insertion in *roa1-2*. To identify the gene responsible for the *roa1-1* and *roa1-2* mutant phenotypes, we prepared mapping populations by crossing *roa1-1* and *roa1-2* mutants with Landsberg *erecta* wild-type plants and performed map-based cloning for the *ROA1* locus. Genetic mapping pinpointed *ROA1* to a genomic region that contains the *At1g60200* gene. In the *roa1-1* mutant, we identified a single nucleotide substitution in *At1g60200*, and this substitution changes alanine at position 899 to valine in the deduced polypeptide (Fig. 2a,b). The *roa1-2* mutation also has a single nucleotide substitution in *At1g60200*, which results in the change of glutamine at position 570 to a premature stop codon in the deduced polypeptide (Fig. 2a,b). We confirmed the *ROA1* gene is *At1g60200* by functional complementation analysis. The wild-type *At1g60200* gene driven by its native promoter complemented the *roa1-1* and *roa1-2* mutant phenotype (Fig. 2d,e). *ROA1* expression is severely disrupted in *roa1-1*, possibly due to the T-DNA insertion downstream of *ROA1* (Fig. 2f).

*ROA1* encodes a putative PWI and RRM motif-containing protein with similarities to the human RNA-binding protein 25 (HsRBM25) (Fig. 2b,c). Hereafter, we refer to *ROA1* as *Arabidopsis* RBM25 and refer to *roa1-1* and *roa1-2* mutant alleles as *rbm25-1* and *rbm25-2*, respectively. *Arabidopsis* RBM25 shares substantial sequence similarity (50% over the RRM motif and 71% over the PWI motif) with the HsRBM25, which is known as a splicing factor<sup>30</sup>. Database searches revealed that RBM25 orthologues are present in other plant species including monocots (such as rice and maize) and dicots (such as soybean and tomato) (Fig. 2c; Supplementary Fig. 2). The last amino acid of RBM25 (changed to valine in *rbm25-1*) is conserved in nearly half of the plant RBM25 orthologues (Supplementary Fig. 2). As expected of a putative splicing factor, the RBM25–green fluorescent protein (GFP) fusion protein driven by the *RBM25* native promoter is localized in the nucleus of *Arabidopsis* root cells (Fig. 2g).

### **RBM25 negatively controls ABA responses in early development**

Seed germination is sensitive to ABA and we were interested whether RBM25 plays a role in this process. *rbm25-1* is slightly hypersensitive to exogenous ABA at the stage of seed germination (Fig. 3a) and is substantially hypersensitive to ABA in post-germination seedling development (Fig. 3b,c). This hypersensitivity to ABA was suppressed in three independent complementation lines of *rbm25-1* (Fig. 3a,b,c), indicating that loss-of-function of RBM25 results in this phenotype. ABA is also known to induce stomatal closure on drought stress. We performed water loss assays with detached shoots of soil-grown *rbm25* mutant plants to determine whether *rbm25* mutations alter plant transpirational water loss. Transpirational water loss rates are the same between the wild-type and *rbm25* mutants with

or without exogenous application of ABA (Fig. 3d). These results suggest that *rbm25* mutations do not alter stomatal responses to ABA.

### The *rbm25-1* mutation alters transcript accumulation profiles

We performed RNA-seq experiments to determine whether the *rbm25-1* mutation affects transcript accumulation profiles and whether altered transcript accumulation profiles may help explain the increased sensitivity of *rbm25-1* mutant to ABA treatment. We used the Illumina HiSeq 2500 System to sequence mRNA-seq libraries prepared from wild-type and *rbm25-1* seedlings with three biological replicates and obtained a minimum of 14.7 million of pair-end clean sequence reads (Supplementary Data 2). Compared with those in the wild-type, 106 genes displayed higher transcript levels (by at least 2-fold and with  $P < 0.05$ ), while 97 genes showed lower transcript levels in *rbm25-1* under control conditions (Fig. 4a,b; Supplementary Data 2). The RNA-seq analysis also revealed that 185 genes displayed at least a 2-fold increase in transcript levels (relative to the wild type and with  $P < 0.05$ ), while 267 genes showed at least a 2-fold decrease in transcripts levels under 100  $\mu\text{M}$  ABA treatment for 6 h (Fig. 4a,b; Supplementary Data 2). Furthermore, we found that there are 37 genes whose transcript levels are increased in *rbm25-1* plants under control and ABA treatment conditions (Fig. 4a; Supplementary Data 2). Similarly, there are 37 genes whose transcript levels are reduced in *rbm25-1* plants under control and ABA treatment conditions (Fig. 4a; Supplementary Data 2). The differentially expressed genes in *rbm25-1* encode proteins with diverse functions, and a substantial number of these proteins have predicted functions in stress responses (Fig. 4c; Supplementary Data 2). For example, under ABA treatment, 62 genes with annotated functions associated with response to abiotic or biotic stimulus or response to stress showed increased transcript levels in *rbm25-1* mutant seedlings (Supplementary Data 2). This observation of large and preferential change in the expression of stress-related genes is consistent with the obviously more 'stressed' appearance of the mutant plants under ABA treatment (Figs 1a and 3b) as a result of their impaired ability to cope with ABA treatment. With quantitative reverse transcription-PCR (qRT-PCR) analysis, we confirmed that the expression of *At5g65080* and *At1g53490* is substantially reduced in *rbm25* mutants, while expression of *At4g33720* is increased in *rbm25* mutants (Fig. 4d). *At5g65080* encodes a MADS-box transcription factor. *At1g53490* encodes a RING/U-box superfamily protein. *At4g33720* encodes a CAP (cysteine-rich secretory proteins, Antigen 5 and Pathogenesis-related 1 protein) superfamily protein.

### RBM25 regulates the splicing of pre-mRNAs

RBM25 has similar functional domains as the human splicing factor HsRBM25. To determine whether RBM25 functions in pre-mRNA splicing, we examined our RNA-seq data sets to look for potential effects of *rbm25-1* mutation on pre-mRNA splicing. The analysis revealed that 359 genes have splicing defects in untreated *rbm25-1*, while 416 genes show splicing defects in ABA-treated *rbm25-1* (Fig. 5a; Supplementary Data 3). Our RNA-seq data analysis further revealed that the *rbm25-1* mutation causes defects in splicing of gene transcripts in eight different alternative splicing categories: alternative acceptor, alternative donor, cassette exon, intron retention (IR), coordinate cassette exon, alternative first exon, alternative last exon and mutually exclusive exon (Fig. 5b; Supplementary Data 3). We designed primers unique to the intron of genes that are retained in the *rbm25-1* plants

and carried out semi-qRT-PCR assays to validate the IR results from the RNA-seq analysis. For RT-PCR analysis, we selected *At4g16143*, *At5g09330*, *At2g42010*, *At3g53340*, *At4g32040* and *At4g35800*. *At4g16143* encodes importin alpha isoform 2. *At5g09330* encodes NAC domain-containing protein 82. *At4g42010* encodes phospholipase D beta 1. *At3g53340* encodes nuclear factor Y, subunit B10. *At4g32040* encodes KNOTTED1-like homeobox gene 5. *At4g35800* encodes the unique largest subunit of nuclear DNA-dependent RNA II. The RT-PCR assays found that transcripts of *At4g16143*, *At5g09330*, *At2g42010*, *At3g53340*, *At4g32040* and *At4g35800* contain an intron in *rbm25-1* and *rbm25-2* with or without ABA treatment (Fig. 5c,e; Supplementary Fig. 3). Thus, the RT-PCR results confirmed the RNA-seq data of IR events in *rbm25-1*. RT-PCR analysis confirmed that exon 4 of *At2g48120* is skipped in *rbm25* mutants under both control and ABA treatment conditions (Fig. 5c; Supplementary Figs 3 and 4). *At2g48120* encodes the pale cress protein involved in chloroplast mRNA maturation<sup>31</sup>.

The genes carrying defective alternative splicing patterns in *rbm25-1* plants under control or ABA treatment conditions encode proteins with diverse functions in many biological processes, and the predicted roles of a good portion of these genes involve responses to abiotic or biotic stresses (Supplementary Data 3). It is possible that altered activities of mis-spliced genes in unstressed *rbm25-1* mutant account for differentially expressed genes in this mutant under control conditions as determined by RNA-seq analyses. There are 31 genes with a predicted role in DNA-dependent transcription in each category of abnormal splicing in unstressed *rbm25-1* mutant (Supplementary Data 3). These genes include promoter-specific transcription factors, general transcription factors and RNA polymerase II subunits.

We reasoned that disrupted function of the mis-spliced genes in *rbm25-1* plants should at least partly explain the impaired responses of *rbm25* mutants to ABA treatment including alterations in gene expression and increased sensitivity of *rbm25* mutants to ABA in terms of root growth and leaf yellowing. Thirty genes involved in transcription were found to have splicing defects in their transcripts in *rbm25-1* mutant under ABA treatment (Supplementary Data 3). These genes include the largest subunit of DNA-dependent RNA polymerase II, TATA-binding proteins, a nuclear factor, promoter-specific transcription factors and the subunits of the mediator complex. The mediators are involved in regulation of transcription from RNA polymerase II<sup>32</sup>. Loss-of-function or reduced activities of these genes will certainly lead to altered expression of genes in *rbm25-1* mutant under ABA treatment. Although genes carrying splicing defects in *rbm25-1* mutant under ABA treatment encode proteins involved in diverse biological processes, a substantial portion of the genes encode proteins that are predicted to function in biotic or abiotic stress-response pathways (Supplementary Data 3). In addition, the calcium-sensing receptor and HAB1 are known regulators of ABA responses<sup>33-35</sup>. There are additional PP2Cs whose transcripts are mis-spliced in ABA-treated *rbm25-1* mutant. Like HAB1, these PP2Cs may function as negative regulators of ABA signalling. Finally, impaired functions of four splicing factors (SC35-like splicing factor 33 encoded by *At1g55310*, CC1-like splicing factor encoded by *At2g16940*, U2 snRNP splicing factor encoded by *At1g60900* and splicing endonuclease 1 encoded by *At3g45590*) may contribute to defects of alternative splicing of gene transcripts in *rbm25-1*.

ABA represses the expression of genes involved in photosynthesis including the small subunit of ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) (*rbcS* genes) and chlorophyll a/b-binding proteins<sup>36,37</sup>. Related to this aspect of cellular function, we found mis-splicing of a group of genes that are involved in biogenesis and functionality of chloroplasts in ABA-treated *rbm25-1* mutant. These genes include pale cress protein (encoded by *At2g48120*; the mutant of this gene has reduced chlorophyll content<sup>31</sup>), the PsbQ subunit of the oxygen-evolving complex of photosystem II (encoded by *At4g21280*)<sup>38</sup>, starch branching enzyme 2.2 (encoded by *At5g03650*; it is transcriptionally regulated by the ABA-insensitive 4 (ABI4)<sup>39</sup>), UDP-glycosyltransferase superfamily protein involved in flavonoid biosynthetic process (encoded by *At4g09500*) and plastid transcriptionally active 5 (PTAC5, encoded by *At4g13670* (ref. 40)) (Supplementary Data 3). Mis-splicing for the transcripts of these proteins in *rbm25-1* mutant may be associated or even contribute to the leaf yellowing phenotype of the mutant. Finally, we observed that a cyclin-dependent protein kinase (CDKC2 encoded by *At5g64960*) is mis-spliced in *rbm25-1* under ABA treatment (Supplementary Data 3). CDKC2 colocalizes with spliceosomal components in a manner dependent on the transcriptional status of the cells and on CDKC2-kinase activity. Expression of CDKC2 modifies the location of spliceosomal components<sup>41</sup>. Thus, disruption of *CDKC2* may exacerbate the splicing defects caused by the *rbm25-1* mutation.

### RBM25 controls the alternative splicing of *HAB1* transcripts

Our RNA-seq analysis revealed that the last intron of *At1g72770* (which encodes *HAB1*) is retained in *rbm25-1* plants (Supplementary Data 3). *HAB1* is one of the clade A PP2Cs that are co-receptors of ABA and important negative regulators of ABA signalling<sup>14,33,42,43</sup>. *HAB1* transcripts are known to exist in three alternatively spliced forms (The *Arabidopsis* Information Resource, <http://www.arabidopsis.org>). In our efforts to confirm intron-retention event in *rbm25-1* and *rbm25-2* with RT-PCR and qRT-PCR analyses, we found additional splicing defects of *HAB1* in the *rbm25* mutants. The *rbm25* mutations cause altered accumulation of four alternatively spliced transcripts of *HAB1* under control and ABA treatment conditions (Fig. 6a,b). Because *HAB1* is upregulated by ABA, the defects of *HAB1* are more pronounced in these two mutants under ABA treatment. In *rbm25-1* and *rbm25-2* plants, *HAB1.1* (*At1g72770.1*) displayed reduced expression level, while *HAB1.2* (*At1g72770.2*) and *HAB1.4* (*At1g72770.4*, a previously unannotated splicing variant) showed increased expression level (Fig. 6a,b). The *HAB1.2* and *HAB1.4* isoforms are the intron-retained *HAB1* transcripts identified in our RNA-seq experiments (Supplementary Data 3). The expression level of *HAB1.3* (*At1g72770.3*) is very low and there is no difference between the wild type and *rbm25* mutants for the accumulation of *HAB1.3* transcripts (Fig. 6b). Transcripts of *HAB1.2* and *HAB1.4* will likely produce non-functional, truncated polypeptides of *HAB1* (Fig. 6c). Therefore, the overall manifestation of the defect in *HAB1* transcripts in *rbm25-1* and *rbm25-2* is the substantial reduction of *HAB1.1*, the major transcript of *HAB1* in wild-type plants (Fig. 6b,c).

### Ectopic expression of *HAB1* suppresses *rbm25-1*

Loss-of-function mutations in *HAB1* cause ABA hypersensitivity in early seedling growth and development<sup>14,33,44</sup>. Consistent with previous reports, the *hab1-1* mutant is hypersensitive to ABA during seed germination and early seedling development (Fig. 3b).

We observed that in response to ABA treatment *hab1-1* mutant seedlings exhibit a leaf yellowing phenotype similar to that of *rbm25* mutants, although the yellowing is not as severe as in *rbm25* (Fig. 2d). Considering the central role of HAB1 in ABA signalling and the seedling growth and development phenotypes of *hab1-1* mutant plants in response to ABA treatment, we hypothesized that the defects in *HAB1* pre-mRNA splicing, particularly the dramatic reduction in *HAB1.1*, may partially explain the leaf yellowing phenotype of ABA-treated *rbm25* mutant plants. To test this hypothesis, we ectopically expressed full-length *HAB1.1* complementary DNA (cDNA) under the CaMV 35S promoter in *rbm25-2* mutant plants. As a control, we also expressed *HAB1.2* cDNA under the CaMV 35S promoter in *rbm25-2* mutant plants. Two independent lines of T<sub>2</sub> transgenic *rbm25-2* seedlings were tested on media supplemented with 25 μM ABA. The results show that the ectopic expression of *HAB1.1* but not *HAB1.2* could partially rescue the leaf yellowing phenotype of *rbm25-2* in response to ABA treatment (Fig. 6d). These results indicated that the *HAB1.1* but not the *HAB1.2* spliced form is functional in controlling leaf development under ABA treatment and the reduction of the *HAB1.1* form contributes significantly to the leaf yellowing phenotype of *rbm25* mutant plants.

### ABA affects splicing of transcripts

We also examined the effect of ABA on the splicing pattern of transcripts in wild-type plants. ABA induces alternative splicing events in the transcripts of 27 genes in wild-type plants (Supplementary Data 3). Among these alternatively spliced genes in ABA-treated wild-type plants, two of them (*At3g28670* and *At3g24170*) showed more severe splicing defects in ABA-treated *rbm25-1* mutant. These results show that the ABA treatment causes defects in alternative splicing of pre-mRNAs in wild-type plants, and the splicing defects become much more severe when the splicing factor RBM25 is impaired.

### ABA regulates *RBM25* transcript level and post-translationally

*ROA1* expression is upregulated by ABA treatment (Fig. 7a). This is consistent with publicly available gene expression profiling data on *ROA1* under ABA treatment obtained using whole genome tiling arrays<sup>45</sup> (Supplementary Fig. 5). This is also consistent with the ABA-induced mutant phenotypes and more severe splicing defects in *rbm25* mutants under ABA treatment (Supplementary Data 3), and suggests a need for more RBM25 to strengthen pre-mRNA splicing under ABA treatment.

Protein phosphorylation is central to ABA signalling<sup>18,20,23</sup>. ABA increases the phosphorylation levels of several splicing factors including arginine/serine-rich splicing factor 41, a PWI domain-containing protein, splicing factor U2af small subunit A and serine/arginine-rich protein splicing factor RSZ32, while ABA also represses the phosphorylation levels of several splicing factors<sup>26</sup>. Because *RBM25* is upregulated by ABA, we examined whether phosphorylation status of RBM25 may also be regulated by ABA. We used a novel mass-spectrometry-based label-free quantitation method that facilitates systematic profiling of plant phosphoproteome changes with high efficiency and accuracy<sup>46</sup>. This method employs synthetic peptide libraries tailored specifically as internal standards for complex phosphopeptide samples and accordingly, a local normalization algorithm, LAXIC, which calculates phosphopeptide abundance normalized locally with co-eluting library peptides.



Our mass-spectrometry studies revealed that RBM25 is phosphorylated and ABA represses the phosphorylation level of RBM25 (Fig. 7b,c). Based on this data, we suggest that dephosphorylation of RBM25 may be important for its splicing function under ABA treatment. Alternatively, dephosphorylation of RBM25 may be important for its protein stability because phosphorylation often leads to protein degradation through ubiquitin-mediated proteolysis<sup>47,48</sup>.

## Discussion

In our unique genetic screen for regulators of ABA responses at seedling developmental stage, we identified a splicing factor, RBM25. The *rbm25* mutant plants do not exhibit obvious growth and developmental defects under normal growth conditions, and the mutants are not altered in their responses to heat stress, ethylene or auxin. We established the identity of *RBM25* through a map-based cloning strategy followed by a gene complementation analysis. Our genetic evidence demonstrated that RBM25 is required for proper ABA responses in plants by maintaining proper patterns of splicing of gene transcripts. Although splicing is important for all cellular processes involving gene expression, the splicing factor RBM25 is essential for root growth and leaf development in response to ABA but appears not to be required for normal growth and development. The *rbm25* mutants are hypersensitive to ABA in the growth medium, indicating that RBM25 may be involved in mediating signal transduction events downstream of ABA. Indeed, our RNA-seq analyses revealed that RBM25 controls the splicing of one of the central ABA signalling molecules, *HAB1* (Fig. 6). Ectopic expression of *HAB1.1* alone can partially suppress the hypersensitive phenotype of *rbm25-2* to ABA in growth medium (Fig. 6). These results indicate that RBM25 is critical for ABA signalling.

Because gene splicing is an essential cellular process, it is difficult to obtain and study null alleles of splicing factor genes due to their lethality. The *rbm25-1* and *rbm25-2* alleles appear to be null alleles since *RBM25* is not expressed in *rbm25-1* and the PWI domain, which is presumably important for splicing function, is truncated in *rbm25-2* (Fig. 2b,f). To our knowledge, this is the first time null alleles of a conserved splicing factor are found and characterized in plants. The conditional phenotype (that is, under ABA treatment only) of *rbm25* mutants implies that RBM25 is not essential for splicing for all or most plant genes. However, there are some splicing defects in *rbm25* mutant plants even under control conditions, but the splicing defects become more severe under ABA treatment such that growth and developmental phenotypes are manifested by the mutants under ABA treatment. Because *rbm25* mutants lack obvious growth and developmental defects in the absence of ABA or in responses to heat stress and two additional hormones (ethylene and auxin), but have severe phenotypic defects in the presence of ABA (Fig. 1 and Supplementary Fig. 1), we propose that plants may have a particular requirement for splicing in the presence of ABA. Indeed, this splicing factor (RBM25) shows increased expression under ABA treatment, which is consistent with the notion of an exaggerated requirement for splicing in the presence of ABA. On the other hand, we speculate that ABA signalling also regulates splicing in at least two ways: to increase the level of *RBM25* transcript; and regulate the phosphorylation status of RBM25 (Fig. 7; Supplementary Fig. 5) and other splicing factors<sup>26</sup>. The phosphorylation levels of several splicing factors are regulated by ABA in

wild-type plants<sup>26</sup>, which perhaps strengthen splicing so as to help plant cope with ABA and related environmental stress. In this study, we showed that the phosphorylation level of RBM25 is downregulated by ABA and we propose that reduced phosphorylation of RBM25 might be important for its activity as a splicing factor or its protein stability under ABA treatment although the physiological significance of RBM25 phosphorylation remains to be determined. Because orthologues of RBM25 exist in various plant species (Fig. 2c; Supplementary Fig. 2), the molecular function of RBM25 may be conserved in other plant species including crops.

Our RNA-seq experiments detected many genes whose splicing is controlled by RBM25. We observed that the *rbm25-1* mutant plants carry splicing defects in alternative splicing of pre-mRNAs (Supplementary Data 3). Our observation that there are more genes with defective splicing patterns when *rbm25-1* mutant was treated with ABA is consistent with ABA upregulation of *ROA1* (Fig. 7a; Supplementary Fig. 5; Supplementary Data 3). Intron-retained transcripts in *rbm25-1* mutant tend to include premature stop codons and therefore produce truncated or even inactive forms of proteins. In addition, many IR events may lead to the generation of nonsense-mediated decay substrates so that the overall level of the transcript would decrease and no translation would take place. Other abnormal, alternatively spliced transcripts in *rbm25-1* mutant probably produce nonfunctional proteins, and such splicing events were seldom detected in wild-type plants in our RNA-seq experiments. Several previous studies have shown that mutants with altered sensitivity to ABA are affected in pre-mRNA splicing under normal developmental conditions<sup>49–55</sup> or under abiotic stress conditions including drought and salinity<sup>56–60</sup>. In the previous studies, the various mutants were defective in ABA-regulated seed germination and/or stomatal regulation, but not in seedling leaf greening. In addition, the previously published mutants are defective in not only splicing but also other RNA processing events, and the published work did not link the mis-splicing of a particular gene(s) to the mutant phenotypes. In contrast, the *rbm25* mutants are primarily defective in leaf greening in response to ABA, and we have identified the mis-splicing of *HABI* as a critical mis-spliced gene contributing to the mutant phenotype (Fig. 6), although the defective splicing in many other genes may also contribute to the mutant phenotypes of *rbm25*.

## Methods

### Plant materials

T-DNA insertional mutant pool with known T-DNA insertion sites including *rbm25-1* (SALK\_064472) were obtained from *Arabidopsis* Biological Resource Center (ABRC; Columbus, OH). *rbm25-2* (containing *pTCH4:LUC* reporter gene<sup>29</sup>; seeds of *pTCH4:LUC* line was kindly provided by Dr Janet Braam) was isolated from a T-DNA-mutagenized *Arabidopsis* T<sub>2</sub> population. Seeds were surface sterilized and sown in germination medium (1 × Murashige–Skoog (MS) salts, 2% sucrose, 1.2% agar, pH 5.7). For ABA-sensitivity screening, 5-day-old seedlings were transferred to new MS agar plates supplemented with 5 μM ABA and were allowed to grow vertically for an additional 15–20 days. Seeds of *hab1-1* (SALK\_002104) were obtained from the ABRC. Plants at all developmental stages were

grown in a growth chamber at  $22\pm 1$  °C under cool, white light ( $\sim 100\text{--}120 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) with a long-day photoperiod (16-h light/8-h dark).

### Map-based cloning and gene complementation

The *rbm25-1* mutant was crossed with the Landsberg *erecta* accession, and a total of 896 plants homozygous for the *rbm25-1* phenotype in response to ABA in MS medium were selected from the F<sub>2</sub> population. A separate F<sub>2</sub> population was also generated from a cross between *rbm25-2* and Landsberg *erecta*, and 896 plants homozygous for the *rbm25-2* phenotype in response to ABA in MS medium were selected. Simple sequence length polymorphism markers were designed according to the information in the Cereon *Arabidopsis* Polymorphism Collection and were used to analyse recombination events. Mapping indicated that *rbm25-1* and *rbm25-2* mutations were located on the BAC clones T30E16 and F8A5. Candidate genes within this region were sequenced from the *rbm25* mutants and compared with those in GenBank to find the *rbm25-1* and *rbm25-2* mutations. For gene complementation of *roa1* mutants, a 7.4-Kb genomic fragment of *At1g60200* including 2.2 Kb upstream of the translation initiation codon and 0.4 Kb downstream of the translation initiation codon was amplified by PCR with T13D8 as a template (primers were listed in Supplementary Data 1). The amplified fragment was first cloned via Gateway technology (Invitrogen) into the pENTR1A Dual Selection Vector. The *RBM25* gene was then introduced into the pGWB501 vector, resulting in plasmid pGWB501-ROA1. The pGWB501-ROA1 construct was transferred into *Agrobacterium tumefaciens* (strain GV3101), and *rbm25-1* and *rbm25-2* plants were transformed by the floral dip method. Transgenic plants were selected on MS agar plates containing  $50 \text{ mg l}^{-1}$  hygromycin.

### Subcellular localization

The  $\sim 2$ -Kb genomic DNA fragment upstream of the start codon of *RBM25* (*At1g60200*) was amplified by PCR with T13D8 as a template using the primer pair At1g60200P-F5 and At1g60200P-R2, resulting in DNA fragment I. The coding region of *At1g60200* (from the start codon to the last codon without stop codon or any intron sequences) was amplified by PCR with the primer pairs At1g60200CDS-F2 and At1g60200CDS-R2, resulting in DNA fragment II. The DNA fragments I and II were joined together by PCR amplification using the primer pairs At1g60200P-F5 and At1g60200CDS-R2 with DNA fragments of I and II as a template. The resulting DNA fragment III was cloned into a vector pENTR1A initially and subsequently into the binary vector pGWB504, resulting in the plasmid RBM25:RBM25–GFP. The RBM25:RBM25–GFP construct was then introduced into *Arabidopsis* wild-type (Col-0) plants by floral dip transformation with *A. tumefaciens* strain GV3101. The subcellular localization of RBM25:RBM25–GFP protein in roots of transgenic plants (T<sub>2</sub> generation) was determined with a Leica SP5X confocal microscope (Leica Microsystems).

### Construction of the 35S:HAB1.1 and 35S:HAB1.2 plasmids

The cDNA fragments were amplified by PCR with primer pairs HAB1CD-F1 plus HAB1CD-R1 (for *HAB1.1*) and HAB1CD-F1 plus HAB1CD-R2 (for *HAB1.2*). The amplified cDNA fragments were initially cloned into the vector pENTR-D/TOPO and subsequently into the binary vector pGWB2 to generate plasmids 35S:HAB1.1 and

35S:HAB1.2. These plasmids were then introduced into *Arabidopsis* wild-type (Col-0) plants into *rbm25-2* plants by floral dip transformation with *A. tumefaciens* strain GV3101.

### RNA-seq data analyses and validation experiments

Fourteen-day-old wild-type and *rbm25-1* seedlings grown on MS medium (1 × MS salts, 2% sucrose, 0.6% agar, pH 5.7) were treated with 0 or 100 μM ABA for 6 h and were used for total RNA extraction. Total RNA was isolated with the Universal Plant Total RNA Extraction Kit (BioTeke) and treated with TURBO DNA-free Kit (Ambion) to remove any genomic DNA contaminants. mRNA-seq libraries were constructed following the standard Illumina protocols. There are three biological replicates per genotype. Illumina sequencing was performed in the Shanghai Center for Plant Stress Biology with an Illumina HiSeq 2500 System.

For each sample, RNA-seq raw reads were trimmed using Trimmomatic v0.32 and PRINSEQ v0.20.4 (refs<sup>61,62</sup>). Trimmomatic was used to remove the potential Illumina adapter contamination and conduct read trimming and clipping the low quality bases. PRINSEQ was subsequently employed to mainly remove low-complexity reads. The remained reads were then aligned to *A. thaliana* genome sequence and the reference-annotated genes (TAIR10) using TopHat v2.0.13 programme<sup>63</sup> with customized parameters specially for our RNA-seq libraries from plant *A. thaliana* (`--read-edit-dist 3--read-realign-edit-dist 0--b2-very-sensitive-r 150--mate-std-dev 200-a 6-i 8-I 10000--min-segment-intron 8--max-segment-intron 10000--microexon-search`).

According to RNA-seq-mapped reads and the reference-annotated transcripts, transcriptomes were reconstructed for each sample by Cufflinks v2.2.1 (ref. 64). Given variable efficiency of mRNA enrichments and rRNA depletion kits in samples, these transcripts were masked in transcriptome constructions to improve the overall robustness of transcript abundance estimates. To obtain a high confidence of transcriptomes, the novel constructed transcript was filtered out when the abundance was <20% (default is 10%) of the most abundant isoform for each gene. Then all of the constructed transcriptomes were merged with the reference-annotated transcripts using Cuffmerge<sup>64</sup> to yield comprehensive re-annotated gene transcripts including known and novel annotated transcripts in our RNA-seq samples. Subsequently, significantly differentially expressed genes were predicted by Cuffdiff<sup>64</sup> between the controls and tested samples, using twofold change and multiple test *P* value <0.05.

Given that IR is the most frequent alternative splicing (AS) type in plants<sup>65,66</sup> and juncBASE especially improves the reliability of the detection for such AS type<sup>67</sup>, we adopted juncBASE v0.6 to identify AS events within genes for our samples. First, two different databases were constructed, one is based on comprehensive re-annotated gene transcripts (as described above) to identify all internal AS events (not alternative first or last exon events), and the other is derived from just the reference-annotated transcripts to define alternative first and last exons. Then, a series of python scripts from juncBASE were employed to identify exon-exon junctions, exon-intron junctions and AS events, qualify the events and give significantly differentially splicing events between samples. Accordingly, juncBASE-detected AS events ranged from ~15,500 to ~20,500, and among the events 358

to 1,464 ones showed significantly differentially AS events ( $P$  value  $<0.05$  and abs (delta\_val)  $>5$ ) between pair-wise samples. Moreover, a module from CASH (comprehensive alternative splicing hunting, the renewed version of ASD (alternative splicing detector<sup>68</sup>) was used to rescue 19 to 50 differential IR events. Only genes with alternative splicing patterns in all three biological replicates of *rbm25-1* mutant plants are listed as differentially alternatively spliced genes.

For RT-PCR and qRT-PCR analyses, 5  $\mu$ g of total RNA was used for synthesis of the first-strand cDNA. Each experiment had three to four biological replicates and was repeated at least three times. The comparative cycle threshold (ct) method was applied for calculating gene expression levels, and *ACT2* was used as a reference gene.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

We thank Dr Shingo Nagawa, Youling Zeng, Sanjib Kumar Panda, Blanca E Barrera-Figueroa and Viswanathan Chinnusamy for technical assistance. This work was supported by the National Science Foundation Grant IOS0919745 to J.Z., by NIH grant R01GM059138 to J.-K.Z. and by the Chinese Academy of Sciences.

## References


1. Zhu JK. Salt and drought stress signal transduction in plants. *Annu Rev Plant Biol.* 2002; 53:247–273. [PubMed: 12221975]
2. Nambara E, Marion-Poll A. Abscisic acid biosynthesis and catabolism. *Annu Rev Plant Biol.* 2005; 56:165–185. [PubMed: 15862093]
3. Cutler SR, Rodriguez PL, Finkelstein RR, Abrams SR. Abscisic acid: emergence of a core signaling network. *Annu Rev Plant Biol.* 2010; 61:651–679. [PubMed: 20192755]
4. Zeevaart, JAD. *Biochemistry and Molecular Biology of Plant Hormones.* Hooykaas, PJJ.Hall, MA., Libbenga, KR., editors. Elsevier Science; 1999. p. 189-207.
5. Qin XQ, Zeevaart JAD. The 9-cis-epoxycarotenoid cleavage reaction is the key regulatory step of abscisic acid biosynthesis in water-stressed bean. *Proc Natl Acad Sci USA.* 1999; 96:15354–15361. [PubMed: 10611388]
6. Wang ZY, Xiong L, Li W, Zhu JK, Zhu J. The plant cuticle is required for osmotic stress regulation of abscisic acid biosynthesis and osmotic stress tolerance in *Arabidopsis*. *Plant Cell.* 2011; 23:1971–1984. [PubMed: 21610183]
7. Ober ES, Sharp RE. Proline accumulation in maize (*Zea mays* L.) primary roots at low water potentials (I. Requirement for increased levels of abscisic acid). *Plant Physiol.* 1994; 105:981–987. [PubMed: 12232259]
8. Bray EA. Classification of genes differentially expressed during water-deficit stress in *Arabidopsis thaliana*: an analysis using microarray and differential expression data. *Ann Bot.* 2002; 89:803–811. [PubMed: 12102506]
9. Neill SJ, Burnett EC. Regulation of gene expression during water deficit stress. *Plant Growth Regul.* 1999; 29:23–33.
10. Fujii H, et al. In vitro reconstitution of an ABA signaling pathway. *Nature.* 2009; 462:660–664. [PubMed: 19924127]
11. Ma Y, et al. Regulators of PP2C phosphatase activity function as abscisic acid sensors. *Science.* 2009; 324:1064–1068. [PubMed: 19407143]
12. Park SY, et al. Abscisic acid inhibits type 2C protein phosphatases via the PYR/PYL family of START proteins. *Science.* 2009; 324:1068–1071. [PubMed: 19407142]

13. Gosti F, et al. ABI1 protein phosphatase 2C is a negative regulator of abscisic acid signaling. *Plant Cell*. 1999; 11:1897–1910. [PubMed: 10521520]
14. Saez A, et al. Gain-of-function and loss-of-function phenotypes of the protein phosphatase 2C HAB1 reveal its role as a negative regulator of abscisic acid signalling. *Plant J*. 2004; 37:354–369. [PubMed: 14731256]
15. Umezawa T, et al. Type 2C protein phosphatases directly regulate abscisic acid-activated protein kinases in *Arabidopsis*. *Proc Natl Acad Sci USA*. 2009; 106:17588–17593. [PubMed: 19805022]
16. Vlad F, et al. Protein phosphatases 2C regulate the activation of the Snf1-related kinase OST1 by abscisic acid in *Arabidopsis*. *Plant Cell*. 2009; 21:3170–3184. [PubMed: 19855047]
17. Fujii H, Zhu JK. *Arabidopsis* mutant deficient in 3 abscisic acid-activated protein kinases reveals critical roles in growth, reproduction, and stress. *Proc Natl Acad Sci USA*. 2009; 106:8380–8385. [PubMed: 19420218]
18. Fujii H, Verslues PE, Zhu JK. Identification of two protein kinases required for abscisic acid regulation of seed germination, root growth, and gene expression in *Arabidopsis*. *Plant Cell*. 2007; 19:485–494. [PubMed: 17307925]
19. Geiger D, et al. Activity of guard cell anion channel SLAC1 is controlled by drought-stress signaling kinase-phosphatase pair. *Proc Natl Acad Sci USA*. 2009; 106:21425–21430. [PubMed: 19955405]
20. Nakashima K, et al. Three *Arabidopsis* SnRK2 protein kinases, SRK2D/SnRK2.2, SRK2E/SnRK2.6/OST1 and SRK2I/SnRK2.3, involved in ABA signaling are essential for the control of seed development and dormancy. *Plant Cell Physiol*. 2009; 50:1345–1363. [PubMed: 19541597]
21. Sato A, et al. Threonine at position 306 of the KAT1 potassium channel is essential for channel activity and is a target site for ABA-activated SnRK2/OST1/SnRK2.6 protein kinase. *Biochem J*. 2009; 424:439–448. [PubMed: 19785574]
22. Geiger D, et al. Guard cell anion channel SLAC1 is regulated by CDPK protein kinases with distinct Ca<sup>2+</sup> affinities. *Proc Natl Acad Sci USA*. 2010; 107:8023–8028. [PubMed: 20385816]
23. Lee SC, Lan W, Buchanan BB, Luan S. A protein kinase-phosphatase pair interacts with an ion channel to regulate ABA signaling in plant guard cells. *Proc Natl Acad Sci USA*. 2009; 106:21419–21424. [PubMed: 19955427]
24. Vahisalu T, et al. Ozone-triggered rapid stomatal response involves the production of reactive oxygen species, and is controlled by SLAC1 and OST1. *Plant J*. 2010; 62:442–453. [PubMed: 20128877]
25. Umezawa T, et al. Genetics and phosphoproteomics reveal a protein phosphorylation network in the abscisic acid signaling pathway in *Arabidopsis thaliana*. *Sci Signal*. 2013; 6:rs8. [PubMed: 23572148]
26. Wang P, et al. Quantitative phosphoproteomics identifies SnRK2 protein kinase substrates and reveals the effectors of abscisic acid action. *Proc Natl Acad Sci USA*. 2013; 110:11205–11210.
27. Munns R, Cramer GR. Is coordination of leaf and root growth mediated by abscisic acid? *Opinion. Plant Soil*. 1996; 185:33–49.
28. Trewavas, AJ., Jones, HG. *Abscisic Acid: Physiology and Biochemistry*. Davies, WJ., Jones, HG., editors. Bios Scientific Publishers; 1991. p. 169-188.
29. Iliiev EA, et al. Transcriptional and posttranscriptional regulation of *Arabidopsis* TCH4 expression by diverse stimuli. Roles of cis regions and brassinosteroids. *Plant Physiol*. 2002; 130:770–783. [PubMed: 12376643]
30. Zhou A, Ou AC, Cho A, Benz EJ Jr, Huang SC. Novel splicing factor RBM25 modulates Bcl-x pre-mRNA 5' splice site selection. *Mol Cell Biol*. 2008; 28:5924–5936. [PubMed: 18663000]
31. Reiter RS, Coomber SA, Bourett TM, Bartley GE, Scolnik PA. Control of leaf and chloroplast development by the *Arabidopsis* gene pale cress. *Plant Cell*. 1994; 6:1253–1264. [PubMed: 7919990]
32. Björklund S, Kim YJ. Mediator of transcriptional regulation. *Trends Biochem Sci*. 1996; 21:335–337. [PubMed: 8870496]
33. Saez A, et al. Enhancement of abscisic acid sensitivity and reduction of water consumption in *Arabidopsis* by combined inactivation of the protein phosphatases type 2C ABI1 and HAB1. *Plant Physiol*. 2006; 141:1389–1399. [PubMed: 16798945]

34. Han S, Tang R, Anderson LK, Woerner TE, Pei ZM. A cell surface receptor mediates extracellular Ca<sup>2+</sup> sensing in guard cells. *Nature*. 2003; 425:196–200. [PubMed: 12968184]
35. Wang WH, et al. Regulation of the calcium-sensing receptor in both stomatal movement and photosynthetic electron transport is crucial for water use efficiency and drought tolerance in *Arabidopsis*. *J Exp Bot*. 2014; 65:223–234. [PubMed: 24187420]
36. Bartholomew DM, Bartley GE, Scolnik PA. Abscisic acid control of rbcS and cab transcription in tomato leaves. *Plant Physiol*. 1991; 96:291–296. [PubMed: 16668167]
37. Weatherwax SC, Ong MS, Degenhardt J, Bray EA, Tobin EM. The interaction of light and abscisic acid in the regulation of plant gene expression. *Plant Physiol*. 1996; 111:363–370. [PubMed: 8787022]
38. Allahverdiyeva Y, et al. *Arabidopsis* plants devoid of PsbQ and PsbR subunits of the oxygen-evolving complex show altered PSII supercomplex organization and short-term adaptive mechanisms. *Plant J*. 2013; 75:671–684. [PubMed: 23647309]
39. Bossi F, et al. The *Arabidopsis* ABA-INSENSITIVE (ABI) 4 factor acts as a central transcription activator of the expression of its own gene, and for the induction of ABI5 and SBE2.2 genes during sugar signaling. *Plant J*. 2009; 59:359–374. [PubMed: 19392689]
40. Zhong L, et al. Chloroplast small heat shock protein HSP21 interacts with plastid nucleoid protein pTAC5 and is essential for chloroplast development in *Arabidopsis* under heat stress. *Plant Cell*. 2013; 25:2925–2943. [PubMed: 23922206]
41. Kitsios G, Alexiou KG, Bush M, Shaw P, Doonan JH. A cyclin-dependent protein kinase, CDKC2, colocalizes with and modulates the distribution of spliceosomal components in *Arabidopsis*. *Plant J*. 2008; 54:220–235. [PubMed: 18208522]
42. Robert N, Merlot S, N'guyen V, Boisson-Dernier A, Schroeder JI. A hypermorphic mutation in the protein phosphatase 2C HAB1 strongly affects ABA signaling in *Arabidopsis*. *FEBS Lett*. 2006; 580:4691–4696. [PubMed: 16876791]
43. Rubio S, et al. Triple loss of function of protein phosphatases type 2C leads to partial constitutive response to endogenous abscisic acid. *Plant Physiol*. 2009; 150:1345–1355. [PubMed: 19458118]
44. Leonhardt N, et al. Microarray expression analyses of *Arabidopsis* guard cells and isolation of a recessive abscisic acid hypersensitive protein phosphatase 2C mutant. *Plant Cell*. 2004; 16:596–615. [PubMed: 14973164]
45. Zeller G, et al. Stress-induced changes in the *Arabidopsis thaliana* transcriptome analyzed using whole genome tiling arrays. *Plant J*. 2009; 58:1068–1082. [PubMed: 19222804]
46. Xue L, et al. Quantitative measurement of phosphoproteome response to osmotic stress in *Arabidopsis* based on Library-Assisted eXtracted Ion Chromatogram (LAXIC). *Mol Cell Proteomics*. 2013; 12:2354–2369. [PubMed: 23660473]
47. Glickman MH, Ciechanover A. The ubiquitin-proteasome proteolytic pathway: destruction for the sake of construction. *Physiol Rev*. 2002; 82:373–428. [PubMed: 11917093]
48. Huang H, Tindall DJ. Regulation of FOXO protein stability via ubiquitination and proteasome degradation. *Biochim Biophys Acta*. 2011; 1813:1961–1964. [PubMed: 21238503]
49. Hugouvieux V, Kwak JM, Schroeder JI. An mRNA cap binding protein, ABH1, modulates early abscisic acid signal transduction in *Arabidopsis*. *Cell*. 2001; 106:477–487. [PubMed: 11525733]
50. Laubinger S, et al. Dual roles of the nuclear cap-binding complex and SERRATE in pre-mRNA splicing and microRNA processing in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA*. 2008; 105:8795–8800. [PubMed: 18550839]
51. Raczynska KD, et al. Involvement of the nuclear cap-binding protein complex in alternative splicing in *Arabidopsis thaliana*. *Nucleic Acids Res*. 2010; 38:265–278. [PubMed: 19864257]
52. Perea-Resa C, Hernández-Verdeja T, López-Cobollo R, del Mar Castellano M, Salinas J. LSM proteins provide accurate splicing and decay of selected transcripts to ensure normal *Arabidopsis* development. *Plant Cell*. 2012; 24:4930–4947. [PubMed: 23221597]
53. Golisz A, Sikorski PJ, Kruszka K, Kufel J. *Arabidopsis thaliana* LSM proteins function in mRNA splicing and degradation. *Nucleic Acids Res*. 2013; 41:6232–6249. [PubMed: 23620288]
54. Zhang XN, Mount SM. Two alternatively spliced isoforms of the *Arabidopsis* SR45 protein have distinct roles during normal plant development. *Plant Physiol*. 2009; 150:1450–1458. [PubMed: 19403727]


55. Jayaweera T, et al. Alternative splicing of *Arabidopsis* IBR5 pre-mRNA generates two IBR5 isoforms with distinct and overlapping functions. PLoS ONE. 2014; 9:e102301. [PubMed: 25144378]
56. Xiong L, et al. Modulation of abscisic acid signal transduction and biosynthesis by an Sm-like protein in *Arabidopsis*. Dev Cell. 2001; 1:771–781. [PubMed: 11740939]
57. Papp I, Mur LA, Dalmadi A, Dulai S, Koncz C. A mutation in the *Cap Binding Protein 20* gene confers drought tolerance to *Arabidopsis*. Plant Mol Biol. 2004; 55:679–686. [PubMed: 15604709]
58. Carvalho RF, Carvalho SD, Duque P. The plant specific SR45 protein negatively regulates glucose and ABA signaling during early seedling development in *Arabidopsis*. Plant Physiol. 2010; 154:772–783. [PubMed: 20699397]
59. Zhang Z, et al. Arabidopsis floral initiator SKB1 confers high salt tolerance by regulating transcription and pre-mRNA splicing through altering histone H4R3 and small nuclear ribonucleoprotein LSM4 methylation. Plant Cell. 2011; 23:396–411. [PubMed: 21258002]
60. Remy E, et al. A major facilitator superfamily transporter plays a dual role in polar auxin transport and drought stress tolerance in *Arabidopsis*. Plant Cell. 2013; 25:901–926. [PubMed: 23524662]
61. Bolger A, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics. 2014; 30:2114–2120. [PubMed: 24695404]
62. Schmieder R, Edwards R. Quality control and preprocessing of metagenomic datasets. Bioinformatics. 2011; 27:863–864. [PubMed: 21278185]
63. Kim D, et al. TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. Genome Biol. 2013; 14:R36. [PubMed: 23618408]
64. Trapnell C, et al. Differential analysis of gene regulation at transcript resolution with RNA-seq. Nat Biotechnol. 2013; 31:46–53. [PubMed: 23222703]
65. Syed NH, Kalyna M, Marquez Y, Barta A, Brown JWS. Alternative splicing in plants—coming of age. Trends Plant Sci. 2012; 17:616–623. [PubMed: 22743067]
66. Reddy ASN, Marquez Y, Kalyna M, Barta A. Complexity of the Alternative Splicing Landscape in Plants. Plant Cell. 2013; 25:3657–3683. [PubMed: 24179125]
67. Brooks AN, et al. Conservation of an RNA regulatory map between *Drosophila* and mammals. Genome Res. 2011; 21:193–202. [PubMed: 20921232]
68. Zhou X, et al. Transcriptome analysis of alternative splicing events regulated by SRSF10 reveals position-dependent splicing modulation. Nucleic Acids Res. 2014; 42:4019–4030. [PubMed: 24442672]
69. Tamura K, Stecher G, Peterson D, Filipinski A, Kumar S. MEGA6: molecular evolutionary genetics analysis version 6.0. Mol Biol Evol. 2013; 30:2725–2729. [PubMed: 24132122]






**Figure 1. Sensitivity of *roa1* mutants to ABA, ethylene precursor ACC and auxin**

Five-day-old seedlings grown in MS medium were transferred to MS medium containing 0 or 5 μM ABA and grown for an additional 14-day period (for **a–d**). (**a**) *roa1* mutants are hypersensitive to ABA. (**b**) Root elongation of seedlings shown in **a**. Root elongation was measured and is shown as a percentage relative to growth on normal MS medium. (**c**) Chlorophyll content of seedlings shown in **a**. (**d**) Sensitivity of F<sub>1</sub> seedlings from pair-wise crosses between *roa1-1* and *roa1-2* to ABA. Hypocotyl elongation of wild-type, *roa1-1* and *roa1-2* seedlings (for **e–f**). Seeds were germinated directly on medium containing various levels of ACC (in **e**; in darkness) or indole-3-acetic acid (in **f**; 8-h light (80 μmol m<sup>-2</sup> s<sup>-1</sup>) and 16-h dark cycle). Seedlings were allowed to grow for an additional 3-day (in **e**) or 5-day period (in **f**) after germination. Values represent mean ± s.d. (*n* = 20 in **b**, 8–10 pools of 4–5 seedlings in **c**, 49–135 in **e** and 89–125 in **f**).




**Figure 2. Molecular cloning of *ROA1***

(a) Map-based cloning of *ROA1*. Structure of the *ROA1* gene and positions of the *roa1-1* and *roa1-2* mutations are indicated. Filled boxes and empty boxes indicate exons and untranslated regions, respectively; the solid lines between boxes indicate introns. Positions are relative to translation start sites. (b) Structure of *ROA1*. Motifs are predicted by the SMART database. (c) Phylogenetic tree of *RBM25* and its close orthologues from other organisms. The protein identities are the same as defined in Supplementary Fig. 2. The Bootstrap consensus tree was generated with MEGA6 (ref. 69). Protein sequences of *RBM25* and its orthologues were aligned by MUSCLE and the Bootstrap consensus phylogenetic tree was reconstructed by the statistical method maximum likelihood using the Jones–Taylor–Thornton (JTT) model with a Bootstrap of 1000. Cut-off value for consensus tree is 50%. (d) Gene complementation analysis. (e) Chlorophyll content in plants shown in d. (f) Relative transcript levels of *ROA1* in WT and *roa1-1* plants as determined by qRT–PCR analysis. (g) Nuclear localization of the *RBM25*–GFP fusion protein. Scale bars, 2.5  $\mu\text{m}$ . Values represent mean  $\pm$  s.d. ( $n = 8$  in e; 4 in f).




**Figure 3. The *rbm25-1* mutation alters sensitivity to ABA in seed germination and post-germination seedling development, but does not affect water loss rates**

(a) Seed germination of WT, *rbm25-1* and *rbm25-1* complementation lines. Seeds (50–60 per genotype) were sown on surface of filter paper soaked with various levels of ABA. Seeds were considered germinated when radicles penetrate the seed coats. The percentage of seeds that had germinated 6 days after sowing was determined. (b) Post-germination seedling development of WT, *rbm25-1* and *rbm25-1* complementation lines subjected to 0 or 0.5  $\mu\text{M}$  ABA. Photograph was taken 16 days after sowing seeds. (c) The percentage of green seedlings in b. Seedlings with fully developed green cotyledons and at least one true green leaf were considered green. (d) Shoots from *rbm25-1*, *rbm25-2* and wild-type plants at the rosette stage were placed on weighing trays and allowed to dry slowly at constant temperature (25  $^{\circ}\text{C}$ ) and humidity (~50%). Weights of shoots were determined over a 4-h period. ABA, 100  $\mu\text{M}$  ABA for 3 h. Values are mean  $\pm$  s.d. ( $n = 8$  in a, 4 (number of MS medium plates containing 0 or 0.5  $\mu\text{M}$  ABA; each plate has 30–40 seedlings per genotype in b, 8 in d).



**Figure 4. The *rbm25-1* mutation causes disrupted gene expression as determined in RNA-seq experiments**

(a) Summary of differentially expressed genes in *rbm25-1*. (b) Heat map of differentially expressed genes in *rbm25-1*. Samples (rows) and genes (column) are hierarchically clustered via Pearson correlation. (c) GO term distribution of differentially expressed genes in *rbm25-1*. (d) Validation of differential gene expression portion of the RNA-seq results by qRT-PCR analysis. Values represent mean $\pm$ s.d. ( $n = 3$ ).



**Figure 5. The *rbm25-1* mutation causes defects in alternative splicing of gene transcripts**  
**(a)** Summary of genes whose transcripts were abnormally spliced in *rbm25-1* as determined by RNA-seq experiments. **(b)** Genes with defects in different types of alternative splicing patterns in *rbm25-1* as determined by RNA-seq experiments. **(c)** Validation of intron-retention and exon-skipping events in WT, *rbm25-1* and *rbm25-2* plants as determined by RT-PCR analysis. **(d)** Visualization with the integrative genomic viewer (IGV) of intron-retention events in WT and *rbm25-1* mutant plants. Annotated gene structures are shown


(bottom), with thick lines representing exons and thin lines representing introns. Wiggle plots represent the normalized read coverage on an autoscale.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



**Figure 6. RBM25 controls alternative splicing of *HAB1***

(a) Different alternative splicing variants of *HAB1* in WT and *rbm25-2* plants as determined by RT-PCR analysis followed by sequence analyses of cloned PCR products. *ACT8* was used as the loading control. In the close-up view of the portion corresponding to most parts of the last two exons of *At1g72770.1*, grey-shaded and orange-shaded boxes represent sequences in an exon, and folded lines between two boxes indicate introns in gene models of *At1g72770.1*, *At1g72770.3* and *At1g72770.4*. (b) Relative transcript levels of different alternative splicing variants of *HAB1* (expression level of *At1g72770.1* in unstressed WT was set to 1) in WT, *rbm25-1* and *rbm25-2* plants subjected to 100  $\mu$ M ABA for 0 or 6 h as determined by qRT-PCR analysis. Values are mean  $\pm$  s.d. ( $n = 4$ ). (c) Deduced C-terminal portion of HAB1.1 and HAB1.2. The predicted domain structures of HAB1.1 and HAB1.2 (HAB1.4 has the same structure as HAB1.2) are also presented. (d) Effect of *HAB1* on ABA sensitivity of *rbm25-1* mutant plants. Five-day-old seedlings grown in MS medium were

transferred to MS medium containing 0 or 25  $\mu\text{M}$  ABA and grown for an additional 20-day period.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



