

All letters are subject to editing and may be shortened. General letters can be sent to bjgpdisc@rcgp.org.uk (please include your postal address for publication), and letters responding directly to *BJGP* articles can be submitted online via **eLetters**. We regret we cannot notify authors regarding publication. For submission instructions visit: bjgp.org/letters

Multidisciplinary care

'Mate I'm getting seriously black poo – definitely getting my iron intake, lol!'

I received the above text message from a friend and, unsurprisingly, it raised a few alarm bells. And not just because a grown man had written 'lol' in a text message.

'I spoke to the pharmacist, he said nothing to worry about, it's all the vitamins I'm taking.'

And herein lies my concerns about the ways in which GP work is increasingly being shifted to other healthcare professionals.¹

In the case of my friend, although I'm not his doctor, I know his medical history well. I knew he suffers from indigestion. I knew he recently had a tooth removed. I knew he likes a few beers after football and that he smokes five cigarettes a day.

It emerged he had been taking high doses of ibuprofen to control his toothache. The multivitamin he was taking contained just 14 mg of iron. He went straight to A&E, and by the following morning had been diagnosed with a gastric ulcer. Thankfully, he is now on the road to recovery.

This is in no way meant to be disparaging towards pharmacists, colleagues whom I respect immensely. But it does concern me. The RCGP's *The 2022 GP* talks about working in federated organisations, with patients receiving routine care from a variety of healthcare professions.² But who will take overall responsibility for patients' care?

I fear that in our need to alleviate pressure, we run the risk of going too far in asking other professionals to take on some of our duties. With this we not only lose part of what makes our job the one we chose, but also risk doing our patients harm.

As GPs we have a unique and privileged role. We get to know our patients well. We take on all aspects of their clinical care, and have the breadth of training to enable us to do so, safely and effectively. Although multidisciplinary teams are no doubt effective, we must remember to ensure that someone is looking at the wider picture.

Simon Lynes,
GP Registrar, Severn Deanery.
E-mail: simon_lynnes@hotmail.com

REFERENCES

1. Butterworth J, Sansom A, Sims L, *et al*. Pharmacists' perceptions of their emerging general practice roles in UK primary care: a qualitative interview study. *Br J Gen Pract* 2017; DOI: <https://doi.org/10.3399/bjgp17X691733>.
2. Royal College of General Practitioners. *The 2022 GP: a vision for general practice in the future NHS*. London: RCGP, 2013. <http://www.rcgp.org.uk/policy/rcgp-policy-areas/general-practice-2022.aspx> (accessed 5 Jul 2017).

DOI: <https://doi.org/10.3399/bjgp17X691805>

Bad Medicine: Defensiveness

Dr Spence makes some very pertinent points about our current NHS.¹ There are long waits for GP appointments, for referrals, for scans, for test results. He suggests less prescribing, fewer referrals, fewer investigations. As a patient who has sustained brain damage due to benzodiazepine withdrawal after 40 years' consumption, I am deeply dismayed by my recent experience of the NHS. I have suffered beyond imagination for 4 years, tortured by my brain and body, and am physically and cognitively disabled as a consequence. My doctors seem totally incapable of grasping the extent of my suffering; instead, they translate my symptoms into something more palatable. I have been given several misleading and inaccurate diagnoses. This has now become a deliberate defensive strategy. I am compelled to seek further opinions, thereby wasting further NHS resources. I am part of a large and growing online community populated by the many other patients harmed by mind-altering prescription drugs, mainly benzodiazepines and antidepressants. We share our similar experiences of being dismissed by our doctors, misdiagnosed, and referred inappropriately.

We are shocked as the prescribing of antidepressants increases year on year despite mounting evidence of their lack of effectiveness and the possibilities of significant harm, including permanent damage to the central nervous system. We discuss the ways our withdrawal symptoms are dismissed as signs of mental illness and how we are misdiagnosed with a variety of diagnostic labels such as

depression, anxiety, CFS/ME, neurological functional impairment, somatic disorders, and medically unexplained symptoms. We wonder why we are offered inappropriate help or no help at all. We wonder why there are no appropriate services for patients who have become unwittingly dependent on prescription drugs. We wonder how a medical profession can get things so very wrong (benzodiazepines) and how they can repeat the same mistakes (antidepressants).

Fiona French,
Retired, Aberdeen.
E-mail: fionahfrench@hotmail.com

REFERENCE

1. Spence D. Bad Medicine: Defensiveness. *Br J Gen Pract* 2017; DOI: <https://doi.org/10.3399/bjgp17X691217>.

DOI: <https://doi.org/10.3399/bjgp17X691817>

Gluten-free food prescriptions for children with coeliac disease: should families have to pay?

The Department of Health consulted (until 22 June)¹ on withdrawing gluten-free food (GFF) prescriptions for patients with coeliac disease (CD), aiming to save £25 million. Many CCGs have already stopped or restricted prescribing, stating that families should buy their own GFF instead.

CD affects 1% of the population, putting them at risk of long-term consequences (including subfertility, osteoporosis, lymphoma, and poor growth), and the only treatment is a strict GF diet. Children are entitled to free NHS prescriptions, and dietary staples (bread, flour, pasta) have been available within regulated limits since the 1960s.² Since 'gluten intolerance' (functional symptoms exacerbated by gluten intake) emerged in the 1980s, GFF is more evident in supermarkets but remains costly. Availability is poor in convenience and budget stores, where low-income families and those without transport shop more frequently.³ One patient questionnaire found that 28%