



Energetic redistribution in allostery to execute protein function

Jin Liu^{a,1} and Ruth Nussinov^{b,c,1}

A perturbation at one site of the protein could cause an effect at a distant site. This important biological phenomenon, termed the “allosteric effect,” is essential for protein regulation and cell signaling, playing an important role in cellular function. Its fundamental functional significance has inspired numerous works aiming to understand how allostery works. Allostery can involve large, or unobserved, subtle (mainly side-chain) conformational changes (1). Conformational changes are driven by enthalpy. The term “dynamic allostery” was coined by Cooper and Dryden in the early 1980s to describe allostery “even in the absence of a macromolecular conformational change” (2). Cooper and Dryden argued that dynamic allostery is primarily an entropy effect. However, numerous works have been published over the last 20 y taking “dynamic allostery” to imply a complete absence of conformational change because the authors did not observe such changes (1). Importantly, “dynamic allostery” without observable conformational changes is still ruled by a population shift between two “distinct” states where a new energetic redistribution favorable for the allosteric (functional) state is either dominated by entropy, enthalpy, or both. Few studies questioned whether enthalpy plays a role in dynamic allostery as well (1). In PNAS, Kumawat and Chakrabarty (3) demonstrate that indeed even in dynamic allostery enthalpy plays a role by redistributing internal energies, especially electrostatic interaction energies, among residues upon perturbation (Fig. 1).

Electrostatics is an established player in function. Decades ago, Warshel (4) and Warshel et al. (5) demonstrated its critical role in protein catalysis and Perutz (6) in allostery. Recently, rearrangements of electrostatic networks were reported for conformationally driven allostery (7, 8). The report of hidden electrostatic interactions in dynamic allostery in PNAS (3) argues that redistribution of electrostatic interaction energy could be universal in allosteric proteins, with or without significant conformational change.

Proteins embrace redistribution of electrostatic interaction energies to execute their functions. For

example, myosin, a motor protein, binds to the negatively charged ATP, which redistributes the electrostatic bond network, thus altering the charge of the actin-binding site, leading to myosin’s dissociation from actin (7). Thus, myosin redistributes electrostatic interaction energy in allostery to execute its function of dissociation from actin.

The PDZ domain protein is a well-known cell signaling protein. The major functions of PDZ domains include recognition of the disordered C-terminal peptide motifs of hundreds of receptors and ion-channel proteins, dimerization with other modular domains, and phospholipid binding. As a dynamic regulator of cell signaling, PDZ function can be modulated allosterically (9). In the PDZ3 domain protein, the peptide recognition site is located at the hydrophobic $\beta 2$ – $\alpha 2$ region. Mutations of distal residues on the PDZ3 domain or deletion of the distal $\alpha 3$ helix (10) allosterically affect the binding affinity at the recognition site without causing significant conformational change. By contrast, binding of the peptide at the recognition site changes the dynamics of the PDZ3 domain, albeit still without large conformational change. Using molecular dynamics simulations and analysis of protein interaction energy, Kumawat and Chakrabarty (3) show that the perturbation of the binding of a peptide in the $\beta 2$ – $\alpha 2$ region leads to the side-chain rearrangement of certain charged residues to form new favorable electrostatic interactions. As a result, some of the previously favorable electrostatic interactions in the unbound state become unfavorable. Like a domino effect, additional side-chain rearrangements occur to release this energetic stress, altering the electrostatic and hydrogen bond network, especially in the N- and C-terminal regions, consistent with the known allosteric sites. How does this energetic redistribution affect PDZ3 function? PDZ domains are usually sequential on the protein chain. The energetic redistribution in one PDZ domain can propagate through an allo-network (11) to a second PDZ domain, thus affecting cell signaling. It is difficult to detect this dramatic redistribution of energy in experiments, because the energetic redistribution does not

^aDepartment of Pharmaceutical Sciences, University of North Texas Systems College of Pharmacy, University of North Texas Health Science Center, Fort Worth, TX 76107; ^bCancer and Inflammation Program, Leidos Biomedical Research, Inc., Frederick National Laboratory for Cancer Research, National Cancer Institute, Frederick, MD 21702; and ^cSackler Institute of Molecular Medicine, Department of Human Genetics and Molecular Medicine, Sackler School of Medicine, Tel Aviv University, Tel Aviv 69978, Israel

Author contributions: J.L. and R.N. designed research; and J.L. wrote the paper.

The authors declare no conflict of interest.

See companion article on page E5825.

¹To whom correspondence may be addressed. Email: jin.liu@unthsc.edu or NussinoR@mail.nih.gov.

the perturbation of external energy. The Kumawat and Chakrabarty (3) observations are thus important in refocusing current views. Rather than classifying allostery as entropic or dynamic, they contend that for the PDZ3 domain protein it should be viewed in terms of internal redistribution and population shift of specific electrostatic interactions. Future studies should explore the

generality of these findings and their predictions in cell signaling and drug discovery.

Acknowledgments

This was supported in whole or in part with federal funds from the Frederick National Laboratory for Cancer Research, National Institutes of Health Contract HHSN261200800001E.

- 1 Nussinov R, Tsai CJ (2015) Allostery without a conformational change? Revisiting the paradigm. *Curr Opin Struct Biol* 30:17–24.
- 2 Cooper A, Dryden DT (1984) Allostery without conformational change. A plausible model. *Eur Biophys J* 11:103–109.
- 3 Kumawat A, Chakrabarty S (2017) Hidden electrostatic basis of dynamic allostery in a PDZ domain. *Proc Natl Acad Sci USA* 114:E5825–E5834.
- 4 Warshel A (1978) Energetics of enzyme catalysis. *Proc Natl Acad Sci USA* 75:5250–5254.
- 5 Warshel A, et al. (2006) Electrostatic basis for enzyme catalysis. *Chem Rev* 106:3210–3235.
- 6 Perutz MF (1978) Electrostatic effects in proteins. *Science* 201:1187–1191.
- 7 Sato T, Ohnuki J, Takano M (2016) Dielectric allostery of protein: Response of myosin to ATP binding. *J Phys Chem B* 120:13047–13055.
- 8 Ohnuki J, Sato T, Takano M (2016) Piezoelectric allostery of protein. *Phys Rev E Stat Nonlin Soft Matter Phys* 94:012406.
- 9 Ivarsson Y (2012) Plasticity of PDZ domains in ligand recognition and signaling. *FEBS Lett* 586:2638–2647.
- 10 Petit CM, Zhang J, Sapienza PJ, Fuentes EJ, Lee AL (2009) Hidden dynamic allostery in a PDZ domain. *Proc Natl Acad Sci USA* 106:18249–18254.
- 11 Nussinov R, Tsai CJ, Csermely P (2011) Allo-network drugs: Harnessing allostery in cellular networks. *Trends Pharmacol Sci* 32:686–693.
- 12 Rao RR, Lawson CL (2014) Structure of catabolite activator protein with cobalt(II) and sulfate. *Acta Crystallogr F Struct Biol Commun* 70:560–563.
- 13 McKay DB, Steitz TA (1981) Structure of catabolite gene activator protein at 2.9 Å resolution suggests binding to left-handed B-DNA. *Nature* 290:744–749.
- 14 Sharma H, Yu S, Kong J, Wang J, Steitz TA (2009) Structure of apo-CAP reveals that large conformational changes are necessary for DNA binding. *Proc Natl Acad Sci USA* 106:16604–16609.
- 15 Townsend PD, et al. (2015) The role of protein-ligand contacts in allosteric regulation of the Escherichia coli catabolite activator protein. *J Biol Chem* 290:22225–22235.
- 16 Kornev AP, Taylor SS (2015) Dynamics-driven allostery in protein kinases. *Trends Biochem Sci* 40:628–647.
- 17 Kim J, et al. (2017) A dynamic hydrophobic core orchestrates allostery in protein kinases. *Sci Adv* 3:e1600663.
- 18 Hart JR, et al. (2015) The butterfly effect in cancer: A single base mutation can remodel the cell. *Proc Natl Acad Sci USA* 112:1131–1136.
- 19 O'Rourke KF, Gorman SD, Boehr DD (2016) Biophysical and computational methods to analyze amino acid interaction networks in proteins. *Comput Struct Biotechnol J* 14:245–251.
- 20 Kalescky R, Zhou H, Liu J, Tao P (2016) Rigid residue scan simulations systematically reveal residue entropic roles in protein allostery. *PLoS Comput Biol* 12:e1004893.
- 21 Kalescky R, Liu J, Tao P (2015) Identifying key residues for protein allostery through rigid residue scan. *J Phys Chem A* 119:1689–1700.
- 22 Law AB, Sapienza PJ, Zhang J, Zuo X, Petit CM (2017) Native state volume fluctuations in proteins as a mechanism for dynamic allostery. *J Am Chem Soc* 139:3599–3602.