



HHS Public Access

Author manuscript

Plant J. Author manuscript; available in PMC 2017 July 27.

Published in final edited form as:

Plant J. 2015 May ; 82(3): 523–531. doi:10.1111/tpj.12780.

***Chlamydomonas* as a model for biofuels and bio-products production**

Melissa A. Scranton, Joseph T. Ostrand, Francis J. Fields, and Stephen P. Mayfield*

California Center for Algae Biotechnology, Division of Biological Sciences, University of California, San Diego, 9500 Gilman Dr, La Jolla, CA 92093-0368, USA

SUMMARY

Developing renewable energy sources is critical to maintaining the economic growth of the planet while protecting the environment. First generation biofuels focused on food crops like corn and sugarcane for ethanol production, and soybean and palm for biodiesel production. Second generation biofuels based on cellulosic ethanol produced from terrestrial plants, has received extensive funding and recently pilot facilities have been commissioned, but to date output of fuels from these sources has fallen well short of what is needed. Recent research and pilot demonstrations have highlighted the potential of algae as one of the most promising sources of sustainable liquid transportation fuels. Algae have also been established as unique biofactories for industrial, therapeutic, and nutraceutical co-products. *Chlamydomonas reinhardtii*'s long established role in the field of basic research in green algae has paved the way for understanding algal metabolism and developing genetic engineering protocols. These tools are now being utilized in *C. reinhardtii* and in other algal species for the development of strains to maximize biofuels and bio-products yields from the lab to the field.

Keywords

Chlamydomonas reinhardtii; recombinant proteins; biofuels; bio-products; molecular engineering

INTRODUCTION

Access to affordable and environmentally sustainable fuels and energy sources may be the greatest challenge of this century. With demand continuing to increase and new supplies costing ever more to extract, the availability of fossil fuels will inevitably shrink, resulting in rising energy prices worldwide. With the rising cost of energy also comes the rising cost of food, as food and fuel prices are closely linked. Recently, algae-based biofuels have been highlighted as one of the best current alternative source of renewable energy (Merchant *et al.*, 2011; Georgianna and Mayfield, 2012; Leite *et al.*, 2013; Oncel, 2013). Algae do not compete for arable land, have fast generation times, can grow in salt and waste water, and have the potential to produce more oil per acre than land plants (Dismukes *et al.*, 2008; Demirbas and Demirbas, 2011). While most focus has been on the production of biodiesel,

*For correspondence (smayfield@ucsd.edu).

algae can also be a source of other fuels, such as bioethanol, biohydrogen, and biogas (Jones and Mayfield, 2011; Oncel, 2013).

While algal biofuels hold significant promise to meet future energy demands, improvements are needed at all levels of production in order to realize this potential. Algal biofuels are not economically viable on their own at current production levels (Brownbridge *et al.*, 2013; Nagarajan *et al.*, 2013). One near-term solution is to couple biofuel production with high-value co-products to increase the commercial value of the entire algal biomass. Current high-value bio-products produced in algae include industrial and therapeutic proteins as well as nutraceuticals and other high-value small molecules (Almaraz-Delgado *et al.*, 2013; Barrera and Mayfield, 2013; Rasala and Mayfield, 2014).

As the most characterized algal species with the largest set of genetic tools and techniques, *Chlamydomonas reinhardtii* is an excellent model organism to understand and improve biofuels and bio-products production in algae. *Chlamydomonas reinhardtii* has led the field in the development of molecular tools for strain selection and engineering for green alga. By far, more recombinant proteins have been expressed in *C. reinhardtii* than all other algal species combined (Almaraz-Delgado *et al.*, 2013; Barrera and Mayfield, 2013; Rasala and Mayfield, 2014). Studies in *C. reinhardtii* have also helped elucidate the molecular mechanisms behind algal lipid and hydrogen metabolism (Merchant *et al.*, 2011; Torzillo and Seibert, 2013). *C. reinhardtii* is also among the first of the engineered algal species to be studied in commercial settings, which allows academic researchers to begin to understand the challenges of bringing transgenic algae to commercial-scale production (Scoma *et al.*, 2012; Gimpel *et al.*, 2014; Schoepp *et al.*, 2014).

BIOENGINEERING ALGAE

Chloroplast engineering

Over the last 70 years, *C. reinhardtii* has become the flagship alga for laboratory studies and genetic manipulation. The eukaryotic green alga has three modifiable genomes and is capable of producing a wide variety of protein products (Rosales-Mendoza *et al.*, 2012; Barrera and Mayfield, 2013; Rasala *et al.*, 2013b). The efficient manufacture of these products at commercial viability will require a myriad of genetic tools to enhance protein accumulation and bioactivity (Figure 1). To date, the chloroplast genome has been the primary target for engineering protein production, predominantly because it readily performs homologous recombination and is easily transformed (Boynnton *et al.*, 1988). Typically, recombinant protein expressed from the chloroplast accounts for 1–10% of total protein (Almaraz-Delgado *et al.*, 2013; Barrera and Mayfield, 2013; Rasala and Mayfield, 2014).

One current limitation of chloroplast engineering in *C. reinhardtii* is the use of native promoters to achieve high levels of protein accumulation. The best exogenous gene expression achieved utilizes the endogenous *psbA* promoter (Manuell *et al.*, 2007). However, the effective use of the *psbA* promoter requires that the endogenous *psbA* regulatory region be removed due to a phenomenon known as auto-attenuation (Minai *et al.*, 2006; Rasala and Mayfield, 2011). This renders the organism non-photosynthetic, eliminating one of the unique attributes of algae. One way to alleviate this issue is to reintroduce the *psbA* gene

under the control of a different endogenous promoter (Gimpel *et al.*, 2014). This method results in photosynthetic *C. reinhardtii* that can produce recombinant proteins, albeit at reduced levels compared with the non-photosynthetic versions. Exogenous transcriptional regulation machinery have been used successfully in *C. reinhardtii* (Kato *et al.*, 2007), however exogenous promoters remain an underutilized resource for chloroplast engineering. Synthetic regulatory elements, such as UTRs, that modify gene expression have also been used in the *C. reinhardtii* chloroplast (Specht and Mayfield, 2012), although the successful use of fully synthetic promoters has yet to be described. A full understanding of synthetic promoter synthesis would allow for fine-tuning of gene expression, which is particularly useful for metabolic engineering.

In addition to engineering strains to produce maximal amounts of recombinant protein, chloroplast engineering is necessary for developing strains for robust growth, particularly in outdoor settings. Light intensity and availability is a major concern for maximizing growth rates and cell density. Excessive solar irradiation leads to photoinhibition, while low light flux due to weather or cell culture density can lead to reduced photosynthetic activity and slower growth. Chloroplast engineering for improved function under varying light conditions would allow maximum production of algae throughout the year. Early data have shown that the light antenna of *C. reinhardtii* can be modified to alter chlorophyll content and subsequently adjust its absorbance spectrum, altering photosynthetic productivity (Kirst *et al.*, 2012).

An alternative method of chloroplast engineering to alter photosynthetic activity has focused on the protein subunits of the photosystem itself. Photosystem II is a highly conserved protein complex, which performs the initial rate limiting step of photosynthesis. It is known that PSII is in part responsible for enhanced fitness in various light conditions (Mulo *et al.*, 2012). Recently, the D1 protein of *C. reinhardtii* was replaced with the homologous D1 proteins from *Synechococcus sp. 7942* (Vinyard *et al.*, 2013, 2014). These studies showed that complementation of the native D1 protein with natural variants from other species could increase photosynthetic efficiencies under varying light conditions, and also that this increased efficiency led to improved overall growth. In addition, due to *C. reinhardtii*'s unique low fluorescence background, these studies were the first to elucidate the mechanisms behind the optimized photosynthetic activity. Thus, not only were improved strains generated, but *C. reinhardtii* proved to be an ideal model for understanding PSII function.

Nuclear engineering

Although the chloroplast can effectively accumulate high levels of recombinant protein, nuclear transformation technology is required to fully engineer algae. Nuclear expression allows for organelle targeting or secretion of proteins, and enables more complex post-translational modifications of proteins such as glycosylation. In particular, proper protein localization is required for metabolic engineering, which relies on compartmentalization for some steps in syntheses. The nuclear genome of *C. reinhardtii* has been much more difficult to develop for recombinant protein production than the chloroplast. Low protein accumulation levels due in part to transgene silencing and insertion positional effects have

limited the use of the nuclear genome from a protein manufacturing standpoint (Cerutti *et al.*, 1997; De Wilde *et al.*, 2000; Wu-Scharf *et al.*, 2000). However, recently it was shown that fusing recombinant proteins to the Sh Ble antibiotic resistance gene, which requires high protein accumulation, could substantially increase recombinant protein accumulation (Rasala *et al.*, 2012). In addition, *C. reinhardtii* lacks strong constitutive promoters comparable to the viral promoters used in plants. While hybrid promoters have had some success in increasing transgene expression in *C. reinhardtii* (Schroda *et al.*, 2000), truly synthetic promoters may hold great promise for tight regulation of robust recombinant protein expression (Venter, 2007).

Homologous recombination

Homologous recombination (HR) allows for targeted gene knock-outs as well as targeted insertion for reducing positional effects for recombinant gene expression. This mechanism has enabled sophisticated engineering in the *C. reinhardtii* chloroplast; however at present, HR does not occur at sufficient levels in the nuclear genome to be of practical use. Knockdowns have been created by artificial microRNAs, which is also a valuable tool for genetic engineering (Molnar *et al.*, 2009; Zhao *et al.*, 2009; Moellering and Benning, 2010; Schmollinger *et al.*, 2010). Alternatively, a knockout strain library is currently being developed by insertional mutagenesis. Because of the scale of this project, a high-throughput screen known as ChlaM-meSeq has been established to identify genes disrupted by insertional mutagenesis. In a pilot screen, a pool of insertional mutants was shown to cover 39% of known protein coding genes (Zhang *et al.*, 2014). Although each of these methods have been valuable for progress of the field, a rapid method for developing novel knock-outs and targeted gene introduction is essential for future development of *C. reinhardtii*.

To date, methods of identifying HR in *C. reinhardtii* have not been reliable, nor have there been ways to significantly increase the low basal rate of HR in Chlamydomonas (Sodeinde and Kindle, 1993; Gumpel *et al.*, 1994; Nelson and Lefebvre, 1995). However, there have been many advances in targeted gene delivery across other platforms in the past decade. Transcription-activator-like effectors (TALEs) allow one to design a unique DNA-binding target sequence and the addition of a nuclease can achieve targeted restriction digestion and subsequent homology directed repair (Li *et al.*, 2011). Successful gene targeting by a TALE has been shown by a subsequent increase in endogenous ARS expression (Gao *et al.*, 2014). However, to this point, TALENs have not successfully been utilized for targeted gene disruption in *C. reinhardtii*. The CRISPR/Cas9 system is an alternative method for inducing targeted gene delivery. Here, a guide RNA will direct restriction by the Cas9 nuclease to a specific sequence of DNA. The CRISPR/Cas9 system has been successful in mammals, fish, fungus, and plants (tobacco, Arabidopsis, sorghum and rice; Jiang *et al.*, 2013). Cas9 has recently been transiently expressed in *C. reinhardtii*, although stable cell lines have not been established (Jiang *et al.*, 2014a). Successful transient expression shows that Cas9 is capable of modifying the *C. reinhardtii* genome.

The *C. reinhardtii* nuclear genome generally performs repairs of double-stranded breaks almost exclusively by non-homologous end joining (NHEJ) and errors are often introduced during the process. Unfortunately, because *Chlamydomonas* relies so heavily on NHEJ as a

repair mechanism, even if TALENs or the CRISPR/Cas9 systems are developed in *C. reinhardtii*, it is unlikely that a large percentage of gene delivery will occur by targeted homology repair. NHEJ machinery is comprised of three main subunits: Ku70, Ku80, and DNA ligase IV. In yeast, Ku70 and Ku80 form a heterodimer that binds to double stranded breaks and recruits DNA ligase IV to mediate NHEJ (Kanaar *et al.*, 1998). The *C. reinhardtii* genome contains homologues to each of these genes. Based on a study in *Pichia pastoris*, one strategy may be to stifle expression of these repair proteins in *Chlamydomonas* in order to increase the level of HR activity (Näätsaari *et al.*, 2012).

High-throughput screening

High-throughput screening allows detection of rare phenotypes including those following gene insertion or mutagenesis. One of the fastest quantitative methods of screening mutants is fluorescence-activated cell sorting (FACS). This technique has been utilized for detection of high lipid accumulating mutants by a process utilizing Nile Red, a lipid dye, to analyze lipid levels in a set of mutants generated using EMS (Xie *et al.*, 2014). This method successfully isolated *C. reinhardtii* strains that accumulated 23–58% higher fatty acid content by dry mass when compared to their parent strain. Alternatively, Terashima *et al.* (2014) developed a high lipid-accumulating mutant pool by insertional mutagenesis, a strategy termed *Chlamydomonas* High Lipid Sorting (CHiLiS). This method generates a detectable insertion that can be used to rapidly identify the disrupted gene(s). Characterization of the strains isolated by this method showed a significant increase in triacylglycerol accumulation. These papers demonstrate that rapid screening can identify mutants with altered metabolism for the production of biofuels, which can easily be translated beyond *C. reinhardtii*. In addition to fluorescent stains, fluorescent protein reporters in *C. reinhardtii* have been shown to effectively sort using FACs (Rasala *et al.*, 2013a). By tagging recombinant proteins with fluorescent protein tags, production strains can be rapidly screened and genes that affect recombinant protein accumulation can be identified. Fluorescent reporters fused to antibodies have also been successfully used to sort for wild isolates with specific cell wall components (Jiang *et al.*, 2014b). Development of techniques using these reporters will be necessary to rapidly isolate and engineer production strains.

Ultimately, high-throughput screening technologies will be an essential tool for generating strains for outdoor production. High-throughput screening allows one to select exclusively based on desirable traits, rather than validating transformation with linkage to an antibiotic marker. Horizontal gene transfer of antibiotic resistance cassettes to environmental microbes poses a theoretical threat to human health and safety. Therefore, high-throughput screening mechanisms that avoid generation of transgenic algae with drug resistant markers, and that can be scaled up to outdoor ponds for economically feasible biofuels production, are an area where more research is required.

ALGAL BIOFUELS

Biodiesel

Chlamydomonas reinhardtii has been a key model in understanding complex algae lipid metabolism. Current efforts have focused on the use of triacylglycerols (TAGs) as a first generation biodiesel. TAGs from algae are also of interest as alternatives to plant-based edible oils (Klok *et al.*, 2014). Traditionally, *C. reinhardtii* has not been seen as an oleaginous species (James *et al.*, 2011). However, studies have shown that natural levels of lipids can vary widely in *C. reinhardtii* strains (Siaut *et al.*, 2011). In addition, wild-type and mutant *C. reinhardtii* can accumulate significant amounts of TAGs in response to nitrogen or salt stress (Wang *et al.*, 2009; Li *et al.*, 2010; Work *et al.*, 2010; Goodson *et al.*, 2011; James *et al.*, 2011; Siaut *et al.*, 2011). In fact, *C. reinhardtii* studies have helped elucidate the mechanisms behind lipid accumulation in response to nitrogen deprivation (Boyle *et al.*, 2012; Msanne *et al.*, 2012; Goodenough *et al.*, 2014). *Chlamydomonas reinhardtii* has also been an excellent model to identify algal genes involved in TAG metabolism (Merchant *et al.*, 2011; Klok *et al.*, 2014).

Targeted overexpression of putative TAG metabolic genes in *C. reinhardtii* has been met with mixed success. Overexpression of type 2 diacylglycerol acyltransferases (DGATs) DGAT2-1 and DGAT2-5 led to increased lipid content, while DGAT2-a,b,c overexpression had no effect (Deng *et al.*, 2012; La Russa *et al.*, 2012). Overexpression of acyl-ACP (acyl carrier protein) esterase (AAE) led to an altered lipid profile but not an increase in lipid content (Blatti *et al.*, 2011). It is clear that even in *C. reinhardtii*, a better understanding of lipid metabolism is required in order to fully utilize algae's potential as a source of TAGs for biodiesel or edible oil production.

Biohydrogen

As an alternative to liquid fuels, *C. reinhardtii* has also been extensively studied as a model for the photoproduction of biohydrogen. Renewed interest in *C. reinhardtii* biohydrogen production began when significant and sustained H₂ production was demonstrated in sulfur starved strains (Melis *et al.*, 2000). The theoretical maximum light conversion efficiency for H₂ production is 13.4%, which is higher than the 11.2% limit for biodiesel (Torzillo and Seibert, 2013). In addition, purities of up to 98% can be achieved without the need of extraction (Torzillo *et al.*, 2009). However, several factors significantly limit the actual production of H₂ in algae. The most significant limitation is the sensitivity of hydrogenase to O₂, which inhibits hydrogenase activity by affecting transcription and protein maturation (Ghirardi *et al.*, 2007). Additional constraints include competition for electrons from alternative pathways, inefficient light conversion of large light-harvesting antennae, and inherent limitations of anaerobic growth (Torzillo and Seibert, 2013).

Studies in *C. reinhardtii* to address these limitations have included the generation of improved strains through either mutagenesis or targeted genetic engineering (Melis *et al.*, 2007; Esquivel *et al.*, 2011; Torzillo and Seibert, 2013; Baltz *et al.*, 2014; Xu *et al.*, 2014). Studies have also shown significantly improved H₂ production through the optimization of growth conditions including periodic sulfur starvation (Melis *et al.*, 2000) and optimized

light, media and photobioreactor conditions (Torzillo and Seibert, 2013; Oncel *et al.*, 2014). However, despite these optimizations, the best light conversion efficiency reached has been approximately 3%. Even if these optimal conditions could be perfectly scaled to outdoor growth, the cost of algal H₂ would be well over US\$8 per gallon gasoline equivalent (gge) (James *et al.*, 2009). However, continued research holds promise to optimize algae strains and growth conditions to make biohydrogen a competitive fuel in the future.

HIGH-VALUE BIO-PRODUCTS

Production of recombinant proteins in *C. reinhardtii* has been a fruitful area of research in recent years. Several recent articles have thoroughly reviewed the plethora of protein products that have been successfully expressed within *C. reinhardtii* (Almaraz-Delgado *et al.*, 2013; Barrera and Mayfield, 2013; Rasala and Mayfield, 2014). To date, 34 different protein co-products have been produced in algae, 29 of which were produced in *C. reinhardtii*.

The vast majority of targeted proteins have been therapeutic proteins. With a current market value of US\$140 billion, therapeutic proteins provide an attractive co-product target to increase the economic value of algal biomass (Walsh, 2014). Therapeutic proteins produced include the UV-protectant metallothionein (Zhang *et al.*, 2006), antibody mimics 14FN3 and SAA-10FN3 (Rasala *et al.*, 2010), and anti-cancer proteins TNF-Related Apoptosis Inducing Ligand (TRAIL; Yang *et al.*, 2006), and allophycocyanin (Su *et al.*, 2005). The glycoprotein hormone erythropoietin has also been produced from the *C. reinhardtii* nucleus, demonstrating the ability to generate recombinant proteins with glycosylation, the most common post-translational modification found on protein therapeutics (Eichler-Stahlberg *et al.*, 2009; Walsh, 2014).

The largest class of therapeutic proteins that have been produced in *C. reinhardtii* are subunit vaccines (Almaraz-Delgado *et al.*, 2013; Barrera and Mayfield, 2013; Rasala and Mayfield, 2014). Vaccines are a particularly attractive target because algae are generally regarded as safe (GRAS) by the FDA. Therefore, if orally available vaccines can be produced, these vaccines may be stored and administered as lyophilized algal pellets. Edible algal vaccines would reduce the cost of vaccines by orders of magnitude and vastly increase their availability since algal vaccine pellets do not need to be purified, do not require needle injection, and can be stored for up to 20 months at RT (Dreesen *et al.*, 2010; Gregory *et al.*, 2013).

Chlamydomonas reinhardtii has also been shown to be able to produce a novel type of antibody therapeutic. Monoclonal antibodies (mAbs) are the largest growing segment of protein therapeutics with almost 27% of the new approvals and six of the top 10 selling biotherapeutics products in 2013 (Walsh, 2014). In particular, antibody-drug conjugates (ADCs) are gaining in popularity as more effective cancer treatment. The most significant drawback of ADCs is the requirement for chemical linkage of the toxic molecule, which leads to increased production costs and potential off-target toxicity of free toxins. In contrast, the unique environment of the *C. reinhardtii* chloroplast is able to produce immunotoxins in which toxic proteins are genetically linked to the targeting antibody (Tran

et al., 2013a,b). This reduces the cost of production since the immunotoxin is produced as a single protein and this production method can also help eliminate off-target toxicity.

Protein co-products expressed in *C. reinhardtii* also include a range of nutraceuticals for use as feed additives and/or human nutritional supplements. Feed additives include phytase, xylanase, and flounder growth hormone (Kim *et al.*, 2002; Yoon *et al.*, 2011; Georgianna *et al.*, 2013). Mammary-associated serum amyloid A (MAA) is one of the best expressed bio-products in *C. reinhardtii*, accumulating to over 5% of total soluble protein (Manuell *et al.*, 2007). MAA can be used as a prophylaxis against enteric bacterial infections in humans as well as replacement for antibiotics in animal feed (Larson *et al.*, 2003; Mack *et al.*, 2003). Recombinant proteins have also been engineered to increase high-value nutritional compounds in *C. reinhardtii* such as organic selenium (Hou *et al.*, 2013), carotenoids (Baldo *et al.*, 2011; Couso *et al.*, 2011), and triacylglycerols (La Russa *et al.*, 2012). Carotenoids and triacylglycerols have the added benefit of being used in biofuels formulations (Merchant *et al.*, 2011; Peralta-Yahya *et al.*, 2011).

OUTDOOR CULTIVATION SYSTEMS

Large-scale microalgae cultivation can occur in either open or closed systems (Ugwu *et al.*, 2008; Singh *et al.*, 2011). Open systems range from small artificial ponds to large open bodies of water; however the most productive open systems currently implemented are shallow raceway ponds (Singh *et al.*, 2011). Although these open systems require large areas of land, they can be constructed in arid environments and avoid competition with preexisting agriculture (Georgianna and Mayfield, 2012). Raceways can be constructed relatively cheaply and require little maintenance (Singh *et al.*, 2011; Schoepp *et al.*, 2014). Open systems are inherently susceptible to contamination and thus are predominantly used to culture only robust photoautotrophic species (Ugwu *et al.*, 2008; Rasala and Mayfield, 2014). Heterotrophic growth in open systems is not a viable option due to the heightened susceptibility of contamination. Greenhouses can be used to cover smaller open systems in order to deter contamination and potentially increase productivity. Additionally, greenhouses allow for the growth of genetically engineered strains in contained 'open' systems, but these structures will add to the initial cost of a facility.

Closed systems are highly variable in design but overall provide a more productive environment while protecting the culture from contamination (Ugwu *et al.*, 2008; Singh *et al.*, 2011). More strains of algae can be cultivated in these closed systems, including mixotrophic and hydrogen-producing species of microalgae (Scoma *et al.*, 2012; Gimpel *et al.*, 2014). The productivity and sterility benefits of a closed system are countered by their heightened costs as they may cost up to 10 times that of open systems due to the additional infrastructure and operational costs (Del Campo *et al.*, 2007).

Recently, pilot facilities containing a variety of these systems have been built to test strains in a setting more comparable to conditions encountered at commercial facilities (Scoma *et al.*, 2012; Gimpel *et al.*, 2014; Schoepp *et al.*, 2014). Experiments with *C. reinhardtii* have been conducted in these outdoor systems to test the viability of the strain as a producer of biomass, biohydrogen and even high-value bio-products. These studies are invaluable for

understanding and addressing the challenges of bringing laboratory strains into commercial production.

OUTDOOR CULTIVATION OF *C. REINHARDTII*

Researchers at the University of California, San Diego have developed one of the first systems to measure research-scale production of algal strains in closed bags and open ponds. Biomass production ($\text{g L}^{-1} \text{ day}^{-1}$) was reported for several algae and cyanobacteria species (Table 1; Schoepp *et al.*, 2014). *Chlamydomonas reinhardtii* (CC-1690) out produced eight other species in a 100-L closed system, but production was reduced by 80% when grown in an open 800-L pond. Lipid content was not measured in this study; however, reported average lipid content in *C. reinhardtii* is comparable to other species (Table 1; Griffiths and Harrison, 2009). This suggests that *C. reinhardtii* is competitive with other algal species in closed systems. While closed systems may be economical for high-value products, the cost and size limitations of closed systems may hinder sufficient production of lipid-accumulating strains for biofuel production (Figure 1).

Chlamydomonas reinhardtii was also tested in an outdoor system for its ability to produce biohydrogen gas. Scoma *et al.* cultivated *C. reinhardtii* heterotrophically in a 50-L tubular closed system. By starving the culture of sulfur, photosystem II was down-regulated to the point where oxygen consumption was greater than oxygen evolution, creating the necessary anaerobic conditions for biohydrogen production. However, the combination of uncontrolled solar intensities and nutrient deprivation can lead to photo-damage and the production of reactive oxygen species in microalgae, thus hindering production. To address this photoinhibition, Scoma *et al.* acclimated their cultures to natural light for 1 week before experimentation to induce physiological changes such as reduced chlorophyll, higher levels of xanthophylls, and increased photosynthetic and respiratory rates. When cells were allowed to acclimate to solar light, a more robust photoprotection system was observed, but biohydrogen production was still only equal to 20% of the laboratory results.

Recombinant protein production in *C. reinhardtii* has been demonstrated in a pilot commercial setting. Bovine Milk Amyloid A (MAA), a protein with anti-microbial properties, has previously been produced to high levels in *C. reinhardtii* in the laboratory (Manuell *et al.*, 2007). High accumulation of MAA in this strain required removal of the *psbA* gene, thus making the strain non-photosynthetic and unfit for outdoor growth. Recently, *psbA* was rescued using the alternative regulatory regions to eliminate issues of auto-attenuation (Gimpel *et al.*, 2014). The best rescued strain was then grown in 100-L hanging bags and accumulated MAA at approximately 3 mg L^{-1} of culture for a maximum yield of $11.8 \text{ g MAA kg}^{-1}$ dry weight biomass. Even with lower photosynthetic efficiency and lower total soluble protein production versus the knockout laboratory strain, the production of MAA in *C. reinhardtii* has the potential to be highly profitable since it can be administered as a solid algal pellet at a cost 60 times lower than traditional purified bovine colostrum MAA.

CONCLUSION

Chlamydomonas reinhardtii with its extensive research history and genetic tool infrastructure has been an excellent system to begin to understand algal metabolism and strain development. *Chlamydomonas reinhardtii*'s ease of transformation, large toolset, and natural variety has led to strains with significantly increased biofuels and bio-product yields. The biggest successes with *C. reinhardtii* have been the production of high-value products at laboratory scale in closed systems (Almaraz-Delgado *et al.*, 2013; Barrera and Mayfield, 2013; Gimpel *et al.*, 2014; Rasala and Mayfield, 2014). However, *C. reinhardtii* remains limited by its naturally low oil content, sensitivity to high solar irradiation and its weak growth in outdoor ponds, thus making it a relatively poor choice for large-scale biofuels production (James *et al.*, 2011; Scoma *et al.*, 2012; Schoepp *et al.*, 2014). While genetic engineering may help to overcome some of *C. reinhardtii*'s limitations, it will be essential to also expand the knowledge and tools developed in *C. reinhardtii* to other algal species.

Algae are the most diverse organisms on the planet, able to grow in almost any environment, from marine systems to hot springs to desert soil crusts and even sewer drains (Norton *et al.*, 1996; Dufresne *et al.*, 2008; Parker *et al.*, 2008; Tirichine and Bowler, 2011; Blunt *et al.*, 2012). Through these adaptations, strains have naturally developed favorable traits for specific manufacturing applications. Many wild-type species of algae will be economically viable as production platforms for bio-products with minimal strain optimization. For instance, cold-tolerant microorganisms have to maintain higher levels of polyunsaturated fatty acids to maintain membrane fluidity at colder temperatures. These extremophiles make obvious choices for sources of omega-3 fatty acids. To date, several natural species have already demonstrated robust growth in commercial-scale systems, mostly for nutraceuticals (Oncel, 2013). In addition, over the past 20 years, we have expanded the use of genetic tools to a wide variety of algal species, albeit with varying success. These tools will be needed to engineer an arsenal of pest-resistant strains with optimal growth rates and production capabilities as well as to utilize algae as a protein biomanufacturing platform. The combination of improved strain selection and engineering in more diverse species is our best chance of developing algal strains for large-scale renewable energy and sustainable recombinant protein production in the future.

Acknowledgments

This work is supported by the US Department of Energy (DEEE0003373) and the California Energy Commission Initiative for Large Molecule Sustainable Fuels (CILMSF 500-10-039).

References

- Almaraz-Delgado AL, Flores-Uribe J, Pérez-España VH, Salgado-Manjarrez E, Badillo-Corona JA. Production of therapeutic proteins in the chloroplast of *Chlamydomonas reinhardtii*. *AMB Express*. 2013; 4:57.
- Baldo FC, Inmaculada C, Rosa L, Herminia R, Vargas MÁ. Enhancement of carotenoids biosynthesis in *Chlamydomonas reinhardtii* by nuclear transformation using a phytoene synthase gene isolated from *Chlorella zofingiensis*. *Appl Microbiol Biotechnol*. 2011; 91:341–351. [PubMed: 21519934]
- Baltz A, Dang KV, Beyly A, Auroy P, Richaud P, Cournac L, Peltier G. Plastidial expression of type II NAD(P)H dehydrogenase increases the reducing state of plastoquinones and hydrogen

- photoproduction rate by the indirect pathway in *Chlamydomonas reinhardtii*. *Plant Physiol.* 2014; 165:1344–1352. [PubMed: 24820024]
- Barrera, DJ., Mayfield, SP. High-value recombinant protein production in microalgae. In: Richmond, A., Hu, Q., editors. *Handbook of Microalgal Culture: Applied Phycology and Biotechnology*. Oxford, UK: John Wiley & Sons, Ltd; 2013. p. 532-544.
- Blatti JL, Beld J, Behnke CA, Mendez M, Mayfield SP, Burkart MD. Manipulating fatty acid biosynthesis in microalgae for biofuel through protein-protein interactions. *PLoS One.* 2011; 7:e42949.
- Blunt JW, Copp BR, Keyzers RA, Munro MH, Prinsep MR. Marine natural products. *Nat Prod Rep.* 2012; 29:144–222. [PubMed: 22193773]
- Boyle N, Page M, Liu B, et al. Three acyltransferases and nitrogen-responsive regulator are implicated in nitrogen starvation-induced triacylglycerol accumulation in *Chlamydomonas*. *J Biol Chem.* 2012; 287:15811–15825. [PubMed: 22403401]
- Boynton JE, Gillham NW, Harris EH, et al. Chloroplast transformation in *Chlamydomonas* with high-velocity microprojectiles. *Science.* 1988; 240:1534–1538. [PubMed: 2897716]
- Brownbridge G, Azadi P, Smallbone A, Bhawe A, Taylor B, Kraft M. The future viability of algae-derived biodiesel under economic and technical uncertainties. *Bioresour Technol.* 2013; 151:166–173. [PubMed: 24220544]
- Cerutti H, Johnson AM, Gillham NW, Boynton JE. Epigenetic silencing of a foreign gene in nuclear transformants of *Chlamydomonas*. *Plant Cell.* 1997; 9:925–945. [PubMed: 9212467]
- Couso I, Vila M, Rodriguez H, Vargas MA. Overexpression of an exogenous phytoene synthase gene in the unicellular alga *Chlamydomonas reinhardtii* leads to an increase in the content of carotenoids. *Biotechnol Prog.* 2011; 27:54–60. [PubMed: 21312355]
- De Wilde C, Van Houdt H, De Buck S, Angenon G, De Jaeger G, Depicker A. Plants as bioreactors for protein production: avoiding the problem of transgene silencing. *Plant Mol Biol.* 2000; 43:347–359. [PubMed: 10999415]
- Del Campo JA, Garcia-Gonzalez M, Guerrero MG. Outdoor cultivation of microalgae for carotenoid production: current state and perspectives. *Appl Microbiol Biotechnol.* 2007; 74:1163–1174. [PubMed: 17277962]
- Demirbas A, Demirbas FM. Importance of algae oil as a source of biodiesel. *Energy Convers Manage.* 2011; 52:163–170.
- Deng XD, Gu B, Li YJ, Hu XW, Guo JC. The roles of acyl-CoA: diacylglycerol acyltransferase 2 genes in the biosynthesis of triacylglycerols by the green alga *Chlamydomonas reinhardtii*. *Mol Plant.* 2012; 5:945–947. [PubMed: 22511604]
- Dismukes CG, Carrieri D, Bennette N, Ananyev GM, Posewitz MC. Aquatic phototrophs: efficient alternatives to land-based crops for biofuels. *Curr Opin Biotechnol.* 2008; 19:235–240. [PubMed: 18539450]
- Dreesen IA, Charpin-El Hamri G, Fussenegger M. Heat-stable oral alga-based vaccine protects mice from *Staphylococcus aureus* infection. *J Biotechnol.* 2010; 145:273–280. [PubMed: 19995584]
- Dufresne A, Ostrowski M, Scanlan DJ, et al. Unraveling the genomic mosaic of a ubiquitous genus of marine cyanobacteria. *Genome Biol.* 2008; 9:R90. [PubMed: 18507822]
- Eichler-Stahlberg A, Weisheit W, Ruecker O, Heitzer M. Strategies to facilitate transgene expression in *Chlamydomonas reinhardtii*. *Planta.* 2009; 229:873–883. [PubMed: 19127370]
- Esquivel MG, Amaro HM, Pinto TS, Fevreiro PS. Efficient H₂ production via *Chlamydomonas reinhardtii*. *Trends Biotechnol.* 2011; 29:595–600. [PubMed: 21794941]
- Gao H, Wright DA, Li T, Wang YJ, Horken K, Weeks DP, Yang B, Spalding MH. TALE activation of endogenous genes in *Chlamydomonas reinhardtii*. *Algal Res.* 2014; 5:52–60.
- Georgianna DR, Mayfield SP. Exploiting diversity and synthetic biology for the production of algal biofuels. *Nature.* 2012; 488:329–335. [PubMed: 22895338]
- Georgianna DR, Michael JH, Marina M, Shuiqin W, Kyle B, Alex JL, James H, Michael M, Stephen PM. Production of recombinant enzymes in the marine alga *Dunaliella tertiolecta*. *Algal Res.* 2013; 2:2–9.
- Ghirardi ML, Posewitz MC, Maness PC. Hydrogenases and hydrogen photoproduction in oxygenic photosynthetic organisms*. *Annu Rev Plant Biol.* 2007; 58:71–91. [PubMed: 17150028]

- Gimpel JA, Hyun JS, Schoepp NG. Production of recombinant proteins in microalgae at pilot greenhouse scale. *Biotechnol Bioeng.* 2014; 112:339–345. [PubMed: 25116083]
- Goodenough U, Blaby I, Casero D, et al. The path to triacylglyceride obesity in the sta6 strain of *Chlamydomonas reinhardtii*. *Eukaryot Cell.* 2014; 13:591–613. [PubMed: 24585881]
- Goodson C, Roth R, Wang ZT, Goodenough U. Structural correlates of cytoplasmic and chloroplast lipid body synthesis in *Chlamydomonas reinhardtii* and stimulation of lipid body production with acetate boost. *Eukaryot Cell.* 2011; 10:1592–1606. [PubMed: 22037181]
- Gregory JA, Topol AB, Doerner DZ, Mayfield SP. Alga-produced cholera toxin-Pfs25 fusion proteins as oral vaccines. *Appl Environ Microbiol.* 2013; 79:3917–3925. [PubMed: 23603678]
- Griffiths MJ, Harrison STL. Lipid productivity as a key characteristic for choosing algal species for biodiesel production. *J Appl Phycol.* 2009; 21:493–507.
- Gumpel NJ, Rochaix JD, Purton S. Studies on homologous recombination in the green alga *Chlamydomonas reinhardtii*. *Curr Genet.* 1994; 26:438–442. [PubMed: 7702662]
- Hou QT, Qiu S, Liu Q, Tian J, Hu ZL, Ni JZ. Selenoprotein-transgenic *Chlamydomonas reinhardtii*. *Nutrients.* 2013; 5:624–636. [PubMed: 23443677]
- James, BD., Baum, GN., Perez, J., Baum, KN. Technoeconomic boundary analysis of biological pathways to hydrogen production. U.S. Dept. of Energy, Washington, DC; 2009. p. 207US Department of Energy, Office of Energy Efficiency & Renewable Energy ed. Subcontract report NREL/SR-560-46674
- James GO, Hocart CH, Hillier W, Chen H, Kordbacheh F, Price DG, Djordjevic MA. Fatty acid profiling of *Chlamydomonas reinhardtii* under nitrogen deprivation. *Bioresour Technol.* 2011; 102:3343–3351. [PubMed: 21146403]
- Jiang W, Zhou H, Bi H, Fromm M, Yang B, Weeks DP. Demonstration of CRISPR/Cas9/sgRNA-mediated targeted gene modification in *Arabidopsis*, tobacco, sorghum and rice. *Nucleic Acids Res.* 2013; 41:e188. [PubMed: 23999092]
- Jiang W, Brueggeman AJ, Horken KM, Plucinak TM, Weeks DP. Successful transient expression of Cas9 and gingle guide RNA genes in *Chlamydomonas reinhardtii*. *Eukaryot Cell.* 2014a; 13:1465–1469. [PubMed: 25239977]
- Jiang W, Cossey S, Rosenberg JN, Oyler GA, Weeks D. A rapid live-cell ELISA for characterizing antibodies against cell surface antigens of *Chlamydomonas reinhardtii* and its use in isolating algae from natural environments with related cell wall components. *BMC Plant Biol.* 2014b; 14:244. [PubMed: 25252698]
- Jones CS, Mayfield SP. Algae biofuels: versatility for the future of bioenergy. *Curr Opin Biotechnol.* 2011; 23:346–351. [PubMed: 22104720]
- Kanaar R, Hoeijmakers JHJ, van Gent DC. Molecular mechanisms of DNA double-strand break repair. *Trends Cell Biol.* 1998; 8:483–489. [PubMed: 9861670]
- Kato K, Marui T, Kasai S, Shinmyo A. Artificial control of transgene expression in *Chlamydomonas reinhardtii* chloroplast using the *lac* regulation system from *Escherichia coli*. *J Biosci Bioeng.* 2007; 104:207–213. [PubMed: 17964485]
- Kim DH, Kim YT, Cho JJ, Bae JH, Hur SB, Hwang I, Choi TJ. Stable integration and functional expression of flounder growth hormone gene in transformed microalga, *Chlorella ellipsoidea*. *Mar Biotechnol.* 2002; 4:63–73. [PubMed: 14961289]
- Kirst H, Garcia-Cerdan JG, Zurbriggen A, Melis A. Assembly of the light-harvesting chlorophyll antenna in the green alga *Chlamydomonas reinhardtii* requires expression of the *TLA2-CpFTSY* gene. *Plant Physiol.* 2012; 158:930–945. [PubMed: 22114096]
- Klok AJ, Lamers PP, Martens DE, Draaisma RB, Wijffels RH. Edible oils from microalgae: insights in TAG accumulation. *Trends Biotechnol.* 2014; 32:521–528. [PubMed: 25168414]
- La Russa M, Bogen C, Uhmeyer A, Doebbe A, Filippone E, Kruse O, Mussgnug JH. Functional analysis of three type-2 DGAT homologue genes for triacylglycerol production in the green microalga *Chlamydomonas reinhardtii*. *J Biotechnol.* 2012; 162:13–20. [PubMed: 22542934]
- Larson MA, Wei SH, Weber A, Mack DR, McDonald TL. Human serum amyloid A3 peptide enhances intestinal MUC3 expression and inhibits EPEC adherence. *Biochem Biophys Res Commun.* 2003; 300:531–540. [PubMed: 12504116]

- Leite GB, Abdelaziz AEM, Hallenbeck PC. Algal biofuels: challenges and opportunities. *Bioresour Technol.* 2013; 145:134–141. [PubMed: 23499181]
- Li Y, Han D, Hu G, Dauvillee D, Sommerfeld M, Ball S, Hu Q. *Chlamydomonas* starchless mutant defective in ADP-glucose pyrophosphorylase hyper-accumulates triacylglycerol. *Metab Eng.* 2010; 12:387–391. [PubMed: 20172043]
- Li T, Huang S, Zhao X, Wright DA, Carpenter S, Spalding MH, Weeks DP, Yang B. Modularly assembled designer TAL effector nucleases for targeted gene knockout and gene replacement in eukaryotes. *Nucleic Acids Res.* 2011; 39:6315–6325. [PubMed: 21459844]
- Mack DR, McDonald TL, Larson MA, Wei S, Weber A. The conserved TFLK motif of mammary-associated serum amyloid A3 is responsible for up-regulation of intestinal MUC3 mucin expression *in vitro*. *Pediatr Res.* 2003; 53:137–142. [PubMed: 12508093]
- Manuell AL, Beligni MV, Elder JH, Siefker DT, Tran M, Weber A, McDonald TL, Mayfield SP. Robust expression of a bioactive mammalian protein in *Chlamydomonas* chloroplast. *Plant Biotechnol J.* 2007; 5:402–412. [PubMed: 17359495]
- Melis A, Zhang L, Forestier M, Ghirardi ML. Sustained photobiological hydrogen gas production upon reversible inactivation of oxygen evolution in the green alga *Chlamydomonas reinhardtii*. *Plant Physiol.* 2000; 122:127–136. [PubMed: 10631256]
- Melis, A., Seibert, M., Ghirardi, ML. Hydrogen fuel production by transgenic microalgae. In: León, R.Galván Cejudo, A., Fernández, E., editors. *Transgenic Microalgae as Green Cell Factories*. New York: Springer New York; 2007. p. 110-121.
- Merchant SS, Kropat J, Liu B, Shaw J, Warakanont J. TAG, You're it! *Chlamydomonas* as a reference organism for understanding algal triacylglycerol accumulation. *Curr Opin Biotechnol.* 2011; 23:352–363. [PubMed: 22209109]
- Minai L, Wostrikoff K, Wollman F. Chloroplast biogenesis of photosystem II cores involves a series of assembly-controlled steps that regulate translation. *Plant Cell.* 2006; 18:159–175. [PubMed: 16339851]
- Moellering ER, Benning C. RNA interference silencing of a major lipid droplet protein affects lipid droplet size in *Chlamydomonas reinhardtii*. *Eukaryot Cell.* 2010; 9:97–106. [PubMed: 19915074]
- Molnar A, Bassett A, Thuenemann E, Schwach F, Karkare S, Ossowski S, Weigel D, Baulcombe D. Highly specific gene silencing by artificial microRNAs in the unicellular alga *Chlamydomonas reinhardtii*. *Plant J.* 2009; 58:165–174. [PubMed: 19054357]
- Msanne J, Xu D, Konda AR, Casas-Mollano JA. Metabolic and gene expression changes triggered by nitrogen deprivation in the photoautotrophically grown microalgae *Chlamydomonas reinhardtii* and *Coccomyxa sp. C-169*. *Phytochemistry.* 2012; 75:50–59. [PubMed: 22226037]
- Mulo P, Sakurai I, Aro EM. Strategies for *psbA* gene expression in cyanobacteria, green algae and higher plants: from transcription to PSII repair. *Biochim Biophys Acta.* 2012; 1817:247–257. [PubMed: 21565160]
- Näätäsaari L, Mistlberger B, Ruth C, Hajek T, Hartner FS, Glieder A. Deletion of the *Pichia pastoris* KU70 homologue facilitates platform strain generation for gene expression and synthetic biology. *PLoS One.* 2012; 7:e39720. [PubMed: 22768112]
- Nagarajan S, Chou SK, Cao S, Wu C, Zhou Z. An updated comprehensive techno-economic analysis of algae biodiesel. *Bioresour Technol.* 2013; 145:150–156. [PubMed: 23260269]
- Nelson JA, Lefebvre PA. Targeted disruption of the *NIT8* gene in *Chlamydomonas reinhardtii*. *Mol Cell Biol.* 1995; 15:5762–5769. [PubMed: 7565729]
- Norton TA, Melkonian M, Andersen RA. Algal biodiversity. *Phycologia.* 1996; 35:308–326.
- Oncel SS. Microalgae for a macroenergy world. *Renew Sustain Energy Rev.* 2013; 26:241–264.
- Oncel SS, Kose A, Faraloni C, Imamoglu E, Elibol M, Torzillo G, Sukan FV. Biohydrogen production using mutant strains of *Chlamydomonas reinhardtii*: the effects of light intensity and illumination patterns. *Biochem Eng J.* 2014; 92:47–52.
- Parker MS, Mock T, Armbrust EV. Genomic insights into marine microalgae. *Annu Rev Genet.* 2008; 42:619–645. [PubMed: 18983264]
- Peralta-Yahya PP, Ouellet M, Chan R. Identification and microbial production of a terpene-based advanced biofuel. *Nat Commun.* 2011; 2:483. [PubMed: 21952217]

- Rasala BA, Mayfield SP. The microalga *Chlamydomonas reinhardtii* as a platform for the production of human protein therapeutics. *Bioeng Bugs*. 2011; 2:50–54. [PubMed: 21636988]
- Rasala BA, Mayfield SP. Photosynthetic biomanufacturing in green algae; production of recombinant proteins for industrial, nutritional, and medical uses. *Photosynth Res*. 2014; 123:227–239. [PubMed: 24659086]
- Rasala BA, Muto M, Lee PA, Jager M, Cardoso RMF, Behnke CA, Kirk P, Hokanson CA, Crea R, Mendez M, et al. Production of therapeutic proteins in algae, analysis of expression of seven human proteins in the chloroplast of *Chlamydomonas reinhardtii*. *Plant Biotechnol J*. 2010; 8:719–733. [PubMed: 20230484]
- Rasala BA, Lee PA, Shen ZX, Briggs SP, Mendez M, Mayfield SP. Robust expression and secretion of Xylanase1 in *Chlamydomonas reinhardtii* by fusion to a selection gene and processing with the FMDV 2A Peptide. *PLoS One*. 2012; 7:e43349. [PubMed: 22937037]
- Rasala B, Barrera D, Ng J, Plucinak T, Rosenberg J, Weeks D, Oyler G, Peterson T, Haerizadeh F, Mayfield S. Expanding the spectral palette of fluorescent proteins for the green microalga *Chlamydomonas reinhardtii*. *Plant J*. 2013a; 74:545–556. [PubMed: 23521393]
- Rasala, BA., Gimpel, JA., Tran, M., Hannon, MJ. Genetic engineering to improve algal biofuels production. In: Borowitzka, MA., Moheimani, NR., editors. *Algae for Biofuels and Energy*. Dordrecht, Netherlands: Springer Netherlands; 2013b. p. 99–113.
- Rosales-Mendoza S, Paz-Maldonado LM, Soria-Guerra RE. *Chlamydomonas reinhardtii* as a viable platform for the production of recombinant proteins: current status and perspectives. *Plant Cell Rep*. 2012; 31:479–494. [PubMed: 22080228]
- Schmollinger S, Strenkert D, Schroda M. An inducible artificial microRNA system for *Chlamydomonas reinhardtii* confirms a key role for heat shock factor 1 in regulating thermotolerance. *Curr Genet*. 2010; 56:383–389. [PubMed: 20449593]
- Schoepp NG, Stewart RL, Sun V, Quigley AJ. System and method for research-scale outdoor production of microalgae and cyanobacteria. *Bioresour Technol*. 2014; 166:273–281. [PubMed: 24926599]
- Schroda M, Blocker D, Beck CF. The *HSP70A* promoter as a tool for the improved expression of transgenes in *Chlamydomonas*. *Plant J*. 2000; 21:121–131. [PubMed: 10743653]
- Scoma A, Giannelli L, Faraloni C, Torzillo G. Outdoor H₂ production in a 50-L tubular photobioreactor by means of a sulfur-deprived culture of the microalga *Chlamydomonas reinhardtii*. *J Biotechnol*. 2012; 157:620–627. [PubMed: 21771618]
- Siaut M, Cuiné S, Cagnon C, Fessler B. Oil accumulation in the model green alga *Chlamydomonas reinhardtii*: characterization, variability between common laboratory strains and relationship with starch reserves. *BMC Biotechnol*. 2011; 11:7. [PubMed: 21255402]
- Singh A, Nigam PS, Murphy JD. Mechanism and challenges in commercialisation of algal biofuels. *Bioresour Technol*. 2011; 102:26–34. [PubMed: 20609580]
- Sodeinde OA, Kindle KL. Homologous recombination in the nuclear genome of *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci USA*. 1993; 90:9199–9203. [PubMed: 8415677]
- Specht E, Mayfield SP. Synthetic oligonucleotide libraries reveal novel regulatory elements in *Chlamydomonas* chloroplast mRNAs. *ACS Synth Biol*. 2012; 2:34–46. [PubMed: 23656324]
- Su ZL, Qian KX, Tan CP, Meng CX, Qin S. Recombination and heterologous expression of allophycocyanin gene in the chloroplast of *Chlamydomonas reinhardtii*. *Acta Biochim Biophys Sin*. 2005; 37:709–712. [PubMed: 16215639]
- Terashima M, Freeman ES, Jinkerson RE, Jonikas MC. A fluorescence-activated cell sorting-based strategy for rapid isolation of high-lipid *Chlamydomonas* mutants. *Plant J*. 2014; 81:147–159. [PubMed: 25267488]
- Tirichine L, Bowler C. Decoding algal genomes: tracing back the history of photosynthetic life on Earth. *Plant J*. 2011; 66:45–57. [PubMed: 21443622]
- Torzillo, G., Seibert, M. Hydrogen production by *Chlamydomonas reinhardtii*. In: Richmond, A., Hu, Q., editors. *Handbook of Microalgal Culture: Applied Phycology and Biotechnology*. 2nd. Oxford, UK: John Wiley & Sons, Ltd; 2013. p. 417–432.

- Torzillo G, Scoma A, Faraloni C, Ena A. Increased hydrogen photoproduction by means of a sulfur-deprived *Chlamydomonas reinhardtii* D1 protein mutant. *Int J Hydrogen Energy*. 2009; 34:4529–4536.
- Tran M, Henry RE, Siefker D, Van C, Newkirk G, Kim J, Bui J, Mayfield SP. Production of anti-cancer immunotoxins in algae: ribosome inactivating proteins as fusion partners. *Biotechnol Bioeng*. 2013a; 110:2826–2835. [PubMed: 23719862]
- Tran M, Van C, Barrera DJ, Pettersson PL, Peinado CD, Bui J, Mayfield SP. Production of unique immunotoxin cancer therapeutics in algal chloroplasts. *Proc Natl Acad Sci USA*. 2013b; 110:E15–E22. [PubMed: 23236148]
- Ugwu CU, Aoyagi H, Uchiyama H. Photobioreactors for mass cultivation of algae. *Bioresour Technol*. 2008; 99:4021–4028. [PubMed: 17379512]
- Venter M. Synthetic promoters: genetic control through cis engineering. *Trends Plant Sci*. 2007; 12:118–124. [PubMed: 17292658]
- Vinyard DJ, Gimpel J, Ananyev GM, Cornejo MA, Golden SS, Mayfield SP, Dismukes GC. Natural variants of photosystem II subunit D1 tune photochemical fitness to solar intensity. *J Biol Chem*. 2013; 288:5451–5462. [PubMed: 23271739]
- Vinyard DJ, Gimpel J, Ananyev GM. Engineered photosystem II reaction centers optimize photochemistry versus photoprotection at different solar intensities. *J Am Chem Soc*. 2014; 136:4048–4055. [PubMed: 24548276]
- Walsh G. Biopharmaceutical benchmarks 2014. *Nat Biotechnol*. 2014; 32:992–1000. [PubMed: 25299917]
- Wang ZT, Ullrich N, Joo S, Waffenschmidt S. Algal lipid bodies: stress induction, purification, and biochemical characterization in wild-type and starchless *Chlamydomonas reinhardtii*. *Eukaryot Cell*. 2009; 8:1856–1868. [PubMed: 19880756]
- Work VH, Radakovits R, Jinkerson RE, Meuser JE. Increased lipid accumulation in the *Chlamydomonas reinhardtii* *sta7–10* starchless isoamylase mutant and increased carbohydrate synthesis in complemented strains. *Eukaryot Cell*. 2010; 9:1251–1261. [PubMed: 20562225]
- Wu-Scharf D, Jeong B, Zhang C, Cerutti H. Transgene and transposon silencing in *Chlamydomonas reinhardtii* by a DEAH-box RNA helicase. *Science*. 2000; 290:1159–1162. [PubMed: 11073454]
- Xie B, Stessman D, Hart JH, Dong H, Wang Y, Wright DA, Nikolau BJ, Spalding MH, Halverson LJ. High throughput fluorescence activated cell sorting for lipid hyperaccumulating *Chlamydomonas reinhardtii* mutants. *Plant Biotechnol J*. 2014; 12:872–882. [PubMed: 24702864]
- Xu L, Wang Q, Wu S, Li D. Improvement of hydrogen yield of *Iba*-transgenic *Chlamydomonas reinhardtii* caused by increasing respiration and impairing photosynthesis. *Int J Hydrogen Energy*. 2014; 39:13347–13352.
- Yang ZQ, Li YN, Chen F, Li D, Zhang ZF, Liu YX, Zheng DX, Wang Y, Shen GF. Expression of human soluble TRAIL in *Chlamydomonas reinhardtii* chloroplast. *Chin Sci Bull*. 2006; 51:1703–1709.
- Yoon SMM, Kim SY, Li KF, Yoon BH, Choe S, Kuo MM. Transgenic microalgae expressing *Escherichia coli* AppA phytase as feed additive to reduce phytate excretion in the manure of young broiler chicks. *Appl Microbiol Biotechnol*. 2011; 91:553–563. [PubMed: 21533578]
- Zhang YK, Shen GF, Ru BG. Survival of human metallothionein-2 transplastomic *Chlamydomonas reinhardtii* to ultraviolet B exposure. *Acta Biochim Biophys Sin (Shanghai)*. 2006; 38:187–193. [PubMed: 16518543]
- Zhang R, Patena W, Armbruster U, Gang SS, Blum SR, Jonikas MC. High-throughput genotyping of green algal mutants reveals random distribution of mutagenic insertion sites and endonucleolytic cleavage of transforming DNA. *Plant Cell*. 2014; 26:1398–1409. [PubMed: 24706510]
- Zhao T, Wang W, Bai X, Qi Y. Gene silencing by artificial microRNAs in *Chlamydomonas*. *Plant J*. 2009; 58:157–164. [PubMed: 19054364]




Figure 1. Steps to engineering a production strain of green algae. Genetic manipulation using advanced tools will result in a desired phenotype. The isolated strain will be tested for scale-up and reanalyzed for characteristics such as growth rate, population density, recombinant protein accumulation, and lipid profile. If these traits do not overcome economic constraints of production, additional genetic manipulation in the laboratory is required. Once an ideal strain is identified, fuels and co-products can be extracted from scaled-up cultures.

Table 1

Biomass production of various algal species in an outdoor setting compared to average reported lipid content

Species	Closed system production (g L ⁻¹ day ⁻¹)	Open system production (g L ⁻¹ day ⁻¹)	Lipid content (nutrient-replete, % dry weight)
<i>Scenedesmus dimorphus</i>	0.095	0.090	26
<i>Chlorella vulgaris</i>	0.047	0.035	25
<i>Chlamydomonas reinhardtii</i>	0.078	0.015	21
<i>Arthrospira platensis</i>	0.040	0.018	13
<i>Anabaena</i> sp.	0.055	–	5
<i>Porphyridium purpureum</i>	0.074	0.036	11
<i>Nannochloropsis salina</i>	0.043	0.028	27
<i>Dunaliella tertiolecta</i>	0.039	0.031	15
<i>Phaeodactylum tricorutum</i>	0.021	–	21

Biomass production data from Schoepp *et al.* (2014) and lipid content from Griffiths and Harrison (2009).