

Emerging pharmaceutical therapies for COPD

Sowmya P Lakshmi^{1,2}

Aravind T Reddy^{1,2}

Raju C Reddy^{1,2}

¹Department of Medicine, Division of Pulmonary, Allergy and Critical Care Medicine, University of Pittsburgh School of Medicine, ²Veterans Affairs Pittsburgh Healthcare System, Pittsburgh, PA, USA

Abstract: COPD, for which cigarette smoking is the major risk factor, remains a worldwide burden. Current therapies provide only limited short-term benefit and fail to halt progression. A variety of potential therapeutic targets are currently being investigated, including COPD-related proinflammatory mediators and signaling pathways. Other investigational compounds target specific aspects or complications of COPD such as mucus hypersecretion and pulmonary hypertension. Although many candidate therapies have shown no significant effects, other emerging therapies have improved lung function, pulmonary hypertension, glucocorticoid sensitivity, and/or the frequency of exacerbations. Among these are compounds that inhibit the CXCR2 receptor, mitogen-activated protein kinase/Src kinase, myristoylated alanine-rich C kinase substrate, selectins, and the endothelin receptor. Activation of certain transcription factors may also be relevant, as a large retrospective cohort study of COPD patients with diabetes found that the peroxisome proliferator-activated receptor γ (PPAR γ) agonists rosiglitazone and pioglitazone were associated with reduced COPD exacerbation rate. Notably, several therapies have shown efficacy only in identifiable subgroups of COPD patients, suggesting that subgroup identification may become more important in future treatment strategies. This review summarizes the status of emerging therapeutic pharmaceuticals for COPD and highlights those that appear most promising.

Keywords: pulmonary, PPAR, phosphodiesterase, emphysema, cigarette, mucus

Introduction

COPD affects ~200 million people worldwide¹ and is the third leading cause of death in the US, claiming ~150,000 lives in 2013.² Approximately 6.3% of US adults aged ≥ 18 years have COPD,³ and its combined indirect and direct costs in 2010 surpassed \$52 billion. More effective treatments for COPD are needed to address this serious health problem. This review summarizes the status of potential therapies currently or recently in clinical trials and highlights those that appear most promising.

COPD pathology

COPD, a complex and heterogeneous chronic progressive inflammatory disease of the distal airways characterized by persistent airflow limitations, typically results from inhaling noxious gases and particles, especially cigarette smoke.⁴⁻⁶ The resulting inflammatory response involves increased numbers of neutrophils, macrophages, B cells, and CD4⁺ and CD8⁺ T lymphocytes in the airways.⁷ Histopathological changes include edema, loss of alveoli, and tissue remodeling involving smooth muscle hypertrophy and fibrosis, which are associated with bronchoconstriction and airway mucus hypersecretion exacerbated by reduced clearance.⁵⁻⁸ The accompanying systemic inflammation and increased reactive oxygen species (ROS) in patients may contribute to such manifestations as cardiovascular complications, loss of skeletal muscle, osteoporosis, and psychosocial effects.

Correspondence: Raju C Reddy
Department of Medicine, Division of Pulmonary, Allergy and Critical Care Medicine, University of Pittsburgh School of Medicine, 3459 Fifth Avenue, Pittsburgh, PA 15213, USA
Tel +1 412 360 6823
Fax +1 412 360 1919
Email reddyrc@upmc.edu

COPD exacerbations

COPD exacerbations can significantly accelerate lung function decline and overall health. In patients with severe COPD, acute exacerbations requiring hospitalization account for up to 50% of COPD health care costs. Exacerbation risk appears to be associated with serum eosinophilia: in patients with clinical COPD and serum eosinophils above 0.34×10^9 cells/L, risk of severe and moderate exacerbations were increased compared to those in patients with fewer eosinophils.⁹ Causes of COPD exacerbations appear multifactorial and elusive but may include bacterial or viral infections, exposure to environmental pollutants, and unidentified factors.⁵ Exacerbations are accompanied by a rapid rise of proinflammatory cytokines and chemokines that trigger and sustain rapid influx of neutrophils and their products such as neutrophil elastase into the airways.^{10,11} Reduction in the frequency of COPD exacerbation is a primary or secondary end point of several current clinical trials of emerging therapies.^{12,13}

Treatment of stable COPD

The American Thoracic Society and other leading pulmonary disease organizations recommend that COPD patients cease smoking and use one or more inhaled bronchodilators.¹⁴ The wide choice of inhaled bronchodilators includes short-acting beta₂-agonists (SABA) and long-acting beta₂-agonists (LABA), short-acting muscarinic antagonists (SAMA) and long-acting muscarinic antagonists (LAMA), and combinations of β -agonists and antimuscarinic agents. Inhaled corticosteroids (ICS) and combinations of corticosteroids with other agents may also be used. Oral medications, including methylxanthines and a phosphodiesterase 4 (PDE4) inhibitor, may also be used as adjunctive agents or, in the case of systemic corticosteroids, for acute exacerbations.

The Global Initiative for Chronic Obstructive Lung Disease (GOLD) recently categorized COPD patients into four groups, in which the familiar spirometry-based grades (GOLD I–IV) are supplemented by patients' symptom burden (assessed by the modified British Medical Research Council questionnaire or COPD Assessment Test) and exacerbation history: spirometry and exacerbation history are jointly used to determine the risk of adverse health events.⁵ Groups A and C have relatively mild symptoms, while Groups A and B have relatively low risk. GOLD recommends that Groups A and B receive SABA, LABA, SAMA, or LAMA bronchodilators. It recommends that Groups C and D also receive ICS or possibly the PDE4 inhibitor roflumilast. Despite

symptomatic relief and reduced rates of exacerbations with current medications, long-term studies of treated COPD patients suggest that medications do not halt the decline of lung function⁴ and resulting mortality, although there is limited evidence that fluticasone and/or salmeterol may slow the rate of decline.^{15,16} Several groups are pursuing identification of biomarkers for COPD subgroups based on either the etiological agent/event triggering the exacerbation or the physiological targets that may indicate response to a given therapy.^{6,17} For example, COPD patients with higher levels of exhaled nitric oxide (FeNO) had a greater probability of the enhanced eosinophilic infiltration¹⁸ that is associated with a greater response to corticosteroids.^{19–21} These studies carry significant implications both for current treatment and for the development of novel therapies that may be targeted to specific groups of patients.

Inflammation in COPD

Smoke, from cigarettes, biomass fuel, or air pollution, is a common inducer of inflammation in COPD. The inhaled irritants provoke significant oxidative stress and activate inflammatory cells.^{22,23} Excess neutrophils and macrophages, along with T cells, B cells, and dendritic cells, infiltrate the peripheral airways,^{24–26} concurrent with the narrowing or obliteration of small bronchioles.²⁷ A number of inflammatory markers are significantly elevated in the serum of patients with COPD, notably including CD40 ligand, epidermal growth factor (EGF), brain-derived neurotrophic factor (BDNF), a variety of acute-phase proteins, and neutrophil-associated proteins.²⁸ These molecules may represent targets for future COPD therapies.

Glucocorticoid (GC) resistance

COPD is relatively resistant to modulation by corticosteroids, as even high doses of GCs do not delay or inhibit COPD progression. Histological analysis of lung tissue removed during lung volume reduction surgery of COPD patients indicated that GC treatment significantly reduced the number of airways with lymphoid follicles but had no long-term effect on luminal occlusion or airway wall thickening.²⁹

Corticosteroids suppress inflammation both by activating transcription of anti-inflammatory and suppressing transcription of proinflammatory genes.³⁰ Binding of corticosteroids to the classic glucocorticoid receptor (GR α) leads to activation, which allows release from its cytoplasmic anchor. The receptor is then acetylated, allowing binding to response elements in the promoter elements of target

genes, and recruitment of coactivators such as CREB-binding protein, with consequent transcriptional activation. GC-mediated gene repression requires histone deacetylase 2 (HDAC2) at two distinct steps.³¹ First, this acetyl group must be removed by HDAC2 to allow binding to proinflammatory transcription factors such as nuclear factor- κ B (NF- κ B). Second, activated GR recruits HDAC2, which then deacetylates histones and thus inactivates transcription. Alveolar macrophages (AMs), lungs, and bronchi of COPD patients express lower levels of HDAC2 than those in healthy controls,³² however, and theophylline treatment, which enhances HDAC activity in AMs, also restores AM GC sensitivity.³³ These effects may underlie the reported ability of theophylline treatment (200–400 mg/day, depending on weight), added to the LABA formoterol and the GC budesonide in a 50-patient (study completion) single-blinded, placebo-controlled study, to improve 6-minute walk distance and forced expiratory volume in 1 second (FEV₁).³⁴ However, a double-blind, placebo-controlled study in 46 per-protocol patients found no effect of low-dose (100 mg bid) theophylline added to fluticasone plus salmeterol on HDAC activity, inflammatory biomarkers, or frequency of exacerbation.³⁵

Observed COPD-associated reductions in HDAC2 are believed to result from oxidative and nitrosative stress,³² and levels of the antioxidant transcription factor NF-E2-related factor 2 (Nrf2) are similarly reduced in AMs of COPD patients.³⁶ Treatment with the Nrf2 activator sulforaphane, similar to theophylline treatment, increases HDAC2 activity and restores GC sensitivity.³⁷

A number of other mechanisms of GC resistance have also been suggested.³⁰ Among these may be increased expression of the translationally inactive GR β , although this remains controversial³⁰ and GR expression has been found to be reduced in lungs of healthy smokers and COPD patients.³⁸ Another possibility is posttranslational modifications of GR α by phosphorylation or nitrosylation.³⁹ Such phosphorylation may be due to p38 mitogen-activated protein kinase (MAPK) activation, and p38 MAPK inhibitors have been shown to reduce the GC resistance seen in some patients with severe asthma.⁴⁰

Restoration of GC sensitivity is an attractive strategy for the development of novel pharmaceutical therapies. As previously noted, the ability of theophylline treatment to accomplish this is controversial.^{34,35} However, roflumilast has been shown *ex vivo* to improve GC sensitivity of neutrophils from COPD patients,⁴¹ and subgroup analysis of two clinical trials showed that patients receiving ICS were among those particularly likely to experience a reduction of exacerbations in roflumilast treatment.⁴² A subsequent 1-year trial showed that roflumilast treatment reduced exacerbations in patients also receiving ICS and LABA.¹³

Proinflammatory cytokines and chemokines

Numerous proinflammatory cytokines and chemokines are significantly higher in COPD patients than in healthy controls,⁴³ and multiple biologics and small molecules that target these mediators are under investigation (Tables 1 and 2). Chemokines significantly contribute to inflammation

Table 1 Developmental status of chemokine receptor inhibitors for COPD

Mediator	Role in COPD	Drug	Clinical development	References
CCR1	Receptor for CCL3 (MIP-1 α) and chemoattractant for inflammatory cells. CCR1 is also among receptors binding CCL5 and CCL7.	AZD4818	AZD4818 (4-week treatment) provided no significant benefit to COPD patients (NCT00629239).	46
CCR2	Receptor for CCL2 (MCP-1), which recruits inflammatory cells to lungs in COPD. Increases synthesis of MUC5AC and MUC5B.	AZD2423	AZD2423 (28-day treatment) in DBPCRT (NCT01215279); study has completed but statistical analysis not released.	47, 48
CXCR2 (IL8RB)	Activated by CXCL8 (IL-8), which is higher in BAL and sputum of COPD patients. CXCL8 is chemotactic for neutrophils.	Navarixin (MK-7123); AZD5069	MK-7123 (navarixin; 6-month treatment) in DBPCRT showed statistically significant improvement in postbronchodilator FEV ₁ (NCT01006616 and NCT00441701). AZD5069 (4-week treatment) in DBPCRT in patients with moderate-to-severe COPD reduced blood neutrophil counts with no increased infection (NCT01233232).	44, 45

Abbreviations: DBPCRT, double-blind, placebo-controlled, randomized trial; IL-8, interleukin 8.

Table 2 Developmental status of cytokine inhibitors for COPD

Mediator	Role in COPD	Drug	Clinical development	References
IL-1	Promotes proinflammatory responses. Elevated in stable COPD and further increased in exacerbations.	Canakinumab, a human anti-IL-1 β monoclonal antibody	A phase I/II RDBPCES of canakinumab (NCT00581945) (45-week treatment), no statistical analysis provided for lung function changes.	49, 50
IL-5	Mediates eosinophil maturation and mobilization; eosinophils increased during some exacerbations.	Mepolizumab (MAB against IL-5), benralizumab (MEDI-563; MAB against IL-5R α)	Mepolizumab (26–52-week treatment) tested as adjunct in DBPCRT targeting COPD exacerbation rate, studies completed, but results not posted (NCT02105948, NCT01463644, and NCT02105961). Benralizumab (\leq 56-week treatment) has completed a trial for moderate-to-severe COPD (NCT01227278), but no evidence of efficacy was observed; studies for exacerbation reduction and other effectiveness measures (NCT02155660 and NCT02138916) are ongoing.	9, 51
IL-13	Plasma but not sputum concentrations inversely correlated with FEV ₁ in COPD. IL-13 induces goblet cell hyperplasia and mucus hypersecretion.	Lebrikizumab, a humanized anti-IL-13 MAB	There is a study of lebrikizumab (24-week treatment) for decline in frequency of COPD exacerbations and lung function (NCT02546700).	100–102
IL-17A	One study found IL-17 reduced in sputum of severe COPD patients but another found numbers of CD4 ⁺ Th17 cells in the airways correlated with airflow limitations.	CNTO6785	CNTO6785 (12-week treatment) is being investigated in moderate-to-severe COPD in DBPCRT (NCT01966549). No results reported yet.	43, 53, 58
Tumor necrosis factor	Higher levels in sputum and serum of COPD patients; augments inflammation.	Infliximab, etanercept	Infliximab (6-month treatment) (NCT00056264) showed no clinical benefit but toxicity – higher rate of pneumonia and malignancies; however, difference in malignancy rate diminished greatly on long-term follow-up (NCT00380796), making the results difficult to interpret. Etanercept (90-day treatment) (NCT00789997) was not more efficacious than prednisone for the treatment of exacerbations.	28, 103–105

Abbreviations: IL, interleukin; RDBPCES, randomized, double-blind, placebo-controlled, exploratory study; MAB, monoclonal antibody; DBPCRT, double-blind, placebo-controlled, randomized trial; FEV₁, forced expiratory volume in 1 second; Th17, T helper 17.

by attracting neutrophils and other inflammation-related cells, and an antagonist of the CXCR2 receptor for interleukin (IL)-8 and other chemokines⁴⁴ has been found beneficial in COPD patients. A different CXCR2 inhibitor reduced blood neutrophil counts with no increased rate of infection,⁴⁵ but compounds blocking CCR1⁴⁶ and CCR2^{47,48} (NCT01215279) had no effect.

IL-1 expression was significantly higher in lung and sputum from COPD patients compared to non-COPD controls.⁴⁹ However, treatment with a monoclonal antibody (MAB) inhibiting IL-1 β (canakinumab)⁵⁰ remains an open question in COPD patients (Table 2).

IL-5 is largely produced by eosinophils and is therefore low in typical COPD. A subset of COPD patients have elevated eosinophil numbers and IL-5 levels in their blood and airways; these patients are at increased risk for acute exacerbations.^{9,51} Two MABs targeted to IL-5 have been investigated for their ability to reduce exacerbation rate in COPD patients, but one showed no evidence of efficacy and results for the other are not yet available (Table 2).

Secretion of IL-17A is a canonical marker of T helper 17 (Th17) lymphocytes, which are distinct from Th1 and Th2 cells. However, most of the cells secreting IL-17 in patients with very severe COPD were identified as mast

cells.⁵² This study also found that levels of IL-17A in lungs of COPD patients correlated with functional decline, with elevations becoming significant in patients with severe and very severe disease. Another study found that the numbers of CD4⁺IL-17⁺ cells in the alveolar walls and small airways of COPD patients and CD4⁺IL-17⁺ numbers in small airways positively correlated with airflow limitations.⁵³ These results are compatible with the finding that Th17 cells were present in the lungs of patients with emphysema but not normal controls and that IL-17A-induced secretion of the elastin-degrading enzyme matrix metalloprotease 12 by lung macrophages.⁵⁴ The levels of IL-17A were also elevated in the sputum of patients with acute COPD exacerbations associated with *Haemophilus influenzae* infection but not other acute exacerbations.⁵⁵ Other studies have not similarly distinguished among causes of acute exacerbations, likely accounting for findings that the ratio of regulatory T cells to IL-17 levels in peripheral blood was similar in COPD patients with and without current acute exacerbations, although exacerbations significantly increased levels of transforming growth factor β (TGF- β).⁵⁶ Indeed, one study was unable to detect IL-17 in sputum or serum of COPD patients with or without exacerbations.⁵⁷ Another study found that the sputum of patients with severe COPD had significantly higher levels of IL-8 but 4.8-fold lower levels of IL-17 compared to that of patients with mild COPD and healthy controls.⁵⁸ T cells from many COPD patients have also been reported to produce less IL-17A and IL-22 (also a signature cytokine of Th17 cells) than those of most normal smokers.⁵⁹ These complex and apparently contradictory findings underline the uncertainty of the role of IL-17 in COPD. Nevertheless, an IL-17 modulator is currently in clinical trials for COPD (Table 2).

Based on an IL-18-overexpressing transgenic mouse model that develops emphysema and airway remodeling,⁶⁰ Kang et al⁶⁰ and Nakajima and Owen⁶¹ proposed that IL-18 is a master regulator of lung pathology in COPD. A phase I safety and tolerability clinical trial (NCT01322594) of the MAb MEDI2338 (targeted to IL-18) in COPD patients found no serious adverse events, but there have been no efficacy studies.⁶²

Tumor necrosis factor α (TNF- α) plays multiple roles in COPD inflammatory pathology, and the levels of interferon γ (IFN γ) and TNF- α in the intraepithelial T cells from bronchi of COPD patients with GOLD II–III disease showed a significant negative correlation with FEV₁.⁶³ Nevertheless, studies with infliximab showed no clinical benefits on FEV₁, dyspnea, or exacerbations and were associated with higher rates of pneumonia and malignancy (Table 2). Similarly, treatment with etanercept, a fusion protein that competitively

binds TNF- α , was not superior to prednisone in COPD exacerbations and in fact was less effective among patients with eosinophilia (Table 2).

Taken together, these data show that an increased level of a specific cytokine or chemokine during COPD exacerbations or stable COPD does not necessarily predict the efficacy of its specific inhibitor in COPD patients. Whether modulators of specific cytokines or chemokines can provide improved efficacy in a subgroup of patients is a possibility and warrants further investigation.⁶

Signaling molecules

Multiple signaling molecules help regulate inflammation and airway remodeling and represent plausible targets for the development of therapeutic candidates. Candidate drugs include inhibitors of p38 MAPK and related kinases, phosphoinositide kinase δ (PI3K δ), leukotriene B₄, selectins, and vasoactive intestinal peptide (Table 3). Although several oral and inhaled p38 MAPK inhibitors have been discontinued, the inhaled narrow spectrum kinase (p38 α + Src family) inhibitor JNJ49095397 (previously RV568) shows promising activity in COPD patients; conference presentations have indicated that RV568 significantly increased FEV₁ and inhibited IL-1 β (90% at 800 μ g dose) and CXCL8 expression (73%).⁶⁴ However, a recent conference report performed in over 200 COPD patients (half placebo, half 400 μ g dose) showed no benefit with RV568 on lung function or EXACT-PRO.⁶⁵ PI3K δ participates in many functions of lymphoid and myeloid cells: B-cell development, migration and activation of natural killer (NK) cells and T cells, neutrophil oxidative burst, macrophage activation triggered by immune complexes, and degranulation and maturation of mast cells.⁶⁶ Specific PI3K δ inhibitors are being developed,⁶⁷ and studies on the effects of such inhibitors in COPD are in progress. Efficacy data remain limited, however (Table 3). Selectins are essential for migration of inflammatory cells from the bloodstream into pulmonary tissue; the selectin modulator bimosiamose reduced inflammation in COPD patients and thus warrants further testing (Table 3).⁶⁸

Similar to the glucocorticoid receptor, peroxisome proliferator-activated receptor γ (PPAR γ), also a member of the nuclear hormone receptor superfamily, exerts potent antioxidant and anti-inflammatory effects via multiple mechanisms, including downregulation of NF- κ B and other proinflammatory transcription factors. Lung tissue and bronchial epithelial cells from COPD patients express significantly lower levels of PPAR γ than those of nonsmoking controls.⁶⁹ In vitro treatment of COPD patient and control bronchial epithelial cells

Table 3 Developmental status of proinflammatory signaling pathway inhibitors for COPD

Mediator	Role in COPD	Drug	Clinical development	References
MAPK (p38 mitogen-activated protein kinase)	Higher MAPK in lungs activates proinflammatory response in AMs and lymphocytes.	Acumapimod, dilmapimod (SB-681323), losmapimod, PH-797804, GSK-610677, AZD-7624	Acumapimod (BCT197A2201) reportedly remains in active development, although phase II results have not been reported and there are no current clinical trials. Dilmapimod (SB-681323) significantly reduced TNF- α production in COPD (NCT00144859) but its development was discontinued. Losmapimod did not induce a significant improvement in exercise tolerance or lung function (NCT01218126) and was discontinued. PH-797804 (6-week treatment) (NCT00559910) significantly improved lung function and dyspnea in moderate-to-severe COPD in DBPCRT but was discontinued. Development of GSK-610677 was discontinued following an unreported phase I trial (NCT00694902). Results of a phase II study of AZD-7624 (NCT02238483) have not been reported.	64, 106–109
Narrow spectrum kinase inhibitors	p38 α and Src family kinases such as Hck are involved in the production of proinflammatory cytokines from macrophages, smooth muscle cells, and human airway epithelial cells. Cigarette smoke activates c-Src and augments airway inflammation and destruction of lung tissue.	JNJ49095397/RV568	RV568 (14-day inhaled treatment) significantly increased FEV ₁ and reduced sputum malondialdehyde and serum myeloperoxidase in COPD patients. A recent conference report, however, in over 200 COPD patients showed no benefit with RV568 for 12 weeks with respect to lung function or EXACT-PRO (NCT01867762, NCT01475292, and NCT01661244).	64, 66, 110
PI3K δ (phosphoinositide-3-kinase δ)	Involved in maturation and effector functions of B cells and other leukocytes.	GSK2269557, RV1729	GSK2269557 (treatment up to 84 days in NCT02522299 or 28 days in NCT02294734) DBPCRTs in patients with acute exacerbations of COPD in progress. RV1729 (treatment up to 28 days) is being tested in NCT02140346 with limited efficacy data being gathered in a predominantly phase I study.	66, 67
IKK (inhibitor of nuclear factor kappa-B kinase)	IKK is an upstream activator of the proinflammatory transcription factor NF- κ B. IKK α and IKK β activity is elevated in patients with COPD.	IMD-1041, an IKK β inhibitor	IMD-1041 in DBPCRT (NCT00883584) has no follow-up information posted since April 2009; unclear whether study was performed.	111
LTB ₄ receptor	LTB ₄ levels are elevated in sputum, breath, and BAL of COPD patients; highest LTB ₄ levels seen in exacerbations. LTB ₄ is chemotactic for neutrophils and T cells. AMs bearing the BLT1 receptor are more common in COPD patients.	BIIL 284	BIIL 284 (12-week treatment) was assessed for effects on lung function, exercise endurance, sputum, and safety in COPD patients (NCT02249247); a 14-day study assessed effects on biomarkers (NCT02249338) – results of neither study have yet been published. Other LTB ₄ receptor antagonists have not demonstrated beneficial results.	4, 112

(Continued)

Table 3 (Continued)

Mediator	Role in COPD	Drug	Clinical development	References
5-LO (5-lipoxygenase)	Involved in synthesis of leukotrienes.	5-Lipoxygenase inhibitor (zileuton) used clinically for asthma	Zileuton (14-day treatment) reduced urinary LTE ₄ levels in hospitalized COPD patients with acute exacerbations in DBPCRT (NCT00493974) but did not significantly shorten stay or reduce treatment failure.	113
Prostaglandin D ₂ receptor or chemoattractant receptor-homologous molecule expressed on Th2 (CRTh2) receptor	Highly expressed on eosinophils, basophils, Th2 (not Th1) lymphocytes, and subset of monocytes. Blocking this receptor inhibits chemotaxis of these cells. CRTh2 is expressed on mucosal epithelia and mononuclear infiltrates from COPD lungs.	AZD1981	AZD1981 (4-week treatment) did not induce significant differences in lung function, quality of life variables, nor use of reliever medication in COPD patients in DBPCRT (NCT00690482).	114
Adenosine A _{2A} receptor	Inhibits neutrophil superoxide production, phagocytosis, adhesion, and cytokine release.	UK-432097 (agonist)	UK-432097 (6-week inhaled treatment) in DBPCRT (NCT00430300) showed no significant differences in FEV ₁ , use of rescue bronchodilator, or quality of life parameters.	115
Selectins	Involved in migration of leukocytes from blood to surrounding tissues. Overexpressed in lung tissue of COPD patients.	Bimosiamose	Inhalation of bimosiamose (28-day treatment) attenuated inflammation by significantly reducing numbers of macrophages and concentrations of CXCL8 in sputum of COPD patients in DBPCRT (NCT01108913). Adverse events were similar between the groups.	68
Vasoactive intestinal peptide (VIP)	Bronchodilatory and immunomodulatory effects in the lungs. Anti-inflammatory activity requires activation of both VPAC1 and VPAC2 receptors, With VPAC1 being particularly elevated in AMs of COPD patients.	VIP; available derivatives have not been tested in human COPD	VIP (3-month inhaled treatment) was tested in DBPCRT in severe COPD patients (NCT00464932). Study was completed in 2006 but no results are available.	116–119

Abbreviations: AM, alveolar macrophage; TNF- α , tumor necrosis factor α ; DBPCRT, double-blind, placebo-controlled, randomized trial; FEV₁, forced expiratory volume in 1 second; NF- κ B, nuclear factor- κ B; Th, T helper.

with PPAR γ agonists including 10-nitro-oleic acid (a possible endogenous ligand) or rosiglitazone (a thiazolidinedione) blocked cigarette smoke-induced production of cytokines, chemokines, and ROS, and accompanying suppression of HDAC2 levels.⁶⁹ The PPAR γ agonist rosiglitazone dose dependently inhibited LPS-induced production of TNF- α and CCL5 by AMs of COPD patients, smokers, and never-smokers, shifting them toward an anti-inflammatory M2 phenotype, while both rosiglitazone and pioglitazone attenuated pulmonary inflammation in a tobacco smoke mouse model.⁷⁰ PPAR γ also appears to play a central role in the development of emphysema. PPAR γ is downregulated in lung myeloid dendritic cells of smokers with emphysema and endogenous PPAR γ agonist activity is reduced in plasma of those with emphysema, while treatment with the thiazolidinedione ciglitazone reverses emphysema in a

mouse model.⁷¹ A retrospective epidemiological study of veterans with both diabetes and COPD indicated that the 7,887 veterans treated with a PPAR γ agonist for their diabetes (97.1% rosiglitazone) significantly reduced the risk of COPD exacerbations in comparison to those receiving other diabetes medications (n=42,347; incidence rate ratio [IRR]=0.85; CI: 0.80–0.91) (Table 4).⁷² Taken together, these data support further studies of PPAR γ agonists in COPD patients with early signs of emphysema.

COPD also involves progressive increase in pulmonary arterial pressure, and 20%–91% (depending on definition, COPD severity, and method of measurement) of COPD patients have progressed to pulmonary hypertension.⁷³ Endothelin signaling plays a major role in vascular remodeling and hence in development of pulmonary hypertension.^{74,75} An 18-month treatment with the

endothelin antagonist bosentan improved measures of pulmonary hypertension compared to those at baseline, especially in GOLD grade III and IV COPD patients, while pulmonary hypertension worsened in placebo-treated patients (Table 4).⁷⁴

However, bosentan can actually worsen hypoxemia in COPD patients without pulmonary hypertension.⁷⁶

Although statins are known to have anti-inflammatory properties, simvastatin treatment (40 mg/day) of COPD

Table 4 Developmental status of miscellaneous inflammatory modulators for COPD

Mediator	Role in COPD	Drug	Clinical development	References
PPAR γ (peroxisome proliferator-activated receptor γ)	Cigarette smoke downregulates PPAR γ . Reduced PPAR γ expression and activity seen in COPD.	Thiazolidinediones: rosiglitazone and pioglitazone	Retrospective cohort study of patients with diabetes and COPD showed that patients who filled ≥ 2 thiazolidinedione prescriptions (97.1% rosiglitazone) had a significantly lower number of COPD exacerbations than those receiving other diabetic medications.	69, 72
IgE activity	The high affinity IgE receptor is overexpressed on myeloid and plasmacytoid dendritic cells (DCs) of current smokers. Expression on plasmacytoid DCs correlates with COPD stage.	Omalizumab	The clinical trial of omalizumab (NCT00851370) was withdrawn due to lack of patients meeting inclusion criteria (elevated IgE and positive skin prick test to environmental allergens).	120
RAR γ (retinoic acid receptor γ)	Regulates function of multiple cells of the immune system.	Palovarotene	RAR γ agonist (2 year treatment) was tested for ability to improve lung function in patients with emphysema in DBPCRT (NCT00413205); in patients with lower lobe emphysema, palovarotene significantly reduced the decline in lung function (from conference report). In another study, over 1 year, palovarotene failed to show a significant benefit on lung density in moderate-to-severe emphysema secondary to severe alpha-1 antitrypsin deficiency.	121, 142, 143
Angiotensin receptor	Angiotensin II receptor blockers reduce in-hospital mortality during first COPD exacerbation.	Losartan	Open-label clinical trial of losartan (4-week treatment) (NCT02416102) to assess effects on mucociliary dysfunction (nasal potential difference, IL-8 and TGF- β in nasal discharge) in COPD patients is recruiting; a 4-year study of losartan for prevention of emphysema progression (NCT02696564) is not yet recruiting.	122
Endothelin	Vasoconstrictor, contributes to pulmonary hypertension in COPD.	Bosentan	Bosentan (18-month treatment) halted progression of PH and led to improvements in most patients, especially those in GOLD grades III and IV. A study of bosentan effects on acute exacerbations and lung function in patients with GOLD III or IV COPD and pulmonary hypertension was initiated but status is unknown (NCT02093195).	73–75
Statins	Statins exert anti-inflammatory effects by several mechanisms independent of cholesterol lowering.	Simvastatin, rosuvastatin	Although retrospective studies suggested that statins may reduce frequency of exacerbations, hospitalization, and mortality in COPD patients, a recent prospective large randomized trial of simvastatin in COPD patients (NCT01061671) did not detect significant differences.	77, 78, 123, 124

(Continued)

Table 4 (Continued)

Mediator	Role in COPD	Drug	Clinical development	References
T cells	T cells infiltrate airways, and are key mediators in the immune response; numbers of senescent cells, which produce increased amounts of proinflammatory and cytotoxic mediators and are relatively resistant to GC treatment, are elevated in blood and lungs of COPD patients.	Cyclosporine, an immunosuppressant affecting T-cell responses	Another 3-month study of simvastatin found no effect in inflammatory biomarkers. Rosuvastatin (12-week treatment) in DBPCRT (NCT00929734) reduced biomarkers of systemic inflammation and improved endothelial function in a prespecified subgroup (patients with supramedian circulating hsCRP levels). A phase I dose-escalation study of inhaled cyclosporine (28-day treatment) in severe COPD patients (NCT00783107) also evaluated inflammatory biomarkers; no results of 2009 study are available. An ongoing phase II DBPCRT (16-week treatment) (NCT00974142) is evaluating oral cyclosporine in severe COPD.	125, 126

Abbreviations: Ig, immunoglobulin; IL, interleukin; TGF- β , transforming growth factor β ; GOLD, Global Initiative for Chronic Obstructive Lung Disease; DBPCRT, double-blind, placebo-controlled, randomized trial; hsCRP, high-sensitivity C-reactive protein.

patients at high risk for exacerbation did not increase the time to first exacerbation nor reduce the number of exacerbations.⁷⁷ However, rosuvastatin (12-week treatment) improved endothelium-dependent vascular function in a prespecified subgroup (patients with supramedian circulating high-sensitivity C-reactive protein [hsCRP]) but not in the total COPD population in a double-blind, placebo-controlled trial.⁷⁸ This study found no statistically significant effect on pulmonary function parameters, however.

PDE inhibitors

The 11-membered phosphodiesterase enzyme family (PDE1-11) differentially hydrolyzes cyclic adenosine monophosphate (cAMP) and cyclic guanosine monophosphate (cGMP), which regulate many cellular processes including release of inflammatory mediators and relaxation of smooth muscles.¹⁰ PDE3 appears to be involved in bronchoconstriction since its inhibitors induce bronchodilation in humans.¹⁰ PDE4 is expressed in most inflammatory cell types and is a main target for emerging COPD therapies (Table 5), with the PDE4 inhibitor roflumilast having been approved by the FDA as a COPD treatment. In vitro studies of low-dose combinations of PDE4 and PI3K δ inhibitors significantly reduced cigarette smoke extract-induced apoptosis of lung epithelial cells, neutrophil elastase production, and macrophage secretion of TNF- α , phosphorylated protein kinase B, and matrix metalloproteinase 9 (MMP-9).⁷⁹

Gastrointestinal adverse effects can be a significant problem with roflumilast, but PDE4 inhibitors with inhaled or nebulized (eg, RPL554, GSK-256066) formulations

appear to have more tolerable side effect profiles than orally administered roflumilast.^{80,81}

The severity of emphysema and small airways disease is associated with higher expression of multiple MMPs.⁸² However, despite promising results in an animal model,⁸³ an inhibitor of MMP-9 and -12 has not shown significant benefit for COPD patients in a clinical trial (Table 6).

Oxidative stress and antioxidants

Oxidants, both from inhaled pollutants and produced as part of the inflammatory response, are major contributors to COPD pathophysiology. Furthermore, inflammation-associated production of ROS and reactive nitrogen species (RNS) is accompanied by downregulation of the antioxidant transcription factor Nrf2. ROS activates the proinflammatory transcription factors activator protein 1 (AP-1) and NF- κ B, with consequent production of inflammatory proteins and mediators. Excess ROS in COPD patients also directly contributes to airspace epithelial injury and inactivates anti-proteases that help prevent emphysema.⁸⁴ Therapeutic agents that activate Nrf2 and thus neutralize the excess oxidants may prove beneficial to COPD treatment.⁸⁵

The antioxidant transcription factor Nrf2, the primary mechanism for limiting oxidative stress, is reduced in COPD patients.³⁶ Kelch-like ECH-associated protein 1 (Keap1) sequesters Nrf2 in the cytoplasm and under healthy conditions targets Nrf2 to Cullin-3 for ubiquitination and degradation.⁸⁶ Keap1 monitors oxidative stress through its multiple cysteines with distinct stressors binding one or more

Table 5 Developmental status of cAMP and cGMP phosphodiesterase inhibitors

Mediator	Role in COPD	Drug	Clinical development	References
PDE4 (phosphodiesterase subtype 4)	Hydrolyzes cAMP, an inhibitor of inflammatory pathways; expressed in a wide variety of cells.	Roflumilast, a selective PDE4 inhibitor; GSK-256066; CHF6001; MK0359; MK-0873; tofomilast; UK-500,001; tetomilast (OPC-6535, PDE4 inhibitor with modest PDE3 inhibitory activity); oglemilast; QAK423A; TPI 1100.	Roflumilast is only US FDA approved PDE4 inhibitor; it reduced exacerbation frequency and also produced clinically significant improvements in dyspnea. GSK-256066 (4-week inhaled treatment) in DBPCRT (NCT00549679) improved residual volume and showed a nonsignificant trend toward augmenting postbronchodilator FEV ₁ . Preclinical testing of CHF6001 (inhaled) shows efficacy and low toxicity in several rat models of pulmonary inflammation. It is in clinical testing (28-day treatment) (NCT01730404) but no results have been reported. Numerous other PDE inhibitors are in clinical testing, including MK-0359 (NCT00482235); MK-0873 (NCT00132730); tofomilast (NCT00219622); UK-500,001 (NCT00263874); tetomilast (OPC-6535) (NCT00874497), terminated, NCT00917150); oglemilast (NCT00671073); QAK423A (NCT01197287); and TPI 1100 (NCT00914433).	10, 42, 80, 99, 127–129
PDE3/PDE4	PDE3 degrades both cAMP and cGMP. It is expressed on airway smooth muscle cells and acts as a bronchoconstrictor. Combined PDE3/PDE4 inhibition is often additive or synergistic.	RPL554	RPL554 (up to 94-day treatment) is being investigated as an adjunct to salbutamol and ipratropium in COPD patients in DBPCRT (NCT02542254).	10
PDE5	PDE5 promotes pulmonary arterial vasoconstriction and vessel wall hypertrophy.	Tadalafil (inhibits PDE5)	Tadalafil, which is approved for pulmonary arterial hypertension, in DBPCRT (12-week treatment) (NCT01197469) did not improve exercise capability or quality of life. Another study (NCT01862536) is in progress.	130

Abbreviations: FEV₁, forced expiratory volume in 1 second; DBPCRT, double-blind, placebo-controlled, randomized trial; FDA, Food and Drug Administration.

specific cysteines. Following binding of these stress-related compounds, release of Nrf2 from Keap1 occurs, which transfers to the nucleus and activates multiple antioxidant enzymes and phase II proteins that counteract oxidative stress.⁸⁶ This makes the Nrf2/Keap1 system an attractive therapeutic target in COPD and other inflammatory diseases. The natural product sulforaphane activated Nrf2 in AMs isolated from COPD patients, denitrosylated HDAC2, and restored sensitivity to the glucocorticoid dexamethasone in a glutathione-dependent manner.³⁷ A study of the effect of two doses of sulforaphane on Nrf2 expression in 89 COPD patients was recently completed, and sulforaphane administered for four weeks to patients with COPD did not induce the expression of Nrf2 target genes or have an effect on oxidative stress, airway inflammation, or lung function. (NCT01335971).⁸⁷

Erdosteine and N-acetylcysteine directly scavenge ROS via their thiol groups (of the metabolite in the case

of erdosteine) and also have mucolytic activity. High-dose (900 mg/day) erdosteine increased the ability of salbutamol to improve %FEV₁ reversibility,⁸⁸ and long-term treatment reduces exacerbations and improves quality of life.⁸⁹ High-dose (600 mg bid) N-acetylcysteine reduces the number of exacerbations patients experience;^{12,90,91} reported effects with a lower dose (600 mg/day) have been inconsistent.^{92–94}

Mucus hypersecretion

Mucus hypersecretion can be modulated by blocking its overproduction and/or by inhibiting its secretion (Table 7). Hypothetically, a rebound effect after cessation of an inhibitor of mucin secretion may involve a rapid release of produced but unsecreted mucin. In contrast, the rebound effect after cessation of an inhibitor of mucin production may involve a lag phase for mucin production and a more gradual increase of mucin secretion. Multiple mucins comprise the gel-forming layer of normal airway mucus. Numerous signals can promote

Table 6 Developmental status of elastin-degrading protease inhibitors for COPD

Mediator	Role in COPD	Drug	Clinical development	References
Neutrophil elastase	Abundant in neutrophils; can degrade extracellular matrix and damage/destroy lung parenchyma; affects mucus secretion.	AZD9668 AZD6553	AZD9668 (12-week treatment) showed no effect on pulmonary function or quality of life when combined with tiotropium (NCT00949975) or budesonide/formoterol (NCT01023516); there was likewise no effect on airway remodeling (NCT01054170) and studies found no decrease in degradation as assessed by urinary desmosine. AZD6553 clinical trial (NCT01068184) was terminated due to PK inconsistent with pharmaceutical properties.	131, 132
Matrix metalloproteinases (MMPs)	Higher levels of multiple MMPs in lungs of COPD patients; involved in matrix breakdown and tissue remodeling.	AZD1236 (anti-MMP-9 and -12); GS-5745 (anti-MMP-9)	In a 6-week DBPCRT (NCT00758706) of AZD1236 (anti-MMP-9 and -12) in moderate-to-severe COPD patients, reduction in urinary desmosine did not reach statistical significance and there was no effect on COPD clinical symptoms. Another study (NCT00758459) has completed but statistical analysis not released. A 28-day safety and PK study of GS-5745 in COPD patients (NCT02077465) has been completed.	82, 133

Abbreviations: PK, pharmacokinetics; DBPCRT, double-blind, placebo-controlled, randomized trial.

mucus secretion: bacterial products, cytokines, cholinergic agonists, elastases, matrix metalloproteases, and activation of epidermal growth factor receptor (EGFR).^{95,96} Cigarette smoke induces EGFR- and hypoxia inducible factor-1 (HIF-1)-mediated signaling⁹⁵ and thus can promote hyperplasia of mucin-producing goblet cells.⁹⁷ Supporting this concept, nuclear HIF-1 α was expressed in the majority of goblet cells in areas of remodeled airway tissues showing goblet cell hyperplasia from COPD patients but not in subjects without COPD.⁹⁷ IL-13, an essential component of COPD-associated inflammation, promotes goblet cell production of mucins.⁹⁸ After the mucins are expressed, glycosylated, and packaged in mucin granules, myristoylated alanine-rich C kinase substrate (MARCKS) mediates movement of intracellular mucin granules to the goblet cell apical membrane and is therefore essential for mucin exocytosis and secretion.⁹⁸

Summary

The heterogeneity of COPD presentation augments the challenges in identifying and developing therapeutic compounds for the treatment of COPD patients. It also emphasizes the importance of tailoring therapy to individual patients and their disease status, which may involve considerations beyond the standard GOLD categories. Several distinct emerging therapies have shown efficacy in at least some COPD patients. The length of treatment (>3 months) and the inhaled route of administration can be associated with a

higher probability of observing a positive effect on COPD variables and a reduced side effect profile, respectively. Although some of the emerging compounds showed significant activity in the total COPD population tested,^{45,68,74,88,99} sometimes only in connection with assessed biomarkers rather than clinically significant outcomes, subgroup analysis showed that most compounds were significantly affected in one or more of the following subgroups: smokers or ex-smokers,^{44,99} patients with chronic bronchitis or emphysema,^{80,99} use of standard COPD therapies,⁹⁹ patients with alteration of relevant biomarkers,⁷⁸ or at different COPD grades.⁷⁴ These data suggest that subgroup analyses and the possibility of individualized therapy can benefit developers of emerging therapies by identification of the patients most likely to benefit from a new therapy.

Acknowledgments

This work was supported by a Merit Review award from the US Department of Veterans Affairs and the National Institutes of Health grants HL093196 and AI125338 (RCR). We gratefully acknowledge the scientific and writing expertise of KL Molnar-Kimber, PhD. The contents in this article do not represent the views of the US Department of Veterans Affairs or the United States Government.

Disclosure

The authors report no conflicts of interest in this work.

Table 7 Developmental status of modulators of mucus-mediated airway obstruction for COPD

Mediator	Role in COPD	Drug	Clinical development	References
Epidermal growth factor receptor (EGFR)	EGFR regulates mucin stores in airway epithelium, which are significantly increased in COPD.	BIBW 2948 (inhibits EGFR autophosphorylation)	Inhalation of BIBW 2948 (4-week treatment) in DBPCRT (NCT00423137) reduced internalization of EGFR but did not reduce mucin stores; BIBW 2948 treatment was associated with higher discontinuation rate (24%) than placebo (4.3%). FEV ₁ in the higher dose group significantly declined by visit 5 but returned to baseline by visit 7.	134
Myristoylated alanine-rich C kinase substrate (MARCKS)	Mediates movement of mucin granules to the apical membrane as part of mucin exocytosis.	BIO-11006	A 21-day phase II DBPCRT of BIO-11006 (inhaled) in COPD (NCT00648245) has been completed; a 2011 abstract reported improved lung function and reduced mucus hypersecretion.	98, 135, 136
Epithelial sodium channel	Role in homeostasis of mucus hydration, ciliary beating, and clearance of mucus.	GS-5737; compound A	Study of effects of GS-5737 on ciliary action in healthy controls (NCT01793649) was terminated. Preclinical study of compound A shows improved ciliary movement, mucus clearance, and airway hydration.	137
Multiple mechanisms				
Anti-inflammatory and mucolytic	Inflammation, oxidative stress and mucus hypersecretion are well-established in COPD.	N-acetylcysteine (NAC)	The 1-year DBPCRT PANTHEON trial found 600 mg bid NAC reduced exacerbations in patients with GOLD II–III COPD (Chinese Clinical Trials Registry TRC-09000460), as did a smaller study (NCT01136239) that found a reduction only in high-risk patients but also observed improvement in airway function. However, two lower dose studies (600 mg/day) (NCT00184977; not registered) found no benefit while another (not registered) did.	12, 90–94, 138, 139
Mucolytic, anti-inflammatory, antioxidant, promotes activity of antibiotics	Inflammation, oxidative stress, and mucus hypersecretion are well-established in COPD.	Erdosteine	DBPCRT (NCT00338507) to test daily erdosteine for 28 days. After 4 weeks, erdosteine treatment significantly reduced plasma oxidant levels and increased %FEV ₁ reversibility by salbutamol treatment. In other reported studies, addition of erdosteine for 7–10 days reduced duration of acute exacerbations, while long-term treatment in stable COPD reduced exacerbations and improved quality of life.	88, 89
Cystic fibrosis transmembrane conductance regulator (CFTR)	One study found that CFTR is downregulated in smokers with and without COPD; another found that expression of CFTR inversely correlated with emphysema severity.	Ivacaftor potentiates chloride transport	A pilot DBPCRT of ivacaftor (NCT02135432) (treatment up to 2 weeks) with outcome assessed by sweat chloride has been completed.	140, 141

Abbreviations: DBPCRT, double-blind, placebo-controlled, randomized trial; FEV₁, forced expiratory volume in 1 second; GOLD, Global Initiative for Chronic Obstructive Lung Disease.

References

- Ferkol T, Schraufnagel D. The global burden of respiratory disease. *Ann Am Thorac Soc*. 2014;11(3):404–406.
- Heron M. Deaths: leading causes for 2013. *Natl Vital Stat Rep*. 2015; 65(2):95.
- Centers for Disease Control and Prevention (CDC). Chronic obstructive pulmonary disease among adults – United States, 2011. *MMWR Morb Mortal Wkly Rep*. 2012;61(46):938–943.
- Barnes PJ. New anti-inflammatory targets for chronic obstructive pulmonary disease. *Nat Rev Drug Discov*. 2013;12(7):543–559.
- Vestbo J, Hurd SS, Agusti AG, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med*. 2013;187(4): 347–365.
- Woodruff PG, Agusti A, Roche N, Singh D, Martinez FJ. Current concepts in targeting chronic obstructive pulmonary disease pharmacotherapy: making progress towards personalised management. *Lancet*. 2015;385(9979):1789–1798.
- Caramori G, Casolari P, Barczyk A, Durham AL, Di Stefano A, Adcock I. COPD immunopathology. *Semin Immunopathol*. 2016;38(4):497–515.

8. Wiegman CH, Michaeloudes C, Haji G, et al. Oxidative stress-induced mitochondrial dysfunction drives inflammation and airway smooth muscle remodeling in patients with chronic obstructive pulmonary disease. *J Allergy Clin Immunol*. 2015;136(3):769–780.
9. Vedel-Krogh S, Nielsen SF, Lange P, Vestbo J, Nordestgaard BG. Blood eosinophils and exacerbations in chronic obstructive pulmonary disease. The Copenhagen General Population Study. *Am J Respir Crit Care Med*. 2016;193(9):965–974.
10. Abbott-Banner KH, Page CP. Dual PDE3/4 and PDE4 inhibitors: novel treatments for COPD and other inflammatory airway diseases. *Basic Clin Pharmacol Toxicol*. 2014;114(5):365–376.
11. Pavord ID, Jones PW, Burgel PR, Rabe KF. Exacerbations of COPD. *Int J Chron Obstruct Pulmon Dis*. 2016;11(Spec Iss):21–30.
12. Tse HN, Raiteri L, Wong KY, Ng LY, Yee KS, Tseng CZ. Benefits of high-dose N-acetylcysteine to exacerbation-prone patients with COPD. *Chest*. 2014;146(3):611–623.
13. Martinez FJ, Calverley PM, Goehring UM, Brose M, Fabbri LM, Rabe KF. Effect of roflumilast on exacerbations in patients with severe chronic obstructive pulmonary disease uncontrolled by combination therapy (REACT): a multicentre randomised controlled trial. *Lancet*. 2015;385(9971):857–866.
14. Celli BR, MacNee W. Standards for the diagnosis and treatment of patients with COPD: a summary of the ATS/ERS position paper. *Eur Respir J*. 2004;23(6):932–946.
15. Celli BR, Thomas NE, Anderson JA, et al. Effect of pharmacotherapy on rate of decline of lung function in chronic obstructive pulmonary disease: results from the TORCH study. *Am J Respir Crit Care Med*. 2008;178(4):332–338.
16. Vestbo J, Anderson JA, Brook RD, et al. Fluticasone furoate and vilanterol and survival in chronic obstructive pulmonary disease with heightened cardiovascular risk (SUMMIT): a double-blind randomised controlled trial. *Lancet*. 2016;387(10030):1817–1826.
17. Bafadhel M, McKenna S, Terry S, et al. Acute exacerbations of chronic obstructive pulmonary disease: identification of biologic clusters and their biomarkers. *Am J Respir Crit Care Med*. 2011;184(6):662–671.
18. Soter S, Barta I, Antus B. Predicting sputum eosinophilia in exacerbations of COPD using exhaled nitric oxide. *Inflammation*. 2013;36(5):1178–1185.
19. Brightling CE, Monteiro W, Ward R, et al. Sputum eosinophilia and short-term response to prednisolone in chronic obstructive pulmonary disease: a randomised controlled trial. *Lancet*. 2000;356(9240):1480–1485.
20. Chanez P, Vignola AM, O'Shaughnessy T, et al. Corticosteroid reversibility in COPD is related to features of asthma. *Am J Respir Crit Care Med*. 1997;155(5):1529–1534.
21. Pizzichini E, Pizzichini MM, Gibson P, et al. Sputum eosinophilia predicts benefit from prednisone in smokers with chronic obstructive bronchitis. *Am J Respir Crit Care Med*. 1998;158(5 pt 1):1511–1517.
22. Pace E, Giarratano A, Ferraro M, et al. TLR4 upregulation underpins airway neutrophilia in smokers with chronic obstructive pulmonary disease and acute respiratory failure. *Hum Immunol*. 2011;72(1):54–62.
23. Mortaz E, Adcock IM, Ito K, Kraneveld AD, Nijkamp FP, Folkerts G. Cigarette smoke induces CXCL8 production by human neutrophils via activation of TLR9 receptor. *Eur Respir J*. 2010;36(5):1143–1154.
24. Hogg JC, Chu F, Utokaparch S, et al. The nature of small-airway obstruction in chronic obstructive pulmonary disease. *N Engl J Med*. 2004;350(26):2645–2653.
25. Burgel PR, Bourdin A, Chanez P, et al. Update on the roles of distal airways in COPD. *Eur Respir Rev*. 2011;20(119):7–22.
26. Chung KF, Adcock IM. Multifaceted mechanisms in COPD: inflammation, immunity, and tissue repair and destruction. *Eur Respir J*. 2008;31(6):1334–1356.
27. McDonough JE, Yuan R, Suzuki M, et al. Small-airway obstruction and emphysema in chronic obstructive pulmonary disease. *N Engl J Med*. 2011;365(17):1567–1575.
28. Loza MJ, Watt R, Baribaud F, Barnathan ES, Rennard SI. Systemic inflammatory profile and response to anti-tumor necrosis factor therapy in chronic obstructive pulmonary disease. *Respir Res*. 2012;13:12.
29. Hogg JC, Chu FS, Tan WC, et al. Survival after lung volume reduction in chronic obstructive pulmonary disease: insights from small airway pathology. *Am J Respir Crit Care Med*. 2007;176(5):454–459.
30. Barnes PJ. Mechanisms and resistance in glucocorticoid control of inflammation. *J Steroid Biochem Mol Biol*. 2010;120(2–3):76–85.
31. Barnes PJ. Corticosteroid resistance in patients with asthma and chronic obstructive pulmonary disease. *J Allergy Clin Immunol*. 2013;131(3):636–645.
32. Barnes PJ. Role of HDAC2 in the pathophysiology of COPD. *Annu Rev Physiol*. 2009;71:451–464.
33. Cosio BG, Tsaprouni L, Ito K, Jazrawi E, Adcock IM, Barnes PJ. Theophylline restores histone deacetylase activity and steroid responses in COPD macrophages. *J Exp Med*. 2004;200(5):689–695.
34. Subramanian, Ragulan, Jindal A, Viswambhar V, Arun Babu V. The study of efficacy, tolerability and safety of theophylline given along with formoterol plus budesonide in COPD. *J Clin Diagn Res*. 2015;9(2):OC10–OC13.
35. Cosio BG, Shafiek H, Iglesias A, et al. Oral low-dose theophylline on top of inhaled fluticasone-salmeterol does not reduce exacerbations in patients with severe COPD: a pilot clinical trial. *Chest*. 2016;150(1):123–130.
36. Suzuki M, Betsuyaku T, Ito Y, et al. Down-regulated NF-E2-related factor 2 in pulmonary macrophages of aged smokers and patients with chronic obstructive pulmonary disease. *Am J Respir Cell Mol Biol*. 2008;39(6):673–682.
37. Malhotra D, Thimmulappa RK, Mercado N, et al. Denitrosylation of HDAC2 by targeting Nrf2 restores glucocorticosteroid sensitivity in macrophages from COPD patients. *J Clin Invest*. 2011;121(11):4289–4302.
38. Marwick JA, Caramori G, Stevenson CS, et al. Inhibition of PI3Kdelta restores glucocorticoid function in smoking-induced airway inflammation in mice. *Am J Respir Crit Care Med*. 2009;179(7):542–548.
39. Boardman C, Chachi L, Gavrilu A, et al. Mechanisms of glucocorticoid action and insensitivity in airways disease. *Pulm Pharmacol Ther*. 2014;29(2):129–143.
40. Mercado N, Hakim A, Kobayashi Y, et al. Restoration of corticosteroid sensitivity by p38 mitogen activated protein kinase inhibition in peripheral blood mononuclear cells from severe asthma. *PLoS One*. 2012;7(7):e41582.
41. Milara J, Lluch J, Almudever P, Freire J, Xiaozhong Q, Cortijo J. Roflumilast N-oxide reverses corticosteroid resistance in neutrophils from patients with chronic obstructive pulmonary disease. *J Allergy Clin Immunol*. 2014;134(2):314–322.
42. Rennard SI, Calverley PM, Goehring UM, Bredenkroter D, Martinez FJ. Reduction of exacerbations by the PDE4 inhibitor roflumilast – the importance of defining different subsets of patients with COPD. *Respir Res*. 2011;12:18.
43. Caramori G, Adcock IM, Di Stefano A, Chung KF. Cytokine inhibition in the treatment of COPD. *Int J Chron Obstruct Pulmon Dis*. 2014;9:397–412.
44. Rennard SI, Dale DC, Donohue JF, et al. CXCR2 antagonist MK-7123. A phase 2 proof-of-concept trial for chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2015;191(9):1001–1011.
45. Kirsten AM, Forster K, Radeckzy E, et al. The safety and tolerability of oral AZD5069, a selective CXCR2 antagonist, in patients with moderate-to-severe COPD. *Pulm Pharmacol Ther*. 2015;31:36–41.
46. Kerstjens HA, Bjermer L, Eriksson L, Dahlstrom K, Vestbo J. Tolerability and efficacy of inhaled AZD4818, a CCR1 antagonist, in moderate to severe COPD patients. *Respir Med*. 2010;104(9):1297–1303.
47. Bai J, Song H, Cai C, Zhang M, Xu S, Tan J. The association of monocyte chemoattractant protein-1 and CC chemokine receptor 2 gene variants with chronic obstructive pulmonary disease. *DNA Cell Biol*. 2012;31(6):1058–1063.
48. Monzon ME, Forteza RM, Casalino-Matsuda SM. MCP-1/CCR2B-dependent loop upregulates MUC5AC and MUC5B in human airway epithelium. *Am J Physiol Lung Cell Mol Physiol*. 2011;300(2):L204–L215.

49. Botelho FM, Bauer CM, Finch D, et al. IL-1alpha/IL-1R1 expression in chronic obstructive pulmonary disease and mechanistic relevance to smoke-induced neutrophilia in mice. *PLoS One*. 2011; 6(12):e28457.
50. Rogliani P, Calzetta L, Ora J, Matera MG. Canakinumab for the treatment of chronic obstructive pulmonary disease. *Pulm Pharmacol Ther*. 2015;31:15–27.
51. Varricchi G, Bagnasco D, Borriello F, Heffler E, Canonica GW. Interleukin-5 pathway inhibition in the treatment of eosinophilic respiratory disorders: evidence and unmet needs. *Curr Opin Allergy Clin Immunol*. 2016;16(2):186–200.
52. Roos AB, Sanden C, Mori M, Bjermer L, Stampfli MR, Erjefalt JS. IL-17A is elevated in end-stage chronic obstructive pulmonary disease and contributes to cigarette smoke-induced lymphoid neogenesis. *Am J Respir Crit Care Med*. 2015;191(11):1232–1241.
53. Zhang J, Chu S, Zhong X, Lao Q, He Z, Liang Y. Increased expression of CD4+IL-17+ cells in the lung tissue of patients with stable chronic obstructive pulmonary disease (COPD) and smokers. *Int Immunopharmacol*. 2013;15(1):58–66.
54. Shan M, Cheng HF, Song LZ, et al. Lung myeloid dendritic cells coordinately induce TH1 and TH17 responses in human emphysema. *Sci Transl Med*. 2009;1(4):4ra10.
55. Roos AB, Sethi S, Nikota J, et al. IL-17A and the promotion of neutrophilia in acute exacerbation of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2015;192(4):428–437.
56. Jin Y, Wan Y, Chen G, et al. Treg/IL-17 ratio and Treg differentiation in patients with COPD. *PLoS One*. 2014;9(10):e111044.
57. Freeman CM, Martinez CH, Todt JC, et al. Acute exacerbations of chronic obstructive pulmonary disease are associated with decreased CD4+ & CD8+ T cells and increased growth & differentiation factor-15 (GDF-15) in peripheral blood. *Respir Res*. 2015;16:94.
58. Zhang X, Zheng H, Zhang H, et al. Increased interleukin (IL)-8 and decreased IL-17 production in chronic obstructive pulmonary disease (COPD) provoked by cigarette smoke. *Cytokine*. 2011; 56(3):717–725.
59. Freeman CM, McCubbrey AL, Crudgington S, et al. Basal gene expression by lung CD4+ T cells in chronic obstructive pulmonary disease identifies independent molecular correlates of airflow obstruction and emphysema extent. *PLoS One*. 2014;9(5):e96421.
60. Kang MJ, Choi JM, Kim BH, et al. IL-18 induces emphysema and airway and vascular remodeling via IFN-gamma, IL-17A, and IL-13. *Am J Respir Crit Care Med*. 2012;185(11):1205–1217.
61. Nakajima T, Owen C. Interleukin-18: the master regulator driving destructive and remodeling processes in the lungs of patients with chronic obstructive pulmonary disease? *Am J Respir Crit Care Med*. 2012;185(11):1137–1139.
62. MedImmune LLC. A study to evaluate the safety of MEDI2338 in subjects with chronic obstructive pulmonary disease; 2013. Available from: <https://clinicaltrials.gov/ct2/show/results/NCT01322594?term=MEDI2338&rank=1>. NLM identifier: NCT01322594. Accessed January 15, 2016.
63. Hodge G, Reynolds PN, Holmes M, Hodge S. Differential expression of pro-inflammatory cytokines in intra-epithelial T cells between trachea and bronchi distinguishes severity of COPD. *Cytokine*. 2012; 60(3):843–848.
64. Norman P. Investigational p38 inhibitors for the treatment of chronic obstructive pulmonary disease. *Expert Opin Investig Drugs*. 2015;24(3): 383–392.
65. Robinson C, Xia K, Russell P, et al. A 12 week, randomized, double-blind, placebo controlled study of the efficacy of RV568 (JNJ49095397), a narrow spectrum kinase inhibitor, in COPD patients. *Am J Respir Crit Care Med*. 2016;193:A6843.
66. Fung-Leung WP. Phosphoinositide 3-kinase delta (PI3Kdelta) in leukocyte signaling and function. *Cell Signal*. 2011;23(4):603–608.
67. Down K, Amour A, Baldwin IR, et al. Optimization of novel indazoles as highly potent and selective inhibitors of phosphoinositide 3-kinase delta for the treatment of respiratory disease. *J Med Chem*. 2015;58(18): 7381–7399.
68. Watz H, Bock D, Meyer M, et al. Inhaled pan-selectin antagonist bimosiamose attenuates airway inflammation in COPD. *Pulm Pharmacol Ther*. 2013;26(2):265–270.
69. Lakshmi SP, Reddy AT, Zhang Y, et al. Down-regulated peroxisome proliferator-activated receptor gamma (PPARgamma) in lung epithelial cells promotes a PPARgamma agonist-reversible proinflammatory phenotype in chronic obstructive pulmonary disease (COPD). *J Biol Chem*. 2014;289(10):6383–6393.
70. Lea S, Plumb J, Metcalfe H, et al. The effect of peroxisome proliferator-activated receptor-gamma ligands on in vitro and in vivo models of COPD. *Eur Respir J*. 2014;43(2):409–420.
71. Shan M, You R, Yuan X, et al. Agonistic induction of PPARgamma reverses cigarette smoke-induced emphysema. *J Clin Invest*. 2014;124(3): 1371–1381.
72. Rinne ST, Liu CF, Feemster LC, et al. Thiazolidinediones are associated with a reduced risk of COPD exacerbations. *Int J Chron Obstruct Pulmon Dis*. 2015;10:1591–1597.
73. Shujaat A, Bajwa AA, Cury JD. Pulmonary hypertension secondary to COPD. *Pulm Med*. 2012;2012:203952.
74. Valerio G, Bracciale P, Grazia D'Agostino A. Effect of bosentan upon pulmonary hypertension in chronic obstructive pulmonary disease. *Ther Adv Respir Dis*. 2009;3(1):15–21.
75. Zakynthinos E, Daniil Z, Papanikolaou J, Makris D. Pulmonary hypertension in COPD: pathophysiology and therapeutic targets. *Curr Drug Targets*. 2011;12(4):501–513.
76. Stolz D, Rasch H, Linka A, et al. A randomised, controlled trial of bosentan in severe COPD. *Eur Respir J*. 2008;32(3):619–628.
77. Criner GJ, Connett JE, Aaron SD, et al. Simvastatin for the prevention of exacerbations in moderate-to-severe COPD. *N Engl J Med*. 2014;370(23): 2201–2210.
78. Neukamm A, Hoiseth AD, Einvik G, et al. Rosuvastatin treatment in stable chronic obstructive pulmonary disease (RODEO): a randomized controlled trial. *J Intern Med*. 2015;278(1):59–67.
79. Dinavahi SS, Nyayapathy S, Perumal Y, Dharmarajan S, Viswanadha S. Combined inhibition of PDE4 and PI3Kdelta modulates the inflammatory component involved in the progression of chronic obstructive pulmonary disease. *Drug Res (Stuttg)*. 2014;64(4):214–219.
80. Mulhall AM, Droegge CA, Ernst NE, Panos RJ, Zafar MA. Phosphodiesterase 4 inhibitors for the treatment of chronic obstructive pulmonary disease: a review of current and developing drugs. *Expert Opin Investig Drugs*. 2015;24(12):1597–1611.
81. De Savi C, Cox RJ, Warner DJ, et al. Efficacious inhaled PDE4 inhibitors with low emetic potential and long duration of action for the treatment of COPD. *J Med Chem*. 2014;57(11):4661–4676.
82. Ostridge K, Williams N, Kim V, et al. Relationship between pulmonary matrix metalloproteinases and quantitative CT markers of small airways disease and emphysema in COPD. *Thorax*. 2016;71(2): 126–132.
83. Churg A, Wang R, Wang X, Onnervik PO, Thim K, Wright JL. Effect of an MMP-9/MMP-12 inhibitor on smoke-induced emphysema and airway remodelling in guinea pigs. *Thorax*. 2007;62(8):706–713.
84. MacNee W. Oxidants/antioxidants and COPD. *Chest*. 2000;117(5 suppl 1): 303S–317S.
85. Bernardo I, Bozinovski S, Vlahos R. Targeting oxidant-dependent mechanisms for the treatment of COPD and its comorbidities. *Pharmacol Ther*. 2015;155:60–79.
86. Bocci V, Valacchi G. Nrf2 activation as target to implement therapeutic treatments. *Front Chem*. 2015;3:4.
87. Wise RA, Holbrook JT, Criner G, et al. Lack of effect of oral sulforaphane administration on Nrf2 expression in COPD: a randomized, double-blind, placebo controlled trial. *PLoS One*. 2016;11(11): e0163716.
88. Dal Negro RW, Visconti M, Turco P. Efficacy of erdosteine 900 versus 600 mg/day in reducing oxidative stress in patients with COPD exacerbations: results of a double blind, placebo-controlled trial. *Pulm Pharmacol Ther*. 2015;33:47–51.
89. Moretti M. Erdosteine: its relevance in COPD treatment. *Expert Opin Drug Metab Toxicol*. 2009;5(3):333–343.

90. Tse HN, Raiteri L, Wong KY, et al. High-dose N-acetylcysteine in stable COPD: the 1-year, double-blind, randomized, placebo-controlled HIACE study. *Chest*. 2013;144(1):106–118.
91. Zheng JP, Wen FQ, Bai CX, et al. Twice daily N-acetylcysteine 600 mg for exacerbations of chronic obstructive pulmonary disease (PANTHEON): a randomised, double-blind placebo-controlled trial. *Lancet Respir Med*. 2014;2(3):187–194.
92. Decramer M, Rutten-van Molken M, Dekhuijzen PN, et al. Effects of N-acetylcysteine on outcomes in chronic obstructive pulmonary disease (Bronchitis Randomized on NAC Cost-Utility Study, BRONCUS): a randomised placebo-controlled trial. *Lancet*. 2005;365(9470):1552–1560.
93. Pela R, Calcagni AM, Subiaco S, Isidori P, Tubaldi A, Sanguinetti CM. N-acetylcysteine reduces the exacerbation rate in patients with moderate to severe COPD. *Respiration*. 1999;66(6):495–500.
94. Schermer T, Chavannes N, Dekhuijzen R, et al. Fluticasone and N-acetylcysteine in primary care patients with COPD or chronic bronchitis. *Respir Med*. 2009;103(4):542–551.
95. Yu H, Li Q, Kolosov VP, Perelman JM, Zhou X. Regulation of cigarette smoke-mediated mucin expression by hypoxia-inducible factor-1 α via epidermal growth factor receptor-mediated signaling pathways. *J Appl Toxicol*. 2012;32(4):282–292.
96. Rubin BK, Priftis KN, Schmidt HJ, Henke MO. Secretory hyperresponsiveness and pulmonary mucus hypersecretion. *Chest*. 2014;146(2):496–507.
97. Polosukhin VV, Cates JM, Lawson WE, et al. Hypoxia-inducible factor-1 signalling promotes goblet cell hyperplasia in airway epithelium. *J Pathol*. 2011;224(2):203–211.
98. Curran DR, Cohn L. Advances in mucous cell metaplasia: a plug for mucus as a therapeutic focus in chronic airway disease. *Am J Respir Cell Mol Biol*. 2010;42(3):268–275.
99. Rennard SI, Sun SX, Tourkodimitris S, et al. Roflumilast and dyspnea in patients with moderate to very severe chronic obstructive pulmonary disease: a pooled analysis of four clinical trials. *Int J Chron Obstruct Pulmon Dis*. 2014;9:657–673.
100. Grubek-Jaworska H, Plapinska M, Hermanowicz-Salamon J, et al. IL-6 and IL-13 in induced sputum of COPD and asthma patients: correlation with respiratory tests. *Respiration*. 2012;84(2):101–107.
101. Lee JS, Rosengart MR, Kondragunta V, et al. Inverse association of plasma IL-13 and inflammatory chemokines with lung function impairment in stable COPD: a cross-sectional cohort study. *Respir Res*. 2007;8:64.
102. Tanabe T, Kanoh S, Tsumura K, Yamazaki Y, Kubo K, Rubin BK. Clarithromycin inhibits interleukin-13-induced goblet cell hyperplasia in human airway cells. *Am J Respir Cell Mol Biol*. 2011;45(5):1075–1083.
103. Rennard SI, Flavin SK, Agarwal PK, Lo KH, Barnathan ES. Long-term safety study of infliximab in moderate-to-severe chronic obstructive pulmonary disease. *Respir Med*. 2013;107(3):424–432.
104. Aaron SD, Vandemheen KL, Maltais F, et al. TNF α antagonists for acute exacerbations of COPD: a randomised double-blind controlled trial. *Thorax*. 2013;68(2):142–148.
105. Chung KF. Inflammatory mediators in chronic obstructive pulmonary disease. *Curr Drug Targets Inflamm Allergy*. 2005;4(6):619–625.
106. Betts JC, Mayer RJ, Tal-Singer R, et al. Gene expression changes caused by the p38 MAPK inhibitor diltapimod in COPD patients: analysis of blood and sputum samples from a randomized, placebo-controlled clinical trial. *Pharmacol Res Perspect*. 2015;3(1):e00094.
107. MacNee W, Allan RJ, Jones I, De Salvo MC, Tan LF. Efficacy and safety of the oral p38 inhibitor PH-797804 in chronic obstructive pulmonary disease: a randomised clinical trial. *Thorax*. 2013;68(8):738–745.
108. Singh D, Smyth L, Borrill Z, Sweeney L, Tal-Singer R. A randomized, placebo-controlled study of the effects of the p38 MAPK inhibitor SB-681323 on blood biomarkers of inflammation in COPD patients. *J Clin Pharmacol*. 2010;50(1):94–100.
109. Watz H, Barnacle H, Hartley BF, Chan R. Efficacy and safety of the p38 MAPK inhibitor losmapimod for patients with chronic obstructive pulmonary disease: a randomised, double-blind, placebo-controlled trial. *Lancet Respir Med*. 2014;2(1):63–72.
110. Geraghty P, Hardigan A, Foronjy RF. Cigarette smoke activates the proto-oncogene c-src to promote airway inflammation and lung tissue destruction. *Am J Respir Cell Mol Biol*. 2014;50(3):559–570.
111. Gagliardo R, Chanez P, Profita M, et al. IkappaB kinase-driven nuclear factor-kappaB activation in patients with asthma and chronic obstructive pulmonary disease. *J Allergy Clin Immunol*. 2011;128(3):635–645. e631–e632.
112. Marian E, Baraldo S, Visentin A, et al. Up-regulated membrane and nuclear leukotriene B4 receptors in COPD. *Chest*. 2006;129(6):1523–1530.
113. Woodruff PG, Albert RK, Bailey WC, et al. Randomized trial of zileuton for treatment of COPD exacerbations requiring hospitalization. *COPD*. 2011;8(1):21–29.
114. Snell N, Foster M, Vestbo J. Efficacy and safety of AZD1981, a CRTH2 receptor antagonist, in patients with moderate to severe COPD. *Respir Med*. 2013;107(11):1722–1730.
115. Mantell S, Jones R, Trevethick M. Design and application of locally delivered agonists of the adenosine A(2A) receptor. *Expert Rev Clin Pharmacol*. 2010;3(1):55–72.
116. Mathioudakis A, Chatzimavridou-Grigoriadou V, Evangelopoulou E, Mathioudakis G. Vasoactive intestinal Peptide inhaled agonists: potential role in respiratory therapeutics. *Hippokratia*. 2013;17(1):12–16.
117. Burian B, Storka A, Marzluf BA, et al. Vasoactive intestinal peptide (VIP) receptor expression in monocyte-derived macrophages from COPD patients. *Peptides*. 2010;31(4):603–608.
118. Onoue S, Aoki Y, Matsui T, et al. Formulation design and in vivo evaluation of dry powder inhalation system of new vasoactive intestinal peptide derivative ([R(15, 20, 21), L(17), A(24, 25), des-N(28)]-VIP-GRR) in experimental asthma/COPD model rats. *Int J Pharm*. 2011;410(1–2):54–60.
119. Wu D, Lee D, Sung YK. Prospect of vasoactive intestinal peptide therapy for COPD/PAH and asthma: a review. *Respir Res*. 2011;12:45.
120. Stoll P, Bahker A, Ulrich M, et al. The dendritic cell high-affinity IgE receptor is overexpressed in both asthma and severe COPD. *Clin Exp Allergy*. 2016;46(4):575–583.
121. Raverdeau M, Mills KH. Modulation of T cell and innate immune responses by retinoic acid. *J Immunol*. 2014;192(7):2953–2958.
122. Ho TW, Tsai YJ, Ruan SY, et al. In-hospital and one-year mortality and their predictors in patients hospitalized for first-ever chronic obstructive pulmonary disease exacerbations: a nationwide population-based study. *PLoS One*. 2014;9(12):e114866.
123. Kaczmarek P, Sladek K, Skucha W, et al. The influence of simvastatin on selected inflammatory markers in patients with chronic obstructive pulmonary disease. *Pol Arch Med Wewn*. 2010;120(1–2):11–17.
124. Lawes CM, Thornley S, Young R, et al. Statin use in COPD patients is associated with a reduction in mortality: a national cohort study. *Prim Care Respir J*. 2012;21(1):35–40.
125. Hodge G, Jersmann H, Tran HB, et al. Lymphocyte senescence in COPD is associated with decreased histone deacetylase 2 expression by pro-inflammatory lymphocytes. *Respir Res*. 2015;16:130.
126. Podolin PL, Foley JP, Carpenter DC, et al. T cell depletion protects against alveolar destruction due to chronic cigarette smoke exposure in mice. *Am J Physiol Lung Cell Mol Physiol*. 2013;304(5):L312–L323.
127. Moretto N, Caruso P, Bosco R, et al. CHF6001 I: a novel highly potent and selective phosphodiesterase 4 inhibitor with robust anti-inflammatory activity and suitable for topical pulmonary administration. *J Pharmacol Exp Ther*. 2015;352(3):559–567.
128. Villetti G, Carnini C, Battipaglia L, et al. CHF6001 II: a novel phosphodiesterase 4 inhibitor, suitable for topical pulmonary administration – in vivo preclinical pharmacology profile defines a potent anti-inflammatory compound with a wide therapeutic window. *J Pharmacol Exp Ther*. 2015;352(3):568–578.

129. Watz H, Mistry SJ, Lazaar AL; Investigators IPC. Safety and tolerability of the inhaled phosphodiesterase 4 inhibitor GSK256066 in moderate COPD. *Pulm Pharmacol Ther.* 2013;26(5):588–595.
130. Goudie AR, Lipworth BJ, Hopkinson PJ, Wei L, Struthers AD. Tadalafil in patients with chronic obstructive pulmonary disease: a randomised, double-blind, parallel-group, placebo-controlled trial. *Lancet Respir Med.* 2014;2(4):293–300.
131. Kuna P, Jenkins M, O'Brien CD, Fahy WA. AZD9668, a neutrophil elastase inhibitor, plus ongoing budesonide/formoterol in patients with COPD. *Respir Med.* 2012;106(4):531–539.
132. Henriksen PA. The potential of neutrophil elastase inhibitors as anti-inflammatory therapies. *Curr Opin Hematol.* 2014;21(1):23–28.
133. Dahl R, Titlestad I, Lindqvist A, et al. Effects of an oral MMP-9 and -12 inhibitor, AZD1236, on biomarkers in moderate/severe COPD: a randomised controlled trial. *Pulm Pharmacol Ther.* 2012; 25(2):169–177.
134. Woodruff PG, Wolff M, Hohlfeld JM, et al. Safety and efficacy of an inhaled epidermal growth factor receptor inhibitor (BIBW 2948 BS) in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 2010;181(5):438–445.
135. Damera G, Jester WF, Jiang M, et al. Inhibition of myristoylated alanine-rich C kinase substrate (MARCKS) protein inhibits ozone-induced airway neutrophilia and inflammation. *Exp Lung Res.* 2010; 36(2):75–84.
136. Ha EV, Rogers DF. Novel therapies to inhibit mucus synthesis and secretion in airway hypersecretory diseases. *Pharmacology.* 2016;97(1–2): 84–100.
137. Astrand AB, Hemmerling M, Root J, et al. Linking increased airway hydration, ciliary beating, and mucociliary clearance through ENaC inhibition. *Am J Physiol Lung Cell Mol Physiol.* 2015;308(1): L22–L32.
138. Black PN, Morgan-Day A, McMillan TE, Poole PJ, Young RP. Randomised, controlled trial of N-acetylcysteine for treatment of acute exacerbations of chronic obstructive pulmonary disease [ISRCTN21676344]. *BMC Pulm Med.* 2004;4:13.
139. Ayfer Aytemur Z, Baysak A, Ozdemir O, Kose T, Sayiner A. N-acetylcysteine in patients with COPD exacerbations associated with increased sputum. *Wien Klin Wochenschr.* 2015;127(7–8):256–261.
140. Dransfield MT, Wilhelm AM, Flanagan B, et al. Acquired cystic fibrosis transmembrane conductance regulator dysfunction in the lower airways in COPD. *Chest.* 2013;144(2):498–506.
141. Bodas M, Min T, Mazur S, Vij N. Critical modifier role of membrane-cystic fibrosis transmembrane conductance regulator-dependent ceramide signaling in lung injury and emphysema. *J Immunol.* 2011; 186(1):602–613.
142. Jones PW, Rames AD. TESRA (treatment of emphysema with a selective retinoid agonist) study results. *Am J Respir Crit Care Med.* 2011;183:A6418.
143. Stolk J, Stockley RA, Stoel BC, et al. Randomised controlled trial for emphysema with a selective agonist of the γ -type retinoic acid receptor. *Eur Respir J.* 2012;40(2):306–312.

International Journal of COPD

Publish your work in this journal

The International Journal of COPD is an international, peer-reviewed journal of therapeutics and pharmacology focusing on concise rapid reporting of clinical studies and reviews in COPD. Special focus is given to the pathophysiological processes underlying the disease, intervention programs, patient focused education, and self management protocols.

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-chronic-obstructive-pulmonary-disease-journal>

Dovepress

This journal is indexed on PubMed Central, MedLine and CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.