



OPEN ACCESS

# 'WNT-er is coming': WNT signalling in chronic lung diseases

H A Baarsma,<sup>1</sup> M Königshoff<sup>1,2</sup>

<sup>1</sup>Comprehensive Pneumology Center, Helmholtz Center Munich, Ludwig Maximilians University Munich, University Hospital Grosshadern, Member of the German Center for Lung Research (DZL), Munich, Germany

<sup>2</sup>Division of Pulmonary Sciences and Critical Care Medicine, Department of Medicine, University of Colorado School of Medicine, Aurora, Colorado, USA

## Correspondence to

Professor M Königshoff, Division of Pulmonary Sciences and Critical Care Medicine, Department of Medicine, University of Colorado School of Medicine, Anschutz Medical Campus M/S C 272, 12700 E 19th Avenue, RC2, 9th Floor, Aurora, CO 80045-2581, USA; Melanie.Koenigshoff@ucdenver.edu

Received 20 November 2016

Revised 1 March 2017

Accepted 16 March 2017

Published Online First

17 April 2017

## ABSTRACT

Chronic lung diseases represent a major public health problem with only limited therapeutic options. An important unmet need is to identify compounds and drugs that target key molecular pathways involved in the pathogenesis of chronic lung diseases. Over the last decade, there has been extensive interest in investigating Wntless/integrase-1 (WNT) signalling pathways; and WNT signal alterations have been linked to pulmonary disease pathogenesis and progression. Here, we comprehensively review the cumulative evidence for WNT pathway alterations in chronic lung pathologies, including idiopathic pulmonary fibrosis, pulmonary arterial hypertension, asthma and COPD. While many studies have focused on the canonical WNT/ $\beta$ -catenin signalling pathway, recent reports highlight that non-canonical WNT signalling may also significantly contribute to chronic lung pathologies; these studies will be particularly featured in this review. We further discuss recent advances uncovering the role of WNT signalling early in life, the potential of pharmacologically modulating WNT signalling pathways and highlight (pre)clinical studies describing promising new therapies for chronic lung diseases.

## THE BASICS: WNT SIGNALLING

The Wntless/integrase-1 (WNT) signalling pathways represent classical developmentally active pathways required for proper organ development. WNT ligands comprise a family of secreted glycoproteins that instruct cells in the respiratory system to adopt particular fates throughout lung development as well as tissue homeostasis in adulthood. In humans, the WNT ligand family is composed of 19 distinct ligands, which are historically classified based on their amino acid sequence rather than their functional properties. Classically, WNT signalling has been separated into canonical and non-canonical signalling. WNT signalling that relies on the activation of the transcriptional coactivator  $\beta$ -catenin is designated as canonical WNT signalling (figure 1) and pathways activated by WNT ligands independently of  $\beta$ -catenin are classified as non-canonical WNT pathways (figure 2). We start off with an explanation of the well described canonical WNT signalling pathway, in which WNT ligands activate  $\beta$ -catenin-mediated gene transcription. In the absence of specific WNT ligands, cytosolic  $\beta$ -catenin is tightly regulated by the so-called ' $\beta$ -catenin destruction complex', a multiprotein complex that targets  $\beta$ -catenin via phosphorylation and ubiquitination for proteasomal degradation (figure 1). The core of the ' $\beta$ -catenin destruction complex' is composed of the proteins axin, adenomatous polyposis coli, casein kinase-1 and glycogen synthase kinase-3 $\beta$  (GSK-3 $\beta$ ). The latter is the primary kinase

involved in the phosphorylation and subsequent degradation of  $\beta$ -catenin. Binding of a specific WNT ligand (eg, WNT-3A) to one of the Frizzled receptors (FZD<sub>1</sub> through FZD<sub>10</sub>) and subsequent activation of the low-density lipoprotein receptor-related proteins 5 and 6 (LRP5/6) co-receptors triggers an intracellular signalling cascade, which results in inactivation of the ' $\beta$ -catenin destruction complex'. Hence, cytosolic  $\beta$ -catenin can accumulate, translocate to the nucleus and, in association with T cell factor/lymphoid enhancer factor-1 (TCF/LEF) family of transcription factors, induce specific gene expression (figure 1).<sup>1</sup>

Activation of non-canonical WNT signalling also relies on the binding of specific WNT ligands (eg, WNT-4 or WNT-5A) to FZD receptors; however, it appears to be independent of LRP5/6 co-receptors. Non-canonical WNT signalling results in the activation of intracellular signalling molecules involved in planar cell polarity (PCP pathway), calcium/calmodulin-dependent protein kinase II (Ca<sup>2+</sup>/CAMKII) signalling and/or various less well defined downstream effector molecules (figure 2). Notably, some classically defined non-canonical WNT ligands are able to negatively influence canonical WNT/ $\beta$ -catenin signalling. Moreover, single WNT ligands can activate multiple signalling pathways suggesting that WNT ligands are not intrinsically canonical or non-canonical. Selectivity in receptor-ligand binding (eg, FZD-WNT interaction) likely dictates the outcome of downstream signalling.<sup>2,3</sup> Indeed, a biochemical study demonstrated that WNT ligands can selectively bind to specific FZD receptors, and that respective WNT-FZD pairs exert functional selectivity in downstream signalling.<sup>4</sup> These data emphasise the interconnectivity and complexity of canonical and non-canonical WNT signalling<sup>1,5-7</sup> (figure 2). The dynamics of WNT and FZD expression in complex biological systems in vivo is currently unknown, thus a better understanding of receptor-ligand interactions in WNT signalling is required to decipher how exactly WNT ligands function. As such, the separation of WNT signalling in purely canonical and non-canonical signalling pathways appears to be outdated and certainly oversimplifies the complexity of this signalling pathway; however for uniformity reasons we maintain this nomenclature in this review. Whenever possible, we mention which WNT ligands, receptors and/or downstream signalling molecules are involved when we refer to canonical or non-canonical WNT signalling.


Over the last decade, there has been extensive interest in investigating WNT signalling pathways in chronic lung diseases. Several components of the WNT pathways serve as potent oncogenes and



CrossMark

To cite: Baarsma HA, Königshoff M. *Thorax* 2017;**72**:746–759.






**Figure 1** Schematic representation of canonical WNT/β-catenin signalling. *Left side:* cytosolic β-catenin is rapidly degraded by the β-catenin destruction complex in the absence of extracellular WNT ligands. The core of the β-catenin destruction complex is composed of: adenomatous polyposis coli (APC), axin, casein kinase-1 (CK-1) and glycogen synthase kinase-3 (GSK-3). GSK-3 is the primary kinase involved in the degradation of β-catenin. *Right side:* an extracellular WNT ligand binds and activates Frizzled (FZD) and the low density lipoprotein receptor-related proteins 5 and 6 (LRP5/6), which results in the activation of and intercellular signalling cascade that leads to the inhibition of the β-catenin destruction complex. Hence, β-catenin can accumulate and translocate to the nucleus to induce gene transcription. In the nucleus β-catenin can associate with various transcriptional coactivators, including T cell factor (TCF) and lymphoid enhancer factor (LEF).

WNT signalling has been linked to lung cancer, which has been extensively reviewed previously and will not be included in this review.<sup>5 8–13</sup> Here, we aimed to comprehensively review cumulative evidence for WNT pathway alterations in chronic lung pathologies, including idiopathic pulmonary fibrosis (IPF), pulmonary arterial hypertension (PAH), asthma and COPD. Early studies have largely focused on the canonical WNT/β-catenin signalling pathway and only recently several reports suggest that non-canonical WNT signalling might also contribute significantly to chronic lung pathologies. These studies will be highlighted in this review. We further discuss recent advances in our knowledge on the role of WNT signalling in early life, and feature novel developments and the potential application of WNT signalling modulation for drug development and (pre) clinical studies.

### WNT SIGNALLING IN EARLY LIFE

Despite intensive research efforts, the aetiology of major chronic lung diseases in children and adults remains elusive. Several lines of evidence indicate that prenatal and/or early postnatal lung injuries will have important implications for future lung function and increase risk for development of chronic lung diseases later in life.<sup>14</sup> Several reports highlight the functional importance of canonical and non-canonical WNT signalling in lung morphogenesis and postnatal development and this has been reviewed previously.<sup>8 15–19</sup> Ectopic expression of specific WNT ligands during lung development, either those involved in

canonical or non-canonical signalling, can result in severe lung phenotypes, which partially resemble lung diseases observed during adulthood.<sup>5 20–23</sup> Moreover, deletion of β-catenin in epithelial cells of embryonic lungs results in disrupted lung morphogenesis.<sup>24</sup> In contrast, overexpression of a truncated, constitutively active form of β-catenin in Clara cell secretory protein (CCSP) positive cells (Club cells) does not influence lung morphogenesis before birth, but leads postnatally to goblet cell hyperplasia, pulmonary tumour development and airspace enlargement.<sup>25</sup> Collectively, these studies highlight the importance of strict spatiotemporal control of WNT signalling for proper lung development and lung physiology early in life and thereafter into adulthood. More recently, studies have focused on the impact of environmental factors on prenatal and postnatal lung development. Epidemiological studies have shown that maternal smoking is a risk factor for the development of several chronic lung diseases.<sup>26 27</sup> It was demonstrated in mice that maternal smoking negatively affects the mRNA expression of the WNT pathway components *frizzled7* (*Fzd7*; receptor) and *Ctmb1* (gene symbol of β-catenin) as well as the WNT target gene *Fn* (fibronectin) in lung tissue of the offspring.<sup>28</sup> Although not investigated in humans, impairment of these WNT pathway components may have implications for lung development, and fitness later in life, as these WNT components regulate neoangiogenesis and lung branching morphogenesis.<sup>28</sup> Thus, the various WNT pathways are part of a signalling network essential for proper lung development and physiology,




**Figure 2** Schematic representation of signalling cascades involved in non-canonical WNT signalling. An extracellular WNT ligand binds to the Frizzled (FZD) receptor, which can subsequently activate a variety of downstream signalling cascades involved in gene transcription, intercellular actin organisation and/or inhibition of the transcriptional coactivator  $\beta$ -catenin. AC, adenylyl cyclase; PKA, protein kinase A; CREB, cAMP responsive element binding protein; DVL, dishevelled; FZD, Frizzled receptor; JNK, c-Jun-N terminal kinase; PLC, phospholipase C; NF-AT, nuclear factor of activated T cells; PKC, protein kinase C; PXN, paxillin; MRLC, myosin regulatory light chain; RAP1, RAS-related protein 1; CAMKII, calcium/calmodulin-dependent kinase II; NLK, Nemo-like-kinase.

which may contribute to (age-associated) chronic lung diseases when impaired at early stages of life.<sup>29</sup>

### WNT SIGNALLING IN PULMONARY FIBROSIS

Impairment of the WNT signalling pathways and their potential role as therapeutic targets in chronic lung disease was first identified and established in IPF. IPF is a devastating, progressive disease characterised by lung epithelial injury and reprogramming, fibroblasts activation and excessive extracellular matrix (ECM) remodelling resulting in distorted lung architecture and a progressive loss of functional lung tissue.<sup>30</sup> Although our understanding of IPF pathogenesis has significantly improved over the recent years, we currently have only limited pharmacological approaches available to treat the disease.<sup>25 31 32</sup> The concept that developmental pathways (such as WNT, Sonic Hedgehog, Notch pathways) are altered in IPF, arose largely from unbiased gene expression profiling of models of experimental lung fibrosis as well as human IPF lung tissue specimens, which particularly indicated a 'WNT signature' within the IPF lung.<sup>33–36</sup> It has been proposed that IPF lungs show remarkable resemblances to the developing lung in terms of pathology. This further implies that particular developmental signalling pathways may be reactivated in adult tissues following injury and contribute to IPF pathogenesis. Indeed, canonical WNT/ $\beta$ -catenin signalling is active in various cell types in human and experimental pulmonary fibrosis<sup>34 35</sup> (figure 3). More specifically, increased gene expression of *WNT-1*, *WNT-7B*, *WNT-10B*, *FZD2*, *FZD3*, *CTNNB1* ( $\beta$ -catenin) and *LEF1* was observed in lung tissue of individuals with IPF compared with individuals

without IPF (donor).<sup>34</sup> Immunohistochemical analysis localised the classically defined canonical ligands WNT-1 and WNT-3A as well as the WNT effector protein  $\beta$ -catenin largely to bronchial and alveolar epithelium, although increased nuclear  $\beta$ -catenin expression has also been observed in (myo)fibroblasts in fibrotic foci.<sup>34 35 37</sup> Increased expression of WNT target genes and, in particular, the increase in WNT1-inducible signalling protein-1 (WISP1) protein expression suggests that functional WNT/ $\beta$ -catenin signalling activity is enhanced in lung tissue of individuals with IPF.<sup>34 35</sup> Antibody-mediated inhibition of WISP1 results in decreased lung pathology in bleomycin-induced lung fibrosis in mice in vivo and WISP1 has been reported to regulate alveolar epithelial cell function and reprogramming as well as (myo)fibroblast activation, thereby impairing lung function.<sup>35</sup> Successive studies demonstrated that WISP1 is a WNT target gene, and at the same time a common downstream mediator of several profibrotic factors, including (1) miR-92A, a micro-RNA that is downregulated in human and experimental lung fibrosis, and (2) Transforming growth factor  $\beta$  (TGF $\beta$ ) and Tumor necrosis factor  $\alpha$  (TNF $\alpha$ ) signalling in human lung fibroblasts, in which WISP1 regulates cell proliferation in an interleukin 6 (IL-6)-dependent fashion.<sup>38 39</sup> Altogether, these findings corroborate WISP1 as a therapeutic target in IPF. With regard to upstream WNT/ $\beta$ -catenin signalling; extensive research has focused on preventing excessive activation of  $\beta$ -catenin signalling by targeting this developmental pathway at various molecular levels (figure 3). Several studies demonstrate that pharmacological and genetic inhibitors of  $\beta$ -catenin signalling attenuate fibrosis in various organs,



**Figure 3** WNT signalling in idiopathic pulmonary fibrosis (IPF) pathogenesis. Increased pulmonary expression of WNT-1, WNT-7B, WNT-10B, Frizzled receptor (FZD)2 and FZD3 in individuals with IPF. Enhanced expression of transcriptionally active  $\beta$ -catenin in pulmonary epithelial cells (top), as a consequence of Wnt-3A and/or TGF- $\beta$  signalling.  $\beta$ -catenin signalling induces mRNA expression of inflammatory and remodelling markers (eg, IL-1 $\beta$  and WNT1-inducible signalling protein-1 (WISP1)) and regulates alveolar epithelial type II cells (ATII)-to-ATI-cell transdifferentiation, a process implicated in wound healing and tissue regeneration. The profibrotic action of WISP1 can be diminished by neutralising antibodies. In pulmonary fibroblasts (bottom), WNT-5B by activating FZD8, in conjunction with TGF- $\beta$  signalling, causes upregulation of mRNA expression of other WNT signal components, extracellular matrix (ECM) components, and myofibroblast markers. Both the expression of ECM components and markers of myofibroblast differentiation are dependent on activation of transcriptionally active  $\beta$ -catenin. Accumulation of transcriptionally active  $\beta$ -catenin can be prevented by small molecule inhibitors of Tankyrases (eg, XAV939), whereas the interaction of  $\beta$ -catenin with specific transcription factors can be inhibited by ICG-001 ( $\beta$ -catenin/cAMP response element-binding protein (CBP)), PKF115–584 ( $\beta$ -catenin/TCF (T cell factor)) or IQ-1 ( $\beta$ -catenin/p300). WNT-5A, via a yet unknown FZD, induces proliferation and protects cells from oxidative-stress-induced apoptosis. Lipoprotein receptor-related protein 5 (LRP5) and TGF- $\beta$  signalling are indispensable for activation of  $\beta$ -catenin signalling in response to a fibrotic insult. Moreover, LRP5 in macrophages contributes to disease progression. See main text for further details.

including the lung.<sup>40–46</sup> Noteworthy, activation of  $\beta$ -catenin in alveolar epithelial cells at the early stages of injury may reflect an attempt to repair and regenerate, however sustained activation of the pathway may drive inflammation and fibrotic changes in the lung.<sup>9 42 47–53</sup>

The alveolar epithelium, consisting of alveolar epithelial type I and II cells (ATI and ATII cells, respectively), represents a major site of tissue damage during lung injury. ATII cells are capable of self-renewal and exert progenitor function for ATI cells upon alveolar epithelial injury, a process dependent on autocrine WNT/ $\beta$ -catenin signalling.<sup>48 50 54 55</sup> In physiological conditions, mature ATII cells exhibit a low degree of constitutive  $\beta$ -catenin signalling *in vivo*; however, the pathway is activated when the lungs are subjected to bleomycin-induced injury.<sup>48 50</sup> Activation of the WNT/ $\beta$ -catenin signalling pathway is also observed during ATII-to-ATI cell transdifferentiation *in vitro*. The transdifferentiation process is characterised by

increased expression of several WNT ligands, enhanced phosphorylation of WNT signalling intermediates (ie, LRP6 and DVL3), increased transcriptionally active  $\beta$ -catenin and an increase in  $\beta$ -catenin-driven target gene expression.<sup>48 50</sup> Collectively, these data indicate that  $\beta$ -catenin activation during alveolar epithelial cell transdifferentiation is dependent on endogenous, autocrine canonical WNT signalling. This was confirmed by experiments in which Dickkopf-1 (DKK-1), an endogenous inhibitor of canonical WNT signalling, was overexpressed. These experiments demonstrated that DKK-1 attenuates  $\beta$ -catenin-driven target gene expression (ie, *Axin2*) in transdifferentiating ATII cells.<sup>48</sup> Furthermore, inhibition of canonical WNT signalling by either the small molecule PKF115–584 (disruption  $\beta$ -catenin/TCF interaction),  $\beta$ -catenin siRNA or by ectopic expression of *inhibitor of  $\beta$ -catenin and TCF* (ICAT) attenuates the time-dependent increase in ATI cell marker expression in these primary cell cultures.<sup>48 50</sup> Ectopic

expression of ICAT also impairs wound closure after scratch-induced injury.<sup>48</sup> Taken together, these data indicate that WNT/ $\beta$ -catenin signalling regulates alveolar epithelial cell transdifferentiation and repair processes, which might initially be beneficial upon lung injury. Nevertheless, persistent WNT/ $\beta$ -catenin activation in the alveolar epithelium may be detrimental in fibrotic lung diseases, like IPF. An unbiased gene expression screen to identify cell-specific mediators of canonical WNT signalling in primary murine ATII cells identified the proinflammatory cytokine IL-1 $\beta$  as one of the highest upregulated genes induced by WNT-3A stimulation.<sup>47</sup> Subsequent experiments confirmed increased protein level of IL-1 $\beta$  in ATII cells in response to WNT-3A, as well as in an animal model of experimental lung fibrosis and, most importantly, in human IPF. These findings indicate that the alveolar epithelium is a relevant source of proinflammatory/profibrotic cytokines induced by active WNT/ $\beta$ -catenin signalling in pulmonary fibrosis.<sup>47</sup> Among many tested, only four cytokines/growth factors (ie, IL-1 $\beta$ , TNF $\alpha$ , Granulocyte macrophage colony stimulating factor (GM-CSF) and TGF- $\beta$ ) have thus far been identified to be able to drive lung fibrosis.<sup>56</sup> Transient overexpression of IL-1 $\beta$  initially results in inflammatory response with tissue injury followed by a progressive fibrotic response mediated in part by TGF- $\beta$ .<sup>57</sup> Thus, sustained canonical WNT signalling activity in alveolar epithelial cells may contribute via IL-1 $\beta$  to lung fibrosis. Furthermore, alveolar epithelial cell reprogramming and epithelial-to-mesenchymal transition (EMT) potentially contributes to chronic lung diseases, like IPF or COPD.<sup>58–62</sup> TGF- $\beta$  is the most commonly used growth factor for promoting epithelial cell reprogramming and its interaction with WNT signalling is important for cell fate determination during development and in the adult. Zhou *et al.*,<sup>63</sup> reported that TGF- $\beta$  induces the expression of the mesenchymal marker  $\alpha$ -smooth muscle-actin ( $\alpha$ -sm-actin) with concomitant upregulation of transcriptionally active  $\beta$ -catenin in lung epithelial cells. Inhibition of  $\beta$ -catenin diminished TGF- $\beta$ -induced  $\alpha$ -sm-actin expression in these cells, whereas enhanced activation of  $\beta$ -catenin via GSK-3 $\beta$  inhibition (ie, LiCl) augmented this process.<sup>63</sup> Subsequent experiments showed that  $\alpha$ -sm-actin is transcriptionally regulated by a complex consisting of SMAD3/ $\beta$ -catenin/cAMP response element-binding protein binding protein (CBP).<sup>63</sup> Furthermore, Ulsamer *et al.*<sup>45</sup> showed that the EMT process, characterised by increased expression of collagen,  $\alpha$ -sm-actin and the transcription factor twist, is largely dependent on activation of a specific form of  $\beta$ -catenin (ie, pY654- $\beta$ -catenin). Accumulation of pY654- $\beta$ -catenin can be attenuated in vitro and in vivo by inhibition of tankyrase-1 (TNKS1), which partially prevents the mesenchymal changes both in primary alveolar epithelial cells and in an animal model of lung fibrosis.<sup>45</sup> Tankyrases (TNKS1 and 2, respectively) are a class of enzymes that facilitate Axin proteolysis, which results in decreased assembly of the ' $\beta$ -catenin destruction complex' and subsequently increases  $\beta$ -catenin signalling.<sup>1 64</sup> Hence, inactivation of tankyrases attenuates canonical WNT/ $\beta$ -catenin signalling. The therapeutic potential of using tankyrase inhibitors in established experimental lung fibrosis was examined by Wang *et al.*<sup>46</sup> Administration of the tankyrase inhibitor XAV939 starting 10 days after the administration of bleomycin to the mice results in attenuation of  $\beta$ -catenin signalling, improved survival of the mice and a significant reduction in lung pathology.<sup>46</sup> Thus, inhibition of this family of enzymes is of potential interest to target  $\beta$ -catenin signalling and to reduce profibrotic responses in the lung. The advantageous effects of XAV939 may be attributed to reduced fibroblasts proliferation, impaired myofibroblast differentiation and,

potentially, increased differentiation of bone marrow-derived mesenchymal stem cells into epithelial-like cells.<sup>46</sup> However, more extensive in vitro and in vivo analyses still have to be performed to unambiguously support the latter finding. Nevertheless, this combined body of evidence suggests that the large repertoire of  $\beta$ -catenin regulated genes is decisive of cellular fate. In the nucleus, gene transcription induced by  $\beta$ -catenin is facilitated by the recruitment and activation of various factors, including transcription factors, proteins involved in histone methylation and acetylation (ie, MLL1/2, CBP and p300), histone deacetylases and histone modifiers (eg, BRG1).<sup>65</sup>  $\beta$ -Catenin as a transcriptional-coactivator has a plethora of binding partners and its interaction with distinct transcription factors (eg, TCFs, SMADs, forkhead box (FOXO) or HIF1 $\alpha$ ) has important implications for its transcriptional activity and determines the functional output.<sup>66–68</sup> One elegant and promising therapeutic option for IPF is to 're-direct'  $\beta$ -catenin signalling by using specific inhibitors that target the interaction of  $\beta$ -catenin with distinct transcriptional cofactors. This approach has been applied in a study by Henderson *et al.*<sup>42</sup> in which the authors demonstrated that selective inhibition of the  $\beta$ -catenin/CBP interaction by the ICG-001 led to reversal of established bleomycin-induced pulmonary fibrosis and improved epithelial cell integrity, whereas this was not achieved by the food and drugs administration (FDA) approved drug pirfenidone. Based on these results it was proposed that the interaction of  $\beta$ -catenin with CBP drives gene transcription critical for the maintenance of cells in an undifferentiated/proliferative state, whereas the interaction of  $\beta$ -catenin with p300 (a transcriptional coactivator homologous to CBP) initiates cellular differentiation. A subsequent study by Rieger *et al.*<sup>69</sup> dissected the role of p300 and CBP in adult progenitor cell differentiation focusing on ATII-to-ATI cell transdifferentiation. Transdifferentiation of rat ATII cells to ATI-like cells was dependent on the interaction of  $\beta$ -catenin with p300, but independent of the interaction with CBP. Pharmacological or genetic disruption of the  $\beta$ -catenin/p300 interaction resulted in decreased expression of aquaporin5 (AQP5; ATI cell marker) and partial stabilisation of surfactant protein C (ATII cell marker) expression. Interestingly, the interaction between  $\beta$ -catenin and p300 in this context was facilitated by the classically defined non-canonical WNT ligand WNT-5A via a molecular mechanism involving protein kinase C-mediated phosphorylation of p300.<sup>69</sup> This study and various other studies demonstrate that targeting  $\beta$ -catenin-dependent signalling may be beneficial for progenitor cell function and consequently the treatment of fibrosis. The question remains if and which secreted WNT ligands actually represent the driving force for the activation of  $\beta$ -catenin in human and experimental pulmonary fibrosis. A better understanding and determination of (cell)-specific ligand/receptor complexes is of high interest, as it will foster the identification of potential therapeutic targets. To elucidate if canonical WNT ligands contribute to disease pathogenesis, Lam *et al.*<sup>44</sup> investigated the role of the indispensable WNT co-receptors LRP5 and LRP6 in murine and human pulmonary fibrosis. Notably, *LRP5* transcript levels were associated with IPF progression, disease severity at presentation, and negatively correlated with clinical parameters like diffusion capacity of the lung for carbon monoxide (DL<sub>co</sub>) and composite physical index (CPI). This study furthermore demonstrated that LRP5-deficient fibroblasts exhibit reduced capacity to activate  $\beta$ -catenin and that mice lacking LRP5 (*LRP5*<sup>-/-</sup> mice) were protected against bleomycin-induced lung fibrosis.<sup>44</sup> In a subsequent study by the same group, microarray analysis revealed that *LRP5*<sup>-/-</sup> mice that were exposed to bleomycin showed enrichment for pathways

related to ECM processing and the innate immune response, suggesting that the immune cell-ECM remodelling axis is important in fibrosis. Macrophage-specific deletion of  $\beta$ -catenin did not have an effect on the development of bleomycin-induced lung fibrosis; however, the resolution of fibrosis in this model was accelerated.<sup>70</sup> Similarly, in a murine model of asbestos-induced, non-resolving lung fibrosis was shown that loss of LRP5 does not influence development of fibrosis, but does result in delayed progression of fibrosis.<sup>70</sup> These data indicate that macrophages with activated WNT/ $\beta$ -catenin signalling play an important role in sustaining pulmonary fibrosis.

In addition to LRP5/6, only a few WNT receptors have been investigated thus far in chronic lung diseases. The previously mentioned study also reported high *FZD8* expression in individuals with IPF, which correlated to more rapid disease progression.<sup>44</sup> Indeed, cellular signalling mediated via *FZD8* is involved in fibrotic responses in the lung.<sup>71</sup> *FZD8* is the most profound upregulated *FZD* receptor in primary human lung fibroblasts in response to TGF- $\beta$  stimulation. Moreover, genetic silencing of *FZD8*, in part, prevents the profibrotic action of TGF- $\beta$  in pulmonary fibroblasts in vitro and in bleomycin-induced lung fibrosis in vivo. Importantly, profibrotic signalling via *FZD8* appears to be mediated by WNT-5B and is independent of  $\beta$ -catenin signalling.<sup>71</sup> These findings suggest that non-canonical WNT signalling also contributes to IPF pathogenesis. In line with this, increased WNT-5A expression has been demonstrated in IPF.<sup>72–74</sup> Vuga *et al.*,<sup>74</sup> reported that this non-canonical WNT ligand mediates ECM deposition by pulmonary fibroblasts and protects these cells against oxidative stress-induced apoptosis, thereby potentially contributing to IPF. Collectively, cumulative evidence strongly suggests that the developmental WNT signalling pathways are able to drive fibrotic changes in the lung and are of great therapeutic interest for the treatment of IPF (figure 3).

### WNT SIGNALLING IN PULMONARY ARTERIAL HYPERTENSION

PAH is a well recognised complication of interstitial lung diseases like IPF, although PAH can also develop without underlying parenchymal lung disease(s). Both idiopathic PAH (IPAH), as well as common acquired forms of PAH, are characterised by a progressive increase in pulmonary vascular resistance, due to (1) loss of peripheral pulmonary arteries as a consequence of endothelial cell apoptosis and (2) occlusion of larger proximal arteries by increased smooth muscle cell proliferation and enhanced ECM deposition. The pathomechanisms in PAH are multifactorial and not fully understood, but include chronic hypoxia and hypoxic vasoconstriction, mechanical lung stress, smoking effects and inflammation.<sup>75</sup> Aberrant growth factor signalling has also been implicated in the pathogenesis of PAH. Heterozygous loss of function mutations in bone morphogenetic protein receptor II (BMPRII) is observed in both sporadic and familial cases of IPAH.<sup>76–77</sup> These findings imply that dysfunction of the developmentally active Bone morphogenetic protein (BMP) signalling is involved in PAH development. Interestingly, BMP-2 via BMPRII recruits canonical and non-canonical WNT pathways to promote pulmonary arterial endothelial proliferation, survival and migration.<sup>78</sup> Activation of canonical WNT/ $\beta$ -catenin signalling, as well as non-canonical WNT/RHOA/RAC1 signalling, are necessary for vascular growth in vivo, suggesting that WNT signalling pathways are required for BMP-2-mediated angiogenesis and therefore might be beneficial in PAH.<sup>78</sup> Nevertheless, an independent study investigating genetic signatures common across multiple cell lines that are

associated with pathological processes in PAH identified WNT signalling as a target pathway, which was validated further in vitro.<sup>79</sup> The data from this study suggest that alterations in WNT signalling are a common molecular defect in both heritable PAH and IPAH, which is linked to decreased BMPRII signalling.<sup>79</sup> Several components of the PCP pathway (non-canonical WNT signalling) are upregulated in pulmonary resistance vessels in IPAH.<sup>80</sup> Accordingly, WNT ligand expression (eg, WNT-5A) positively correlated to pulmonary arterial pressure in pulmonary arterial smooth muscle cells from patients with PAH.<sup>81</sup> Furthermore, Wu *et al.*<sup>82</sup> investigated the expression and molecular target(s) of several microRNAs in the pathogenesis of PAH and discovered that Mir-199b-5p was overexpressed in PAH and negatively regulated the expression of GSK-3 $\beta$  resulting in enhanced  $\beta$ -catenin-driven gene transcription. These findings suggest enhanced WNT signalling in PAH. Thus, WNT signalling pathways have a crucial role in the regulation of pulmonary angiogenesis and vascular remodelling, and therapies that modulate WNT pathway activity could be of use in patients with PAH. The complexity of WNT pathway signalling, the extent of crosstalk with other (growth factor-dependent) signalling pathways and the plethora of signalling components, remain challenges to identify safe and effective therapies that specifically target WNT signalling in PAH.

### WNT SIGNALLING IN ASTHMA

Asthma is a chronic airway disease with high prevalence in children as well as adults. The hallmark pathological features of asthma include eosinophilic airway inflammation and structural changes within the airways (airway remodelling), which are associated with an irreversible loss in lung function.<sup>83</sup> Identifying genetic determinants for lung function is important in providing insight into the pathophysiology of asthma and other chronic lung diseases. One of the first studies suggesting a genetic link between WNT signalling and asthma was performed by Sharma *et al.*,<sup>84</sup> who demonstrated that WNT signalling genes are associated with impaired lung function in asthmatic children with *WISP1* associating with FEV<sub>1</sub> and FVC, whereas WNT inhibitory factor-1 (*WIF-1*) associated with FVC and FEV<sub>1</sub>/FEV (not with FEV<sub>1</sub> alone). Furthermore, a single nucleotide polymorphism (SNP; rs2929973) in the *WISP1* gene is significantly associated with FEV<sub>1</sub> in asthmatic children, whereas an SNP in *WIF-1* does influence lung function (ie, FEV<sub>1</sub>).<sup>85</sup> Moreover, a recent publication, which used a genome-wide association study (GWAS)-based analysis to prioritise asthma susceptibility genes, revealed two biological pathways to be enriched in asthma pathogenesis: (1) cytokine-cytokine receptor interactions and (2) WNT signalling pathway(s). The same study identified a novel susceptibility locus near WNT signalling genes, in particular SNPs located near *FZD3* and *FZD6*.<sup>86</sup>


Airway inflammation in allergic asthma is mainly characterised by a T helper 2 cell (Th2) signature and eosinophilia.<sup>87</sup> In human asthmatic airways, gene expression of multiple WNT ligands is positively (*WNT-3A*, *WNT-5A*, *WNT-6* and *WNT-10A*) or negatively (*WNT-5B*) associated with a Th2 signature. In addition, *FZD5* (WNT receptor) is upregulated in patients with Th2-high asthma.<sup>87</sup> Recently, a study investigated whether eosinophil-induced airway remodelling in asthma is associated with alterations in WNT expression. Eosinophils derived from patients with asthma enhanced the expression of *WNT-5A*, *TGF- $\beta$*  and gene expression of several ECM components in human airway smooth muscle cells. In addition, these eosinophils induced airway smooth muscle cell proliferation, which is

potentially mediated by WNT-5A. This study links WNT signalling to inflammatory processes, and airway remodelling; however, this was not directly addressed experimentally.<sup>88</sup> Furthermore, these studies suggest that WNT signalling potentially modulates allergic responses in the lung. By using the mouse model of ovalbumin (OVA)-induced allergic asthma, Reuter *et al*<sup>89</sup> examined the impact of WNT-1 on development of allergic lung disease. Overexpression of this specific WNT ligand in CCSP-positive cells (Club cells) had beneficial effects on allergic airway disease, demonstrated by a reduction in airway hyper-responsiveness (AHR), decreased eosinophilia in the bronchoalveolar lavage and a reduced number of mucus cells in comparison to animals that did not overexpress the ligand. The beneficial effects of WNT-1 expression were due to impairment of dendritic cell-dependent activation of T cells.<sup>89</sup> In addition to inflammation, the impact of WNT signalling on airway remodelling has been investigated using animal models of allergic asthma, in which increased expression of specific WNT ligands (ie, *Wnt-5A* and *Wnt-7B*),  $\beta$ -catenin and the canonical WNT target gene *Axin2* was observed.<sup>90–91</sup> Genetic silencing of  $\beta$ -catenin, by siRNA, attenuated inflammation and airway remodelling in asthmatic mice in vivo.<sup>90</sup> Airway remodelling may be, in part, a consequence of a process in which pulmonary epithelial cells acquire a mesenchymal phenotype (EMT). In accordance, the aeroallergen house dust mite, together with TGF- $\beta$ , induces mesenchymal transition of bronchial epithelial cells. This process was associated with loss of E-cadherin, a binding partner of  $\beta$ -catenin at the plasma membrane. This potentially leads to dissociation of  $\beta$ -catenin from the membrane and the subsequent translocation of  $\beta$ -catenin to the nucleus, where it assists the transcription of mesenchymal genes.<sup>92</sup> Furthermore, in a rat model of allergic asthma, airway remodelling in response to OVA exposure was partially prevented by antibody-mediated inhibition of the WNT target gene *WISP1*.<sup>93</sup> In a more recent study, no differences were observed in  $\beta$ -catenin expression in the airway smooth muscle layer in either an animal model of allergic asthma or in smooth muscle biopsies from asthmatic individuals compared with non-asthmatic donors.<sup>94</sup>  $\beta$ -Catenin-driven gene expression (*Axin2*, *Wisp1* and *c-Myc*), however, was highly increased in the used mouse model of OVA-induced asthma. Moreover, pharmacological inhibition of  $\beta$ -catenin/CBP interaction by the small molecule ICG-001 prevented smooth muscle remodelling and ECM expression in vitro and in vivo models of asthma.<sup>94</sup> In agreement,  $\beta$ -catenin signalling is required for TGF- $\beta$ -induced expression of ECM proteins by human airway smooth muscle cells and pulmonary fibroblasts. Furthermore, enhanced synthesis of the ECM protein fibronectin by human airway smooth muscle cells was observed after ectopic expression of constitutively active  $\beta$ -catenin (ie, S33Y- $\beta$ -catenin; resistant to GSK-3-mediated proteasomal degradation).<sup>95–96</sup> On the other hand, while overexpression of a non-degradable form of  $\beta$ -catenin (S37A- $\beta$ -catenin) in fibroblasts is insufficient to induce TGF- $\beta$ , profibrotic marker expression and/or myofibroblast differentiation, it does increase fibroblast migration and proliferation.<sup>43</sup> Of note,  $\beta$ -catenin might further play an important role in airway smooth muscle proliferation and contraction independently of WNT signalling. The latter is presumably due to its role in cell-cell contacts as a component of cell adherens junctions.<sup>97–98</sup> This may represent an additional mechanism by which  $\beta$ -catenin contributes to asthma pathology. By and large, these findings indicate that  $\beta$ -catenin is a potential therapeutic target in asthma, which is discussed in more detail in a comprehensive review by Kumawat *et al*.<sup>17</sup>

Remarkably, among all the WNT ligands detected in human airway smooth muscle cells, *WNT-5A* is the most abundant WNT ligand expressed.<sup>99</sup> Moreover, *WNT-5A* protein expression is approximately twofold higher in airway smooth muscle of asthmatics compared with healthy donors. These data suggest a potential role for non-canonical WNT signalling as well in asthma.<sup>100</sup> Stimulation of airway smooth muscle cells with solely *WNT-5A* was not sufficient to induce ECM deposition; however, siRNA-mediated silencing of *WNT-5A* largely blunted TGF- $\beta$ -induced ECM deposition (collagen 1 $\alpha$ 1 and fibronectin) by airway smooth muscle cells.<sup>99</sup> Furthermore, subsequent in vitro studies demonstrated that *WNT-5A* as well as *WNT-11* may contribute to AHR by inducing actin polymerisation and the upregulation of  $\alpha$ -sm-actin in human airway smooth muscle, respectively.<sup>101–102</sup> Whether this pathway is also involved in asthma pathogenesis in vivo is not yet elucidated. In summary, these data indicate that specific WNT ligands, by acting on various cell types, potentially contribute to pathogenic processes in asthma, including airway inflammation and remodelling, and highlight (canonical WNT-driven)  $\beta$ -catenin as a potential therapeutic target in asthma.

### WNT SIGNALLING IN COPD

COPD is characterised by chronic airflow limitation caused by (1) small airway disease, which is composed of (small) airway remodelling and chronic bronchitis, and (2) parenchymal destruction, which leads to emphysema. The relative contribution of each of these pathological features varies in individual patients who suffer from COPD. Given the importance for WNT/ $\beta$ -catenin signalling for lung development and growth, this pathway represents a prime target for initiation of lung repair and regeneration. A tremendous challenge in emphysema is the restoration of functional lung tissue and one current concept is that the progressive destruction of tissue is partially due to the inability of the lung to activate self-repair mechanisms in COPD. Similarly to IPF, no therapy is currently available for COPD that stabilises or reverses disease progression.<sup>103–104</sup> Cigarette smoke, the main risk factor for developing COPD, reduces canonical WNT signalling in vitro and in vivo in human bronchial epithelial cells.<sup>105–106</sup> Other studies report that bronchial epithelial cells of smokers and individuals with COPD might exhibit enhanced nuclear  $\beta$ -catenin expression and that components of cigarette smoke (eg, nicotine) can induce EMT in a WNT-3A/ $\beta$ -catenin-dependent manner.<sup>62–107</sup> Nevertheless, transcript levels of several components of canonical WNT signalling (eg, *CTNNB1*, *GSK-3 $\beta$*  and *TCF4*) are significantly lower in peripheral lung tissue of individuals with COPD compared with smokers without COPD, indicating that processes involved in COPD pathogenesis affect components of canonical WNT signalling independently of smoking.<sup>105</sup> These data confirm previous findings in which expression profiles obtained from lung tissue specimens from smokers without obstruction and individuals with COPD were compared.<sup>108</sup> In this study, several biological pathways that may be relevant to COPD pathogenesis independently of smoking were identified, including TGF- $\beta$  signalling, proteins involved in focal adhesion and WNT signalling pathways.<sup>108</sup> Notably, in experimental and human COPD/emphysema canonical WNT signalling is silenced in the alveolar epithelium as observed by decreased nuclear expression of  $\beta$ -catenin, a surrogate marker for canonical WNT signalling activity (figure 5).<sup>109</sup> Several recent studies investigated potential underlying mechanism(s) for reduced  $\beta$ -catenin signalling in COPD.<sup>110–112</sup> family with sequence similarity 13 member A (*FAM13A*) was identified in a GWAS as a gene that predisposes




**Figure 4** WNT signalling in asthma. In airway smooth muscle cells, WNT-5A activates Frizzled receptor (FZD)8, which together with TGF- $\beta$  stimulation results in increased mRNA expression of extracellular matrix (ECM) components. In addition, WNT-5A, via a not further specified FZD, enhances actin polymerisation and contractile capacity of the smooth muscle cell. Eosinophil-driven airway inflammation stimulates the smooth muscle cells to increase *WNT-5A* expression. Increased expression of WNT-1 prevents dendritic cell-mediated activation of T cells, thereby attenuating airway hyper-responsiveness (AHR) and airway remodelling. WNT-11, via an unspecified FZD, conjointly with TGF- $\beta$  stimulation causes upregulation of the contractile protein  $\alpha$ -sm-actin. Activation of transcriptionally active  $\beta$ -catenin in response to WNT-7B, TGF- $\beta$  and/or (aero) allergens results in augmented mRNA expression of ECM components and genes involved in cell proliferation. Similarly, ectopic expression of a non-degradable form of  $\beta$ -catenin (S33Y- $\beta$ -catenin) enhances expression of ECM components. Transcriptional activity of  $\beta$ -catenin can be inhibited by PKF115-584 or ICG-001, whereas the canonical WNT target gene WNT1-inducible signalling protein-1 (WISP1) can be inhibited with neutralising antibodies. See main text for further details.

to increased susceptibility to COPD.<sup>113 114</sup> A subsequent study investigated the functional role of FAM13A in COPD pathogenesis and demonstrated that FAM13A induces emphysema by targeting canonical WNT signalling.<sup>111</sup> Increased expression of FAM13A was observed in several cell types in the lungs of patients with COPD and FAM13A-deficient mice (FAM13A<sup>-/-</sup>) were protected against cigarette smoke-induced as well as elastase-induced emphysema development. The resistance to elastase-induced emphysema in FAM13A<sup>-/-</sup> mice was abrogated by coadministration of a  $\beta$ -catenin inhibitor (PKF118-310). Biochemical experiments revealed that FAM13A associates with protein phosphatase 2A to facilitate GSK-3 $\beta$ -mediated degradation of  $\beta$ -catenin. Collectively, this study demonstrated that FAM13A is increased in lungs of patients with COPD and might contribute to COPD susceptibility by promoting  $\beta$ -catenin degradation.<sup>111</sup> An extensive analysis of FZD receptor expression in experimental and human COPD, identified FZD4 as an important regulator of canonical WNT/ $\beta$ -catenin signalling and potential candidate for therapeutic intervention in this chronic lung disease.<sup>112</sup> In physiological conditions, the FZD4 receptor is expressed on alveolar epithelial cells, however the expression of this receptor is diminished in individuals with moderate and very severe COPD (Global initiative for chronic obstructive lung diseases (GOLD) stages II and III/IV). Moreover, *FZD4*

expression correlated positively to lung function (%FEV<sub>1</sub>, post-bronchodilator) and negatively with smoking pack-years in the analysed cohort of patients. In vivo and in vitro studies demonstrated that cigarette smoke was able to directly attenuate FZD4 expression, which was accompanied by a reduction in active  $\beta$ -catenin expression.<sup>112</sup> This suggests that FZD4 is a positive regulator of  $\beta$ -catenin signalling in alveolar epithelial cells. Pharmacological inhibition of FZD4 via the small molecule FzM1 decreased (WNT-3A-driven)  $\beta$ -catenin signalling in alveolar epithelial cells, whereas overexpression of FZD4 had the opposite effect. Functional studies demonstrated that the receptor is important for wound healing and repair by the alveolar epithelium, as FZD4 facilitates cell proliferation, migration and ATII-to-ATI cell transdifferentiation.<sup>112</sup> Taken together, these findings strongly suggest that FZD4 signalling and  $\beta$ -catenin activation might be a promising route to induce lung repair in COPD. Indeed, preventive as well as therapeutic reactivation of  $\beta$ -catenin signalling via GSK-3 $\beta$  inhibition resulted in alveolar epithelial cell activation, attenuated emphysema pathology, and improved lung function in experimental emphysema in vivo.<sup>109</sup> Notably, the potential therapeutic application of WNT/ $\beta$ -catenin activation in COPD has recently been tested in more detail and translated into human tissue specimen using three-dimensional lung tissue cultures (3D-LTCs).<sup>115</sup> Patient-derived 3D-LTCs are





**Figure 5** WNT signalling in COPD. Neutrophil elastase and cigarette smoke attenuate pulmonary expression of *WNT-2*, *WNT-3A*, *WNT-10B*, *LRP6*, *FZD1*, *AXIN1*, *AXIN2*, *CTNNB1* ( $\beta$ -catenin), *LEF1* and *TCF4* in human and/or animal models of COPD. In bronchial epithelial cells (left side of dashed line), WNT-4, independently of  $\beta$ -catenin, induces expression of extracellular matrix (ECM) components and of genes involved in cell proliferation. WNT-5B together with TGF- $\beta$ /SMAD signalling activates gene transcription of MMP2, MMP9 and FN. Additionally, TGF- $\beta$  inactivates glycogen synthase kinase-3 (GSK-3) $\beta$  via phosphorylation resulting in activation of  $\beta$ -catenin, which facilitates the epithelial-to-mesenchymal transition (EMT) process of bronchial epithelial cells. In alveolar epithelial cells (right side of dashed line),  $\beta$ -catenin is a critical regulator of ATIII-to-ATI-cell transdifferentiation. WNT-5A, which is increased in individuals with COPD and secreted by pulmonary fibroblasts, acts a negative regulator of  $\beta$ -catenin signalling, thereby impairing endogenous tissue repair by alveolar epithelial cells. FAM13, a COPD susceptibility gene, together with GSK-3 $\beta$  contributes to the development of emphysema by enhanced targeting of  $\beta$ -catenin for proteasomal degradation in alveolar epithelial cells. Moreover, cigarette smoke inhibits  $\beta$ -catenin signalling and epithelial cell repair by reducing FZD4 expression (indicated in grey). Pharmacological reactivation of  $\beta$ -catenin signalling via GSK-3 $\beta$  inhibition (by eg, LiCl, CT99021 or SB216763) in experimental emphysema in vivo as well as patient-derived COPD tissue ex vivo results in epithelial cell activation and attenuated emphysema pathology. In pulmonary fibroblasts, expression of proinflammatory genes induced by IL-1 $\beta$  or Epidermal growth factor (EGF) is mediated in part by FZD8. Additionally, IL-1 $\beta$  and EGF induce the expression of FZD8 via a yet unidentified signalling cascade, whereas TGF- $\beta$ -induced expression of FZD8 is dependent on SMAD signalling. Additionally, TGF- $\beta$  induces the expression of WNT-5A and WNT-5B, which in turn induce expression of proinflammatory genes in a FZD2-dependent and/or FZD8-dependent manner. Activated WNT signalling together with TGF- $\beta$  induces mRNA expression of WNT signalling components, ECM components and myofibroblast markers.  $\beta$ -Catenin is required for the expression of ECM components and myofibroblast differentiation. See main text for further details.

generated ex vivo and represent a valuable tool for preclinical target validation and drug testing. Using this set-up, we demonstrated the potential of activating canonical WNT/ $\beta$ -catenin signalling in 3D-LTCs derived from lung tissue of individuals with COPD.<sup>115</sup> Activation of  $\beta$ -catenin by GSK-3 $\beta$  inhibition, by either LiCl or CT99021, led to increased alveolar epithelial cell marker expression, decreased matrix metalloproteinase-12 expression, as well as altered macrophage activity and elastin remodelling in 3D-LTCs.<sup>115</sup> Altogether, these data underline the potential suitability of WNT/ $\beta$ -catenin activators, such as GSK-3 $\beta$  inhibitors, for the treatment of COPD and urge further preclinical studies for this devastating disease (figure 5).

Similarly, to IPF and asthma, WNT signalling independent of  $\beta$ -catenin (non-canonical WNT signalling) may contribute to

COPD pathogenesis. Two independent studies nearly simultaneously showed aberrant WNT-4 expression in bronchial cells of patients with COPD.<sup>116 117</sup> This specific WNT ligand induces cell proliferation and proinflammatory cytokine secretion by bronchial epithelial cells. In addition, two structurally closely related WNT ligands, WNT-5A and WNT-5B, have been recently investigated in COPD.<sup>96 110 118 119</sup> WNT-5B is predominantly expressed in airway epithelial cells with significantly higher WNT-5B staining in bronchial epithelial cells of patients with COPD compared with smokers.<sup>118</sup> In vitro experiments with recombinant WNT-5B revealed that this specific WNT ligand induces expression of genes related to airway remodelling (ie, FN, MMP-2 and MMP-9) independently of  $\beta$ -catenin signalling, but required activation of TGF- $\beta$ /SMAD3 signalling.<sup>118</sup> In

**Table 1** Overview of molecular targets and applied tools/compounds to modify WNT signalling, which have been investigated in chronic lung diseases

Target	Tools/compounds	Disease	Summary of study	References
<b>Ligands</b>				
WNT-1	CCSP-driven WNT-1 overexpression (in vivo)	Asthma	<ul style="list-style-type: none"> <li>▶ WNT-1 overexpression attenuated AHR, eosinophilia and number of mucus producing cells.</li> <li>▶ Reduced dendritic cell-mediated activation of T cells.</li> </ul>	89
WNT-5A	Recombinant protein (in vitro)	IPF/fibrosis	<ul style="list-style-type: none"> <li>▶ Enhances ECM deposition by lung fibroblasts in vitro.</li> <li>▶ Protects lung fibroblasts against oxidative stress-induced apoptosis.</li> </ul>	74
	Recombinant protein (in vitro)	Physiological conditions	<ul style="list-style-type: none"> <li>▶ Facilitates <math>\beta</math>-catenin/p300 interaction in transdifferentiating primary rat alveolar type-II cells.</li> </ul>	69
	Surfactant protein C (SPC)-driven WNT-5A overexpression, recombinant protein, neutralising antibodies (in vitro and in vivo)	COPD	<ul style="list-style-type: none"> <li>▶ Lung specific overexpression aggravates elastase-induced emphysema in vivo.</li> <li>▶ Attenuation of <math>\beta</math>-catenin-driven wound healing by alveolar epithelial cells.</li> <li>▶ In vivo inhibition of WNT-5A attenuated tissue destruction, improved lung function and restoration of alveolar epithelial cell markers expression in two animal models of COPD.</li> </ul>	110
<b>Receptors</b>				
LRP5	LRP5 <sup>-/-</sup> mice and siRNA (in vitro and in vivo)	IPF/fibrosis	<ul style="list-style-type: none"> <li>▶ LRP5<sup>-/-</sup> mice are protected against bleomycin-induced lung fibrosis.</li> <li>▶ Delayed progression of asbestos-induced lung fibrosis in mice lacking LRP5.</li> </ul>	44
FZD4	FzM1, FZD4 siRNA and overexpression (in vitro)	COPD	<ul style="list-style-type: none"> <li>▶ FZD4 is expressed in alveolar epithelial cells and is reduced in COPD.</li> <li>▶ Inhibition of FZD4 by FzM1 decreased (WNT-3A-driven) <math>\beta</math>-catenin signalling in alveolar epithelial cells, whereas overexpression of FZD4 has the opposite effect.</li> <li>▶ FZD4 regulates wound healing and repair by the alveolar epithelium by facilitating cell proliferation, cell migration and ATII-to-ATI cell transdifferentiation.</li> </ul>	112
FZD8	FZD8 <sup>-/-</sup> mice and siRNA (in vitro and in vivo)	IPF/fibrosis	<ul style="list-style-type: none"> <li>▶ FZD8<sup>-/-</sup> mice are partially protected against bleomycin-induced lung fibrosis. FZD8 siRNA attenuates TGF-<math>\beta</math>-induced ECM deposition by human lung fibroblasts.</li> </ul>	71
		COPD	<ul style="list-style-type: none"> <li>▶ Reduced cigarette smoke-induced inflammation in FZD8<sup>-/-</sup> mice compared with Wild type (WT) mice.</li> <li>▶ Association between SNP in FZD8 and chronic mucus hypersecretion in cohort of smokers.</li> <li>▶ Involvement of receptor in cytokine secretion by human lung fibroblasts.</li> </ul>	123
<b>Intercellular proteins</b>				
FAM13A	FAM13A <sup>-/-</sup> mice (in vivo)	COPD	<ul style="list-style-type: none"> <li>▶ FAM13A is a COPD susceptibility gene.</li> <li>▶ Increased expression of FAM13A facilitates <math>\beta</math>-catenin degradation thereby contributing to emphysema development in two murine models of COPD.</li> </ul>	111
GSK-3 $\beta$	LiCl (in vitro)	IPF/fibrosis/COPD	<ul style="list-style-type: none"> <li>▶ Pharmacological inhibition of GSK-3<math>\beta</math> enhances TGF-<math>\beta</math>-induced EMT</li> </ul>	63
	LiCl (in vivo)	COPD	<ul style="list-style-type: none"> <li>▶ Inhibition of GSK-3<math>\beta</math> enhanced <math>\beta</math>-catenin in alveolar epithelial cells in vivo.</li> <li>▶ LiCl prevents the development and progression of elastase-induced emphysema.</li> </ul>	109
	SB216763 (in vivo)	COPD	<ul style="list-style-type: none"> <li>▶ Inhibition of GSK-3<math>\beta</math> prevented right ventricle hypertrophy and small airway remodelling in a guinea pig model of LPS-induced COPD.</li> </ul>	125
	CT99021 or LiCl (ex vivo)	COPD	<ul style="list-style-type: none"> <li>▶ GSK-3<math>\beta</math> inhibition in 3D-LTCs: increased alveolar epithelial cell marker expression, decreased MMP12 expression, and altered elastin remodelling.</li> </ul>	115
<i>Tankyrases</i>	XAV939 (in vivo and in vitro)	IPF/fibrosis	<ul style="list-style-type: none"> <li>▶ Reduced <math>\beta</math>-catenin activation due to Axin stabilisation.</li> <li>▶ XAV939 attenuates bleomycin-induced lung fibrosis.</li> <li>▶ Inhibition of TKNS reduces fibroblast proliferation and myofibroblast differentiation.</li> </ul>	45
$\beta$ -catenin	siRNA (in vivo)	Asthma	<ul style="list-style-type: none"> <li>▶ siRNA against <math>\beta</math>-catenin resulted in reduced inflammation and airway remodelling in a murine model of OVA-induced asthma.</li> </ul>	90
<b><math>\beta</math>-catenin interaction</b>				
$\beta$ -catenin/TCF	ICAT (in vitro)	Physiological conditions	<ul style="list-style-type: none"> <li>▶ Ectopic expression of <i>Inhibitor of <math>\beta</math>-catenin and TCF</i> (ICAT) impairs murine ATII-to-ATI cell transdifferentiation.</li> </ul>	48
$\beta$ -catenin/TCF	PKF115–584 (in vitro)	Physiological conditions	<ul style="list-style-type: none"> <li>▶ Inhibition of <math>\beta</math>-catenin/TCF interaction by PKF115–584 impairs murine ATII-to-ATI cell transdifferentiation.</li> </ul>	50
		Physiological conditions	<ul style="list-style-type: none"> <li>▶ PKF115–584 prevents TGF-<math>\beta</math>-induced ECM deposition by human airway smooth muscle cells.</li> </ul>	95
		Physiological conditions	<ul style="list-style-type: none"> <li>▶ PKF115–584 decreases contractile force generation by airway smooth muscle.</li> </ul>	97
		Physiological conditions	<ul style="list-style-type: none"> <li>▶ Disruption of <math>\beta</math>-catenin/TCF interaction decreases ECM deposition by lung fibroblasts and prevents myofibroblast differentiation.</li> </ul>	96

Continued

Table 1 Continued

Target	Tools/compounds	Disease	Summary of study	References
$\beta$ -catenin/CBP	ICG-001 (in vitro and in vivo)	Asthma	▶ Reduced airway smooth muscle remodelling and ECM expression in a murine model of OVA-induced asthma in isolated smooth muscle cells.	94
	ICG-001 (in vivo)	IPF	▶ Selective inhibition of the $\beta$ -catenin/CBP interaction by the ICG-001 led to reversal of established bleomycin-induced pulmonary fibrosis and improved epithelial cell integrity.	42
$\beta$ -catenin/p300	IQ-1 (in vitro)	Physiological conditions	▶ Disruption of the $\beta$ -catenin/p300 interaction interferes with rat A1II-to-A1I cell transdifferentiation. ▶ IQ-1 partially stabilises SPC (A1II cell marker) expression, whereas the small molecule attenuates aquaporin5 (A1I cell marker) expression.	69
WNT target gene				
WISP1	WISP1 neutralising antibodies (in vitro and in vivo)	IPF	▶ WISP1 is upregulated in murine model of bleomycin-induced fibrosis. ▶ Neutralising mAbs specific for WISP1 reduced the expression of genes characteristic of fibrosis and reversed the expression of genes associated with EMT. ▶ Inhibition of WISP1 attenuates pathological changes in experimental lung fibrosis in vivo.	35
		Asthma	▶ Inhibition of WISP1 by neutralising antibodies results in less airway remodelling in a rat model of allergic asthma.	93

3D-LTCs, three-dimensional lung tissue cultures; AHR, airway hyper-responsiveness; ATI (or II), alveolar epithelial type I (or II) cells; CBP, cAMP response element-binding protein binding protein; CCSP, Clara cell secretory protein; ECM, extracellular matrix; EMT, epithelial-to-mesenchymal transition; FZD, Frizzled receptor; GSK-3, glycogen synthase kinase-3; ICAT, inhibitor of  $\beta$ -catenin and TCF; IPF, idiopathic pulmonary fibrosis; LRP5<sup>-/-</sup>, mice lacking lipoprotein receptor-related protein 5; OVA, ovalbumin; SNP, single nucleotide polymorphism; TCF, T cell factor; WISP1, WNT1-inducible signalling protein-1.

addition, WNT-5A has also been linked to inflammatory processes by pulmonary fibroblasts and COPD pathogenesis. Individual components of cigarette smoke, including nicotine, can induce WNT-5A in cells of the respiratory system and WNT-5A is a direct target of miR-487b, a microRNA that is repressed by cigarette smoke.<sup>110 120 121</sup> Moreover, cigarette smoke can directly activate TGF- $\beta$ , a growth factor implicated in WNT-5A induction in several distinct cells of the lungs.<sup>72 96 99 100 110</sup> These findings indicate WNT-5A as a prime target of COPD-related exposures. Most recently, we reported increased WNT-5A transcript and protein in lung tissue of individuals with COPD compared with non-COPD controls.<sup>110</sup> We furthermore observed altered post-translational modifications (ie, glycosylation, palmitoylation and oligomerisation) of WNT-5A in lung tissue specimens of individuals with COPD compared with tissue of donors.<sup>110</sup> The increased expression of this ligand was recapitulated in a murine model of cigarette smoke-induced COPD as well as in elastase-induced emphysema.<sup>110</sup> Notably, various studies have indicated that specific non-canonical WNT ligands are able to inhibit canonical WNT/ $\beta$ -catenin signalling.<sup>3 122</sup> Accordingly, mature WNT-5A attenuated canonical WNT/ $\beta$ -catenin-driven alveolar epithelial cell wound healing and A1II-to-A1I cell transdifferentiation in vitro. Furthermore, lung-specific overexpression of WNT-5A aggravated elastase-induced emphysema in vivo. Most importantly, both prophylactic as well as therapeutic inhibition of WNT-5A, using neutralising antibodies or a synthetic peptide, resulted in attenuated tissue destruction, improved lung function and restoration of  $\beta$ -catenin-driven transcriptional targets (eg, *Axin2* and *Nkd1*) as well as alveolar epithelial cell marker expression in two animal models of COPD.<sup>110</sup> These findings suggest a canonical to non-canonical WNT signal switch in COPD, which contributes to disease development and progression. Moreover, our study highlights that non-canonical WNT signalling and, in particular targeting WNT-5A, is a promising therapeutic option for the treatment of emphysema/COPD.<sup>110</sup> However, which FZD receptors are mediating the effects of these WNT ligands (ie, WNT-4, WNT-5A and/or WNT-5B) in COPD is largely unknown. Nevertheless, a recent study by

Spanjer *et al*<sup>123</sup> linked the WNT receptor FZD8 to chronic bronchitis and mucus hypersecretion, but did not investigate which WNT ligand(s) might be involved in these processes, and this needs further examination.

In summary, it appears that canonical and non-canonical WNT signalling pathways play opposing or balancing roles in COPD, wherein (re)activation of WNT/ $\beta$ -catenin signalling is a potential remedy for emphysema whereas non-canonical WNT signalling contributes to airway remodelling and cigarette smoke-induced inflammation, thus perpetuating disease development and progression (figure 5).

#### FIGHTING THE COLD WNT-er

An important unfulfilled need is to identify new agents that interact with key molecular pathways involved in the pathogenesis of chronic lung diseases. Since the last decade, our knowledge about deviations in WNT signalling in the aforementioned chronic lung disease continues to grow (figures 3–5 and table 1). The ability to target the WNT signalling pathway offers immense promise in chronic lung diseases; however, substantial risks and concerns remain present with regard to the targeting of such crucial signalling pathways. The challenge for the future will be identifying which branch of WNT signalling to target at what time point and how to specifically target major signalling hubs within these pathways. Inhibition of biogenesis and secretion of WNT ligands as well as blocking the interaction of WNT ligands with the cell-surface receptors can be achieved to a certain extent by pharmacologically active small molecules.<sup>124</sup> While these approaches do not necessarily discriminate between targeting either the canonical or non-canonical WNT signalling pathway and are therefore relatively non-specific, this may also be a potential benefit as both these WNT signalling pathways are involved in pathological processes in IPF and asthma. A major benefit of the canonical WNT signalling pathway over the non-canonical pathways is that  $\beta$ -catenin is a specific downstream effector molecule of canonical signalling, which can be targeted pharmacologically. Activation of  $\beta$ -catenin signalling is of therapeutic potential in emphysema/COPD and can be achieved by various molecular mechanisms. Most commonly,

$\beta$ -catenin signalling is activated via inhibition of GSK-3 $\beta$ , the primary kinase involved in  $\beta$ -catenin degradation.<sup>1</sup> Prudence is advised with artificially activating  $\beta$ -catenin signalling in the lung due to the risk of potential adverse effects, such as EMT, tissue remodelling or tumorigenesis. However, in a guinea pig model of lipopolysaccharide (LPS)-induced COPD airway wall remodelling was substantially reduced by pharmacological inhibition of GSK-3 $\beta$ .<sup>125</sup> Moreover, the GSK-3 $\beta$  inhibitor lithium has been used to treat patients for many years now without apparent increases in the cancer incidence.<sup>126</sup> Taken together, these studies suggest that inhibition of GSK-3 $\beta$  has therapeutic potential in COPD. In contrast to emphysema/COPD, increased  $\beta$ -catenin signalling is a hallmark of IPF and inhibition of this transcriptional coactivator is of great therapeutic interest. Most pharmacological approaches to attenuate canonical WNT/ $\beta$ -catenin signalling focus on blocking the interaction of  $\beta$ -catenin with the transcription factors.<sup>127</sup> A primary consideration, however, is that  $\beta$ -catenin-dependent signalling can also be activated by other growth factors than WNTs and that transcriptional activity of  $\beta$ -catenin is influenced by its interaction with various binding partners.

Finally, the majority of chronic lung diseases affect diverse cellular compartments, and cell-specific targeting of WNT signalling to ensure a beneficial clinical outcome represents a challenge for the future. Our knowledge about (cell)-specific ligand/receptor complexes has just recently begun to increase and this route holds promise for treatment applications, as drugs targeting distinct WNT ligands and/or FZD receptors are currently under investigation.<sup>1 128–132</sup> Taken together, several valuable approaches to target the WNT pathways exist or are under development and thus offer encouraging routes to potentially treat chronic lung diseases in the future.

**Twitter** Follow Melanie Königshoff @m\_konigshoff

**Acknowledgements** The authors thank all members of the Königshoff laboratory, in particular Florian Ciolek, Stephan Klee, Mareike Lehmann, Carlo Mümmeler, Wioletta Skronska-Wasek and Darcy Wagner, for stimulating discussions about the content and the title of this review at ATS 2016.

**Contributors** HAB and MK conceptually designed and wrote the manuscript.

**Funding** HAB is supported by a postdoctoral fellowship from the European Respiratory Society (ERS Fellowship LTRF 79-2012) and a fellowship from Helmholtz Zentrum Germany (PFP PF-135). MK is supported by a European Research Council Starting Grant (ERC-StG-L57, Grant No. 261302) and a W2/3 Professorship Award by the Helmholtz Association.

**Competing interests** None declared.

**Provenance and peer review** Commissioned; externally peer reviewed.

**Open Access** This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

## REFERENCES

- Baarsma HA, Königshoff M, Gosens R. The WNT signaling pathway from ligand secretion to gene transcription: molecular mechanisms and pharmacological targets. *Pharmacol Ther* 2013;138:66–83.
- van Amerongen R, Mikels A, Nusse R. Alternative wnt signaling is initiated by distinct receptors. *Sci Signal* 2008;1:re9.
- Mikels AJ, Nusse R. Purified Wnt5a protein activates or inhibits beta-catenin-TCF signaling depending on receptor context. *PLoS Biol* 2006;4:e115.
- Dijksterhuis JP, Baljinnam B, Stanger K, et al. Systematic mapping of WNT-FZD protein interactions reveals functional selectivity by distinct WNT-FZD pairs. *J Biol Chem* 2015;290:6789–98.
- Li C, Bellusci S, Borok Z, et al. Non-canonical WNT signalling in the lung. *J Biochem* 2015;158:355–65.
- Macdonald BT, Semenov MV, He X. Snapshot: Wnt/beta-catenin signaling. *Cell* 2007;131:1204.
- Semenov MV, Habas R, Macdonald BT, et al. Snapshot: Noncanonical Wnt Signaling Pathways. *Cell* 2007;131:1378.
- De Langhe SP, Reynolds SD. Wnt signaling in lung organogenesis. *Organogenesis* 2008;4:100–8.
- Königshoff M, Eickelberg O. WNT signaling in lung disease: a failure or a regeneration signal? *Am J Respir Cell Mol Biol* 2010;42:21–31.
- Stewart DJ. Wnt signaling pathway in non-small cell lung cancer. *J Natl Cancer Inst* 2014;106:djt356.
- Mazieres J, He B, You L, et al. Wnt signaling in lung cancer. *Cancer Lett* 2005;222:1–10.
- He B, Barg RN, You L, et al. Wnt signaling in stem cells and non-small-cell lung cancer. *Clin Lung Cancer* 2005;7:54–60.
- Tennis M, Van Scoyk M, Winn RA. Role of the wnt signaling pathway and lung cancer. *J Thorac Oncol* 2007;2:889–92.
- Krauss-Etschmann S, Bush A, Bellusci S, et al. Of flies, mice and men: a systematic approach to understanding the early life origins of chronic lung disease. *Thorax* 2013;68:380–4.
- Ota C, Baarsma HA, Wagner DE, et al. Linking bronchopulmonary dysplasia to adult chronic lung diseases: role of WNT signaling. *Mol Cell Pediatr* 2016;3:34.
- Yates LL, Dean CH. Planar polarity: a new player in both lung development and disease. *Organogenesis* 2011;7:209–16.
- Kumawat K, Koopmans T, Gosens R.  $\beta$ -catenin as a regulator and therapeutic target for asthmatic airway remodeling. *Expert Opin Ther Targets* 2014;18:1023–34.
- Beers MF, Morrissey EE. The three R's of lung health and disease: repair, remodeling, and regeneration. *J Clin Invest* 2011;121:2065–73.
- Pongracz JE, Stockley RA. Wnt signalling in lung development and diseases. *Respir Res* 2006;7:15.
- Li C, Hu L, Xiao J, et al. Wnt5a regulates Shh and Fgf10 signaling during lung development. *Dev Biol* 2005;287:86–97.
- Li C, Xiao J, Hormi K, et al. Wnt5a participates in distal lung morphogenesis. *Dev Biol* 2002;248:68–81.
- Caprioli A, Villasenor A, Wylie LA, et al. Wnt4 is essential to normal mammalian lung development. *Dev Biol* 2015;406:222–34.
- Goss AM, Tian Y, Tsukiyama T, et al. Wnt2/2b and beta-catenin signaling are necessary and sufficient to specify lung progenitors in the foregut. *Dev Cell* 2009;17:290–8.
- Mucenski ML, Wert SE, Nation JM, et al. beta-Catenin is required for specification of proximal/distal cell fate during lung morphogenesis. *J Biol Chem* 2003;278:40231–8.
- Mucenski ML, Nation JM, Thitoff AR, et al. Beta-catenin regulates differentiation of respiratory epithelial cells in vivo. *Am J Physiol Lung Cell Mol Physiol* 2005;289:L971–9.
- Banderali G, Martelli A, Landi M, et al. Short and long term health effects of parental tobacco smoking during pregnancy and lactation: a descriptive review. *J Transl Med* 2015;13:327.
- Duijts L, Reiss IK, Brusselle G, et al. Early origins of chronic obstructive lung diseases across the life course. *Eur J Epidemiol* 2014;29:871–85.
- Blacquièrre MJ, Timens W, van den Berg A, et al. Maternal smoking during pregnancy decreases Wnt signalling in neonatal mice. *Thorax* 2010;65:553–4.
- Lehmann M, Baarsma HA, Königshoff M. WNT Signaling in Lung Aging and Disease. *Ann Am Thorac Soc* 2016;13(Suppl\_5):S411–16.
- Gross TJ, Hunninghake GW. Idiopathic pulmonary fibrosis. *N Engl J Med* 2001;345:517–25.
- Behr J. Evidence-based treatment strategies in idiopathic pulmonary fibrosis. *Eur Respir Rev* 2013;22:163–8.
- Raghu G, Collard HR, Egan JJ, et al. An official ATS/ERS/JRS/ALAT statement: idiopathic pulmonary fibrosis: evidence-based guidelines for diagnosis and management. *Am J Respir Crit Care Med* 2011;183:788–824.
- Selman M, Pardo A, Kaminski N. Idiopathic pulmonary fibrosis: aberrant recapitulation of developmental programs? *PLoS Med* 2008;5:e62.
- Königshoff M, Balsara N, Pfaff EM, et al. Functional Wnt signaling is increased in idiopathic pulmonary fibrosis. *PLoS ONE* 2008;3:e2142.
- Königshoff M, Kramer M, Balsara N, et al. WNT1-inducible signaling protein-1 mediates pulmonary fibrosis in mice and is upregulated in humans with idiopathic pulmonary fibrosis. *J Clin Invest* 2009;119:772–87.
- Rosas IO, Richards TJ, Konishi K, et al. MMP1 and MMP7 as potential peripheral blood biomarkers in idiopathic pulmonary fibrosis. *PLoS Med* 2008;5:e93.
- Chilosi M, Poletti V, Zamò A, et al. Aberrant Wnt/beta-catenin pathway activation in idiopathic pulmonary fibrosis. *Am J Pathol* 2003;162:1495–502.
- Berschneider B, Ellwanger DC, Baarsma HA, et al. miR-92a regulates TGF- $\beta$ 1-induced WISP1 expression in pulmonary fibrosis. *Int J Biochem Cell Biol* 2014;53:432–41.
- Klee S, Lehmann M, Wagner DE, et al. WISP1 mediates IL-6-dependent proliferation in primary human lung fibroblasts. *Sci Rep* 2016;6:20547.
- Guo Y, Xiao L, Sun L, et al. Wnt/beta-catenin signaling: a promising new target for fibrosis diseases. *Physiol Res* 2012;61:337–46.

- 41 He W, Dai C, Li Y, et al. Wnt/beta-catenin signaling promotes renal interstitial fibrosis. *J Am Soc Nephrol* 2009;20:765–76.
- 42 Henderson WR Jr, Chi EY, Ye X, et al. Inhibition of Wnt/beta-catenin/CREB binding protein (CBP) signaling reverses pulmonary fibrosis. *Proc Natl Acad Sci USA* 2010;107:14309–14.
- 43 Lam AP, Flozak AS, Russell S, et al. Nuclear  $\beta$ -catenin is increased in systemic sclerosis pulmonary fibrosis and promotes lung fibroblast migration and proliferation. *Am J Respir Cell Mol Biol* 2011;45:915–22.
- 44 Lam AP, Herazo-Maya JD, Sennello JA, et al. Wnt coreceptor Lrp5 is a driver of idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 2014;190:185–95.
- 45 Ulsamer A, Wei Y, Kim KK, et al. Axin pathway activity regulates in vivo pY654- $\beta$ -catenin accumulation and pulmonary fibrosis. *J Biol Chem* 2012;287:5164–72.
- 46 Wang C, Zhu H, Sun Z, et al. Inhibition of Wnt/ $\beta$ -catenin signaling promotes epithelial differentiation of mesenchymal stem cells and repairs bleomycin-induced lung injury. *Am J Physiol Cell Physiol* 2014;307:C234–44.
- 47 Aumiller V, Balsara N, Wilhelm J, et al. WNT/beta-catenin signaling induces IL-1 $\beta$  expression by alveolar epithelial cells in pulmonary fibrosis. *Am J Respir Cell Mol Biol* 2013;49:96–104.
- 48 Flozak AS, Lam AP, Russell S, et al. Beta-catenin/T-cell factor signaling is activated during lung injury and promotes the survival and migration of alveolar epithelial cells. *J Biol Chem* 2010;285:3157–67.
- 49 Marconett CN, Zhou B, Rieger ME, et al. Integrated transcriptomic and epigenomic analysis of primary human lung epithelial cell differentiation. *PLoS Genet* 2013;9:e1003513.
- 50 Mutze K, Vierkotten S, Milosevic J, et al. Enolase 1 (ENO1) and protein disulfide-isomerase associated 3 (PDIA3) regulate Wnt/ $\beta$ -catenin-driven trans-differentiation of murine alveolar epithelial cells. *Dis Model Mech* 2015;8:877–90.
- 51 Lawson WE, Blackwell TS. beta-Catenin and CCNs in lung epithelial repair. *Am J Physiol Lung Cell Mol Physiol* 2013;304:L579–81.
- 52 Tanjore H, Degryse AL, Crossno PF, et al.  $\beta$ -catenin in the alveolar epithelium protects from lung fibrosis after intratracheal bleomycin. *Am J Respir Crit Care Med* 2013;187:630–9.
- 53 Zemans RL, McClendon J, Aschner Y, et al. Role of  $\beta$ -catenin-regulated CCN matricellular proteins in epithelial repair after inflammatory lung injury. *Am J Physiol Lung Cell Mol Physiol* 2013;304:L415–27.
- 54 Desai TJ, Brownfield DG, Krasnow MA. Alveolar progenitor and stem cells in lung development, renewal and cancer. *Nature* 2014;507:190–4.
- 55 Treutlein B, Brownfield DG, Wu AR, et al. Reconstructing lineage hierarchies of the distal lung epithelium using single-cell RNA-seq. *Nature* 2014;509:371–5.
- 56 Kelly M, Kolb M, Bonniaud P, et al. Re-evaluation of fibrogenic cytokines in lung fibrosis. *Curr Pharm Des* 2003;9:39–49.
- 57 Kolb M, Margetts PJ, Anthony DC, et al. Transient expression of IL-1 $\beta$  induces acute lung injury and chronic repair leading to pulmonary fibrosis. *J Clin Invest* 2001;107:1529–36.
- 58 Chapman HA. Epithelial-mesenchymal interactions in pulmonary fibrosis. *Annu Rev Physiol* 2011;73:413–35.
- 59 Kim KK, Kugler MC, Wolters PJ, et al. Alveolar epithelial cell mesenchymal transition develops in vivo during pulmonary fibrosis and is regulated by the extracellular matrix. *Proc Natl Acad Sci USA* 2006;103:13180–5.
- 60 Sohal SS, Reid D, Soltani A, et al. Evaluation of epithelial mesenchymal transition in patients with chronic obstructive pulmonary disease. *Respir Res* 2011;12:130.
- 61 Wang Q, Wang Y, Zhang Y, et al. The role of uPAR in epithelial-mesenchymal transition in small airway epithelium of patients with chronic obstructive pulmonary disease. *Respir Res* 2013;14:67.
- 62 Milara J, Peiró T, Serrano A, et al. Epithelial to mesenchymal transition is increased in patients with COPD and induced by cigarette smoke. *Thorax* 2013;68:410–20.
- 63 Zhou B, Liu Y, Kahn M, et al. Interactions between  $\beta$ -catenin and transforming growth factor- $\beta$  signaling pathways mediate epithelial-mesenchymal transition and are dependent on the transcriptional co-activator cAMP-response element-binding protein (CREB)-binding protein (CBP). *J Biol Chem* 2012;287:7026–38.
- 64 Huang SM, Mishina YM, Liu S, et al. Tankyrase inhibition stabilizes axin and antagonizes Wnt signalling. *Nature* 2009;461:614–20.
- 65 Mosimann C, Hausmann G, Basler K. Beta-catenin hits chromatin: regulation of Wnt target gene activation. *Nat Rev Mol Cell Biol* 2009;10:276–86.
- 66 Labbé E, Letamendia A, Attisano L. Association of Smads with lymphoid enhancer binding factor 1/T cell-specific factor mediates cooperative signaling by the transforming growth factor-beta and wnt pathways. *Proc Natl Acad Sci USA* 2000;97:8358–63.
- 67 Jin T, George Fantus I, Sun J. Wnt and beyond Wnt: multiple mechanisms control the transcriptional property of beta-catenin. *Cell Signal* 2008;20:1697–704.
- 68 Valenta T, Hausmann G, Basler K. The many faces and functions of  $\beta$ -catenin. *EMBO J* 2012;31:2714–36.
- 69 Rieger ME, Zhou B, Solomon N, et al. p300/ $\beta$ -Catenin Interactions Regulate Adult Progenitor Cell Differentiation Downstream of WNT5a/Protein Kinase C (PKC). *J Biol Chem* 2016;291:6569–82.
- 70 Sennello JA, Misharin AV, Flozak AS, et al. Lrp5/beta-catenin Signaling Controls Lung Macrophage Differentiation and Inhibits Resolution of Fibrosis. *Am J Respir Cell Mol Biol* 2017;56:191–201.
- 71 Spanjer AI, Baarsma HA, Oostenbrink LM, et al. TGF- $\beta$ -induced profibrotic signaling is regulated in part by the WNT receptor Frizzled-8. *FASEB J* 2016;30:1823–35.
- 72 Newman DR, Sills WS, Hanrahan K, et al. Expression of WNT5A in idiopathic pulmonary fibrosis and its control by TGF- $\beta$  and WNT7B in human lung fibroblasts. *J Histochem Cytochem* 2016;64:99–111.
- 73 Rydell-Tormanen K, Zhou XH, Hallgren O, et al. Aberrant nonfibrotic parenchyma in idiopathic pulmonary fibrosis is correlated with decreased beta-catenin inhibition and increased Wnt5a/b interaction. *Physiol Rep* 2016;4:e12727.
- 74 Yuga LJ, Ben-Yehudah A, Kovkarova-Naumovski E, et al. WNT5A is a regulator of fibroblast proliferation and resistance to apoptosis. *Am J Respir Cell Mol Biol* 2009;41:583–9.
- 75 Tsangaris I, Tsaknis G, Anthi A, et al. Pulmonary hypertension in parenchymal lung disease. *Pulm Med* 2012;2012:684781.
- 76 Thomson JR, Machado RD, Pauciuolo MW, et al. Sporadic primary pulmonary hypertension is associated with germline mutations of the gene encoding BMPRII, a receptor member of the TGF-beta family. *J Med Genet* 2000;37:741–5.
- 77 Deng Z, Morse JH, Slager SL, et al. Familial primary pulmonary hypertension (gene PPH1) is caused by mutations in the bone morphogenetic protein receptor-II gene. *Am J Hum Genet* 2000;67:737–44.
- 78 de Jesus Perez VA, Alastalo TP, Wu JC, et al. Bone morphogenetic protein 2 induces pulmonary angiogenesis via Wnt-beta-catenin and Wnt-RhoA-Rac1 pathways. *J Cell Biol* 2009;184:83–99.
- 79 West JD, Austin ED, Gaskill C, et al. Identification of a common Wnt-associated genetic signature across multiple cell types in pulmonary arterial hypertension. *Am J Physiol Cell Physiol* 2014;307:C415–30.
- 80 Laumanns IP, Fink L, Wilhelm J, et al. The noncanonical WNT pathway is operative in idiopathic pulmonary arterial hypertension. *Am J Respir Cell Mol Biol* 2009;40:683–91.
- 81 Fantozzi I, Huang W, Zhang J, et al. Divergent effects of BMP-2 on gene expression in pulmonary artery smooth muscle cells from normal subjects and patients with idiopathic pulmonary arterial hypertension. *Exp Lung Res* 2005;31:783–806.
- 82 Wu D, Talbot CC Jr, Liu Q, et al. Identifying microRNAs targeting Wnt/ $\beta$ -catenin pathway in end-stage idiopathic pulmonary arterial hypertension. *J Mol Med* 2016;94:875–85.
- 83 Saglani S, Lloyd CM. Novel concepts in airway inflammation and remodelling in asthma. *Eur Respir J* 2015;46:1796–804.
- 84 Sharma S, Tantisira K, Carey V, et al. A role for Wnt signaling genes in the pathogenesis of impaired lung function in asthma. *Am J Respir Crit Care Med* 2010;181:328–36.
- 85 Wang SH, Xu F, Dang HX, et al. Genetic variations in the Wnt signaling pathway affect lung function in asthma patients. *Genet Mol Res* 2013;12:1829–33.
- 86 Barreto-Luis A, Corrales A, Acosta-Herrera M, et al. A pathway-based association study reveals variants from Wnt signaling genes contributing to asthma susceptibility. *Clin Exp Allergy*. Published Online First 12 Jan 2017. doi: 10.1111/cea.12883.
- 87 Choy DF, Modrek B, Abbas AR, et al. Gene expression patterns of Th2 inflammation and intercellular communication in asthmatic airways. *J Immunol* 2011;186:1861–9.
- 88 Januskevicius A, Vaitkiene S, Gosens R, et al. Eosinophils enhance WNT-5a and TGF- $\beta$ 1 genes expression in airway smooth muscle cells and promote their proliferation by increased extracellular matrix proteins production in asthma. *BMC Pulm Med* 2016;16:94.
- 89 Reuter S, Martin H, Beckert H, et al. The Wnt/ $\beta$ -catenin pathway attenuates experimental allergic airway disease. *J Immunol* 2014;193:485–95.
- 90 Kwak HJ, Park DW, Seo JY, et al. The Wnt/ $\beta$ -catenin signaling pathway regulates the development of airway remodeling in patients with asthma. *Exp Mol Med* 2015;47:e198.
- 91 Cohen ED, Ihida-Stansbury K, Lu MM, et al. Wnt signaling regulates smooth muscle precursor development in the mouse lung via a tenascin C/PDGFR pathway. *J Clin Invest* 2009;119:2538–49.
- 92 Heijink IH, Postma DS, Noordhoek JA, et al. House dust mite-promoted epithelial-to-mesenchymal transition in human bronchial epithelium. *Am J Respir Cell Mol Biol* 2010;42:69–79.
- 93 Yang M, Du Y, Xu Z, et al. Functional Effects of WNT1-Inducible Signaling Pathway Protein-1 on Bronchial Smooth Muscle Cell Migration and Proliferation in OVA-Induced Airway Remodeling. *Inflammation* 2016;39:16–29.
- 94 Koopmans T, Crutzen S, Menzen MH, et al. Selective targeting of CBP/ $\beta$ -catenin inhibits growth of and extracellular matrix remodelling by airway smooth muscle. *Br J Pharmacol*. Published Online First: 15 Sep 2016. doi:10.1111/bph.13620.
- 95 Baarsma HA, Menzen MH, Halayko AJ, et al.  $\beta$ -Catenin signaling is required for TGF- $\beta$ 1-induced extracellular matrix production by airway smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol* 2011;301:L956–65.

- 96 Baarsma HA, Spanjer AI, Haitsma G, *et al.* Activation of WNT/ $\beta$ -catenin signaling in pulmonary fibroblasts by TGF- $\beta$ ; is increased in chronic obstructive pulmonary disease. *PLoS ONE* 2011;6:e25450.
- 97 Jansen SR, Van Ziel AM, Baarsma HA, *et al.*  $\beta$ -Catenin regulates airway smooth muscle contraction. *Am J Physiol Lung Cell Mol Physiol* 2010;299:L204–14.
- 98 Nunes RO, Schmidt M, Dueck G, *et al.* GSK-3/ $\beta$ -catenin signaling axis in airway smooth muscle: role in mitogenic signaling. *Am J Physiol Lung Cell Mol Physiol* 2008;294:L1110–18.
- 99 Kumawat K, Menzen MH, Bos IS, *et al.* Noncanonical WNT-5A signaling regulates TGF- $\beta$ -induced extracellular matrix production by airway smooth muscle cells. *FASEB J* 2013;27:1631–43.
- 100 Kumawat K, Gosens R. WNT-5A: signaling and functions in health and disease. *Cell Mol Life Sci* 2016;73:567–87.
- 101 Koopmans T, Kumawat K, Halayko AJ, *et al.* Regulation of actin dynamics by WNT-5A: implications for human airway smooth muscle contraction. *Sci Rep* 2016;6:30676.
- 102 Kumawat K, Koopmans T, Menzen MH, *et al.* Cooperative signaling by TGF- $\beta$ 1 and WNT-11 drives sm- $\alpha$ -actin expression in smooth muscle via Rho kinase-actin-MRTF-A signaling. *Am J Physiol Lung Cell Mol Physiol* 2016;311:L529–37.
- 103 Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med* 2006;3:e442.
- 104 Vestbo J, Hurd SS, Agustí AG, *et al.* Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* 2013;187:347–65.
- 105 Guo L, Wang T, Wu Y, *et al.* WNT/ $\beta$ -catenin signaling regulates cigarette smoke-induced airway inflammation via the PPAR $\delta$ /p38 pathway. *Lab Invest* 2016;96:218–29.
- 106 Wang R, Ahmed J, Wang G, *et al.* Down-regulation of the canonical Wnt  $\beta$ -catenin pathway in the airway epithelium of healthy smokers and smokers with COPD. *PLoS ONE* 2011;6:e14793.
- 107 Zou W, Zou Y, Zhao Z, *et al.* Nicotine-induced epithelial-mesenchymal transition via Wnt/ $\beta$ -catenin signaling in human airway epithelial cells. *Am J Physiol Lung Cell Mol Physiol* 2013;304:L199–209.
- 108 Ezzie ME, Crawford M, Cho JH, *et al.* Gene expression networks in COPD: microRNA and mRNA regulation. *Thorax* 2012;67:122–31.
- 109 Kneidinger N, Yildirim AO, Callegari J, *et al.* Activation of the WNT/ $\beta$ -catenin pathway attenuates experimental emphysema. *Am J Respir Crit Care Med* 2011;183:723–33.
- 110 Baarsma HA, Skronska-Wasek W, Mutze K. Noncanonical WNT-5A signaling impairs endogenous lung repair in COPD. *J Exp Med* 2017;214:143–63.
- 111 Jiang Z, Lao T, Qiu W, *et al.* A chronic obstructive pulmonary disease susceptibility gene, FAM13A, regulates protein stability of  $\beta$ -catenin. *Am J Respir Crit Care Med* 2016;194:185–97.
- 112 Skronska-Wasek W, Mutze K, Baarsma HA, *et al.* Reduced Frizzled receptor 4 expression prevents WNT/ $\beta$ -catenin-driven alveolar lung repair in COPD. *Am J Respir Crit Care Med*. Published Online First: 28 Feb 2017. doi:10.1164/rccm.201605-0904OC.
- 113 Cho MH, Boutaoui N, Klanderma BJ, *et al.* Variants in FAM13A are associated with chronic obstructive pulmonary disease. *Nat Genet* 2010;42:200–2.
- 114 Ziolkowska-Suchanek I, Mosor M, Gabryel P, *et al.* Susceptibility loci in lung cancer and COPD: association of IREB2 and FAM13A with pulmonary diseases. *Sci Rep* 2015;5:13502.
- 115 Uhl FE, Vierkotten S, Wagner DE, *et al.* Preclinical validation and imaging of Wnt-induced repair in human 3D lung tissue cultures. *Eur Respir J* 2015;46:1150–66.
- 116 Durham AL, McLaren A, Hayes BP, *et al.* Regulation of Wnt4 in chronic obstructive pulmonary disease. *FASEB J* 2013;27:2367–81.
- 117 Heijink IH, de Bruin HG, van den Berge M, *et al.* Role of aberrant WNT signalling in the airway epithelial response to cigarette smoke in chronic obstructive pulmonary disease. *Thorax* 2013;68:709–16.
- 118 Heijink IH, de Bruin HG, Dennebos R, *et al.* Cigarette smoke-induced epithelial expression of WNT-5B: implications for COPD. *Eur Respir J* 2016;48:504–15.
- 119 van Dijk EM, Menzen MH, Spanjer AI, *et al.* Noncanonical WNT-5B signaling induces inflammatory responses in human lung fibroblasts. *Am J Physiol Lung Cell Mol Physiol* 2016;310:L1166–76.
- 120 Whang YM, Jo U, Sung JS, *et al.* Wnt5a is associated with cigarette smoke-related lung carcinogenesis via protein kinase C. *PLoS ONE* 2013;8:e53012.
- 121 Xi S, Xu H, Shan J, *et al.* Cigarette smoke mediates epigenetic repression of miR-487b during pulmonary carcinogenesis. *J Clin Invest* 2013;123:1241–61.
- 122 Nemeth MJ, Topol L, Anderson SM, *et al.* Wnt5a inhibits canonical Wnt signaling in hematopoietic stem cells and enhances repopulation. *Proc Natl Acad Sci USA* 2007;104:15436–41.
- 123 Spanjer AI, Menzen MH, Dijkstra AE, *et al.* A pro-inflammatory role for the Frizzled-8 receptor in chronic bronchitis. *Thorax* 2016;71:312–22.
- 124 Chen B, Dodge ME, Tang W, *et al.* Small molecule-mediated disruption of Wnt-dependent signaling in tissue regeneration and cancer. *Nat Chem Biol* 2009;5:100–7.
- 125 Baarsma HA, Bos S, Meurs H, *et al.* Pharmacological inhibition of GSK-3 in a Guinea pig model of LPS-induced pulmonary inflammation: I. Effects on lung remodeling and pathology. *Respir Res* 2013;14:113.
- 126 McCubrey JA, Steelman LS, Bertrand FE, *et al.* GSK-3 as potential target for therapeutic intervention in cancer. *Oncotarget* 2014;5:2881–911.
- 127 Kahn M. Can we safely target the WNT pathway? *Nat Rev Drug Discov* 2014;13:513–32.
- 128 Chen M, Wang J, Lu J, *et al.* The anti-helminthic niclosamide inhibits Wnt/Frizzled1 signaling. *Biochemistry* 2009;48:10267–74.
- 129 Jenei V, Sherwood V, Howlin J, *et al.* A t-butylloxycarbonyl-modified Wnt5a-derived hexapeptide functions as a potent antagonist of Wnt5a-dependent melanoma cell invasion. *Proc Natl Acad Sci USA* 2009;106:19473–8.
- 130 Laeremans H, Hackeng TM, van Zandvoort MA, *et al.* Blocking of frizzled signaling with a homologous peptide fragment of wnt3a/wnt5a reduces infarct expansion and prevents the development of heart failure after myocardial infarction. *Circulation* 2011;124:1626–35.
- 131 Nagayama S, Fukukawa C, Katagiri T, *et al.* Therapeutic potential of antibodies against FZD 10, a cell-surface protein, for synovial sarcomas. *Oncogene* 2005;24:6201–12.
- 132 Paes KT, Wang E, Henze K, *et al.* Frizzled 4 is required for retinal angiogenesis and maintenance of the blood-retina barrier. *Invest Ophthalmol Vis Sci* 2011;52:6452–61.