

α -Synuclein occurs in lipid-rich high molecular weight complexes, binds fatty acids, and shows homology to the fatty acid-binding proteins

Ronit Sharon[†], Matthew S. Goldberg[†], Ifat Bar-Josef[†], Rebecca A. Betensky[‡], Jie Shen[†], and Dennis J. Selkoe^{†§}

[†]Center for Neurologic Diseases, Harvard Medical School and Brigham and Women's Hospital, Boston, MA 02115; and [‡]Harvard School of Public Health, Boston, MA 02115

Communicated by Laszlo Lorand, Northwestern University Medical School, Chicago, IL, June 14, 2001 (received for review March 23, 2001)

α -Synuclein (α S) is a 140-residue neuronal protein that forms insoluble cytoplasmic aggregates in Parkinson's disease (PD) and several other neurodegenerative disorders. Two missense mutations (A53T and A30P) are linked to rare forms of familial PD. The normal function of α S is unknown, and cultured cell systems that model its modification from soluble monomers to aggregated forms have not been reported. Through a systematic centrifugal fractionation of mesencephalic neuronal cell lines and transgenic mouse brains expressing wild-type or A53T human α S, we observed unusual, previously unrecognized species of α S that migrate well above the 17-kDa monomeric form in denaturing gels. Incubation at 65°C of high-speed cytosols from cells or brains revealed a modified α S species migrating at \approx 36 kDa and an extensive higher molecular mass α S-reactive smear. Extraction of the cytosols with chloroform/methanol or with a resin (Lipidex 1000) that binds fatty acids resulted in a similar pattern of higher molecular mass α S forms. On the basis of this effect of delipidation, we reexamined the primary structure of α S and detected a motif at the N and C termini that is homologous to a fatty acid-binding protein signature. In accord, we found that purified human α S binds oleic acid, with an apparent K_d of 12.5 μ M. We also observed an enhanced association of A53T α S with microsomal membranes in both mesencephalic cells and transgenic mouse brains. We conclude that α S has biochemical properties and a structural motif that suggest it is a novel member of the fatty acid-binding protein family and may thus transport fatty acids between the aqueous and membrane phospholipid compartments of the neuronal cytoplasm.

Progress in characterizing insoluble protein aggregates in human neurodegenerative diseases has led to new insights into the molecular mechanisms of these previously obscure disorders (1). A salient example is the discovery of the protein α -synuclein (α S) as the principal constituent of the Lewy bodies that are the cytopathological hallmark of dopaminergic neuronal degeneration in Parkinson's disease (PD) (2, 3). This finding emerged from the observation of α S missense mutations in rare dominantly inherited forms of PD (4, 5). Lewy bodies, which are ubiquitinated cytoplasmic inclusions, and related neuritic alterations called Lewy neurites contain insoluble filamentous aggregates of the otherwise highly soluble α S protein (2, 6–8). Such aggregates also occur in diffuse Lewy body dementia, multiple systems atrophy, and some cases of Alzheimer's disease and Down's syndrome (9–12). Accordingly, there is now great interest in elucidating the normal structure–function relationships of α S in neurons and their modification in PD and the other “synucleinopathies.”

α S is a 140-residue soluble protein that is abundantly expressed in neurons, where it is localized in part to presynaptic nerve endings (13). Developmental studies in songbirds suggest that its expression in certain neurons is associated with the acquisition of learning (14). It may thus play a role in synaptic plasticity, perhaps by helping to maintain the size of the presynaptic vesicular pool (15). α S can inhibit *in vitro* the enzyme phospholipase D2, which hydrolyzes

phosphatidylcholine to phosphatidic acid and may be involved in vesicle trafficking in the secretory pathway (16–18). Although α S is believed to be principally a soluble cytosolic protein, some portion of α S is associated with membranes (19–22). The N-terminal region of α S may interact reversibly with membranes via 7 repeated motifs of 11 amino acids each that are shared with the class A2 helices of apolipoproteins (9–12). α S appears to associate with membranes via not only this domain but also its C-terminal region, in which no such lipophilic sequence has been observed (21). *In vitro* studies suggest that α S binds acidic phospholipid vesicles in a way that markedly alters its secondary structure (23, 24).

Some transgenic (tg) mouse lines expressing wild-type (wt) human α S develop neuronal cytoplasmic inclusions reminiscent of Lewy bodies, loss of dopaminergic synapses, and associated motoric impairment (25). Expression of wt or mutant human α S in *Drosophila* leads to Lewy body-like inclusions in dopaminergic neurons associated with neuronal loss and impaired crawling (26). *In vitro* aggregation studies of purified α S show that its oligomerization into protofibrillar intermediates is accelerated by both the A53T and A30P missense mutations (8, 27, 28), although only A53T also increases the conversion of such intermediates into amyloid-like fibrils (29). The occurrence of the mutations in the region of homology to the class A2 helices of apolipoproteins suggests possible altered interactions with membrane lipids and mislocalization of α S. However, α S distribution in fractionated brain synaptic vesicles was found to be unaltered in wt vs. A30P tg mice (20), and no differences in membrane association were detectable in transfected primary cortical neurons by using fluorescence resonance energy transfer (21).

Here, we conducted a systematic biochemical fractionation study of α S in non-tg and tg mouse brains and in an α S-transfected rat mesencephalic (MES) neuronal cell line, MES 23.5 (30). The results reveal that α S shares biochemical and structural features with the fatty acid-binding proteins (FABPs) and binds fatty acids (FAs).

Materials and Methods

Cultured Cells. The rodent mesencephalic neuronal cell line MES 23.5, which has dopaminergic properties (30), was stably transfected with wt or A53T α S cDNA in the pCDNA 3.1 vector. An unexpected and striking feature of these α S-transfected MES cells was that all stable clones gradually lost α S expression after being continuously passaged for 2–3 mos or more. The wt and A53T clones selected for the study expressed similar levels of α S.

Abbreviations: α S, α -synuclein; PD, Parkinson's disease; wt, wild type; FABP, fatty acid-binding protein; tg, transgenic; FA, fatty acid; AA, arachidonic acid; DHA, docosahexaenoic acid; OA, oleic acid; MES, rat mesencephalic neuronal cell line.

§To whom reprint requests should be addressed. E-mail: Selkoe@cnd.bwh.harvard.edu.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. §1734 solely to indicate this fact.




Fig. 1. Distribution of α S in subcellular fractions in normal and tg mouse brains and transfected MES cells. (A) Scheme of the centrifugal fractionation procedure. (B) Endogenous α S distribution in normal mouse brain. Whole normal mouse brain was fractionated as in A and subjected to quantitative Western blotting with H3C Ab. Results are presented as the ratio of the amounts of α S monomer (17 kDa) in each fraction to that of total α S monomer in the starting homogenate (100%) run in parallel on each gel (means \pm SD of three brains). (C) Higher portion of A53T than WT α S is associated with P25 microsomes. Results are presented as the ratio of the amounts of α S monomer in P25 to that in S370 analyzed in parallel. [Means \pm SD of $n = 5$ brains and $n = 8$ MES cultures each for wt (empty bars) and A53T (filled bars).]

Tg Mice. Mice expressing human wt or A53T mutant α S were generated routinely, as will be described in detail elsewhere (M.S.G., C. A. Lemere, M. P. Frosh, P. T. Lansbury, and J.S., unpublished work). Briefly, constructs containing the human wt or A53T mutant α S cDNA behind the platelet-derived growth factor (PDGF)-B promoter were injected into pronuclei of C57BL/6J mouse embryos, and resulting tg mice were maintained in the same background. Western and Northern analyses showed brain levels of human α S approximately half those of endogenous mouse α S, primarily in the cerebral cortex and hippocampus, as expected from the known expression pattern of the PDGF-B promoter. A complete description of the ongoing phenotypic analysis of these mice will be reported elsewhere.

Fractionation Protocol. A protocol described previously (31) was modified to gain improved separation of microsomal vesicles (see Fig. 1A). All procedures were at 4°C. Briefly, whole mouse brains or cultured MES cell pellets ($\approx 2 \times 10^7$ cells) were homogenized (20 up-and-down strokes with a Teflon homogenizer, followed by five passages through a 22-gauge needle) in 1:10 (wt/vol) homogenization (H) buffer [20 mM Hepes, pH 7.4/1 mM $MgCl_2$ /0.32 M sucrose/43 mM 2-mercaptoethanol/1 \times protease inhibitor mix (Sigma)]. The homogenate was centrifuged at $170 \times g$ for 10 min. The resultant pellet (P1) was washed in H buffer and respun at $370 \times g$ for 15 min. The pellet (P1*) was resuspended in H buffer, and the suspension was brought to 2.1 M sucrose and spun at $175,000 \times g$ for 1 h. The pellet (consisting of nuclei) and the lipid-rich fraction floating to the top of this sucrose cushion were collected. S1 supernatant was centrifuged at $8,000 \times g$ for 15 min to obtain P2 and S2. S2 supernatant was then subjected to sequential 60-min spins at

$25,000 \times g$, $100,000 \times g$, $230,000 \times g$, and $370,000 \times g$. At each speed, the resultant supernatant (designated S25, S100, etc.) was transferred to a clean tube, and the resultant pellet was washed and then resuspended in homogenization buffer containing 1% Nonidet P-40. The pellets were extracted on ice for 20 min, cleared in a microfuge at $15,000 \times g$ for 10 min, and the detergent-solubilized pellet extract was transferred to a clean tube. Samples were stored at $-70^\circ C$ for up to 6 mos. Protein concentrations were assayed by the Bradford method (32).

Western Blotting for Quantitative Analysis of the Distribution of α S Monomer (17 kDa) in the Above Brain and Cell Fractionations. To reduce experimental variation, experiments were conducted in pairs, i.e., wt and A53T samples were worked up in parallel on the same day, from harvesting the cells or sacrificing the mice through the fractionation, protein assay, and gel electrophoresis steps. Equal amounts of total protein (in duplicate for each fraction) were loaded on 8–16% Tris-glycine gels (NOVEX, San Diego), transferred to poly(vinylidene difluoride) membrane, and blotted by using an enhanced chemiluminescence detection kit (Amersham Pharmacia). The blots were scanned in a Umax Magic Scan (Eastman Kodak) and analyzed for density of the α S monomeric band by using UN-SCAN-IT GEL 3.1 software (Silk Scientific). Results in the various fractions were normalized to the amount of α S monomer either in the corresponding total homogenate or in the S370 fraction run in parallel on each gel, as indicated. Although the stable lines analyzed express closely similar amounts of α S, results are presented as the ratio of α S in a specific fraction to total or S370 α S to eliminate small differences in expression levels. α S Abs and their epitopes were: LB509 (amino acid 115–122) (3); 15G7 (amino acid 116–121) (20); Syn-1 (amino acid 15–123) (Transduction Laboratories, Lexington, KY); H3C (amino acid 126–140) (14); PER4 (amino acid 1–120) (33); affinity purified 7384 (amino acid 123–140) (this study).

Lipid Removal by Chloroform/Methanol Extraction. S370 samples were extracted with chloroform/methanol to remove lipids attached to proteins, as originally described (34). In an Eppendorf tube, 2 vol of chloroform and 1 vol of methanol were mixed, and 0.25 vol of S370 was added. The mixture was vortexed for 30 sec and centrifuged at $10,000 \times g$ for 5 min (room temperature). The resulting aqueous (upper) and organic (lower) phases were removed. The interface, containing all α S proteins (see Results), was lyophilized and resuspended in water or 2 \times Laemmli buffer.

FA-Binding Studies. S370 samples were incubated at 37°C or 65°C overnight with Lipidex 1000 (5% wt/vol). The tube was spun at $10,000 \times g$ for 5 min and the supernatant transferred to a clean tube. Proteins were eluted from the Lipidex pellet by boiling for 10 min in 2 \times Laemmli buffer.

Sequence Homology Search. We searched through the ExPASy web site (<http://expasy.cbr.nrc.ca>) under “proteomic tools” for “pattern and profile searches,” by using scan Prosite. By allowing mismatches of up to 50%, we detected the “cytosolic FABP signature” motif in the α S sequence (see Results) by using the option “PROSITE scan” at the URL <http://expasy.cbr.nrc.ca/tools/#pattern>.

Oleic Acid (OA)-Binding Assays. Assays were carried out as previously described (35). Briefly, purified human recombinant wt α S was resuspended in binding buffer (10 mM Tris-HCl, pH 7.5, 150 mM NaCl), filtered through Microcon YM-100 (Millipore), and its concentration determined both by OD 276 nm and Bradford (32). Samples containing wt α S (5 μ M) and [^{14}C] OA (NEN) (at the indicated concentrations) in 100 μ l of binding buffer were processed as described previously for the Lipidex assay (35). Each value was corrected against a blank processed identically

but without α S protein. Cold FA displacement experiments were performed by using the same assay. The reaction mix contained fixed concentrations of α S (5 μ M) and [14 C] OA (10 μ M) and increasing concentrations of the unlabeled competitors: arachidonic acid (AA), docosahexaenoic acid (DHA), and OA. The radiolabeled OA and the competitors were added simultaneously to the protein. For quantifying FA binding directly to the S370, extracts (5 μ g) from nontransfected or wt α S-transfected MES cells were treated with 0.5% SDS for 10' at 37°C. SDS was diluted 1:10 with binding buffer to final reaction vol of 100 μ l, and the S370 was subjected to the Lipidex assay (above).

Statistical Analyses. The amounts of α S in P25 membrane pellets (Fig. 1A and B) were assumed to follow a Poisson distribution (36) with rate given by $2 \cdot S370 \cdot \mu$. We analyzed the equality of the parameters μ between P25 α S values from A53T vs. wt α S and between tg mice vs. MES cells by using Poisson regression, adjusting for correlation from experiments conducted at the same time by including random effects for time of experiment. This was fit by using SAS software (SAS Institute, Cary, NC), release 8.1.

Results

The initial goals of this study were to determine the relative distribution of soluble and membrane-associated forms of α S in neural cells and to search specifically for forms of the protein that are aggregated or have unusual solubility properties and thus might be relevant to PD. To these ends, we carried out a systematic centrifugal fractionation of α S from two disease-relevant sources: human α S stable MES transfectants and human α S tg mouse brains. We expressed wt or A53T α S in the MES cells and selected stable clones with closely similar α S expression levels. To complement the data obtained in these neuronal cell lines, we examined C57BL/6J mice tg for either wt or A53T human α S. For some studies, we compared the results obtained in the wt and A53T human α S tg mice with the properties of endogenous α S in brains from non-tg littermates.

Fractionation of Brain and MES α S by Differential Centrifugation. The distribution of endogenous α S was first examined in normal (non-tg) mouse brain by using quantitative Western blotting (see *Materials and Methods*). Approximately 50% of total brain α S was found to be cytosolic, i.e., was recovered in the final high-speed (370,000 \times g) supernatant (Fig. 1B). The remaining \approx 50% of brain α S was distributed among several membrane-enriched pellets, with the large majority of this noncytosolic protein being recovered in pellets P1 (170 \times g) (\approx 25% of total) and P2 (8,000 \times g) (\approx 20% of total). No significant α S-reactive protein was detected in the nuclear fraction, which was obtained from the P1 pellet (Fig. 1A). However, the lipid-rich material floating to the top of the sucrose cushion used to pellet the nuclei was highly α S immunoreactive (see below).

Because the primary structure of α S predicts that the N-terminal portion interacts with lipid molecules and thus with membranes, and because the two known PD-associated mutations are localized to this region, we were interested in the portion of total brain α S associated with various sized microsomal membranes. We therefore used differential centrifugation to collect four higher-speed membrane pellets from the S2 sup:P25 (i.e., 25,000 \times g), P100, P230, and P370 (Fig. 1A). Taken together, these four pellets contained about 8% of total brain α S. Among these four higher-speed membrane pellets, the highest percentage of total brain α S was associated with P25 (mean \approx 7%) (Fig. 1B). Western blotting with a panel of membrane protein markers revealed that P25 was particularly enriched in calnexin (an endoplasmic reticulum marker) and several markers of endosomes, including transferrin, syntaxin 6, and syntaxin 13, as well as synaptotagmin, a synaptic vesicle marker (data not shown).

Next, we carried out the same fractionation protocol on the

brains of tg mice expressing wt or A53T human α S and on MES neuronal lines stably expressing these α S forms. Using the human α S-specific Ab, LB509 (3), we found that the overall distribution of both human isoforms among the tg brains and MES cell fractions was indistinguishable from that of endogenous mouse brain α S (Fig. 1B), i.e., was well within the range of experimental variation. Because P25 contained most of the α S immunoreactivity among the four microsomal pellet fractions, we compared the relative amounts of P25 α S immunoreactivity (normalized to the amount of S370 α S in the same fractionation) between wt and A53T expressors. Significantly more α S was associated with P25 microsomes in A53T than in wt α S expressors, both in the MES cells ($P < 0.001$) and in the tg mouse brains ($P < 0.014$) (Fig. 1C).

Identification of Higher Molecular Mass α S-Reactive Polypeptides in the Cytosol of Brain Tissue and Mesencephalic Neuronal Cells. In the quantitative Western blot analysis of α S distribution described above, the vast majority of α S-reactive protein in all of the fractions migrated at \approx 17 kDa, the expected molecular mass for the α S monomer (3, 14, 37). However, we also found variable amounts of α S-reactive protein that remained gel-excluded (i.e., in the sample well). In most fractionations, we further observed variable amounts of an \approx 85-kDa α S-reactive protein specifically labeled by LB509 in the very high-speed pellet (P370) and supernatant (S370) of both the MES transfectants and tg mouse brains (not shown). In view of these small amounts of apparent higher M_r forms of human α S in the cytosol of mesencephalic neurons and brain, we undertook a concerted search for larger α S forms by attempting to dissociate or depolymerize such forms using a variety of chemical and physical treatments on the S370 supernatants.

Treatment of the S370 cytosol with 8 M urea, 2% SDS, 1.5% β -ME, divalent cations (10 mM $MgCl_2$, 10 mM $CaCl_2$ or 1 mM $ZnSO_4$), 5 mM EDTA, 0.1 M NaOH, or 0.1 M HCl did not alter the α S electrophoretic pattern (not shown). We next incubated the S370 at either 37°C, 65°C, or 100°C for increasing time intervals to determine whether heat would alter the pattern of α S immunoreactivity (Fig. 2A–D). Incubation at 37°C for 18 h revealed a α S-reactive 36-kDa band, which was not observed at 0°C (Fig. 2A). A dramatic change in the α S Western blot pattern was observed upon incubation at 65°C for 18 h. The change comprised a modest increase in the amount of α S monomer, the appearance of the 36-kDa band that became increasingly visible between 2 and 24 h of 65°C incubation, and the appearance of an extensive smear of α S immunoreactive material throughout the middle to higher M_r range of the gel (Fig. 2A and B). This pattern was highly reproducible in both mouse brains (non-tg and tg) and MES transfectants. We observed no detectable difference in these changes between wt and A53T samples, both in brains and MES cells. This temperature-dependent change in the α S-reactive pattern of the S370 cytosols was not observed in any of the membrane pellets (P25 to P370), but it was observed in the lipid-rich floating material obtained during the sucrose cushion fractionation of the P1 pellet (not shown).

Incubation of S370 at 100°C for 18 h led to a α S-reactive pattern partially reminiscent of that at 65°C, but it was accompanied by a loss of the α S monomer and the appearance of gel-excluded α S in the sample well (Fig. 2A). Incubation of S370 at 37°C for 18 h did not reveal the higher M_r smear (Fig. 2A); however, longer (24- or 45-h) incubations at 37°C did produce a pattern closely similar to that seen after 65°C incubation (not shown).

To determine the specificity of the new α S-reactive species that appeared with 65°C treatment, we performed Western blotting with six different α S-specific Abs (see *Materials and Methods*) and found that all six Abs recognized the 17-kDa monomer, the 36-kDa band, and the mid-high M_r smear (Fig. 2C and data not shown). Preabsorption of H3C, LB509, or Syn-1 Abs with purified recombinant human α S protein markedly reduced the staining of the 17-kDa monomer and abolished the




Fig. 2. α S forms higher M_r aggregated species in the cytosols of mouse brain and α S-transfected MES cells. (A) The appearance of higher M_r α S immunoreactive forms is temperature-dependent. Aliquots of S370 cytosol (20 μ g) from normal mouse brain were incubated at the indicated temperatures for 18 h, loaded on an 8–16% Tris-Glycine SDS gel, and blotted with H3C Ab. *, Gel-excluded material. (B) Appearance of the higher M_r α S immunoreactive forms is time-dependent. Aliquots (20 μ g) of S370 cytosol from human wt α S transfected MES cells were incubated at 65°C for the indicated times and blotted with human α S-specific Ab LB509. (C) The α S-reactive higher M_r material is immunospecific. The indicated α S Abs were preabsorbed (+) or not (–) at 4°C overnight with 5 μ g purified α S per microliter Ab and used to blot duplicate aliquots (20 μ g) of S370 (from wt human α S-transfected MES cells) that had been incubated either at 0°C or 65°C overnight. (D) α S monomer is shifted upwards by \approx 1 kDa on 18 h incubation at 65°C (also visible in C). S370 aliquots (20 μ g) from wt α S-transfected MES cells were blotted with LB509.

36-kDa band and the higher M_r smear (Fig. 2C). By densitometry, the total amount of α S immunoreactivity detectable by Western blots was 10- to 100-fold higher in the 65°C-treated S370 fractions than in the same cytosol incubated simultaneously at 0°C. On very brief development of the Western blots, we noticed that the α S monomeric band migrating at 17 kDa before 65°C incubation shifted to a slightly higher molecular mass (18 kDa) after the 65°C treatment (Fig. 2D). This subtle but highly consistent shift in apparent M_r was observed in non-tg and tg mouse brains as well as in transfected MES cells. In some samples, only a portion of the α S monomer was up-shifted, resulting in a closely spaced doublet (17 + 18 kDa) containing the up-shifted and the conventional monomers (not shown).

Both Monomeric and Higher M_r α S Species in the Cytosol Are Bound to Lipids. The striking increase in α S immunoreactive material effected by 65°C treatment suggested that heating exposed α S-reactive epitopes by denaturing the protein, releasing lipids or other bound molecules from it, and/or depolymerizing higher molecular weight forms and allowing them to enter the gel. In an attempt to distinguish among these explanations, we assessed the effects of delipidating agents on the α S immunoreactive pattern. Extraction of the S370 cytosol with chloroform/methanol (2:1) resulted in the recovery of α S-reactive monomer and abundant higher M_r species in the interface between the upper (aqueous) and lower (organic) phases (Fig. 3A, lane 2). This α S-reactive pattern after chloroform/methanol extraction was immunospe-




Fig. 3. α S associates with lipids in brain and MES cells. (A) Aliquots (20 μ g) of S370 cytosol from normal mouse brain were untreated (lane 1), treated with chloroform/methanol and the interface between the upper and lower phases collected (lane 2) or incubated at 65°C overnight (lane 3) and then blotted with H3C. (B) Exposure of novel α S species by treatment with a FA-binding resin (Lipidex 1000), without or with heating. Aliquots (20 μ g) of S370 from human wt α S-transfected MES cells were incubated overnight with or without Lipidex at the indicated temperatures, and samples were spun (10,000 \times g for 5 min). The post-Lipidex supernatant (S-Lx) and the sample without Lipidex treatment (S370) were loaded directly, whereas the Lipidex pellet (P-Lx) was boiled for 10 min in 2 \times Laemmli buffer, and the extract was loaded. LB509 Western blots.

cific (not shown) and resembled in part that observed when S370 was incubated at 65°C (Fig. 3A, lane 3).

An alternative method of attempting to delipidate the various α S species is the use of lipid-binding resins, which can extract lipid bound to proteins (38–40). We therefore treated the S370 fraction with Lipidex 1000, a resin that principally binds FAs (see *Materials and Methods*). The Lipidex-treated supernatant again showed a temperature-dependent α S pattern. After Lipidex treatment at 0°C, we saw no effect on the α S-reactive pattern (Fig. 3B). Lipidex treatment at 37°C led to the recovery of some monomeric α S and a small amount of the 36-kDa species in the supernatant. However, greater amounts of both monomeric α S and the 36-kDa band were recovered in the Lipidex pellet (Fig. 3B). At 65°C, we again observed the marked enhancement of all α S species. At this temperature, the post-Lipidex supernatant and the Lipidex pellet contained more α S immunoreactivity than the S370 before Lipidex, and the higher M_r immunoreactive bands appeared more defined, i.e., not just as a smear (Fig. 3B). In this experiment, the post-Lipidex supernatant and Lipidex pellet originate from the same cytosol sample and contain the same amount of starting material as the parallel aliquot incubated without Lipidex (Fig. 3B). Therefore, much more α S-reactive material was recovered in the presence of Lipidex. We conclude that the recovery of novel higher M_r α S species does not depend on temperature alone but also occurs on removal of lipids, particularly FAs given the specificity of Lipidex.

α S Shares Sequence Homology with FABPs and Binds OA Stochiometrically. Taken together, the effects in both brain and MES cells of the 65°C incubation, the chloroform/methanol treatment, and the delipidation with Lipidex all strongly support previous *in vitro* evidence that α S interacts with lipids (19, 23, 24). Further, these data suggest that the lipids mask immunoreactive epitopes in the α S protein. We therefore searched the α S amino acid sequence for motifs that might have homology with known lipid-binding proteins (see *Materials and Methods*). We found a significant regional homology to the family of FABP. We observed an 18-residue sequence at the extreme C terminus of α S that showed 67% homology (or 83% including the intervening space-holding residues) with a signature motif char-




Fig. 4. α S has characteristics of a FABP. (A) The cytosolic FABPs signature. The 18-residue motif comprises nine specific residues and nine undefined residues (represented by X). The homology percentage is calculated either by considering only the specific residues [67% in the amino acid 123–140 stretch (C terminus) and 55% in the amino acid 2–19 stretch (N terminus)] or by including the X residues needed to maintain proper spacing between the specific residues (83% at the C terminus and 78% at the N terminus). Underlined residues are shared with the FABP signature. (B) Purified human α S (5 μ M) was incubated with increasing concentrations of 14 C OA. Protein-bound OA was separated from free OA by using the Lipidex assay (see *Materials and Methods*). Data points are means of triplicates \pm SD. (Inset) Scatchard plot. Estimated K_d and B_{max} values were 12.5 μ M and 1, respectively. (C) FA displacement curve: purified human α S (5 μ M) was incubated with 10 μ M [14 C] OA and increasing concentrations of unlabeled OA (●) AA (■) or DHA (▲), as indicated. (D) Aliquots (5 μ g) of S370 cytosols from untransfected or wt- α S-transfected MES cells were incubated with increasing concentrations of 14 C OA, as in B. Nontransfected cells (●), α S transfected cells (■). (E) Aliquots of S370 (20 μ g) were incubated at 65°C overnight without (–) or with (+) free cold OA (5 μ M) and Western blotted with LB509.

acteristic of the FABP (Fig. 4A). A stretch of 18 residues near the N terminus of α S also showed 55% homology (or 78% including intervening space-holding residues) with this same motif of the FABP. These previously unrecognized homologies suggest that α S shares some structure and properties with the FABP. Consistent with this hypothesis are the observations that the length of α S (140 aa) is closely similar to that of the large majority of FABPs (41) and that α S, like FABPs, is a cytosolic protein that associates with membranes and is subject to regulation by phosphorylation (42).

In view of these similarities, we asked whether purified recombinant α S was capable of incorporating FA. We measured 14 C OA binding by purified human wt α S in the Lipidex assay (see *Materials and Methods*). OA was chosen as a general FA ligand commonly used to characterize FABP and does not indicate the nature of any *in vivo* FA ligand(s). A dose-dependent binding of 14 C OA up to saturation was observed when a constant

amount (5 μ M) of purified recombinant human wt α S was incubated with increasing OA concentrations (2–15 μ M) (Fig. 4B). A Scatchard plot of the binding data yielded an estimate for the apparent K_d of 12.5 μ M (mean of $n = 3$). The predicted molar ratio of α S to OA is 1:1, based on a B_{max} of 1.

The specificity of the OA binding was then tested by displacement with unlabeled DHA, AA, and OA. DHA displaced 14 C OA binding most potently (i.e., 100% at 150 μ M), followed by AA (100% at 175 μ M) and OA (100% at 200 μ M) (Fig. 4C). Next we attempted to relate this binding of FA to purified human α S (Fig. 4B and C) to the effect of delipidation of cytosolic human α S in mouse brains and MES neuronal cells (Figs. 2 and 3). We compared the amounts of 14 C OA incorporation into the S370 cytosols of untransfected vs. human wt α S-transfected MES cells. The S370 cytosols of untransfected cells underwent a dose-dependent incorporation of OA (Fig. 4D). The α S-transfected S370 incubated simultaneously under identical conditions showed greater amounts of OA incorporation (Fig. 4D). To further link these findings to the α S-reactive species revealed by delipidation, we incubated S370 cytosol samples at 65°C without or with unlabeled OA and examined it by Western blotting. OA incubation substantially masked the 36 kDa and higher α S-reactive species, while only slightly decreasing the appearance of the 17-kDa monomer (Fig. 4E). Thus adding FA produces the opposite effect on the α S-reactive high M_r pattern than does removal of FAs by Lipidex 1000.

Discussion

α S is a principal constituent of Lewy bodies (2, 3), and recombinant α S assembles into filaments that are structurally and immunochemically highly similar to those of Lewy bodies (8, 27). These results indicate that α S accumulates abnormally and can aggregate into insoluble forms in PD. However, there have been no biochemical descriptions heretofore of altered or aggregated forms of wt or mutant α S in *in vivo* models, either in cell culture or in the brains of tg α S mice.

During a systematic biochemical fractionation of α S in mesencephalic neuronal cells and mouse brains expressing human α S, we discovered unusual, previously unrecognized species of α S that migrate well above the 17-kDa monomeric form in denaturing gels. A \approx 36-kDa species is invariably present after 65°C incubation of the high-speed cytosol (S370) of mouse brain and mesencephalic neuronal cells. In addition, an extensive immunoreactive smear of α S material appears in the middle and upper M_r regions of the gels. The occurrence of these species not only in tg mice and stable MES transfectants but also endogenously in normal mouse brain suggests that modified and/or aggregated forms of α S occur normally in neurons.

Because the amount of α S monomer (17 kDa) was unchanged or modestly increased between the 0°C and 65°C incubations, it is unlikely that the newly appearing 36-kDa form and the higher smear of α S-reactive material arise from α S polymerization *in vitro* during the incubations. Rather, we hypothesize that this treatment exposes hitherto buried epitopes on an array of α S-reactive species in the cytosol that are not visualized under conventional Western blotting conditions and/or causes partial depolymerization of very high M_r aggregates of α S that cannot otherwise be detected by Western blotting. The reduced recovery of these species after incubating the cytosols with free OA (Fig. 4E) supports the first explanation of epitope exposure. We interpret the small upward shift in electrophoretic migration of the α S monomer consistently induced by 65°C incubation (Fig. 2D) as evidence of the unfolding of a partially SDS-resistant conformer of the α S monomer. Such subtle shifts in the electrophoretic migration of lipid-binding proteins on further denaturation have been described before, including in the brain proteolipid protein (43).

The effect of lipid extraction by using the Lipidex 1000 resin is strongly indicative of the presence of lipid molecules, most

likely FAs, on α S, including on the 36-kDa and higher M_r α S-reactive species. The nature of the endogenous lipids that are attached to α S in mesencephalic neurons and in the brain is currently unknown. However, our unexpected observation of a regional homology between α S and a conserved motif characteristic of the FABP family, coupled with our demonstration that OA is incorporated stoichiometrically into purified α S, leads us to hypothesize that α S may function as a FABP. Indeed two FAs that are abundant in brain tissue, AA and DHA, displace OA from purified α S in a dose-dependent fashion. Moreover, the enhanced binding of OA by the S370 cytosol of α S-overexpressing MES cells (Fig. 4D) is consistent with a role for α S as a FABP.

FABP are small (≈ 15 kDa) cytosolic lipid-transport proteins of diverse primary structure that include a large and growing list of tissue-specific proteins. The 14.5-kDa brain-specific (B)-FABP has been localized to glial cells, in particular to radial glia (44). B-FABP has not been described in neurons and Schwann cells. Because α S is principally expressed in the presynaptic terminals of neurons in the cerebral cortex, hippocampus, basal ganglia, and brainstem (13, 45), it appears that B-FABP and α S do not coexist in brain cells. As α S is expressed in very high abundance in the brain [≈ 0.5 – 1% of total cytosolic proteins (11)], our findings suggest that α S is a novel brain-specific FABP.

The temperature-dependent effects of Lipidex treatment we report (Fig. 3B) are similar to those described previously for other proteins (40). Importantly, the latter study found that Lipidex binds only free FA at 0°C , whereas it binds protein-bound as well as free FA at 37°C (40). On this basis, we propose that the presence of the 36-kDa α S-reactive species bound to the Lipidex pellet after 37°C incubation (Fig. 3B, lanes 4–6) represents a FA-bound form of the protein. The combined treatment of the S370 cytosol with Lipidex at 65°C markedly enhances the

appearance of additional higher M_r α S-reactive species (Fig. 3B, lanes 7–9). We hypothesize that these forms are being visualized because of a dual effect: temperature-dependent partial denaturation exposes not only α S immunoreactive epitopes but also bound FA molecules, and the latter are captured by Lipidex.

The finding of α S association with various microsomal membranes is of special interest in light of its conserved FABP signature and its binding of FAs. FABP have been shown to transport free FAs from the aqueous phase of the cytoplasm to various membranes, during which the FABP transiently associates with the membrane to release a FA to or remove one from the membrane, a process called collisional transfer (41). The significant increase in the amount of α S associated with P25 microsomal membranes in both mouse brains and MES cells expressing the A53T familial PD mutation (Fig. 1C) raises the possibility that the function of α S in the FA transport process is partially altered by this mutation.

The accumulation of modified insoluble forms of wt α S in the Lewy bodies and Lewy neurites of idiopathic PD and in related cytoplasmic inclusions of other synucleinopathies leads us to speculate that a necessary FA transport function of cytosolic α S is altered in these disorders. Our mesencephalic neuronal cell model and tg mice should be useful in further testing this hypothesis.

We thank Drs. Wei-Dong Le and Stanley Appel (Baylor College of Medicine, Houston, TX) for providing the MES 23.5 cell line; Drs. Jean-Christoph Rochet and Peter Lansbury, Jr. (Brigham and Women's Hospital) for providing purified recombinant human α S, and Drs. Judith Storch, Marjorie Lees, and Omanand Koul for many helpful suggestions. This work was supported by the Morris K. Udall PD Center (National Institutes of Health Grant T50NS38375) at the Brigham and Women's Hospital.

- Lansbury, P. T. (1997) *Neuron* **19**, 1151–1154.
- Spillantini, M. G., Schmidt, M. L., Lee, V. M.-Y., Trojanowski, J. Q., Jakes, R. & Goedert, M. (1997) *Nature (London)* **388**, 389–390.
- Baba, M., Nakajo, S., Tu, P.-H., Tomita, T., Nakaya, K., Lee, V. M.-Y., Trojanowski, J. Q. & Iwatsubo, T. (1998) *Am. J. Pathol.* **152**, 879–884.
- Polymeropoulos, M. H., Lavedan, C., Leroy, E., Ide, S. E., Dehejia, A., Dutra, A., Pike, B., Root, H., Rubenstein, J., Boyer, R., et al. (1997) *Science* **276**, 2045–2047.
- Kruger, R., Kuhn, W., Muller, T., Woitalla, D., Graeber, M., Kosel, S., Przuntek, H., Epplen, J. T., Schols, L. & Riess, O. (1998) *Nat. Genet.* **18**, 106–108.
- Spillantini, M. G., Crowther R.A., Jakes, R., Hasegawa, M. & Goedert, M. (1998) *Proc. Natl. Acad. Sci. USA* **95**, 6469–6473.
- Lippa, C. F., Fujiwara, H., Mann, D. M. A., Giasson, B., Baba, M., Schmidt, M. L., Nee, L. E., O'Connell, B., Pollen, D., George-Hyslop, P., et al. (1998) *Am. J. Pathol.* **153**, 1365–1370.
- Conway, K. A., Harper, J. D. & Lansbury, P. T. (1998) *Nat. Med.* **4**, 1318–1320.
- Clayton, D. F. & George, J. M. (1999) *J. Neurosci. Res.* **58**, 120–129.
- Hashimoto, M. & Masliah, E. (1999) *Brain Pathol.* **9**, 707–720.
- Kruger, R., Muller, T. & Riess, O. (2000) *J. Neural. Transm.* **107**, 31–40.
- Duda, J. E., Lee, V. M.-Y. & Trojanowski, J. Q. (2000) *J. Neurosci. Res.* **61**, 121–127.
- Iwai, A., Masliah, E., Ge, N., Flanagan, L., de Silva, H. A. R., Kittel, A. & Saitoh, T. (1995) *Neuron* **14**, 467–475.
- George, J. M., Jin, H., Woods, W. S. & Clayton, D. F. (1995) *Neuron* **15**, 361–372.
- Murphy, D. D., Rueter, S. M., Trojanowski, J. Q. & Lee, V. M.-Y. (2000) *J. Neurosci.* **20**, 3214–3220.
- Jenco, J. M., Rawlins, A., Daniels, B. & Morris, A. J. (1998) *Biochemistry* **37**, 4901–4909.
- Exton, J. H. (1997) *Physiol. Rev.* **77**, 303–320.
- Chen, Y.-G., Siddhanta, A., Austin, C. D., Hammond, S. M., Sung T.-C., Frohman, M. A., Morris, A. J. & Shields, D. (1997) *J. Cell. Biol.* **138**, 495–504.
- Jensen, P. H., Nielsen, M. S., Jakes, R., Dotti, C. G. & Goedert, M. (1998) *J. Biol. Chem.* **273**, 26292–26294.
- Kahle, P. J., Neumann, M., Ozmen, L., Muller, V., Jacobsen, H., Schindzielorz, A., Okochi, M., Leimer, U., van Der Putten, H., Probst, A., et al. (2000) *J. Neurosci.* **20**, 6365–6373.
- McLean, P., Kawamata, H., Ribich, S. & Hyman, B. T. (2000) *J. Biol. Chem.* **275**, 8812–8816.
- Segrest, J., Jones, M., De Loof, H., Brouillette, C., Venkatachalapathi, Y. & Anantharamaiah, G. (1992) *J. Lipid Res.* **33**, 141–166.
- Davidson, W. S., Jonas, A., Clayton, D. F. & George, J. M. (1998) *J. Biol. Chem.* **273**, 9443–9449.
- Perrin, R. J., Woods, W. S., Clayton, D. F. & George, J. M. (2000) *J. Biol. Chem.* **275**, 34393–34398.
- Masliah, E., Rockenstein, E., Veinbergs, I., Mallory, M., Hashimoto, M., Takeda, A., Sagara, Y., Sisk, A. & Mucke, L. (2000) *Science* **287**, 1265–1269.
- Feany, M. B. & Bender, M. W. (2000) *Nature (London)* **404**, 394–398.
- Narhi, L., Wood, S. J., Steavenson, S., Jiang, Y., Wu, G. M., Anafi, D., Kaufmann, S. A., Martin, F., Sitney, K., Denis, P., et al. (1999) *J. Biol. Chem.* **274**, 9843–9846.
- Conway, K. A., Lee, S.-J., Rochet, J.-C., Ding, T. T., Williamson, R. E. & Lansbury, P. T. (2000) *Proc. Natl. Acad. Sci. USA* **97**, 571–576.
- Conway, K. A., Harper, J. D. & Lansbury, P. T. (2000) *Biochemistry* **39**, 2552–2563.
- Crawford, G. D. J., Le, W. D., Smith, R. G., Xie, W. J., Stefani, E. & Appel, S. H. (1992) *J. Neurosci.* **12**, 3382–3398.
- Otto, H., Buchner, K., Beckmann, R., Hilbert, R. & Hucho, F. (1992) *Neurochem. Int.* **21**, 409–414.
- Bradford, M. M. (1976) *Anal. Biochem.* **72**, 248–254.
- Jakes, R., Spillantini, M. G. & Goedert, M. (1994) *FEBS Lett.* **345**, 27–32.
- Folch, J., Lees, M. B. & Sloane, S. G. H. (1957) *J. Biol. Chem.* **226**, 497–509.
- Smith, A. F., Tsuchida, K., Hanneman, E., Suzuki, T. C. & Wells, M. A. (1992) *J. Biol. Chem.* **267**, 380–384.
- Wolfinger, R. D. & O'Connell, M. (1993) *J. Stat. Comp. Simul.* **48**, 233–243.
- Maroteaux, L., Campanelli, J. T. & Scheller, R. H. (1988) *J. Neurosci.* **8**, 2804–2815.
- Beijer, K. & Nyström, E. (1972) *Anal. Biochem.* **48**, 1–8.
- Chojnacki, T., Jankowski, W., Jankowski, T. & Sasak, W. (1975) *Anal. Biochem.* **69**, 114–119.
- Glatz, J. F. C. & Veerkamp, J. H. (1983) *Anal. Biochem.* **132**, 89–95.
- Storch, J. & Thumser, A. E. A. (2000) *Biochem. Biophys. Acta* **1486**, 28–44.
- Veerkamp, J. H., Peeters, R. A. & Maatman, R. G. H. J. (1991) *Biochem. Biophys. Acta* **1081**, 1–24.
- Chan, D. S. & Lees, M. B. (1974) *Biochemistry* **13**, 2704–2712.
- Kurtz, A., Zimmer, A., Schnutgen, F., Bruning, G., Spener, F. & Muller, T. (1994) *Development (Cambridge, U.K.)* **120**, 2637–2649.
- Petersen, K., Olesen, O. F. & Mikkelsen, J. D. (1999) *Neuroscience* **91**, 651–659.